Author's Accepted Manuscript

Suicidal ideation and physical illness: does the link lie with depression?

Livia Sanna, Amanda L. Stuart, Julie A Pasco, Mark A Kotowicz, Michael Berk, Paolo Girardi, Lana J Williams

www.elsevier.com/locate/jad

PII: S0165-0327(13)00737-4
DOI: <http://dx.doi.org/10.1016/j.jad.2013.10.008>
Reference: JAD6423

To appear in: *Journal of Affective Disorders*

Received date: 19 July 2013
Revised date: 1 October 2013
Accepted date: 6 October 2013

Cite this article as: Livia Sanna, Amanda L. Stuart, Julie A Pasco, Mark A Kotowicz, Michael Berk, Paolo Girardi, Lana J Williams, Suicidal ideation and physical illness: does the link lie with depression?, *Journal of Affective Disorders*, <http://dx.doi.org/10.1016/j.jad.2013.10.008>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Suicidal ideation and physical illness: does the link lie with depression?

Livia Sanna^{1,2}: liviasanna@gmail.com

Amanda L Stuart²: amandh@barwonhealth.org.au

Julie A Pasco^{2,4}: juliep@barwonhealth.org.au

Mark A Kotowicz^{2,4}: markk@barwonhealth.org.au

Michael Berk^{2,3,5,6}: mikebe@barwonhealth.org.au

Paolo Girardi¹: paolo.girardi@uniroma1.it

Lana J Williams^{2,3}: lanaw@barwonhealth.org.au

Word count:

Abstract 203; Manuscript 2,520; Tables 2

Affiliations:

¹ Unit of Psychiatry , Neurosciences, Mental Health and Sensory Organs Department (NeSMOS), Faculty of Medicine and Psychology, Sant'Andrea Hospital, Sapienza University of Rome, Italy; ²School of Medicine, Deakin University, Geelong, Australia; ³Department of Psychiatry, The University of Melbourne, Parkville, Australia; ⁴NorthWest Academic Centre, Department of Medicine, The University of Melbourne, Western Health, St Albans, Australia; ⁵Orygen Youth Health Research Centre, Parkville, Australia; ⁶Florey Institute for Neuroscience and Mental Health, Parkville, Australia

Corresponding author and contact for re-print requests:

Dr Lana Williams

Deakin University

School of Medicine

PO Box 281, Geelong 3220, Australia

Tel: +61 3 4215 3303 Fax +61 3 4215 3491

Email: lanaw@barwonhealth.org.au

ABSTRACT

Objective: Medical illness is a risk factor for suicidality; however, disorder-specific risks are not well-known and these relationships are often explained by major depressive disorder (MDD). We aimed to investigate the relationship between suicidal ideation, MDD and medical illnesses in an age-stratified, population-based sample of men participating in the Geelong Osteoporosis Study.

Methods: Suicidal ideation and medical conditions were self-reported. Medical conditions were confirmed by medical records, medication use or clinical data where possible. MDD

was determined using the Structured Clinical Interview for DSM-IV-TR Research Version, Non-patient edition.

Results: Of the 907 men, 8.5% reported suicidal ideation. Thyroid disorders (OR 3.85, 95%CI 1.2-12.1), syncope and seizures (OR 1.96, 95%CI 1.1-3.5), liver disorders (OR 3.53, 95%CI 1.1-11.8; younger men only) and alcoholism (OR 2.15, 95%CI 1.1-4.4) were associated with increased odds of suicidal ideation, independent of age and MDD. Major vascular events doubled the odds of suicidal ideation but this was explained by MDD. No association was evident with high medical burden, musculoskeletal disease, metabolic factors, gastrointestinal disorders, headaches, cardiovascular disease, COPD, cancer and psoriasis.

Conclusion: Health care professionals should focus on identification, assessment and management of suicidal ideation in the medically ill in patients both with and without MDD.

Keywords: comorbidity; depression; medical illness; suicidal ideation

INTRODUCTION

Although the rate of suicide in Australia has been seen to be dropping over time, over 2,000 Australians complete suicide each year (Large and Nielssen, 2010, Australian Bureau of Statistics, 2012). It has been estimated that the lifetime prevalence of suicidal ideation, suicide plan and suicide attempt is 13.3%, 4.0% and 3.2%, respectively (Johnston et al., 2009), with men more likely to have a successful attempt (13.9 per 100 000 males, versus 3.96 in females) (Large and Nielssen, 2010).

It is well-known that physical and mental illness are strongly linked (Liew, 2012, Roy-Byrne et al., 2008). Over the past 15 years, researchers have systematically investigated the association between suicidality and medical conditions, yet study cohorts are usually restricted to elderly populations (Waern et al., 2002, Juurlink et al., 2004, Quan et al., 2002) or psychiatric/inpatient settings (Furlanetto and Stefanello, 2011, Webb et al., 2012, Botega et al., 2010) with few population-based studies (Druss and Pincus, 2000, Scott et al., 2010). There is also controversy surrounding whether suicide is more likely to be associated with general medical burden rather than a specific condition (Webb et al., 2012, Scott et al., 2010, Waern et al., 2002). Understanding the role of medical conditions in suicidal ideation is complex due to frequent comorbidity with major depressive disorder (MDD). Webb and colleagues demonstrated a higher suicide risk among people affected by coronary heart disease (CHD), stroke, chronic obstructive pulmonary disease (COPD) and osteoporosis, but these associations were explained by adjustment for MDD (Webb et al., 2012). On the other hand, Scott and colleagues recently found physical conditions to be a risk factor for suicidal behaviour independent of mental disorders in a large, population based cross-national sample of 37,915 men and women (Scott et al., 2010).

Considering suicidality as a continuum between passive death-related thoughts and self-inflicted death, we aimed to investigate the association between suicidality, MDD, medical conditions and overall medical burden in a population based sample of men spanning the full adult age range.

METHOD

Participants

This study examined data collected from the 5-year follow-up of men participating in the Geelong Osteoporosis Study (GOS) (Pasco et al., 2012), an epidemiological population-based study involving individuals randomly selected from the Commonwealth of Australia Electoral Rolls for the Barwon Statistical Division. A total of 1540 male participants aged 20-97 years were randomly recruited between 2001 and 2006 (67% response), with 978 returning for the 5-year follow up (81% of eligible men). Reasons for non-participation included death (n=141), change of region (n=41), incapacity to provide written informed consent (n=16), inability to be contacted (n=139) and 225 declined participation.

Of the 978 participants, 22 did not provide medical information and 49 participants did not complete the suicidal ideation question and were thus excluded from the current analysis, resulting in a sample of 907 eligible men aged 25-98 years. The study was approved by the Barwon Health Human Research Ethics Committee and written informed consent was obtained from all participants.

Assessments

Outcomevariable

Suicidal ideation was determined from a self-reported question eliciting a binary yes/no response: Have you had for a two-week period, recurrent thoughts of death or thoughts of suicide, or a plan for committing suicide?

Exposure variables

The presence of medical conditions (lifetime) was self-reported and confirmed by medical records, medication use or clinical data where possible. Low bone mineral density (BMD) self-reported osteoarthritis, ankylosing spondylitis, psoriatic arthritis, rheumatoid arthritis, muscle disease or weakness, adult low trauma fracture and self-reported or medicated gout (agents used in gout or hyperuricaemia) were termed Musculoskeletal disease. Low BMD at the lumbar spine (L2-L4) and/or femoral neck was measured by Dual Energy X-ray Absorptiometry (DXA) using GE Lunar Prodigy (Prodigy GE Lunar Madison, WI, USA) and identified as T-score less than -1 (Henry et al., 2010). Thyroid disorders included self-reported or medicated (thyroid hormones and anti-thyroid agents) hyperthyroidism, hypothyroidism, Graves' disease, Hashimoto's disease, thyroiditis and other unspecified thyroid conditions. A diagnosis of diabetes, defined by self-report or current medication use (oral hypoglycaemic agents and insulin preparations), obesity defined by a body mass index (BMI) greater than or equal to 30kg/m^2 , self-reported or medicated hypercholesterolemia

(cholesterol-lowering agents) and hypertension (self-report and current use of antihypertensive medication/Beta-adrenergic blocking agents/diuretics, or systolic blood pressure of >140mmHg or diastolic blood pressure of >90mmHg) were collectively termed Metabolic risk factors. BMI (kg/m^2) was calculated from measurements of height and weight to the nearest 0.1cm and 0.1kg, respectively. Gastrointestinal disorders encompassed peptic ulcer, chronic gastritis and causes of malabsorption such as pancreatitis, gastric surgery, chronic diarrhoea, irritable bowel syndrome, inflammatory bowel syndrome and coeliac disease. Hypertension (as detailed above) or a combination of self-reported or medicated (beta-adrenergic blocking agents, diuretics, anti-arrhythmic, anti-angina, cardiac inotropic agents, adrenergic stimulants, other cardiovascular agents) stroke, ischaemic cardiomyopathy, arrhythmia and valve and vascular diseases were collectively termed Cardiovascular diseases. Major vascular events included stroke and ischaemic cardiomyopathy. Epilepsy, blackouts, fainting and dizzy spells were grouped together as syncope and seizures. Recurrent headaches included migraine and other recurrent headaches. Diagnoses of asthma, emphysema and chronic bronchitis were grouped as pulmonary disease. Psoriasis was defined by self-report. Liver disease included fatty liver, hepatitis, cirrhosis and liver failure. Malignant tumours including non-melanoma skin cancer were grouped as Cancer. A diagnosis of three or more conditions listed above awarded the term high medical burden.

Past and current alcoholism, including alcohol abuse and dependence and MDD were assessed using the Structured Clinical Interview for DSM-IV-TR Research Version, Non-patient edition (SCID-I/NP) (First et al., 2002). Clinical interviewers were trained in administering the SCID-I/NP and held qualifications in psychology.

Information on lifestyle and other health factors was obtained via questionnaire. Participants were asked to bring in a list of medications or containers to assist with accurate recording of details. Cigarette smoking was documented and grouped as current, past or never. Men participating in light to vigorous activity on a regular basis were classed as active. Socio-economic status (SES) was defined based on quintiles of the Index of Relative Socioeconomic Advantage/Disadvantage (IRSAD), derived from the Socio-Economic Index for Areas (SEIFA) (Australian Bureau of Statistics, 2006).

Statistical Analysis

The prevalence of medical conditions in participants with and without suicidal ideation was analysed using chi square analysis and Kruskal-Wallis, where applicable. Age- and age and MDD- adjusted binary logistic regression were performed to evaluate the relationship between lifetime suicidal ideation and lifetime medical conditions. Interaction terms were tested for effect modification. Post hoc analyses incorporating alcoholism into the models were performed. Statistical analysis was performed using Minitab (version 16; Minitab, State College, PA).

RESULTS

Seventy-seven men (8.5%) were identified as having a history of suicidal ideation. Characteristics of men with and without suicidal ideation are listed in Table 1. There were no differences in age, BMI, physical activity, smoking status, SES or alcohol consumption between the groups. Males with MDD were more likely to have suicidal ideation than those without.

Table 2 presents both age- and age/MDD -adjusted odds ratios for medical conditions in men with a lifetime history of suicidal ideation. Thyroid disorders were associated with a 4-fold increased odds of suicidal ideation (age adjusted OR 4.0, 95% CI 1.4-11.6, $p=0.009$), which persisted following adjustment for MDD (OR 3.9, 95% CI 1.2-12.1, $p=0.02$). Self-reported syncope and seizures were associated with a two-fold increased risk of suicidal ideation before (age adjusted OR 2.4, 95% CI 1.4-4.1, $p=0.002$) and after adjustment for MDD (OR 2.0, 95% CI 1.1-3.5, $p=0.02$). Major vascular events were associated with a two-fold increased risk of suicidal ideation (age-adjusted OR 2.1, 95% CI 1.1-4.2, $p=0.03$); however, this was attenuated following adjustments for MDD (OR 1.6, 95% CI 0.8-3.4, $p=0.19$). Age was an effect modifier in the association between liver disorders and suicidal ideation. Liver disorders were associated with a 3-fold increase in risk of suicidal ideation among younger men both before (OR 4.0, 95% CI 1.3-11.9, $p=0.01$) and after (OR 3.5, 95% CI 1.1-11.8, $p=0.04$) adjustment for MDD, but not among those aged over 60 years ($p>0.05$). Alcoholism was associated with an increased risk of suicidal ideation both before (OR 3.2, 95% CI 1.7-6.2, $p<0.001$) and after (OR 2.2, 95% CI 1.1-4.4, $p=0.04$) adjusting for MDD.

Further adjustments for BMI, physical activity, smoking status, SES, alcohol consumption and alcoholism did not impact on the odds ratios.

There was no association between suicidal ideation and musculoskeletal disease, metabolic risk factors, gastrointestinal disorders, recurrent headaches, cardiovascular diseases, pulmonary disease, cancer, psoriasis and high medical burden (all $p > 0.05$).

DISCUSSION

Our data demonstrate that suicidality is associated with several medical conditions; thyroid disorders, syncope and seizures, alcoholism and liver disorders among younger men only, independent of age, MDD, sociodemographic factors and other lifestyle factors. There were no associations detected between suicidality and musculoskeletal disease, metabolic risk factors, gastrointestinal disorders, recurrent headaches, cardiovascular diseases, pulmonary disease, cancer, psoriasis and high medical burden and the significant relationship between suicidality and major vascular events was lost following adjustment for MDD.

The high odds of suicidality in men with thyroid dysfunction and young men with liver disorders, corroborate recent data (Duval et al., 2010, Kristiansen et al., 2011). Similarly, our results confirm previous observations of a positive association between suicidality and neurological disorders such as blackouts, dizzy spells and epilepsy (Christensen et al., 2007), and alcohol abuse and dependence (Clark, 2003), regardless of a diagnosis of depression. It is notable that Clark suggested that alcoholism can lead to suicidal behaviour in an indirect

way by affecting mood, but also in a direct way by enhancing impulsivity (Clark, 2003). However, our results show that the association between medical conditions and suicidality is independent of alcohol abuse/dependence. The relationship between major vascular events and suicidal ideation was influenced by MDD. This result supports that by Scott and colleagues who reported a 2.7 fold increased risk of planned suicide attempts amongst those with past history of stroke/heart attack was not significant after adjusting for mental disorders in their analysis of World Mental Health surveys (Scott et al., 2010).

There was no evidence of an increased odds of suicidal ideation associated with high medical burden; being in accordance with Webb and colleagues who reported that heightened risk of suicide was associated with multiple physical diseases for women only (Webb et al., 2012). However, in another study, a high overall medical burden has been associated with an almost threefold increase in rate of self-inflicted death in elderly men (Waern et al., 2002). It is noteworthy that cut-off points differed between the studies; high medical burden was categorised as 3 or more and 10 or more medical disorders, respectively.

Our results confirm previous data showing no association between suicidality and metabolic risk factor such as diabetes, hypertension and obesity (Harris and Barraclough, 1994) and gastrointestinal disorders such as peptic ulcer (Quan et al., 2002) and raise questions regarding other known associations with cardiovascular disease (Larsen et al., 2010,

Lossnitzer et al., 2009), COPD (Goodwin, 2011), psoriasis (Gupta and Gupta, 1998), cancer (Scott et al., 2010) and recurrent headaches (Scott et al., 2010).

Diseases were grouped based on organs/systems affected in order to give sufficient power for analyses, therefore examining larger groups rather than individual illness. This may help to explain the differences between our observations and those of others. For example, it has been previously reported that osteoarthritis is protective against suicidal ideation whereas osteoporosis is a risk factor (Webb et al., 2012); however, in our analysis these diseases were pooled reporting no significant association.

Reasons for the relationship between suicide and medical illness are still unknown and explanatory pathways are likely to be different for different conditions. Depression is a possible explanation, however, life-threatening characteristics of the condition, lack of social support/coping capacities (Marusic and Goodwin, 2006), functional status (Kaplan et al., 2007), subjective perception of the illness (Olfson et al., 1996) or chronic pain (Purcell et al., 1999) are also potential mechanisms.

It is necessary to note the methodological characteristics of the study. Strengths include the use of a randomly-selected population-based sample of men spanning the full adult age range, thus allowing generalization of our findings to the adult population and assessment of possible age-related associations. Suicidality has been previously shown to differ between elderly and young physically-ill patients (Viilo et al., 2005). Furthermore, self-reported data

were confirmed by medical records, medication use or clinical data where possible and a gold standard tool was used to assess mood and alcohol use disorders. Limitations of our study include potential recall bias, healthy participant bias and the absence of illness severity and functional disability. It is possible that the relationship between suicidal ideation and medical illness is driven by severity of illness rather than the physical disease per se. Although the total sample size was moderate, the small size of some of the subgroups of medical conditions, albeit likely to be representative of the prevalence of the medical condition in the general population, may have limited the power to detect a difference between the groups. Furthermore, medication effects that may induce self-harm ideation have not been taken into consideration. We evaluated self-reported passive and active death-related thoughts through a question with a binary (yes/no) response, rather than using a detailed rating scale. Furthermore, we investigated suicidal thoughts rather than suicidal behaviour as data on suicide attempt and completed suicides were not available. It is estimated that only a small proportion of individuals who exhibit suicidal ideation go on to attempt suicide. However, desire of death, suicidal ideation and suicide planning are main predictors of suicide behaviour (Baca-Garcia et al., 2011). Finally, the cross-sectional nature of our study does not enable conclusions about causal directionality, but we can speculate that there might be a reciprocal relationship between health status and self-harm ideation.

Past research has shown that many patients who commit suicide have seen their primary care physician within one month (Luoma et al., 2002) and often within the week of their death (Conwell, 1997) with many physicians unaware of their patients' intentions (Murphy,

1975). Suicidality, including in its cognitive expression, is a condition often independent of a depressive syndrome, and thus it is a responsibility of any clinician monitoring a patient's health, both physical and mental. In light of this, primary care physicians should be trained in screening patients for suicidal ideation, both active and passive. In addition, information and awareness programs should be implemented at the level of the general population to educate and reduce stigma surrounding suicidality. Considering desire for death, suicidal ideation and suicidal planning as elements on a continuum, we suggest that a correct assessment and management of these factors could improve patients' quality of life, and have the capacity to prevent the progression to the last step represented by completed suicide. In conclusion, our data suggest that suicidality is associated with several medical conditions outside of hospital or psychiatric settings. Thus, awareness of this relationship and identification and management of suicidal ideation should be a priority.

Acknowledgments

We acknowledge the men who participated in the study. The study was funded by the National Health and Medical Research Council. The funding provider played no role in the design or conduct of the study; collection, management, analysis, and interpretation of the data; or in the preparation, review, or approval of the manuscript.

References

- AUSTRALIAN BUREAU OF STATISTICS (2006) Socio-Economic indexes for Areas (SEIFA) - Technical Paper. IN AUSTRALIAN BUREAU OF STATISTICS (Ed.) Canberra, ABS.
- AUSTRALIAN BUREAU OF STATISTICS (2012) Suicides, Australia 2010. IN AUSTRALIAN BUREAU OF STATISTICS (Ed.) Canberra, ABS.
- BACA-GARCIA, E., PEREZ-RODRIGUEZ, M. M., OQUENDO, M. A., KEYES, K. M., HASIN, D. S., GRANT, B. F. & BLANCO, C. (2011) Estimating risk for suicide attempt: are we asking the right

- questions? Passive suicidal ideation as a marker for suicidal behavior. *J Affect Disord*, 134, 327-32.
- BOTEGA, N. J., DE AZEVEDO, R. C., MAURO, M. L., MITSUUSHI, G. N., FANGER, P. C., LIMA, D. D., GASPAR, K. C. & DA SILVA, V. F. (2010) Factors associated with suicide ideation among medically and surgically hospitalized patients. *Gen Hosp Psychiatry*, 32, 396-400.
- CHRISTENSEN, J., VESTERGAARD, M., MORTENSEN, P. B., SIDENIUS, P. & AGERBO, E. (2007) Epilepsy and risk of suicide: a population-based case-control study. *Lancet Neurol*, 6, 693-8.
- CLARK, D. B. (2003) Serum tryptophan ratio and suicidal behavior in adolescents: a prospective study. *Psychiatry Res*, 119, 199-204.
- CONWELL, Y. (1997) Management of suicidal behavior in the elderly. *Psychiatr Clin North Am*, 20, 667-83.
- DRUSS, B. & PINCUS, H. (2000) Suicidal ideation and suicide attempts in general medical illnesses. *Arch Intern Med*, 160, 1522-6.
- DUVAL, F., MOKRANI, M. C., LOPERA, F. G., DIEP, T. S., RABIA, H. & FATTAH, S. (2010) Thyroid axis activity and suicidal behavior in depressed patients. *Psychoneuroendocrinology*, 35, 1045-54.
- FIRST, M., SPITZER, R., GIBBON, M. & WILLIAMS, J. (2002) *Structured clinical interview for DSM-IVTR axis I disorders, research version, non-patient edition (SCID-I/NP)*, New York, Biometrics Research, New York State Psychiatric Institute.
- FURLANETTO, L. M. & STEFANELLO, B. (2011) Suicidal ideation in medical inpatients: psychosocial and clinical correlates. *Gen Hosp Psychiatry*, 33, 572-8.
- GOODWIN, R. D. (2011) Is COPD associated with suicide behavior? *J Psychiatr Res*, 45, 1269-71.
- GUPTA, M. A. & GUPTA, A. K. (1998) Depression and suicidal ideation in dermatology patients with acne, alopecia areata, atopic dermatitis and psoriasis. *Br J Dermatol*, 139, 846-50.
- HARRIS, E. C. & BARRACLOUGH, B. M. (1994) Suicide as an outcome for medical disorders. *Medicine (Baltimore)*, 73, 281-96.
- HENRY, M. J., PASCO, J. A., KORN, S., GIBSON, J. E., KOTOWICZ, M. A. & NICHOLSON, G. C. (2010) Bone mineral density reference ranges for Australian men: Geelong Osteoporosis Study. *Osteoporos Int*, 21, 909-17.
- JOHNSTON, A. K., PIRKIS, J. E. & BURGESS, P. M. (2009) Suicidal thoughts and behaviours among Australian adults: findings from the 2007 National Survey of Mental Health and Wellbeing. *Aust N Z J Psychiatry*, 43, 635-43.
- JUURLINK, D. N., HERRMANN, N., SZALAI, J. P., KOPP, A. & REDELMEIER, D. A. (2004) Medical illness and the risk of suicide in the elderly. *Arch Intern Med*, 164, 1179-84.
- KAPLAN, M. S., MCFARLAND, B. H., HUGUET, N. & NEWSOM, J. T. (2007) Physical illness, functional limitations, and suicide risk: a population-based study. *Am J Orthopsychiatry*, 77, 56-60.
- KRISTIANSEN, M. G., LOCHEN, M. L., GUTTEBERG, T. J., MORTENSEN, L., ERIKSEN, B. O. & FLORHOLMEN, J. (2011) Total and cause-specific mortality rates in a prospective study of community-acquired hepatitis C virus infection in northern Norway. *J Viral Hepat*, 18, 237-44.
- LARGE, M. M. & NIELSSEN, O. B. (2010) Suicide in Australia: meta-analysis of rates and methods of suicide between 1988 and 2007. *Med J Aust*, 192, 432-7.
- LARSEN, K. K., AGERBO, E., CHRISTENSEN, B., SONDERGAARD, J. & VESTERGAARD, M. (2010) Myocardial infarction and risk of suicide: a population-based case-control study. *Circulation*, 122, 2388-93.
- LIEW, H. P. (2012) Depression and chronic illness: a test of competing hypotheses. *J Health Psychol*, 17, 100-9.
- LOSSNITZER, N., MULLER-TASCH, T., LOWE, B., ZUGCK, C., NELLES, M., REMPPIS, A., HAASS, M., RAUCH, B., JUNGER, J., HERZOG, W. & WILD, B. (2009) Exploring potential associations of

- suicidal ideation and ideas of self-harm in patients with congestive heart failure. *Depress Anxiety*, 26, 764-8.
- LUOMA, J. B., MARTIN, C. E. & PEARSON, J. L. (2002) Contact with mental health and primary care providers before suicide: a review of the evidence. *Am J Psychiatry*, 159, 909-16.
- MARUSIC, A. & GOODWIN, R. D. (2006) Suicidal and deliberate self-harm ideation among patients with physical illness: the role of coping styles. *Suicide Life Threat Behav*, 36, 323-8.
- MURPHY, G. E. (1975) The physician's responsibility for suicide. II. Errors of omission. *Ann Intern Med*, 82, 305-9.
- OLFSON, M., WEISSMAN, M. M., LEON, A. C., SHEEHAN, D. V. & FARBER, L. (1996) Suicidal ideation in primary care. *J Gen Intern Med*, 11, 447-53.
- PASCO, J. A., NICHOLSON, G. C. & KOTOWICZ, M. A. (2012) Cohort Profile: Geelong Osteoporosis Study. *Int J Epidemiol*, 41, 1565-1575.
- PURCELL, D., THRUSH, C. R. & BLANCHETTE, P. L. (1999) Suicide among the elderly in Honolulu County: a multiethnic comparative study (1987-1992). *Int Psychogeriatr*, 11, 57-66.
- QUAN, H., ARBOLEDA-FLOREZ, J., FICK, G. H., STUART, H. L. & LOVE, E. J. (2002) Association between physical illness and suicide among the elderly. *Soc Psychiatry Psychiatr Epidemiol*, 37, 190-7.
- ROY-BYRNE, P. P., DAVIDSON, K. W., KESSLER, R. C., ASMUNDSON, G. J., GOODWIN, R. D., KUBZANSKY, L., LYDIARD, R. B., MASSIE, M. J., KATON, W., LADEN, S. K. & STEIN, M. B. (2008) Anxiety disorders and comorbid medical illness. *Gen Hosp Psychiatry*, 30, 208-25.
- SCOTT, K. M., HWANG, I., CHIU, W. T., KESSLER, R. C., SAMPSON, N. A., ANGERMEYER, M., BEAUTRAIS, A., BORGES, G., BRUFFAERTS, R., DE GRAAF, R., FLORESCU, S., FUKAO, A., HARO, J. M., HU, C., KOVESS, V., LEVINSON, D., POSADA-VILLA, J., SCOCCO, P. & NOCK, M. K. (2010) Chronic physical conditions and their association with first onset of suicidal behavior in the world mental health surveys. *Psychosom Med*, 72, 712-9.
- VIILO, K. M., TIMONEN, M. J., HAKKO, H. H., SARKIOJA, T., MEYER-ROCHOW, V. B. & RASANEN, P. K. (2005) Lifetime prevalences of physical diseases and mental disorders in young suicide victims. *Psychosom Med*, 67, 241-5.
- WAERN, M., RUBENOWITZ, E., RUNESON, B., SKOOG, I., WILHELMSON, K. & ALLEBECK, P. (2002) Burden of illness and suicide in elderly people: case-control study. *Bmj*, 324, 1355.
- WEBB, R. T., KONTOPANTELOS, E., DORAN, T., QIN, P., CREED, F. & KAPUR, N. (2012) Suicide risk in primary care patients with major physical diseases: a case-control study. *Arch Gen Psychiatry*, 69, 256-64.

Table 1. Characteristics between men with or without lifetime history of suicidal ideation. Values are given as median (interquartile range) or n (%).

Medical Condition (ever)	Total n=907	Suicidal ideation (ever)		p value
		Yes n=77	No n=830	
Age at interview(yr), median (IQR)	59.6 (45.7, 73.2)	56.1 (44.4, 68.3)	60.0 (45.7, 73.5)	0.15
Major Depressive Disorder	133 (14.8%)	39 (52.0%)	94 (11.4%)	<0.001
BMI (kg/m ²), median (IQR)	27.1 (24.7, 29.7)	26.8 (24.3, 30.2)	27.1 (24.7, 29.6)	0.86
Physical Activity (active)	650 (71.7%)	54 (70.1%)	596 (71.9%)	0.74
Smoking (current)	98 (10.8%)	10 (13.0%)	88 (10.6%)	0.53
Socioeconomic Status (IRSAD)	Quintile 1 (low)	144 (15.9%)	14 (18.2%)	0.23
	Quintile 2	186 (20.5%)	18 (23.4%)	
	Quintile 3	179 (19.7%)	10 (13.0%)	
	Quintile 4	195 (21.5%)	22 (28.6%)	
	Quintile 5	203 (22.4%)	13 (16.9%)	
Alcohol consumption	0	93 (10.5%)	8 (10.5%)	0.41
	1-2 glasses/day	341 (51.0%)	24 (31.6%)	
	≥3 glasses/day	452 (38.5%)	44 (57.9%)	
Musculoskeletal disease	679 (76.1%)	60 (79.0%)	619 (75.9%)	0.55
Thyroid disorders	21 (2.3%)	5 (6.5%)	16 (1.9%)	0.01
Metabolic risk factors	636 (90.1%)	54 (88.5%)	582 (90.2%)	0.67
Gastrointestinal disorders	158 (17.4%)	14 (18.2%)	144 (17.4%)	0.85
Recurrent headaches	48 (5.3%)	7 (9.1%)	41 (4.9%)	0.12
Syncope and seizures	150 (16.5%)	22 (28.6%)	128 (15.4%)	0.003
Cardiovascular disease	575 (81.8%)	49 (75.4%)	526 (82.5%)	0.16
Major vascular events	130 (14.3%)	15 (19.5%)	115 (13.9%)	0.19
Pulmonary disease	175 (19.3%)	15 (19.5%)	160 (19.3%)	0.97
Liver disorders	55 (6.1%)	8 (10.4%)	47 (5.7%)	0.10

Age <60yo	18/459 (3.9%)	5/44 (11.4%)	13/415 (3.1%)	0.007
Age ≥60yo	37/448 (8.3%)	3/33 (9.1%)	34/415 (8.2%)	0.86
Cancer	157 (17.3%)	16 (20.8%)	141 (17.0%)	0.40
Alcoholism	65 (7.2%)	14 (18.2%)	51 (6.1%)	<0.001
Psoriasis	39 (4.3%)	3 (3.9%)	36 (4.3%)	0.86
High medical burden	358 (48.6%)	35 (52.2%)	323 (48.2%)	0.53

Missing data: MDD n=10, BMI n=27, Physical Activity n=1, Smoking n=3, Musculoskeletal disease n=15, Metabolic risk factors n=201, Cardiovascular diseases n=204, High medical burden n=170

Table 2. Age- and MDD/age-adjusted odds ratios for physical illness in men with a lifetime history of suicidal ideation.

	Model	Suicidal ideation		
		Odds ratio	95% CI	p value
Musculoskeletal disease	I	1.29	0.72 – 2.31	0.40
	II	1.23	0.65 – 2.31	0.53
Thyroid disorders	I	4.04	1.42 – 11.55	0.01
	II	3.85	1.22 – 12.13	0.02
Metabolic risk factors	I	0.95	0.41 – 2.20	0.90
	II	0.91	0.35 – 2.35	0.84
Gastrointestinal disorders	I	1.15	0.62 – 2.13	0.66
	II	1.04	0.54 – 1.99	0.91
Recurrent headaches	I	1.99	0.86 – 4.62	0.11
	II	1.52	0.61 – 3.76	0.37
Syncope and seizures	I	2.38	1.39 – 4.07	0.002
	II	1.96	1.10 – 3.50	0.02
Cardiovascular disease	I	0.79	0.42 – 1.49	0.47
	II	0.76	0.37 – 1.55	0.45
Major vascular events	I	2.11	1.07 – 4.19	0.03
	II	1.63	0.78 – 3.40	0.19
Pulmonary disease	I	1.01	0.56 – 1.82	0.98
	II	0.79	0.41 – 1.51	0.48
Liver disorders <60yo*	I	3.97	1.32 – 11.93	0.01
	II	3.53	1.06 – 11.81	0.04
Liver disorders ≥60yo*	I	1.06	0.30 – 3.71	0.93
	II	0.56	0.14 – 2.27	0.42
Cancer	I	1.54	0.83 – 2.85	0.17
	II	1.80	0.93 – 3.49	0.08
Alcoholism	I	3.23	1.68 – 6.20	<0.001

	II	2.15	1.05 – 4.40	0.04
Psoriasis	I	0.95	0.29 – 3.18	0.94
	II	0.80	0.22 – 2.88	0.74
High medical burden	I	1.53	0.85- 2.77	0.16
	II	1.09	0.57 – 2.06	0.80

Model I: Age-adjusted Model II: Age / MDD-adjusted

* Alcohol consumption-adjusted

Conflict of interest

Livia Sanna, Amanda L Stuart and Mark A Kotowicz have no conflicts of interest, including specific financial interests and relationships and affiliations relevant to the subject matter or materials discussed in the manuscript.

Julie A Pasco has received speaker fees from Amgen, Eli Lilly and Sanofi-Aventis and funding from the Geelong Region Medical Research Foundation, Barwon Health, Perpetual Trustees, the Dairy Research and Development Corporation, The University of Melbourne, the Ronald Geoffrey Arnott Foundation, ANZ Charitable Trust, the American Society for Bone and Mineral Research, Amgen (Europe) GmbH and the NHMRC.

Michael Berk has received Grant/Research Support from the NIH, Simons Foundation, CRC for Mental Health, Stanley Medical Research Institute, MBF, NHMRC, Beyond Blue, Geelong Medical Research Foundation, Bristol Myers Squibb, Eli Lilly, Glaxo SmithKline, Organon, Novartis, Mayne Pharma, Servier and Astra Zeneca. He has been a paid consultant for Astra Zeneca, Bristol Myers Squibb, Eli Lilly, Glaxo SmithKline, Janssen Cilag, Lundbeck and Pfizer

and a paid speaker for Astra Zeneca, Bristol Myers Squibb, Eli Lilly, Glaxo SmithKline, Janssen Cilag, Lundbeck, Organon, Pfizer, Sanofi Synthelabo, Solvay and Wyeth.

Paolo Girardi has in the past three years received research support from Lilly and Janssen, participated in advisory boards for Lilly, Organon, Pfizer, and Schering, and received honoraria from Lilly and Organon.

Lana J Williams has received Grant/Research support from Eli Lilly, Pfizer, The University of Melbourne, Deakin University and the NHMRC.

Author contribution

LS and AS were involved in the conception and design of the study, analysis of data and interpretation of data. JP, MK and LW were involved in the conception and design of the study and interpretation of data. MB and PG were involved in the interpretation of data. All authors were involved in the drafting, revision process and gave final approval of the published version of the manuscript

Acknowledgments

We acknowledge the men who participated in the study. The study was funded by the National Health and Medical Research Council. The funding provider played no role in the design or conduct of the study; collection, management, analysis, and interpretation of the data; or in the preparation, review, or approval of the manuscript.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Sanna, L;Stuart, AL;Pasco, JA;Kotowicz, MA;Berk, M;Girardi, P;Williams, LJ

Title:

Suicidal ideation and physical illness: Does the link lie with depression?

Date:

2014-01-01

Citation:

Sanna, L., Stuart, A. L., Pasco, J. A., Kotowicz, M. A., Berk, M., Girardi, P. & Williams, L. J. (2014). Suicidal ideation and physical illness: Does the link lie with depression?. JOURNAL OF AFFECTIVE DISORDERS, 152 (1), pp.422-426. <https://doi.org/10.1016/j.jad.2013.10.008>.

Persistent Link:

<http://hdl.handle.net/11343/43871>