

1 **Type of contribution:** Research paper

2 **Date of preparation:** 28/3/2017

3

4 **Title: Effects of the nitrification inhibitor acetylene on nitrous oxide emissions and**
5 **ammonia-oxidizing microorganisms of different agricultural soils under laboratory**
6 **incubation conditions**

7 **Authors:** Rui Liu^{a*}, Helen L. Hayden^b, Hangwei Hu^a, Jizheng He^a, Helen Suter^a, Deli Chen^a

8 **Affiliations:**

9 ^a Faculty of Veterinary and Agricultural Sciences, The University of Melbourne, Victoria
10 3010, Australia

11 ^b Department of Economic Development, Jobs, Transport and Resources, Victoria 3083,
12 Australia

13

14 *Corresponding author:

15 Address: Faculty of Veterinary and Agricultural Sciences, The University of Melbourne,
16 Parkville, Victoria 3010, Australia

17 Tel.: +61433990822

18 E-mail address: rliu@unimelb.edu.au

19

20 **Submitted to:** *Applied Soil Ecology*

21

22

23 **Abstract**

24 Acetylene (C_2H_2) is an effective nitrification inhibitor targeting autotrophic ammonia
25 oxidizers, and has shown promise for improving nitrogen use efficiency by mitigating
26 greenhouse gas nitrous oxide (N_2O) emissions and reducing nitrate leaching. Its efficacy,
27 however, varies considerably with edaphic and environmental conditions and remains largely
28 less studied in dryland agricultural soils. Here we conducted two laboratory microcosm
29 incubations to explore the efficacy of C_2H_2 across various agricultural soils and under
30 different conditions. The first incubation was with four agricultural soils at $25^\circ C$ and 60%
31 water-filled pore space (WFPS), and the second incubation included one cropping soil under
32 a range of conditions ($15^\circ C$, $25^\circ C$, $35^\circ C$ and 50%, 70% WFPS). Our results showed that
33 incubation of soil with 1% v/v C_2H_2 resulted in complete or partial inhibition of nitrification,
34 N_2O emission, and AOA or AOB growth under the experimental conditions. Acetylene can
35 totally inhibit nitrification in acidic cropping and dairy pasture soils through retarding both
36 AOA and AOB growth, while C_2H_2 partly inhibited nitrification and N_2O emission in the
37 alkaline vegetable soil through impeding only AOB growth. The highest inhibition effect of
38 C_2H_2 was achieved at $25^\circ C$ and 50% WFPS, while there was no inhibitory effect of C_2H_2
39 when soil was incubated at $15^\circ C$ and 50% WFPS suggesting soil temperature may have a
40 significant influence on C_2H_2 effectiveness. The inhibition of C_2H_2 on cumulative N_2O
41 emission increased with increasing temperature at 50% WFPS. In contrast, at 70% WFPS, the
42 inhibition of C_2H_2 on cumulative N_2O emission decreased with increasing temperature. Since
43 the effect of C_2H_2 varied with soils and environmental conditions, this highlights the
44 assumption that N_2O production and nitrification can be affected by low concentrations of
45 C_2H_2 may be not appropriate in some occasions.

46 **Keywords:** Acetylene; nitrification inhibitor; AOA; AOB; N₂O emission; temperature;
47 moisture

48

49 **1. Introduction**

50 Nitrogen (N) is an essential nutrient for food production, but the amount of applied fertilizer
51 N used by crops rarely exceeds 40% (Chen et al., 2008), and can be as low as 20% in
52 vegetable production systems in Australia (Suter et al., 2014). The majority of applied N is
53 lost from agro-ecosystems through ammonia (NH₃) volatilization, gaseous emission of
54 nitrous oxide (N₂O) and di-nitrogen (N₂) and nitrate (NO₃⁻) leaching. Nitrous oxide is a
55 potent greenhouse gas contributing significantly to global climate change with a 300-fold
56 higher global warming potential than CO₂ (IPCC, 2007). It is also involved in the destruction
57 of the protective ozone layer (Ravishankara et al., 2009). Soil ecosystems are the largest
58 source of N₂O, accounting for approximately 65% of the atmospheric N₂O loading (IPCC,
59 2007). The major pathways of N₂O production in soils include microbial-mediated
60 nitrification and denitrification (Hu et al., 2015; Zhang et al., 2015). Microorganisms, such as
61 ammonia oxidizers and bacterial denitrifiers, involved in the N cycle can directly regulate
62 N₂O production and consumption from soils, and increased abundance and activity of these
63 microorganisms may increase N₂O emissions (Burger et al., 2005).

64 Nitrification inhibitors (NIs) can decelerate the rate of soil nitrification by deactivating the
65 enzyme ammonia monooxygenase (AMO) which catalyses ammonia oxidation, the first and
66 rate-limiting step of nitrification which is encoded by the *amoA* gene within ammonia-
67 oxidizing archaea (AOA) and ammonia-oxidizing bacteria (AOB) (Zhang et al., 2012). NIs
68 can block the growth of AOA and AOB (Di et al., 2010; Hink et al., 2016), and are widely
69 used to improve N fertiliser efficiency, mitigate N₂O emissions and reduce NO₃⁻ leaching in

70 agricultural systems (Chen et al., 2008; Kelly et al., 2008; Chen et al., 2010; Di et al., 2010;
71 Zhang et al., 2012; Hu et al., 2015). Acetylene (C_2H_2) is an effective inhibitor of bacterial
72 ammonia oxidation (Offre et al., 2009), acting with AMO as a suicide substrate. It is usually
73 used as an autotrophic nitrification inhibitor in experimental studies (de Boer and
74 Kowalchuk, 2001) and can inhibit nitrification at a low concentration (e.g. 10 Pa) in most
75 soils under aerobic conditions (Hyman and Wood, 1985; Schmidt and Bock, 1998; De Boer
76 et al., 1991; Offre et al., 2009). In contrast, C_2H_2 does not efficiently inhibit ammonia
77 oxidation by heterotrophic nitrifiers (Moir et al., 1996; Daum et al., 1998). Higher
78 concentrations of C_2H_2 (1–20 kPa) will inhibit the N_2O reductase of denitrifying
79 microorganisms (Davidson et al., 1986; Klemetsson et al., 1988). As a result of this C_2H_2
80 has been used as a routine method to distinguish nitrification-related N_2O and denitrification-
81 related N_2O in soils experiments based on the inhibition of soil ammonia oxidation (Bateman
82 and Baggs, 2005; Butterbach-Bahl et al., 2013).

83 It is generally believed that low concentrations of C_2H_2 (0.1–10 Pa) totally inhibit
84 nitrification (Hynes and Knowles, 1978; Berg et al., 1982) by forming a reactive epoxide
85 which then irreversibly inactivates the AMO enzyme (Hyman and Wood, 1985), however,
86 this does not always occur and its efficacy is considerably varied. While it was previously
87 reported that 10 Pa of C_2H_2 totally inhibited nitrification (Wrage et al., 2004), Bremner and
88 Blackmer (1979) found that 10 Pa only partially inhibited soil nitrification.

89 Acetylene was also widely used to determine the community compositions of nitrifiers in
90 soils through laboratory work (Boyle-Yarwood et al., 2008; Scheer et al., 2014). Recent
91 studies have shown however, that AOA and AOB may have a variety of responses to C_2H_2
92 application. Gubry-Rangin et al. (2010) and Offre et al. (2009) revealed that AOA growth
93 was inhibited in C_2H_2 -containing microcosms but not AOB growth. Liu et al. (2015a)
94 demonstrated that both AOA and AOB were inhibited by C_2H_2 in three Australian

95 agricultural soils. However, these results contrast with those of Jia and Conrad (2009), who
96 found changes in the abundance of AOB *amoA* genes correlated best with nitrification rate
97 rather than AOA *amoA*, and bacterial growth occurred only in actively nitrifying microcosms
98 with added C₂H₂. Considering the various responses of AOA and AOB to C₂H₂ addition, it is
99 therefore necessary to find out how soil factors influence the response of AOA and AOB to
100 C₂H₂ addition and what are the key factors affecting the response of AOA and AOB to C₂H₂.

101 This study was designed to determine the impact of C₂H₂ on N₂O emissions, nitrification
102 rates and the abundance of ammonia oxidizers in different agricultural soils under laboratory
103 conditions. Microcosm incubation experiments were established under a set of controlled
104 environment conditions with the following objectives: (i) to examine the effects of C₂H₂ on
105 N₂O emissions and the abundances of AOA and AOB from different agricultural soils and (ii)
106 to investigate the effects of C₂H₂ on N₂O emissions and the abundances of AOA and AOB
107 under different temperature and soil water contents in one agricultural soil. We hypothesized
108 that (i) C₂H₂ would have significant inhibitory effects on nitrification and N₂O production
109 from soils with different physicochemical traits, and (ii) AOA and AOB would exhibit
110 distinctly different responses to C₂H₂. This study represents comprehensive efforts to
111 examine C₂H₂ efficacy from different soils under controlled conditions, and the findings from
112 the study can improve our understanding of the interactions between soil microbial
113 communities and the nitrification inhibitor C₂H₂ in different agricultural soils under
114 laboratory controlled conditions.

115 **2. Materials and methods**

116 **2.1 Site description and soil sampling**

117 The soils used in this study were collected from four agricultural sites in Australia: vegetable
118 soil at Boneo, VIC (38.3°S, 144.9°E), sugarcane soil at Bundaberg, QLD (24.8°S, 152.3°E),

119 dairy pasture soil at Glenormiston, VIC (38.2°S, 143°E), and cereal cropping soil at
120 Hamilton, VIC (38.3°S, 142.7°E). At each site, 10 replicate samples of the top soil (0–10 cm)
121 were collected, thoroughly homogenized, and transported on ice to the laboratory. Fresh soils
122 were sieved through a 2.0 mm mesh, and root and leaf residues were removed with tweezers
123 prior to the establishment of microcosms. Soil moisture contents were determined by oven-
124 drying three subsamples (10 g of fresh soil) at 105°C for 48 h. Soil texture (sieve and
125 hydrometer procedures), pH (1:5 soil/water), total carbon (Dumas method) and other soil
126 properties were determined and are shown in Table 1.

127 **2.2 Soil microcosm incubations**

128 **2.2.1 The laboratory incubation with different agricultural soils**

129 Soil microcosms were established in 500 ml vials containing 60 g of soils (oven-dry
130 equivalent). Distilled water was added to soil to just under the final moisture content (60%
131 water-filled pore space, WFPS) and the microcosms were pre-incubated at 25°C for three
132 weeks to stabilise soil microbial communities and minimise priming effects associated with
133 wetting events. After pre-incubation, treatment was applied to each incubation vial to reach
134 60% WFPS (Linn and Doran, 1984). The treatments contained 100 mg N kg⁻¹ soil as
135 exchangeable NH₄⁺-N and 50 mg N kg⁻¹ soil as NO₃⁻-N, which were added to the soil as 1)
136 NH₄Cl + KNO₃; and 2) NH₄Cl + KNO₃ + C₂H₂. Five ml of C₂H₂ (1% v/v) was injected into
137 the headspace of the vials using an air-tight syringe. Aerobic conditions, soil moisture and
138 C₂H₂ contents in the vials were maintained every three days by opening microcosms and
139 replenishing. Soil microcosms were incubated at 25°C in the dark for three weeks.

140 **2.2.1.1 Gas sampling and analysis**

141 Gas samples were collected on days 0, 4, 8, 12, 16 and 20 after fertilizer application. Gas
142 samples (20 ml) for N₂O analysis were taken from the 500 ml vials using gas-tight syringes.

143 Prior to collection of gas samples, the vials were opened to ensure that N₂O concentration in
144 the headspace was at ambient levels. During each sampling, gas samples were collected at 0,
145 8, 16, 24, 48 and 72 hours after vials closure. Before gas collection, 20 ml compressed zero
146 air was injected into 500 ml vials to maintain the pressure in the vials and then 20 ml gas
147 samples were collected into a pre-evacuated 12 ml exetainer (Exetainer®, Labco Ltd.,
148 Lampeter, Ceredigion, UK). Samples were analysed for N₂O concentration by a gas
149 chromatograph (Agilent 7890A) using an ECD (N₂O) detector.

150 **2.2.1.2 Soil Sampling and analysis**

151 Soils were destructively sampled for soil mineral N analysis on days 0, 7, 14 and 21
152 immediately after gas sampling. There were four replicates at each sampling day. A
153 subsample (2 g) of soil was taken from each vial for molecular analysis and stored in a -80°C
154 freezer before DNA extraction. The remaining 50 g of soil in the vials was shaken with 250
155 ml of 2 M KCl for 1 h at 200 rpm at room temperature, and the extract filtered through a
156 qualitative filter paper (Whatman 42). The extracts (30 ml) were stored at -20°C prior to
157 measurement of NH₄⁺ and NO₃⁻ concentrations on a segmented-flow analyzer (Skalar,
158 SAN++).

159 **2.2.1.3 Soil DNA extraction and quantitative PCR (qPCR)**

160 The Power Soil DNA Isolation Kit (MO BIO Laboratories Inc., Carlsbad, CA, USA) was
161 used for DNA extraction following the manufacturer's instructions. Extracted DNA was
162 quantified using a NanoDrop ND2000c spectrophotometer (NanoDrop Technologies,
163 Wilmington, DE, USA) and the quality of extracted DNA was checked on a 1% agarose gel.
164 The archaeal and bacterial *amoA* gene copy numbers were quantified from triplicate samples
165 using qPCR. The primer sets were Arch-amoAF/Arch-amoAR (Francis et al., 2005) and
166 amoA1F/amoA2R (Rotthauwe et al., 1997), respectively. Each qPCR reaction for the

167 archaeal *amoA* gene was performed in a 20 µl volume including 0.5 µM of each primer, 10 µl
168 SensiFAST SYBR No-ROX reagent (Bioline, Sydney, Australia), and 2 µl of 10-fold dilution
169 DNA template (1–10 ng). Each qPCR reaction for the bacterial *amoA* gene was performed in
170 a 10 µl volume containing 0.6 µM of each primer, 5 µl iTaq Universal SYBR Green
171 Supermix (Bio-Rad Laboratories, USA), and 2 µl of 10-fold dilution DNA template (1–10
172 ng). Amplification programs for both AOA and AOB were as follows: 3 mins at 95°C, 40
173 cycles of 5 s at 95°C, 30 s at 60°C, and 72°C for 45 s. A known copy number of plasmid
174 DNA was used to create a standard curve for each AOA or AOB assay. For all assays, qPCR
175 efficiency was 92.7–98.4% and r^2 was 0.96–0.99.

176 **2.2.2 Soil microcosm incubation under different temperature and water contents with** 177 **one agricultural soil**

178 The cropping soil used for the second experiment was chosen from one of the four sites and
179 collected from a cropping paddock in Hamilton, VIC (38.3°S, 142.7°E) (Table 1). The study
180 site is in a high rainfall zone (688 mm per annum), and was chosen because the soil was
181 subjected to two different land management practices, cropping and pasture. Field studies
182 have shown that high N₂O emissions occurred when the soil was converted from pasture to
183 cropping. The laboratory soil incubation was carried out in 500 ml vials containing 60 g of
184 soils (oven dry weight equivalent) to investigate the effects of C₂H₂ on N₂O emissions and
185 the abundances of *amoA* genes under different soil temperature and moisture contents. The
186 same treatments were established as the first experiment, though after the pre-incubation the
187 samples were incubated at three temperatures (15°C, 25°C, 35°C) and two soil moisture
188 levels (50% and 70% WFPS). During the 21-day incubation, soil, gas samples and *amoA*
189 genes abundance were analysed as well.

190 **2.3 Calculations**

191 The equation developed by Persson and Wirén (1995) was used to calculate the net
192 nitrification rates over the incubation time (21 days)

$$193 \quad n = [(\text{NO}_3^- \text{-N})_{d21} - (\text{NO}_3^- \text{-N})_{d0}] / 21$$

194 where $(\text{NO}_3^- \text{-N})_{d0}$ and $(\text{NO}_3^- \text{-N})_{d21}$ are the $\text{NO}_3^- \text{-N}$ concentrations in the soil on days 0 and 21,
195 respectively.

196 **2.4 Statistical Analyses**

197 Data were analysed using SPSS 19 and means were compared using one-way analysis of
198 variance (ANOVA) between treatments to test the variance with a level of significance of $P <$
199 0.05. Spearman correlation analysis was performed to test the relationships between N_2O , the
200 abundances of AOA and AOB under different conditions.

201

202 **3. Results**

203 **3.1 Soil microcosm incubation with different land-use agricultural soils**

204 The physical and chemical properties of the soil samples were highly variable across the
205 different sampling sites (Table 1). Briefly, all soils except the vegetable soil were acidic (pH
206 ≤ 6.2). The cropping soil had the highest organic C content (6.2%), while the vegetable soil
207 had the lowest (0.8%). Nitrate-N ($\text{NO}_3^- \text{-N}$) was the dominant inorganic nitrogen ranging from
208 8.8 to 93 mg kg^{-1} soil, with the highest value recorded in the cropping soil. Sugarcane and
209 vegetable soils had a sandy texture, but cropping and dairy pasture soils were loam. The
210 cropping soil has the highest clay content at 19%.

211 The highest cumulative N_2O emission was 3204.5 (± 52.5) mg N kg^{-1} soil in the cropping soil,
212 compared to 122.4 (± 5.1) mg N kg^{-1} soil in the sugarcane soil, which was the lowest emission
213 from fertilizer treatments (Table 2). Acetylene addition significantly reduced the cumulative

214 N₂O emissions by 28.6% to 54.6% (Table 2). The efficacy of C₂H₂ on reducing cumulative
215 N₂O emissions was ordered by dairy pasture (28.6%) < sugarcane (37.6%) < vegetable
216 (44.9%) < cropping (54.6%).

217 Changes in the NO₃⁻-N concentrations of the four soils during the incubation period are
218 shown in Figure 1. The NO₃⁻-N concentrations showed an increasing trend in all soils from
219 fertilized treatments. While C₂H₂ addition significantly reduced the NO₃⁻-N concentrations in
220 all the soils (*P* < 0.05), the efficacy of C₂H₂ varied among soils. For example, the average net
221 nitrification rates over 21 days in the dairy pasture soil were 1.8 (±0.1) mg NO₃-N kg⁻¹ soil
222 day⁻¹ for the fertilized treatment and 0 mg NO₃-N kg⁻¹ soil day⁻¹ for the fertilizer plus C₂H₂
223 treatment, while in the vegetable soil they were 2.1 (±0.4) mg NO₃⁻-N kg⁻¹ soil day⁻¹ for the
224 fertilized treatment and 1.5 (±0.3) mg NO₃⁻-N kg⁻¹ soil day⁻¹ for the fertilizer plus C₂H₂
225 treatment (Table 2).

226 The abundance of AOB before adding fertilizer varied greatly from 2.7 × 10⁵ copies g⁻¹ soil
227 in the dairy pasture soil to 2.6 × 10⁶ copies g⁻¹ soil in the vegetable soil, and was evidently
228 higher in the vegetable soil than in other soils (Figure 2). The addition of fertilizer
229 significantly increased AOB abundance in all soils on day 7, while C₂H₂ significantly
230 decreased the AOB abundance in cropping, sugarcane and vegetable soils on day 7. In the
231 dairy pasture soil, the abundance of AOB significantly decreased compared to the fertilizer
232 treatment on day 14 (*P* < 0.05). Before fertilizer addition, the AOA abundance was highest in
233 the vegetable soil up to 1.5 × 10⁸ copies g⁻¹ soil and much lower in the cropping, sugarcane,
234 and dairy pasture soils ranging between 5.8 × 10⁶ and 1.1 × 10⁷ copies g⁻¹ soil (Figure 3).

235 Addition of fertilizer significantly increased AOA abundance in all soils except the vegetable
236 soil. There was a decreasing trend in AOA abundance during incubation in the vegetable soil.
237 AOA growth was inhibited by C₂H₂ addition in cropping, sugarcane and dairy pasture soils,

238 however, there was no significant difference in the AOA abundance between treatments
239 during the incubation in the vegetable soil (Figure 3).

240 **3.2 Soil microcosm incubation under different temperature and soil water contents**

241 The NO_3^- -N concentrations in fertilized treatments increased under 50% WFPS with
242 increasing soil temperature except at 35°C (Figure 4). At 70% WFPS, the soil NO_3^- -N
243 concentrations decreased with increasing temperature over the incubation period. The
244 concentrations of NO_3^- -N remained largely unchanged in the fertilizer plus C_2H_2 treatments at
245 different conditions except at 35°C with 70% WFPS. Under 35°C with 70% WFPS, NO_3^- -N
246 concentration significantly ($P < 0.05$) decreased after day 14 in the fertilizer plus C_2H_2
247 treatment. Acetylene addition substantially reduced the nitrification rate by 53% - 100%,
248 suggesting C_2H_2 was capable of inhibiting nitrification under all conditions (Table 3), with
249 the highest inhibition effect found at 25°C and 50% WFPS. The efficacy of C_2H_2 on
250 nitrification decreased with increasing soil temperature and moisture except at 15°C.

251 Total N_2O emission increased with increasing temperature and moisture (Table 3). When
252 C_2H_2 was applied into soil microcosms, cumulative N_2O emission was reduced by 0–86.7%.
253 There was no difference in cumulative N_2O emission between the fertilizer and fertilizer plus
254 C_2H_2 treatments at 15°C and 50% WFPS. The inhibition of C_2H_2 on cumulative N_2O
255 emission increased with increasing temperature at 50% WFPS. In contrast, when soil was
256 wetted to 70% WFPS, the inhibition of C_2H_2 on cumulative N_2O emission decreased with
257 increasing temperature. At the same soil temperature, the efficacy of C_2H_2 on total N_2O
258 emission decreased with increasing moisture except at 15°C.

259 The changes in abundances of AOB and AOA *amoA* genes are shown in Figures 5 and 6. In
260 the fertilized treatment, AOA and AOB *amoA* gene copy numbers ranged from 1.0×10^7 to
261 3.3×10^8 and 1.8×10^6 to 1.7×10^7 copies g^{-1} soil, respectively, and increased with

262 increasing soil moisture. However, both AOA and AOB abundance decreased with increasing
263 soil temperature. AOA *amoA* genes at different soil temperatures and moistures were 3.3-107
264 times more abundant than AOB *amoA* genes. AOA *amoA* gene abundance comprised 93–
265 96% and 76–92% of the total *amoA* gene abundance at 15°C to 35°C, respectively. The ratio
266 of AOA to AOB decreased significantly with increasing soil temperature ($P < 0.05$).
267 Acetylene application significantly ($P < 0.05$) decreased AOA abundance by 48% (15°C and
268 70% WFPS) and reduced AOB abundance by 93% (25°C and 70% WFPS) relative to that in
269 the fertilizer treatments on day 21. However, no significant effect of C₂H₂ addition on AOA
270 or AOB *amoA* gene abundances was observed at 35°C, 50% WFPS and 15°C, 70% WFPS,
271 respectively.

272

273 **4. Discussion**

274 This study investigated the changes in N₂O production, nitrification rates, and abundance of
275 ammonia oxidizers after addition of the nitrification inhibitor C₂H₂ at various conditions. Our
276 experiments showed that incubation of soil with 1% v/v C₂H₂ resulted in complete or partial
277 inhibition of nitrification, N₂O emission, and AOA or AOB growth in all conditions.

278 **4.1 Inhibitory effects of C₂H₂ in different soils.**

279 The results demonstrate that C₂H₂ has different efficacy in inhibiting nitrification and N₂O
280 emission in different soils. Nitrification can be inhibited by C₂H₂, implying AMO-dependent
281 (and presumably autotrophic) nitrification, as frequently observed in many acidic soils (De
282 Boer and Kowalchuk, 2001). In this study, C₂H₂ was much more effective in inhibiting
283 nitrification and cumulative N₂O emission in acidic soils than the alkaline soil, and the acidic
284 cropping soil had the highest inhibitory effect among the four soils. Soil pH might be a key
285 factor in C₂H₂ inhibitory effects because pH is a strong environmental determinant of AOA

286 and AOB abundance. A number of studies have shown niche separation based on pH with
287 AOA favoured in acidic soils and AOB in alkaline. The abundances of AOA and AOB in our
288 acidic and alkaline soils supported this observation (as reviewed in He et al., 2012). Our
289 findings are consistent with these previous studies, and supported the clear niche separation
290 between AOA and AOB shaped by soil pH. We suggest that in the alkaline soil, nitrifiers can
291 be protected within microenvironments due to the high amounts of carbonates present in the
292 alkaline condition.

293 Besides soil pH, other soil physicochemical traits may also have the potential effects on C₂H₂
294 inhibitory efficacy. The dairy pasture soil had the lowest inhibition by C₂H₂ on cumulative
295 N₂O emission (28.6%) while there was a complete inhibition (100%) on nitrification. It is
296 possibly that because the dairy pasture soil had the highest ratio of C to NO₃⁻, that there might
297 be a loss in inhibitory efficiency on N₂O production. Previous studies have shown that the use
298 of C₂H₂ to inhibit N₂O production and reduction has been problematic for soils with a very
299 high ratio of C to NO₃⁻ (Davidson et al., 1986). Another potential explanation for low
300 inhibition of C₂H₂ on cumulative N₂O emission is that the release of N₂O was due to
301 processes other than nitrification, such as denitrification and heterotrophic nitrification,
302 rendering the C₂H₂ unable to effect N₂O emitted from either denitrification or heterotrophic
303 nitrification. The dairy pasture soil had a high organic carbon content which might result in
304 high denitrification-related N₂O because of the high availability of soil organic C for
305 denitrifiers. Wan et al. (2009) also demonstrated that more N₂O was released from
306 denitrification than from nitrification in a soil with high organic carbon. Regarding the
307 highest inhibitory effect on nitrification in dairy pasture soil, this may be due to nitrification
308 in the dairy pasture soil being largely autotrophic. This can be supported by our previous
309 study (Liu et al., 2015b) where we indicated that nitrification was primarily autotrophic with
310 heterotrophic nitrification accounting for only 20%. In the alkaline sandy soil, C₂H₂

311 inhibition of NO_3^- production was only 29%, thus 1% v/v of C_2H_2 may be insufficient to
312 inhibit nitrification completely in this soil, nevertheless, most studies have indicated that this
313 partial pressure of C_2H_2 is clearly sufficient for a complete inhibition. This result contrasts
314 with our previous study (Liu et al., 2015a) where we found that C_2H_2 addition produced
315 100% nitrification inhibition in an alkaline clay loam soil. The difference between these two
316 studies may be because the two alkaline soils differed in soil texture.

317 Overall, the four soils chosen from different land uses had very differing physicochemical
318 properties. Of the properties characterised for the four soils, soil pH, texture, organic C and
319 NO_3^- content might be the key factors influencing the effectiveness of C_2H_2 on nitrification
320 and cumulative N_2O emission. Many recent studies also showed that soil physicochemical
321 properties affected the efficacy of other NIs, such as DMPP and DCD, with their efficacy
322 diminished with the addition of soil organic matter (Fisk et al., 2015) and decreased with
323 higher clay content (Marsden et al., 2016). Although it is not possible to clearly discern the
324 effects of land use from this experimental design, we speculate that land use may affect the
325 efficacy of C_2H_2 to inhibit nitrification and N_2O emissions. However, a multiple regression
326 including more soil physicochemical properties is still necessary.

327 C_2H_2 addition was able to inhibit nitrification and N_2O emission in all soils, albeit to varying
328 degrees. It is unclear however if this reduction in N_2O emission and nitrification is linked to
329 soil microbial communities due to adding C_2H_2 and further investigation is needed to
330 examine if C_2H_2 stimulates changes in microbial community population and activity.

331 As observed previously (He et al., 2007; Shen et al., 2008; Levc̃nik-Höfferle et al., 2012;
332 Liu et al., 2015a), AOA grow better in acidic soils, while AOB thrive better in alkaline soils,
333 which was supported by our study. In our study, the acidic sugarcane soil had the highest
334 AOA abundance, while the highest AOB population was found in the alkaline vegetable soil.

335 Growth of AOB, and not of AOA, has been linked to soil nitrification activity with high
336 levels of ammonium (Di et al., 2009; Jia and Conrad, 2009). In contrast, growth of AOA is
337 associated with nitrification in soils with a continual supply of ammonia at low concentration
338 through the mineralisation of organic matter (Offre et al., 2009). The difference in substrate
339 preferences could affect the distribution of AOA and AOB in soils and further affect the
340 efficacy of C₂H₂ across different soils. Our study indicated that the application of C₂H₂ could
341 block AOA and AOB growth but to different extents. In acidic cropping, sugarcane and dairy
342 pasture soils, C₂H₂ addition significantly decreased AOA and AOB abundance, suggesting
343 both AOA and AOB mediated nitrification in these soils and were sensitive to C₂H₂. The
344 observed significant decrease in N₂O emission and nitrification by the addition of C₂H₂ in the
345 cropping and sugarcane soils is mostly likely caused by the inhibitory effect of C₂H₂ on both
346 AOA and AOB growth. However, there was no inhibitory effect on AOA in vegetable soil
347 and AOB on day 7 in dairy pasture soil, suggesting that AOB rather than AOA were involved
348 in nitrification in the alkaline vegetable soil while in the dairy pasture soil AOB might be less
349 important in nitrification than AOA. Moreover, it is possible that C₂H₂ application may also
350 change active strains of ammonia oxidizers and different effectiveness possibly due to
351 different sensitive strains appearing in soils (Belser et al., 1980), and this needs further
352 investigation.

353 **4.2 Inhibitory effects of C₂H₂ at different incubation conditions.**

354 From both previous studies (Kool et al., 2010; Liu et al., 2017) and the current study, soil
355 water content and temperature were the predominant factors regulating N₂O emission from
356 soils. The cumulative N₂O emissions increased with increasing WFPS and temperature in the
357 cropping soil (Table 3). The significant increase in emissions at 35°C between 50% and 70%
358 WFPS is probably because these incubation conditions favoured nitrifying and denitrifying
359 enzyme synthesis (Liu et al., 2017), and 35°C at 70% WFPS is the most suitable condition for

360 microbial activity. The degree of inhibition of cumulative N₂O emission and nitrification by
361 C₂H₂ was shown to vary across the different incubation conditions, indicating that
362 temperature and moisture content significantly affect C₂H₂ efficacy. At lower soil
363 temperature (15°C) and in drier soil (50% WFPS), C₂H₂ addition had no inhibitory effect on
364 the total N₂O emissions, suggesting that nitrification may be heterotrophic under these
365 conditions in this soil. This result is supported by Liu et al. (2015c) where under 15°C and
366 50% WFPS incubation, nitrification was found to be predominantly heterotrophic. The low
367 efficacy of C₂H₂ to inhibit N₂O emission under wetter soils (70% WFPS) is probably because
368 more N₂O was produced from denitrification. Another possibility for the low efficacy of
369 C₂H₂ at the lower soil temperature of 15°C is that C₂H₂ may be unable to form a reactive
370 epoxide to inactivate the AMO enzyme (Hynes and Knowles, 1978).

371 Compared to 70% WFPS, the inhibitory effect of C₂H₂ was much greatest at inhibiting
372 nitrification and cumulative N₂O emission at 50% WFPS except at 15°C. This is possibly
373 because 25°C and 35°C with 50% WFPS are more suitable for nitrification to occur (Garrido
374 et al., 2002; Huang et al., 2014). During incubation, the nitrification rates in the fertilizer
375 treatments were similar at low water content (50% WFPS) regardless of soil temperature,
376 while the effectiveness of C₂H₂ on inhibiting nitrification was different (Table 3). This
377 possibly indicates that concentrations of C₂H₂ may need to be adjusted for optimal inhibition
378 under different soil temperature and moisture.

379 AOA and AOB reduced in abundance with varying degrees after C₂H₂ addition under
380 different environmental conditions, which might be attributed to the differential sensitivity of
381 AOA and AOB to C₂H₂ under different conditions. Under 15°C, 50% and 70% WFPS, C₂H₂
382 showed no effect on AOB populations. One possible explanation is that under lower soil
383 temperature (15°C), the active strains of AOB were less abundant than under higher
384 temperature resulting in less sensitivity to C₂H₂ addition. Another possible explanation is

385 AOA may prefer lower a soil temperature and be more involved in nitrification at 15°C than
386 AOB.

387 The underlying mechanism of how C₂H₂ targets AOA remains largely unknown. Acetylene
388 might suppress the growth of *amoA*-containing archaea by inactivating the archaeal AMO
389 protein, as demonstrated in AOB (Hyman and Wood 1985). However, Offre et al. (2009)
390 demonstrated that the enzymes and metabolic pathways of AOA might differ significantly
391 from those of AOB, in which C₂H₂ interferes with the AMO protein. Future studies based on
392 soil RNA and pure cultures are therefore necessary to elucidate the mechanism by which
393 C₂H₂ blocks the growth of AOA.

394 In the first incubation, we found that different soils have different C₂H₂ efficacy. Therefore,
395 we cannot ascertain whether the findings of the incubation study at different temperatures and
396 moistures can be extended to other soils. Future work with alkaline soils or a larger range of
397 soil properties is definitely needed to clarify this question.

398

399 **5. Conclusions**

400 In conclusion, C₂H₂ was more effective in inhibiting nitrification and N₂O emissions in acidic
401 soils and under drier condition (50% WFPS). Soil pH, C content, texture, temperature,
402 moisture and land use might be important factors affecting the efficacy of C₂H₂. The various
403 C₂H₂ inhibitory effects on nitrification and N₂O production were also linked to different
404 responses of the ammonia oxidizers to C₂H₂. Therefore, since the effect of C₂H₂ varied with
405 soils and environmental conditions, the assumption that N₂O production and nitrification can
406 be affected by low concentrations of C₂H₂ may need verification for specific soils and
407 conditions of interest.

408

409 **Acknowledgements**

410 The authors would like to acknowledge the financial support by Incitec Pivot, the Australian
411 Government Department of Agriculture through the Grains Research and Development
412 Corporation, and Australian Research Council (DE150100870, DP160101028,
413 LP160101134).

414

415 **Reference**

416 Bateman, E.J., and Baggs, E.M., 2005. Contributions of nitrification and denitrification to
417 N_2O emission from soils at different water-filled pore space. *Biol. Fertil. Soils.* 41,
418 379–388.

419 Belser, L.W., and Mays, E.L., 1980. Specific inhibition of nitrite oxidation by chlorate and its
420 use in assessing nitrification in soils and sediments. *Appl. Environ. Microb.* 39, 505–
421 510.

422 Berg, P., Klemmedtsson, L., Rosswall, T., 1982. Inhibitory effect of low partial pressures of
423 acetylene on nitrification. *Soil. Biol. Biochem.* 14, 301–303.

424 Boyle-Yarwood, S.A., Bottomley, P.J., Myrold, D.D., 2008. Community composition of
425 ammonia-oxidising bacteria and archaea in soils under stands of red alder and
426 Douglas fir in Oregon. *Environ. Microbiol.* 10, 2956–2965.

427 Bremner, J., and Blackmer, A.M., 1978. Nitrous oxide: emission from soils during
428 nitrification of fertilizer nitrogen. *Science.* 199, 295–296.

429 Burger, M., Jackson, L.E., Lundquist, E.J., Louie, D.T., Miller, R.L., Rolston, D.E., Scow,
430 K.M., 2005. Microbial responses and nitrous oxide emissions during wetting and

431 drying of organically and conventionally managed soil under tomatoes. *Biol. Fertil.*
432 *Soils.* 42, 109–118.

433 Butterbach-Bahl, K., Baggs, E.M., Dannenmann, M., Kiese, R., Zechmeister-Boltenstern, S.,
434 2013. Nitrous oxide emissions from soils: how well do we understand the processes and
435 their controls?. *Phil. Trans. R. Soc. B.* 368, DOI: 10.1098/rstb.2013.0122.

436 Chen, D.L., Suter, H.C., Islam, A., Edis, R., Freney, J.R., Walker, C.N., 2008. Prospects of
437 improving efficiency of fertilizer nitrogen in Australian agriculture; a review of
438 enhanced efficiency fertilizers. *Aust. J. Soil Res.* 46, 289–301.

439 Chen, D.L., Suter, H.C., Islam, A., Edis, R., 2010. Influence of nitrification inhibitors on
440 nitrification and nitrous oxide (N₂O) emission from a clay loam soil fertilized with
441 urea. *Soil. Biol. Biochem.* 42, 660–664.

442 Daum, M., Zimmer, W., Papen, H., Kloos, K., Nawrath, K., Bothe, H., 1998. Physiological and
443 molecular biological characterization of ammonia oxidation of the heterotrophic
444 nitrifier *Pseudomonas putida*. *Curr. Microbiol.* 37, 281–288.

445 Davidson, E.A., Swank, W.T., Perry, T.O., 1986. Distinguishing between nitrification and
446 denitrification as sources of gaseous nitrogen production in soil. *Appl. Environ.*
447 *Microbiol.* 52, 1280–1286.

448 De Boer, W., and Kowalchuk, G., 2001. Nitrification in acid soils: micro-organisms and
449 mechanisms. *Soil. Biol. Biochem.* 33, 853–866.

450 De Boer, W., Klein Gunnewiek, P.J.A., Veenhuis, M., Bock, E., Laanbroek, H.J., 1991.
451 Nitrification at low pH by aggregated autotrophic bacteria. *Appl. Environ. Microb.* 57,
452 3600–3604.

453 Di, H.J., Cameron, K.C., Sherlock, R., Shen, J.P., He, J.Z., Winefield, C., 2010. Nitrous
454 oxide emissions from grazed grassland as affected by a nitrification inhibitor,
455 dicyandiamide, and relationships with ammonia-oxidising bacteria and archaea. *J.*
456 *Soils Sediments*. 10, 943–954.

457 Di, H., Cameron, K., Shen, J., 2009. Nitrification driven by bacteria and not archaea in
458 nitrogen-rich grassland soils. *Nat. Geosci.* 2, 621–624.

459 Fisk, L.M., Maccarone, L.D., Barton, L., Murphy, D.V., 2015. Nitrapyrin decreased
460 nitrification of nitrogen released from soil organic matter but not *amoA* gene
461 abundance at high soil temperature. *Soil. Biol. Biochem.* 88, 214–223.

462 Francis, C.A., Roberts, K.J., Beman, J.M., Santoro, A.E., Oakley, B.B., 2005. Ubiquity and
463 diversity of ammonia-oxidizing archaea in water columns and sediments of the ocean.
464 *Proc. Natl. Acad. Sci. U.S.A.* 102, 14683–14688.

465 Garrido, F., Henault, C., Gaillard, H., Perez, S., Germon, J.C., 2002. N₂O and NO emissions
466 by agricultural soils with low hydraulic potentials. *Soil. Biol. Biochem.* 34, 559–575.

467 Gubry-Rangin, C., Nicol, G.W., Prosser, J.I., 2010. Archaea rather than bacteria control
468 nitrification in two agricultural acidic soils. *FEMS Microbiol. Ecol.* 74, 566–574.

469 He, J.Z., Hu, H.W., Zhang, L.M., 2012. Current insights into the autotrophic thaumarchaeal
470 ammonia oxidation in acidic soils. *Soil Biol. Biochem.* 55, 146-154.

471 He, J.Z., Shen, J.P., Zhang, L.M., Zhu, Y.G., Zheng, Y.M., Xu, M.G., Di, H., 2007.
472 Quantitative analyses of the abundance and composition of ammonia-oxidizing
473 bacteria and ammonia-oxidizing archaea of a Chinese upland red soil under long-term
474 fertilization practices. *Environ. Microbiol.* 9, 2364–2374.

475 Hink, L., Nicol, G.W., Prosser, J.I., 2016. Archaea produce lower yields of N₂O than bacteria
476 during aerobic ammonia oxidation in soil. *Environ. Microbiol.* doi:10.1111/1462-
477 2920.13282.

478 Hu, H.W., Chen, D.L., He, J.Z., 2015. Microbial regulation of terrestrial nitrous oxide
479 formation: understanding the biological pathways for prediction of emission rates.
480 *FEMS Microbiol. Rev.* 39, 729–49.

481 Huang, T., Gao, B., Hu, X.K., 2014. Ammonia-oxidation as an engine to generate nitrous
482 oxide in an intensively managed calcareous Fluvo-aquic soil. *Sci. Rep.* 4, 3950.

483 Hyman, M.R., Wood, P.M., 1985. Suicidal inactivation and labelling of ammonia
484 monooxygenase by acetylene. *Biochem. J.* 227, 719–725.

485 Hynes, R.K., Knowles, R., 1978. Inhibition by acetylene of ammonia oxidation in
486 *Nitrosomonas europaea*. *FEMS. Microbiol. Lett.* 4, 319–321.

487 IPCC. *Climate change 2007: the physical science basis*. In: Solomon, S., Qin, D., Manning,
488 M., (eds). *Contribution of Working Group I to the Fourth Assessment Report of the*
489 *Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University
490 Press, 2007.

491 Jia, Z., Conrad, R., 2009. Bacteria rather than Archaea dominate microbial ammonia
492 oxidation in an agricultural soil. *Environ. Microbiol.* 11, 1658–1671.

493 Kelly, K.B., Phillips, F.A., Baigent, R., 2008. Impact of dicyandiamide application on nitrous
494 oxide emissions from urine patches in northern Victoria, Australia. *Aust. J. Exp. Agr.*
495 48, 156–159.

496 Klemetsson, L., Svensson, B., Rosswall, T., 1988. A method of selective inhibition to
497 distinguish between nitrification and denitrification as sources of nitrous oxide in soil.
498 *Biol. Fertil. Soils.* 6, 112–119.

499 Kool, D.M., Wrage, N., Zechmeister-Boltenstern, S., 2010. Nitrifier denitrification can be a
500 source of N₂O from soil: a revised approach to the dual-isotope labelling method. *Eur.*
501 *J. Soil. Sci.* 61,759–772.

502 Levic̃nik-Höfferle, S., Nicol, G.W., Ausec, L., Mandic-Mulec, I., Prosser, J.I., 2012.
503 Stimulation of thaumarchaeal ammonia oxidation by ammonia derived from organic
504 nitrogen but not added inorganic nitrogen. *FEMS. Microbiol. Ecol.* 80, 114–123.

505 Linn, D.M., and J.W. Doran., 1984. Effect of water-filled pore space on carbon dioxide and
506 nitrous oxide production in tilled and nontilled soils. *Soil. Sci. Soc. Am. J.* 48,1267-
507 1272.

508 Liu, R., Hayden, H., Suter, H., He, J.Z., Chen, D.L., 2015a. The effect of nitrification
509 inhibitors in reducing nitrification and the ammonia oxidizer population in three
510 contrasting soils. *J. Soils. Sediments.* 15, 1113–1118.

511 Liu, R., Suter, H., Hayden, H., He, J.Z., Chen, D.L., 2015b. Nitrate production is mainly
512 heterotrophic in an acid dairy soil with high organic content in Australia. *Biol. Fertil.*
513 *Soils.* 51, 891.

514 Liu, R., Hayden, H., Suter, H., Hu, H.W., He, J.Z., Pauline, M.M., Chen, D.L., 2017. The
515 effect of temperature and moisture on the source of N₂O and contributions from
516 ammonia oxidizers in an agricultural soil. *Biol. Fertil. Soils.* 53, 141–152.

517 Liu, R., Suter, H., He, J. Hayden, H., Chen, D. 2015c. Influence of temperature and
518 moisture on the relative contributions of heterotrophic and autotrophic nitrification
519 to gross nitrification in an acid cropping soil. *J. Soils. Sediments.* 15, 2304.

520 Marsden, K.A., Marín-Martínez, A.J., Vallejo, A., Hill, P.W., Jones, D.L., Chadwick, D.R.,
521 2016. The mobility of nitrification inhibitors under simulated ruminant urine
522 deposition and rainfall: a comparison between DCD and DMPP. *Biol. Fertil. Soils.*
523 52, 491–503.

524 Moir, J.W.B., Crossman, L.C., Spiro, S., Richardson, D.J., 1996. The purification of
525 ammonia monooxygenase from *Paracoccus denitrificans*. *FEBS. Lett.* 387, 71–74.

526 Offre, P., Prosser, J.I., Nicol, G.W., 2009. Growth of ammonia-oxidizing archaea in soil
527 microcosms is inhibited by acetylene. *FEMS. Microbiol. Ecol.* 70, 99–108.

528 Persson, T., Wirén, A., 1995. Nitrogen mineralization and potential nitrification at different
529 depths in acid forest soils. *Plant. Soil.* 168, 55–65.

530 Ravishankara, A.R., Daniel, J.S., Portmann, R.W., 2009. Nitrous Oxide (N₂O): The
531 Dominant Ozone-Depleting Substance Emitted in the 21st Century. *Science.* 326, 123–
532 125.

533 Rotthauwe, J.H., Witzel, K.P., Liesack, W., 1997. The ammonia monooxygenase structural
534 gene *amoA* as a functional marker: molecular finescale analysis of natural ammonia
535 oxidizing populations. *Environ. Microbiol.* 63, 4704–4712.

536 Scheer, C., Rowlings, D.W., Firrel, M., Deuter, P., Morris, S., Grace, P.R., 2014. Impact of
537 nitrification inhibitor (DMPP) on soil nitrous oxide emissions from an intensive
538 broccoli production system in sub-tropical Australia. *Soil. Biol. Biochem.* 77, 243–
539 251.

540 Shen, J.P., Zhang, L.M., Zhu, Y.G., Zhang, J.B., He, J.Z., 2008. Abundance and composition
541 of ammonia-oxidizing bacteria and ammonia-oxidizing archaea communities of an
542 alkaline sandy loam. *Environ. Microbiol.* 10, 1601–1611.

543 Schmidt, I., Bock, E., 1998. Anaerobic ammonia oxidation by cell-free extracts of
544 *Nitrosomonas eutropha*. *Antonie. Van. Leeuwenhoek.* 73, 271–278.

545 Suter, H, Lam, S.K., Davies, R., 2014. Soil Science Australia National Soil Science
546 Conference, 24-27th November, Melbourne.

547 Wan, Y., Ju, X., Ingwersen J., Schwarz, U., Stange, C.F., Zhang, F., Streck, T., 2009. Gross
548 Nitrogen Transformations and Related Nitrous Oxide Emissions in an Intensively
549 Used Calcareous Soil. *Soil. Sci. Soc. Am. J.* 73, 102-112.

- 550 Wrage, N., Velthof, G.L., Oenema, O., Laanbroek, H.J., 2004. Acetylene and oxygen as
551 inhibitors of nitrous oxide production in *Nitrosomonas europaea* and *Nitrosospira*
552 *briensis*: a cautionary tale. *FEMS Microbiol. Ecol.* 47, 13–18.
- 553 Zhang, L.M., Hu, H.W., Shen, J.P., He, J.Z., 2012. Ammonia-oxidising archaea have more
554 important role than ammonia-oxidising bacteria in ammonia oxidation of strongly
555 acidic soils. *ISME J.* 6, 1032–1045.
- 556 Zhang, J.B., Muller, C., Cai, Z.C., 2015. Heterotrophic nitrification of organic N and its
557 contribution to nitrous oxide emissions in soils. *Soil. Biol. Biochem.* 84, 109–209.



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Liu, R; Hayden, HL; Hu, H; He, J; Suter, H; Chen, D

Title:

Effects of the nitrification inhibitor acetylene on nitrous oxide emissions and ammonia-oxidizing microorganisms of different agricultural soils under laboratory incubation conditions

Date:

2017-10-01

Citation:

Liu, R., Hayden, H. L., Hu, H., He, J., Suter, H. & Chen, D. (2017). Effects of the nitrification inhibitor acetylene on nitrous oxide emissions and ammonia-oxidizing microorganisms of different agricultural soils under laboratory incubation conditions. *APPLIED SOIL ECOLOGY*, 119, pp.80-90. <https://doi.org/10.1016/j.apsoil.2017.05.034>.

Persistent Link:

<http://hdl.handle.net/11343/208822>

File Description:

Accepted version