

3. Singer P, Blaser AR, Berger MM, Alhazzani W, Calder PC, Casaer MP, *et al.* ESPEN guideline on clinical nutrition in the intensive care unit. *Clin Nutr* 2019;38:48–79.
4. Marik PE, Hooper MH. Normocaloric versus hypocaloric feeding on the outcomes of ICU patients: a systematic review and meta-analysis. *Intensive Care Med* 2016;42:316–323.
5. Zusman O, Theilla M, Cohen J, Kagan I, Bendavid I, Singer P. Resting energy expenditure, calorie and protein consumption in critically ill patients: a retrospective cohort study. *Crit Care* 2016;20:367.
6. Allingstrup MJ, Kondrup J, Wiis J, Claudius C, Pedersen UG, Hein-Rasmussen R, *et al.* Early goal-directed nutrition versus standard of care in adult intensive care patients: the single-centre, randomised, outcome assessor-blinded EAT-ICU trial. *Intensive Care Med* 2017;43:1637–1647.

Copyright © 2020 by the American Thoracic Society



Reply to Peçanha Antonio *et al.*

From the Authors:

Dr. Peçanha Antonio and colleagues write that administration of at least 70% of estimated energy requirements to critically ill patients provides a survival advantage. However, randomized clinical trials are the gold-standard method for drawing causal inferences about therapies (1), and there is evidence from three well-conducted multicenter randomized trials involving 1,000, 894, and 3,957 patients, respectively, that mortality is unaffected by targeting a specific energy delivery during the acute phase of critical illness (2–4).

The letter also raises concerns that our study compared “overfeeding” with usual feeding practice, particularly given the proportion of patients with obesity in our cohort. The body mass index of our study participants (29.2 kg/m²) reflects the adult body mass index in our region (5). Those assigned the energy-dense formula received a mean (SD) of 29.1 (6.2) kcal/kg of ideal body weight/d and 23.2 (7.1) kcal/kg of actual body weight/d from the enteral route (*see* Table E4 in the online supplement of Ref. 5). We are not aware of any published data establishing that this dose of energy delivered via the enteral route represents overfeeding and/or is a risk factor for increased death. Moreover, for our primary outcome (Day 90 mortality), body mass index was a predefined subgroup of interest, and the relative risk of death (energy-dense formula vs. standard nutrition for the 1,423 participants with a body mass index \geq 30 kg/m²) was 0.94 (95% confidence interval, 0.77–1.14) (4). Finally, as there is no consensus definition for overfeeding or how to diagnose it, the belief that it has occurred in the subgroup with obesity is speculative at best.

Dr. Peçanha Antonio and colleagues also suggest that our study provides limited additional information, given the 6-month outcome data provided by a previous trial (6). We agree that the EAT-ICU (Early Goal-directed Nutrition in ICU Patients) trial was well conducted but assert that a single-center open-label trial

providing 6-month outcome data from 105 survivors does not diminish the information obtained from analyzing data from 2,492 survivors randomly assigned to receive a blinded intervention at 1 of 46 ICUs across two countries.

We agree that future trials of nutritional therapy, particularly focused on the time after the first 5–7 days, are warranted. However, the conduct of an adequately powered randomized trial to evaluate the effect on patient-centered outcomes of an intervention commencing after 5–7 days of admission to an ICU represents substantial challenges. We maintain that our trial has provided valuable insights into the current population-level effects of acute energy delivery on longer-term outcomes. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Adam M. Deane, Ph.D.*
Rinaldo Bellomo, Ph.D.
The University of Melbourne
Parkville, Victoria, Australia

Marianne J. Chapman, Ph.D.
Kylie Lange, B.Sc.
Sandra L. Peake, Ph.D.
University of Adelaide
Adelaide, South Australia, Australia

Paul Young, Ph.D.
Medical Research Institute of New Zealand
Wellington, New Zealand

Theodore J. Iwashyna, Ph.D.
University of Michigan
Ann Arbor, Michigan

ORCID IDs: 0000-0002-7620-5577 (A.M.D.); 0000-0002-3428-3083 (P.Y.).

*Corresponding author (e-mail: adam.deane@mh.org.au).

References

1. Balshem H, Helfand M, Schünemann HJ, Oxman AD, Kunz R, Brozek J, *et al.* GRADE guidelines: 3. rating the quality of evidence. *J Clin Epidemiol* 2011;64:401–406.
2. Rice TW, Wheeler AP, Thompson BT, Steingrub J, Hite RD, Moss M, *et al.*; National Heart, Lung, and Blood Institute Acute Respiratory Distress Syndrome (ARDS) Clinical Trials Network. Initial trophic vs full enteral feeding in patients with acute lung injury: the EDEN randomized trial. *JAMA* 2012;307:795–803.
3. Arabi YM, Aldawood AS, Haddad SH, Al-Dorzi HM, Tamim HM, Jones G, *et al.*; PermiT Trial Group. Permissive underfeeding or standard enteral feeding in critically ill adults. *N Engl J Med* 2015;372:2398–2408.
4. Chapman M, Peake SL, Bellomo R, Davies A, Deane A, Horowitz M, *et al.*; TARGET Investigators, for the ANZICS Clinical Trials Group. Energy-dense versus routine enteral nutrition in the critically ill. *N Engl J Med* 2018;379:1823–1834.
5. Deane AM, Little L, Bellomo R, Chapman MJ, Davies AR, Ferrie S, *et al.* Outcomes six months after delivering 100% or 70% of enteral calorie requirements during critical illness (TARGET): a randomized controlled trial. *Am J Respir Crit Care Med* 2020;201:814–822.
6. Allingstrup MJ, Kondrup J, Wiis J, Claudius C, Pedersen UG, Hein-Rasmussen R, *et al.* Early goal-directed nutrition versus standard of care in adult intensive care patients: the single-centre, randomised, outcome assessor-blinded EAT-ICU trial. *Intensive Care Med* 2017;43:1637–1647.

Copyright © 2020 by the American Thoracic Society

§This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0 (<http://creativecommons.org/licenses/by-nc-nd/4.0/>). For commercial usage and reprints, please contact Diane Gern (dgern@thoracic.org).

Originally Published in Press as DOI: 10.1164/rccm.202005-1810LE on June 9, 2020



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Deane, AM; Bellomo, R; Chapman, MJ; Lange, K; Peake, SL; Young, P; Iwashyna, TJ

Title:

Too Many Calories for All? Reply

Date:

2020-10-01

Citation:

Deane, A. M., Bellomo, R., Chapman, M. J., Lange, K., Peake, S. L., Young, P. & Iwashyna, T. J. (2020). Too Many Calories for All? Reply. AMERICAN JOURNAL OF RESPIRATORY AND CRITICAL CARE MEDICINE, 202 (7), pp.1060-1060.

<https://doi.org/10.1164/rccm.202005-1810LE>.

Persistent Link:

<http://hdl.handle.net/11343/251488>

File Description:

published version

License:

CC BY-NC-ND