

RESEARCH ARTICLE

Knowledge of risk factors for diabetes or cardiovascular disease (CVD) is poor among individuals with risk factors for CVD

Monique F. Kilkenny^{1,2*}, Libby Dunstan³, Doreen Busingye¹, Tara Purvis^{1,2}, Megan Reyneke¹, Mary Orgill⁴, Dominique A. Cadilhac^{1,2,5}

1 Stroke and Ageing Research, School of Clinical Sciences at Monash Health, Department of Medicine, Monash University, Clayton, Victoria, Australia, **2** Stroke Division: Public Health, the Florey Institute of Neurosciences and Mental Health, Heidelberg, Victoria, Australia, **3** Stroke Foundation, Brisbane, Queensland, Australia, **4** Stroke Foundation, Melbourne, Victoria, Australia, **5** The University of Melbourne, Victoria, Australia

* monique.kilkenny@monash.edu



OPEN ACCESS

Citation: Kilkenny MF, Dunstan L, Busingye D, Purvis T, Reyneke M, Orgill M, et al. (2017) Knowledge of risk factors for diabetes or cardiovascular disease (CVD) is poor among individuals with risk factors for CVD. *PLoS ONE* 12 (2): e0172941. doi:10.1371/journal.pone.0172941

Editor: Gianpaolo Reboldi, Università degli Studi di Perugia, ITALY

Received: March 11, 2016

Accepted: February 13, 2017

Published: February 28, 2017

Copyright: © 2017 Kilkenny et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data are available from the Monash University Ethics Committee. Please contact the Stroke Foundation Manager of the Evaluation (who provided evaluation of the Know your numbers Program) to seek approval email via admin@strokefoundation.com.au

Funding: Dr Monique Kilkenny is supported by a NHMRC Early Career Fellowships (1109426). A/ Prof Dominique Cadilhac is supported by a NHMRC Public Health/National Heart Foundation Future Leader Research Fellowship (1063761). We

Abstract

Background

There is limited evidence on whether having pre-existing cardiovascular disease (CVD) or risk factors for CVD such as diabetes, ensures greater knowledge of risk factors important for motivating preventative behaviours. Our objective was to compare knowledge among the Australian public participating in a health check program and their risk status.

Methods

Data from the Stroke Foundation 'Know your numbers' program were used. Staff in community pharmacies provided opportunistic health checks (measurement of blood pressure and diabetes risk assessment) among their customers. Participants were categorised: 1) CVD ± risk of CVD: history of stroke, heart disease or kidney disease, and may have risk factors; 2) risk of CVD only: reported having high blood pressure, high cholesterol, diabetes or atrial fibrillation; and 3) CVD risk free (no CVD or risk of CVD). Multivariable logistic regression analyses were performed including adjustment for age and sex.

Findings

Among 4,647 participants, 12% had CVD (55% male, 85% aged 55+ years), 47% were at risk of CVD (40% male, 72% 55+ years) and 41% were CVD risk free (33% male, 27% 55+ years). Participants with CVD (OR: 0.66; 95% CI: 0.55, 0.80) or risk factors for CVD (OR: 0.65; 95% CI: 0.57, 0.73) had poorer knowledge of the risk factors for diabetes/CVD compared to those who were CVD risk free. After adjustment, only participants with risk factors for CVD (OR: 0.80; 95% CI: 0.69, 0.93) had poorer knowledge. Older participants (55+ years) and men had poorer knowledge of diabetes/CVD risk factors and complications of diabetes.

acknowledge that the Know your numbers Program was a government funded program in the Australian states of New South Wales Health and Queensland Health.

Competing interests: The authors have declared that no competing interests exist

Conclusions

Participants with poorer knowledge of risk factors were older, more often male or were at risk of developing CVD compared with those who were CVD risk free. Health education in these high risk groups should be a priority, as diabetes and CVD are increasing in prevalence throughout the world.

Introduction

The prevalence of diabetes mellitus (diabetes), and particularly type 2 diabetes, is considered a global epidemic [1]. It is projected to become the leading cause of disease burden for males and the second leading cause for females by 2023 [2]. Diabetes is an independent risk factor for cardiovascular disease (CVD). CVD mainly includes coronary heart disease, heart failure, cardiomyopathy and stroke [3]. About 11.7 million adult Australians (95%) have at least one of the major modifiable risk factors for stroke and heart disease including high blood pressure (BP), diabetes, high cholesterol, smoking, high consumption of alcohol, poor diet, obesity or inadequate physical activity [4]. Furthermore, knowledge of risk factors for CVD and diabetes in the community is poor [5].

The risk factors for diabetes and CVD overlap. Type 2 diabetes results from a combination of genetic and environmental factors. Although there is a strong genetic predisposition to the development of diabetes, the risk is greatly increased when associated with other factors such as high BP, being overweight or obese, physical inactivity and poor diet. It is unclear whether individuals with CVD or risk factors for CVD including diabetes are aware of the risk factors for diabetes/CVD compared to those without CVD or its risk factors. There are limited data on whether individuals with diabetes or who are at high absolute risk of developing diabetes are aware of the complications of diabetes (such as CVD events, amputation, blindness or nerve damage).

Health promotion programs in the community such as the Know your numbers (KYN) program, which involved opportunistic CVD screening assessments in pharmacies, may improve community awareness and knowledge of CVD and associated risk factors such as diabetes [6,7].

Our aim was to compare KYN participants' knowledge of risk factors for diabetes/CVD and complications associated with diabetes based on their risk status. We hypothesised that knowledge about risk factors or complications would be better in people with pre-existing disease or who were at risk of CVD or diabetes than among those who were risk free. This is because these people with known risk or pre-existing disease should have received some health promotion education if the health system is being effective.

Methods

KYN diabetes program

Since 2007, the Stroke Foundation in Australia has undertaken the KYN program which aims to improve public awareness and early detection of individuals at risk of stroke and other CVD [6,7]. The program is designed to facilitate opportunistic CVD and diabetes risk screening assessments in pharmacy and community settings targeting people aged 50+ years. However, subjects aged less than 50 years were still able to participate in the health check program. The features of this program include a standardised BP check, self-reported risk factor checklist,

education brochures and doctor referral letter if required. Since 2011 the KYN program has included diabetes risk assessment and educational information on diabetes [8]. The diabetes risk assessment included the participants completing a self-administered screening questionnaire called the Australian Type 2 Diabetes Risk Assessment Tool (AUSDRISK) [9]. A score is calculated from the results of the questionnaire that determines a level of risk for the person developing type 2 diabetes within the next five years. A score of 12 or more means that the participant is at high risk of developing type 2 diabetes within five years or having undiagnosed type 2 diabetes [9]. Participation in KYN provides an important opportunity for members of the community to learn about their CVD and diabetes risk including health-related behaviours (e.g. physical inactivity) or biomedical factors (e.g. high BP) and receive health promotion education [6–8].

Study design

For this observational study, data from the 2012–2014 KYN program delivered in Australian community pharmacies in the states of Queensland and New South Wales were used. Detailed information was collected from the participant on the day of the health check (called “Participant questionnaire”) from pharmacies who volunteered to collect these data. This questionnaire included information on demographics, self-reported health history, awareness of the KYN health check, diabetes risk assessment and assessment of knowledge of causes of diabetes/CVD and complications of diabetes e.g. stroke or heart attack (Table 1). The questionnaires were developed during the pilot phases of KYN with feedback from the participants [8].

Covariate measures

Risk factor measurements. The AUSDRISK was used as part of the assessment of risk of diabetes within the KYN program [8]. The AUSDRISK score and the BP reading were recorded on the participant questionnaire on the day of the health check. A participant was classified as **High Risk** if they recorded a high BP reading ($\geq 140/90$ mmHg) or a high AUSDRISK score of 12 or more.

Socioeconomic status (SES) was defined using the Index of Relative Socioeconomic Disadvantage (IRSD) [10]. Pharmacy address was used as a surrogate for a participant’s place of

Table 1. Example of self-reported medical history and knowledge questions from the Participant Questionnaire collected at the time of the Know your numbers health check.

Your medical history? (Please cross all that apply)		
<input type="checkbox"/> High Blood pressure (hypertension)	<input type="checkbox"/> Diabetes	<input type="checkbox"/> High cholesterol (Dyslipidemia)
<input type="checkbox"/> Previous stroke/ Transient Ischaemic Attack (TIA)	<input type="checkbox"/> Current smoker	<input type="checkbox"/> Ex-smoker
<input type="checkbox"/> Ischaemic heart disease	<input type="checkbox"/> Previous heart disease or heart attack	<input type="checkbox"/> Atrial Fibrillation (AF)
<input type="checkbox"/> Kidney Disease	<input type="checkbox"/> Peripheral vascular Disease	<input type="checkbox"/> None of the above
What do you think causes diabetes and cardiovascular diseases (e.g. stroke or heart attack)? (Please cross all that apply)		
<input type="checkbox"/> Being overweight	<input type="checkbox"/> Making unhealthy food choices e.g. not eating enough fruit/vegetables	
<input type="checkbox"/> High blood pressure	<input type="checkbox"/> High cholesterol	<input type="checkbox"/> Smoking
<input type="checkbox"/> Chronic kidney disease	<input type="checkbox"/> Diabetes	<input type="checkbox"/> Family history of CVD or diabetes
<input type="checkbox"/> Being physically inactive (e.g.: less than 30 minutes a day on most days of week)		
<input type="checkbox"/> Drinking too much alcohol (e.g.: more than 2 standard drinks per day)		
Diabetes can lead to serious complications. What are these complications? (Please cross all that apply)		
<input type="checkbox"/> Heart disease	<input type="checkbox"/> Developing blindness	<input type="checkbox"/> Nerve damage
<input type="checkbox"/> Stroke	<input type="checkbox"/> Amputation	<input type="checkbox"/> Kidney disease
<input type="checkbox"/> Circulation problems	<input type="checkbox"/> Unsure	

doi:10.1371/journal.pone.0172941.t001

residence. These were matched to local government area (LGA) and then matched to the IRSD and corresponding quintiles were derived. Quintile 1 (IRSD 1) refers to area of most socioeconomic disadvantage and Quintile 5 (IRSD 5) refers to area of least socioeconomic disadvantage. **High SES:** includes quintiles 4 and 5 of the Index of Relative Socioeconomic Disadvantage.

Knowledge assessment: Risk factors for diabetes/CVD. Participants were asked to select from a pre-defined list of ten risk factors for diabetes and CVD (Table 1). Knowledge scores were calculated for each participant based on the number of options selected. Participants received one point for each of the ten response options. Knowledge scores ranged from 0 to 10.

Knowledge assessment: Complications of diabetes. Participants were asked to select the complications of diabetes from a pre-defined list (Table 1). Knowledge scores were calculated for each participant based on the number of options selected as complications of diabetes. Participants received one point for each of the seven complications of diabetes correctly selected. Knowledge scores ranged from 0 to 7.

Definition of subject groups

- **CVD risk status:** Participants were categorised into the following CVD risk groups based on their self-reported medical history (Table 1): 1) **CVD ± risk of CVD** included stroke/transient ischaemic attack (TIA), heart disease/heart attack, peripheral vascular disease, ischaemic heart disease, or kidney disease. They may have also reported risk factors including high BP, high cholesterol, diabetes, or atrial fibrillation; 2) **Risk of CVD** only included high BP, high cholesterol, diabetes or atrial fibrillation; 3) **CVD risk free** if the participant did not report any of these options.
- **Diabetes risk status:** Participants were categorised into the following diabetes risk groups based on their self-reported medical history (Table 1) or their AUSDRISK score. 1) **Diabetes** included self-reported diagnosis of diabetes; 2) **Risk of diabetes** if the participant did not have a self-reported diagnosis of diabetes but recorded a high AUSDRISK score of 12 or more; 3) **Diabetes risk free** if the participant did not have a self-reported diagnosis of diabetes and recorded an AUSDRISK scored less than 12.

Analyses

Chi-square test for categorical variables and Kolmogorov Smirnov test for continuous data were used to compare participant characteristics and knowledge of risk factors and complications. To determine whether history of CVD or diabetes risk was associated with greater knowledge, univariable and multivariable logistic regression analyses were performed using the knowledge score (lesser knowledge group coded as zero and greater knowledge group coded as one) as a dependent variable. The lesser knowledge group were those with a knowledge score below the median score while the greater knowledge group had a knowledge score at or above the median score. The multivariable models were adjusted for risk status, age, sex, socioeconomic status and high risk (based on results of health check). Level of significance was $p < 0.05$. All analyses were performed using Intercool STATA 12.1 for Windows software (Stata Corp PL, 2013).

Ethics

Ethics approval for the evaluation was provided by the Monash University Human Research Ethics Committee (HREC CF12/1103–2012000528). Written informed consent was not

obtained from the participant to collect detailed information on the “Participant questionnaire”. However, consent was implied, as evidenced that they completed the form. Information was collected anonymously from participants and was de-identified in these analyses.

Results

Questionnaire data were completed by 4,647 participants from pharmacies in Queensland and New South Wales (Australia) participating in the KYN program from 2012 to 2014. More than half of the participants (55%) were aged over 54 years and 39% were male. Following the KYN health check, 2,318 participants (50%) were assessed to be at high risk. One third of participants (37%) recorded a high BP and 45% were assessed as having a high 5-year risk of developing diabetes (AUSDRISK score 12+). Of the participants who had CVD, one third reported history of stroke/TIA; 23% ischaemic heart disease; 20% kidney disease; 15% peripheral vascular disease and 49% heart disease/heart attack. The participants who had CVD also reported the following risk factors: 76% had high BP; 47% high cholesterol; 30% diabetes and 14% atrial fibrillation. Of the participants who were categorised as at risk of CVD only: 83% had high BP; 37% high cholesterol; 22% diabetes and 5% atrial fibrillation.

CVD risk status

Univariable analyses of the participant characteristics according to CVD status are provided in [Table 2](#). Participants attending a pharmacy in an area of relative socioeconomic advantage (high SES) were more often CVD risk free. Compared with participants who were CVD risk free, participants with pre-existing or risk of CVD were more often older (55 years or more), male and had recorded a high BP reading or high AUDRISK score on the day of the health check.

Univariable analyses of knowledge according to CVD status on the same day of the health check are provided in [Table 3](#). Compared with participants who were CVD risk free, participants with CVD or risk of CVD more often had poorer knowledge of risk factors for diabetes/CVD such as being physically inactive, or having diabetes. The median knowledge score was 6 (Q1: 3, Q3: 9 [Q1: 25th percentile, Q3: 75th percentile]). To assess factors associated with knowledge of risk factors, participants were divided into two groups: the group with the lesser knowledge (knowledge median score <6) and the group with greater knowledge (knowledge median score 6+) in multivariable analyses ([Table 4](#)).

Multivariable analyses of knowledge of risk factors of diabetes/CVD according to CVD status are provided in [Table 4](#). Interestingly, having CVD or risk factors for CVD was associated with poorer knowledge of the risk factors for diabetes/CVD. However, on adjustment for age, sex, high risk (based on health check) and socioeconomic status, this association remained in individuals with risk factors for CVD only.

Diabetes risk status

Compared with participants who were diabetes risk free, participants with risk of diabetes were more likely to have poor knowledge of complications of diabetes such as stroke or circulation problems. The median knowledge score was 4 (Q1: 2, Q3: 6). Participants were divided into two groups, with those with poor knowledge having a score less than four (the median knowledge score), and the group with greater knowledge having a median knowledge score of 4 or more ([Table 5](#)).

In contrast, having diabetes was not independently associated with good knowledge of the complications of diabetes ([Table 6](#)) compared to those who were diabetes risk free. Individuals with risk factors for diabetes were associated with having poorer knowledge of the

Table 2. Characteristics of participants by CVD risk status.

Characteristics	CVD ¹ (N = 552) n (%)	Risk of CVD ² (N = 2,185) n (%)	CVD risk free ³ (N = 1,910) n (%)	p-value CVD vs CVD Risk free	p-value Risk of CVD vs CVD Risk free
Age group (years)				<0.001	<0.001
<34 years	7 (1)	57 (3)	576 (31)		
35–44	18 (3)	160 (7)	343 (18)		
45–54	57 (10)	388 (18)	436 (23)		
55–64	107 (20)	640 (30)	295 (16)		
65–74	195 (36)	567 (26)	151 (8)		
75+	162 (30)	356 (16)	66 (4)		
Age 55+ years	464 (85)	1,563 (72)	512 (27)	<0.001	<0.001
Patient characteristics					
Male	293 (55)	849 (40)	608 (33)	<0.001	<0.001
Australian	400 (72)	1,470 (67)	1,308 (68)	0.07	0.41
Aboriginal/Torres Strait Islander	15 (3)	37 (2)	38 (2)	0.30	0.48
Socioeconomic status (SES)				0.15	0.002
IRSD 1	66 (12)	359 (17)	233 (13)		
IRSD 2	112 (21)	369 (17)	300 (16)		
IRSD 3	135 (25)	530 (25)	478 (26)		
IRSD 4	142 (26)	539 (25)	511 (27)		
IRSD 5	86 (16)	353 (16)	340 (18)		
High SES (IRSD 4/5)	228 (42)	892 (41)	851 (45)	0.14	0.007
Results of KYN check					
High 5-year risk* diabetes	218 (81)	816 (63)	355 (23)	<0.001	<0.001
High BP reading (\geq 140/90 mmHg)	228 (46)	990 (49)	373 (21)	<0.001	<0.001
High BP/High 5-year risk diabetes*	346 (63)	1,374 (63)	598 (31)	<0.001	<0.001

CVD: Cardiovascular disease

¹CVD ± risk of CVD: heart disease, heart attack, stroke/transient ischaemic attack, ischaemic heart disease, kidney disease, and may have high blood pressure, high cholesterol, diabetes, atrial fibrillation

²Risk of CVD only: high blood pressure, high cholesterol, diabetes atrial fibrillation

³CVD risk free: no CVD or risk of CVD; KYN: Know your numbers program

*based on AUSDRISK score 12+; IRSD: Index of Relative Socioeconomic Disadvantage.

doi:10.1371/journal.pone.0172941.t002

complications of diabetes compared to those who were diabetes risk free. However, this association did not remain after adjustment for age, sex, socioeconomic position and diabetes risk status. In all analyses, being older (aged 55 years or more) and male were consistently associated with poorer knowledge of the complications of diabetes.

Discussion

Evidence about individuals' knowledge of risk factors associated with diabetes/CVD and complications of diabetes is rare. This study contributes new knowledge for developed countries on the association between knowledge of risk factors for diabetes/CVD and complications of diabetes in individuals with pre-existing disease or who are at risk, and those who were risk free. The findings provide evidence that knowledge of risk factors for diabetes/CVD was poorer in participants who were at risk of CVD who were more often male and over 55 years.

Table 3. Univariable analysis of knowledge of risk factors for CVD by CVD risk status.

Risk factors for diabetes/CVD	CVD ¹ (N = 552) n (%)	Risk of CVD ² (N = 2,185) n (%)	CVD Risk free ³ (N = 1,910) n (%)	p-value CVD vs CVD Risk free	p-value Risk of CVD vs CVD Risk free
Being physically inactive	322 (59)	1,263 (58)	1,258 (67)	<0.001	<0.001
Unhealthy foods	345 (63)	1,392 (64)	1,328 (70)	0.001	<0.001
Being overweight	444 (81)	1,840 (85)	1,630 (86)	0.001	0.20
Kidney disease	167 (30)	551 (25)	601 (32)	0.51	<0.001
High blood pressure	397 (72)	1,573 (73)	1,347 (71)	0.70	0.38
Diabetes	228 (41)	911 (42)	964 (51)	<0.001	<0.001
Drinking too much alcohol	279 (51)	1,022 (47)	1,102 (58)	0.001	<0.001
High cholesterol	340 (62)	1,280 (59)	1,233 (65)	0.13	<0.001
Smoking	328 (60)	1,251 (58)	1,287 (68)	<0.001	<0.001
Family history	345 (63)	1,339 (62)	1,325 (70)	0.001	<0.001
Knowledge score				0.001	<0.001
0	32 (6)	111 (5)	96 (5)		
1–3	123 (22)	499 (23)	313 (17)		
4–7	191 (35)	800 (37)	614 (33)		
8–10	203 (37)	755 (35)	863 (46)		
Median score (Q1, Q3)	6 (3, 9)	6 (3, 9)	7 (4, 9)	<0.001	<0.001

CVD: Cardiovascular disease

¹CVD ± risk of CVD: heart disease, heart attack, stroke/transient ischaemic attack, ischaemic heart disease, kidney disease and may have high blood pressure, high cholesterol, diabetes atrial fibrillation

²Risk of CVD only: high blood pressure, high cholesterol, diabetes atrial fibrillation

³CVD risk free: no CVD or risk of CVD. Q1: 25th percentile; Q3: 75th percentile.

doi:10.1371/journal.pone.0172941.t003

The factors associated with poor knowledge of risk factors of CVD/diabetes were older age, being male and those participants who recorded a high BP or risk of diabetes (at the health check).

Table 4. Multivariable analysis of factors associated with knowledge of risk factors for diabetes/CVD.

Median knowledge score of 6+	Univariable analysis OR (95% CI)	p-value	Multivariable analysis OR (95% CI)	p-value
Age (55+ years)	0.59 (0.52, 0.66)	<0.001	0.69 (0.60, 0.79)	<0.001
Male	0.75 (0.66, 0.84)	<0.001	0.80 (0.70, 0.91)	0.001
High BP/High 5-year risk diabetes* (at health check)	0.64 (0.53, 0.76)	<0.001	0.79 (0.70, 0.90)	<0.001
CVD risk status				
CVD risk free ³	1.0 (reference group)		1.0 (reference group)	
Risk of CVD²	0.65 (0.57, 0.73)	<0.001	0.80 (0.69, 0.93)	0.003
CVD¹	0.66 (0.55, 0.80)	<0.001	0.87 (0.70, 1.08)	0.20
High SES ⁴	1.07 (0.95, 1.21)	0.24	1.07 (0.95, 1.21)	0.28

CVD: Cardiovascular disease

¹CVD ± CVD risk: heart disease, heart attack, stroke/transient ischaemic attack, ischaemic heart disease, kidney disease and may have high blood pressure, high cholesterol, diabetes, atrial fibrillation

²Risk of CVD only: high blood pressure, high cholesterol, diabetes atrial fibrillation

³CVD risk free: no CVD or risk of CVD

*based on AUSDRISK score 12+

⁴High SES: quintile 4 and 5 of the Index of Relative Socioeconomic Disadvantage; Multivariable model adjusted for age, sex, high BP/high 5-year risk diabetes and CVD risk status.

doi:10.1371/journal.pone.0172941.t004

Table 5. Univariable analysis of knowledge of complications of diabetes by diabetes risk status.

Complications of diabetes	Diabetes ¹ (N = 634) n (%)	Risk of diabetes ² (N = 1,389) n (%)	Diabetes risk free ³ (N = 1,702) n (%)	p-value Diabetes vs Diabetes risk free	p-value Risk of diabetes vs Diabetes risk free
Kidney disease	315 (50)	636 (46)	923 (54)	0.06	<0.001
Nerve damage	294 (47)	573 (41)	819 (48)	0.48	<0.001
Amputation	188 (30)	432 (31)	526 (31)	0.58	0.88
Developing blindness	392 (62)	819 (59)	1,082 (64)	0.47	0.01
Circulation problems	401 (63)	864 (62)	1,180 (69)	0.006	<0.001
Stroke	398 (63)	814 (59)	1,095 (64)	0.52	0.001
Heart disease	445 (70)	894 (65)	1,150 (68)	0.20	0.07
Unsure	48 (8)	92 (7)	114 (7)	0.45	0.94
Knowledge score				0.28	<0.001
0	52 (8)	127 (9)	152 (9)		
1–3	229 (36)	557 (40)	557 (33)		
4–7	351 (56)	701 (51)	991 (58)		
Median score (Q1, Q3)	4 (2, 6)	4 (2, 6)	4 (2, 6)	0.21	<0.001

¹Diabetes: reported in medical history

²Risk of diabetes: AUSDRISK score recorded 12+

³Diabetes risk free: no diabetes, or AUSDRISK score recorded <12; Q1: 25th percentile; Q3: 75th percentile.

doi:10.1371/journal.pone.0172941.t005

Our study contrasts with a United States study of 4,193 healthy people in which knowledge of CVD risk factors was associated with presence of risk factors [11]. However, this study was conducted in young adults (mean age 30 years), compared to our study where half of the participants were aged 55 years or older. Our findings were similar in univariable analyses to those of a study from China in which knowledge of risk factors for stroke was reported to be poorer in patients with previous stroke or TIA [12]. Although risk of diabetes was not independently associated with knowledge of complications, this does not negate the fact that individuals with pre-existing diabetes were more likely to mention the following complications of diabetes: kidney disease, nerve damage, blindness, circulation problems and stroke. This further highlights the need for public health awareness programs such as KYN, aimed at educating the public about CVD risks and complications, particularly those related to diabetes.

Table 6. Multivariable analysis of factors associated with greater knowledge of complications of diabetes.

Median knowledge score of 4+	Univariable analysis OR (95% CI)	p-value	Multivariable analysis OR (95% CI)	p-value
Age (55+ years)	0.66 (0.58, 0.74)	<0.001	0.72 (0.62, 0.83)	<0.001
Male	0.75 (0.66, 0.84)	<0.001	0.77 (0.67, 0.89)	<0.001
Diabetes status				
Diabetes risk free ¹	1.0 (reference group)		1.0 (reference group)	
Risk of diabetes²	0.73 (0.64, 0.85)	<0.001	0.88 (0.75, 1.03)	0.11
Diabetes³	0.89 (0.74, 1.07)	0.23	1.14 (0.93, 1.40)	0.21
High SES ⁴	0.92 (0.82, 1.04)	0.17	0.98 (0.86, 1.12)	0.80

¹Diabetes risk free: no diabetes, or AUSDRISK score recorded <12

²Risk of diabetes: AUSDRISK score recorded 12+

³Diabetes: reported in medical history

⁴High SES: quintile 4 and 5 of the Index of Relative Socioeconomic Disadvantage; Multivariable model adjusted for age, sex, socioeconomic position and diabetes risk status.

doi:10.1371/journal.pone.0172941.t006

Previous research shows that older age, lower educational status and limited English language are associated with poor knowledge of diabetes/CVD risk factors or complications [13]. In one study, it was reported that age had a negative association with knowledge of risk factors for diabetes in a study conducted in Saudi Arabia [14]. However, in a small Australian study of 181 patients with type 2 diabetes, there was no association found between older age and knowledge of diabetes [13]. In our study, educational status and English fluency were not collected as part of our participant questionnaire, so we were unable to adjust for these factors in our regression models. Nevertheless, there is substantial evidence from other studies that show lower socioeconomic position, measured by level of education, to be associated with poorer knowledge of risk factors [11,13].

Our findings highlight poor knowledge of risk factors for diabetes and CVD in a high risk group. Given that better knowledge of diabetes is associated with ability to perform self-care activities (e.g. proper diet, regular exercise and blood glucose monitoring) [15], and better medication adherence and glycaemic control [16], this underscores the need for patient and community education programs for diabetes. Moreover, educational interventions for diabetes have been shown to improve patients' knowledge about diabetes, clinical, behavioural, and patient-centred outcomes and complications associated with diabetes [17–19]. Perhaps, as we have shown previously, volunteer-led community interventions such as the NSF StrokeSafe Ambassador Program could be adopted to improve knowledge and awareness of diabetes and CVD risk factors and complications of diabetes in high risk groups as well as in communities [20]. The Ambassador program included presentations by volunteers trained to be 'ambassadors' to spread standard information about stroke to the public on stroke risks and warning signs.

The potential limitations of the KYN data have been noted in previous publications as related to the KYN program [6–8]. In brief, the group of participants being tested was a convenience sample and may not be representative of the general population. The questionnaires used were pilot tested and refined based on feedback from participants, KYN staff and the working group to ensure face validity including readability and completeness. Other forms of reliability and validity were not undertaken. We also may have overestimated levels of knowledge by using close-ended questions in which specific risk factors were listed as opposed to using open-ended questions in the participant questionnaire. The strengths of our study include the large sample of pharmacies and community-based participants.

Increasing awareness of CVD and diabetes in the population is important, given the high proportion of participants with pre-existing CVD/diabetes or at risk of CVD/diabetes who were unaware of the risk factors and complications of diabetes. There is a need for further education of people who attend health checks who record a high BP or risk of diabetes, as these people have inadequate knowledge of risk factors for CVD/diabetes. Diabetes Australia have developed education programs related to diabetes and complications [13]. These education programs could be modified to target older individuals and men in these high risk groups to improve knowledge and awareness of diabetes and CVD.

In summary, the findings of this study suggest that individuals who are at risk of CVD, male or over 55 years have poorer knowledge of risk factors for CVD/diabetes. Targeting these groups with tailored educational information may help improve knowledge of risk factors and their understanding of the complications of diabetes.

Acknowledgments

We thank staff in participating pharmacies, volunteers from Rotary and study participants without whose support this program would not be possible. We acknowledge the input from

the Advisory Committee and especially Li Chun Quang for the design of the questionnaires using Teleform and staff at the Florey Institute of Neuroscience and Mental Health who undertook data processing; Dr Erin Lalor (former Chief Executive Officer, Stroke Foundation) for the initial development of the program; Belinda Wilkinson (National Know your numbers Coordinator) and Roslyn Johnson (National Know your numbers Manager) for assisting with the coordination of the program.

Author Contributions

Conceptualization: MK DC.

Data curation: MK LD.

Formal analysis: MK DB MR.

Funding acquisition: LD MO DC.

Investigation: LD MO MK.

Methodology: MK DC TP.

Project administration: LD DC MO MK.

Resources: DC MK LD MO.

Software: MK DC MR.

Supervision: DC MO LD MK.

Validation: DB MR TP.

Visualization: MK DB MR TP.

Writing – original draft: MK DB TP.

Writing – review & editing: DC MK TP MR DB LD MO.

References

1. Guariguata L, Whiting DR, Hambleton I, Beagley J, Linnenkamp U, et al. (2014) Global estimates of diabetes prevalence for 2013 and projections for 2035. *Diabetes research and clinical practice* 103: 137–149. doi: [10.1016/j.diabres.2013.11.002](https://doi.org/10.1016/j.diabres.2013.11.002) PMID: [24630390](https://pubmed.ncbi.nlm.nih.gov/24630390/)
2. Begg SJ, Vos T, Barker B, Stanley L, Lopez AD (2008) Burden of disease and injury in Australia in the new millennium: measuring health loss from diseases, injuries and risk factors. *Med J Aust* 188: 36–40. PMID: [18205562](https://pubmed.ncbi.nlm.nih.gov/18205562/)
3. Alliance NVDP (2012) Guidelines for the management of absolute cardiovascular risk. Melbourne: National Heart Foundation of Australia.
4. Australian Institute of Health and Welfare (2012) Australia's Health 2012. Canberra: AIHW.
5. Slark J, Bentley P, Majeed A, Sharma P (2012) Awareness of stroke symptomatology and cardiovascular risk factors amongst stroke survivors. *J Stroke Cerebrovasc Dis* 21: 358–362. doi: [10.1016/j.jstrokecerebrovasdis.2010.09.010](https://doi.org/10.1016/j.jstrokecerebrovasdis.2010.09.010) PMID: [21111631](https://pubmed.ncbi.nlm.nih.gov/21111631/)
6. Cadilhac DA, Kilkenny MF, Johnson R, Wilkinson B, Amatya B, et al. (2015) The Know Your Numbers (KYN) program 2008 to 2010: impact on knowledge and health promotion behavior among participants. *International journal of stroke: official journal of the International Stroke Society* 10: 110–116.
7. Kilkenny MF, Johnson RR, Cadilhac DA, National Stroke F (2010) Feasibility of a pilot programme to increase awareness of blood pressure as an important risk factor for stroke in Australia. *International journal of stroke: official journal of the International Stroke Society* 5: 344–350.
8. Kilkenny MF, Johnson R, Andrew NE, Purvis T, Hicks A, et al. (2014) Comparison of two methods for assessing diabetes risk in a pharmacy setting in Australia. *BMC public health* 14: 1227. doi: [10.1186/1471-2458-14-1227](https://doi.org/10.1186/1471-2458-14-1227) PMID: [25427845](https://pubmed.ncbi.nlm.nih.gov/25427845/)

9. Chen L, Magliano DJ, Balkau B, Colagiuri S, Zimmet PZ, et al. (2010) AUSDRISK: an Australian Type 2 Diabetes Risk Assessment Tool based on demographic, lifestyle and simple anthropometric measures. *The Medical journal of Australia* 192: 197–202. PMID: [20170456](#)
10. Adams RJ, Howard N, Tucker G, Appleton S, Taylor AW, et al. (2009) Effects of area deprivation on health risks and outcomes: a multilevel, cross-sectional, Australian population study. *International Journal of Public Health* 54: 183–192. doi: [10.1007/s00038-009-7113-x](#) PMID: [19214382](#)
11. Lynch EB, Liu K, Kiefe CI, Greenland P (2006) Cardiovascular disease risk factor knowledge in young adults and 10-year change in risk factors: the Coronary Artery Risk Development in Young Adults (CARDIA) Study. *American Journal of Epidemiology* 164: 1171–1179. doi: [10.1093/aje/kwj334](#) PMID: [17038418](#)
12. Zeng Y, He GP, Yi GH, Huang YJ, Zhang QH, et al. (2012) Knowledge of stroke warning signs and risk factors among patients with previous stroke or TIA in China. *Journal of Clinical Nursing* 21: 2886–2895. doi: [10.1111/j.1365-2702.2012.04118.x](#) PMID: [22985321](#)
13. Fenwick EK, Xie J, Rees G, Finger RP, Lamoureux EL (2013) Factors associated with knowledge of diabetes in patients with type 2 diabetes using the Diabetes Knowledge Test validated with Rasch analysis. *PLoS ONE [Electronic Resource]* 8: e80593.
14. Aljoudi AS, Taha AZ (2009) Knowledge of diabetes risk factors and preventive measures among attendees of a primary care center in eastern Saudi Arabia. *Annals of Saudi Medicine* 29: 15–19. doi: [10.4103/0256-4947.51813](#) PMID: [19139622](#)
15. Persell SD, Keating NL, Landrum MB, Landon BE, Ayanian JZ, et al. (2004) Relationship of diabetes-specific knowledge to self-management activities, ambulatory preventive care, and metabolic outcomes. *Preventive Medicine* 39: 746–752. doi: [10.1016/j.yjmed.2004.02.045](#) PMID: [15351541](#)
16. Al-Qazaz HK, Sulaiman SA, Hassali MA, Shafie AA, Sundram S, et al. (2011) Diabetes knowledge, medication adherence and glycemic control among patients with type 2 diabetes. *International Journal of Clinical Pharmacy* 33: 1028–1035. doi: [10.1007/s11096-011-9582-2](#) PMID: [22083724](#)
17. Lee TI, Yeh YT, Liu CT, Chen PL (2007) Development and evaluation of a patient-oriented education system for diabetes management. *International Journal of Medical Informatics* 76: 655–663. doi: [10.1016/j.ijmedinf.2006.05.030](#) PMID: [16815741](#)
18. Molsted S, Tribler J, Poulsen PB, Snorgaard O (2012) The effects and costs of a group-based education programme for self-management of patients with Type 2 diabetes. A community-based study. *Health Education Research* 27: 804–813. doi: [10.1093/her/cyr053](#) PMID: [21712503](#)
19. Norris SL, Engelgau MM, Narayan KMV (2001) Effectiveness of self management training in type 2 diabetes—A systematic review of randomized controlled trials. *Diabetes Care* 24: 561–587. PMID: [11289485](#)
20. Kilkenny MF, Purvis T, Werner M, Reyneke M, Czerenkowski J, et al. (2016) Improving stroke knowledge through a 'volunteer-led' community education program in Australia. *Prev Med* 86: 1–5. doi: [10.1016/j.yjmed.2016.01.015](#) PMID: [26820114](#)



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Kilkenny, MF; Dunstan, L; Busingye, D; Purvis, T; Reyneke, M; Orgill, M; Cadilhac, DA

Title:

Knowledge of risk factors for diabetes or cardiovascular disease (CVD) is poor among individuals with risk factors for CVD

Date:

2017-02-28

Citation:

Kilkenny, M. F., Dunstan, L., Busingye, D., Purvis, T., Reyneke, M., Orgill, M. & Cadilhac, D. A. (2017). Knowledge of risk factors for diabetes or cardiovascular disease (CVD) is poor among individuals with risk factors for CVD. PLOS ONE, 12 (2), <https://doi.org/10.1371/journal.pone.0172941>.

Persistent Link:

<http://hdl.handle.net/11343/258289>

File Description:

Published version

License:

CC BY