

Correction

Correction: Gravely, S., et al. Awareness, Trial, and Current Use of Electronic Cigarettes in 10 Countries: Findings from the ITC Project. *Int. J. Environ. Res. Public Health* 2014, *11*, 11691–11704

Shannon Gravely¹, **Geoffrey T. Fong**^{1,2,*}, **K. Michael Cummings**³, **Mi Yan**¹, **Anne C. K. Quah**¹, **Ron Borland**⁴, **Hua-Hie Yong**⁴, **Sara C. Hitchman**⁵, **Ann McNeill**⁵, **David Hammond**⁶, **James F. Thrasher**^{7,8}, **Marc C. Willemsen**⁹, **Hong Gwan Seo**¹⁰, **Yuan Jiang**¹¹, **Tania Cavalcante**¹², **Cristina Perez**¹², **Maizurah Omar**¹³ and **Karin Hummel**⁹

¹ Department of Psychology, University of Waterloo, 200 University Avenue West, Waterloo, Ontario, N2L 3G1, Canada; E-Mails: shannon.gravely@uwaterloo.ca (S.G.); m6yan@uwaterloo.ca (M.Y.); ackquah@uwaterloo.ca (A.C.K.Q.)

² Ontario Institute for Cancer Research, MaRS Center, West Tower, 661 University Avenue, Suite 510, Toronto, Ontario, M5G 0A3, Canada

³ Medical University of South Carolina, 171 Ashley Ave, Charleston, SC 29425, USA; E-Mail: cummingk@musc.edu

⁴ The Cancer Council Victoria, 615 St. Kilda Road, Melbourne, Victoria 3004, Australia; E-Mails: Ron.Borland@cancervic.org.au (R.B.); Hua.Yong@cancervic.org.au (H.-H.Y.)

⁵ King's College London, Strand, London, WC2R 2LS, UK; E-Mails: sara.hitchman@kcl.ac.uk (S.C.H.); ann.mcneill@kcl.ac.uk (A.M.)

⁶ School of Public Health and Health Systems, University of Waterloo, 200 University Avenue West, Waterloo, Ontario, N2L 3G1, Canada; E-Mail: dhammond@uwaterloo.ca

⁷ Department of Health Promotion, Education & Behavior, Arnold School of Public Health, University of South Carolina, Columbia, SC 29208, USA; E-Mail: thrasher@mailbox.sc.edu

⁸ National Institute of Public Health, Universidad N. 655, Colonia Stana Maria Ahuacatitlan, C.P. 62100, Cuernavaca, Mexico

⁹ Maastricht University (CAPHRI), P. Debyeplein 1, 6229 HA Maastricht, the Netherlands; E-Mails: marc.willemsen@maastrichtuniversity.nl (M.C.W.); Karin.hummel@maastrichtuniversity.nl (K.H.)

¹⁰ National Cancer Center of Korea, 323 Ilsan-ro, Ilsandong-gu, Goyang-si Gyeonggi-do, 410-769, Korea; E-Mail: hongwan@ncc.re.kr

¹¹ China Center for Disease Control and Prevention, 155 Changbai Road, Changping District, Beijing 102206, China; E-Mail: jiangyuan88@vip.sina.com

¹² National Cancer Institute of Brazil, Praca Cruz Vermelha, 23, Centro, 20230-130, Rio de Janeiro, Brazil; E-Mails: taniac@inca.gov.br (T.C.); cperez@inca.gov.br (C.P.)

¹³ University Sains Malaysia, Jalan Sungai Dua, 11800 Georgetown, Pulau Pinang, Malaysia; E-Mail: maizurahomar@hotmail.com

* Author to whom correspondence should be addressed; E-Mail: gfong@uwaterloo.ca; Tel.: +1-519-888-4567 (ext. 33597); Fax: +1-519-746-8631.

Academic Editor: Paul B. Tchounwou

Received: 21 April 2015 / Accepted: 22 April 2015 / Published:

The authors wish to make the following amendments to their paper published in *International Journal of Environmental Research and Public Health* [1]. These amendments only apply to the data from Brazil:

1. Page 11692, the abstract, line 15, the data collection for “Awareness of e-cigarettes” in Brazil should be 37%, and the data for “Trial of Electronic Cigarettes” in Brazil should be 8%. The correct sentence should be “There was considerable cross-country variation by year of data collection and for awareness of e-cigarettes (Netherlands (2013: 88%), Republic of Korea (2010: 79%), United States (2010: 73%), Australia (2013: 66%), Malaysia (2011: 62%), United Kingdom (2010: 54%), Canada (2010: 40%), Brazil (2013: 37%), Mexico (2012: 34%), and China (2009: 31%)), in self-reports of ever having tried e-cigarettes (Australia, (20%), Malaysia (19%), Netherlands (18%), United States (15%), Republic of Korea (11%), United Kingdom (10%), Brazil (8%), Mexico (4%), Canada (4%), and China (2%)), and in current use (Malaysia (14%), Republic of Korea (7%), Australia (7%), United States (6%), United Kingdom (4%), Netherlands (3%), Canada (1%), and China (0.05%))”.
2. Page 11696, Table 1, the data for Brazil are corrected. There are no changes in the data of any of the other countries. The corrected Table 1 is as follows:

Table 1. Sociodemographic and Smoking Characteristics and Patterns of E-Cigarette Use.

Country	N	Dates of Data Collection (Survey Mid-Point ‡)	Survey Mode	Response Rate (Average Retention Rate)	N (%) Female	Age (Mean ± SD)	% Aware of E-Cigarettes (95% CI)	% Ever-Tried E-Cigarettes (95% CI)	% Currently Using E-Cigarettes (95% CI)
China	5583				297 (5%)	50 ± 13	31% (28.7–34.0)	2% (1.8, 2.9)	0% (0.0–0.1)
<i>Cigarette Smokers</i>	5209	May 2009–Oct 2009 (Jul-2009)	Face-to-Face	52.8% ** (81.0%)	270 (5%)	50 ± 13	31% (28.4–33.8)	2% (1.8–3.0)	~0%
<i>Recent quitters</i>	103				7 (7%)	52 ± 13	40% (29.9–50.8)	2% (0–4.4)	0%
United Kingdom	1325				726 (55%)	49 ± 13	54% (50.9–57.9)	10% (7.1–12.1)	4% (2.5–6.5)
<i>Cigarette Smokers</i>	977	Jul 2010–Jun 2011 (Aug-2010)	Web or Phone	37.8% (72.8%)	544 (56%)	49 ± 13	56% (51.9–60.0)	11% (8.0–13.9)	5% (2.8–7.0)
<i>Recent quitters</i>	77				46 (60%)	47 ± 13	60% (44.4–74.9)	16% (0–32.9)	11% (0–28.5)
United States	1520				805 (53%)	51 ± 13	73% (70.5–76.4)	15% (12.1–17.8)	6% (3.6–7.6)
<i>Cigarette Smokers</i>	1262	Jul 2010–Jun 2011 (Aug-2010)	Web or Phone	25.6% (63.4%)	671 (53%)	52 ± 13	74% (70.5–76.8)	18% (14.2–21.0)	6% (4.0–8.8)
<i>Recent quitters</i>	63				28 (44%)	50 ± 14	72% (54.6–90.0)	10% (1.0–19.9)	7% (0–15.8)

Table 1. Cont.

Country	N	Dates of Data Collection (Survey Mid-Point ‡)	Survey Mode	Response Rate (Average Retention Rate)	N (%) Female	Age (Mean ± SD)	% Aware of E-Cigarettes (95% CI)	% Ever-Tried E-Cigarettes (95% CI)	% Currently Using E-Cigarettes (95% CI)
Canada *	1581				872 (55%)	47 ± 12	40% (36.5–42.6)	4% (2.7–5.3)	1% (0.6–2.1)
<i>Cigarette Smokers</i>	1243	Jul 2010–Jun 2011 (Aug-2010)	Web or Phone	49.5% (73.0%)	683 (55%)	48 ± 12	40% (36.1–43.0)	4% (2.9–6.0)	2% (0.8–2.7)
<i>Recent quitters</i>	74				34 (46%)	47 ± 14	39% (25.2–53.7)	5% (0–12.4)	0% -
Republic of Korea	1753				83 (5%)	49 ± 16	79% (77.0–81.2)	12% (10.4–14.1)	7% (5.4–8.4)
<i>Cigarette Smokers</i>	1560	Oct 2010–Dec 2010 (Nov-2010)	Phone	14.5% (50.3%)	76 (5%)	49 ± 16	80% (77.6–82.0)	13% (10.9–14.9)	7% (5.7–8.9)
<i>Recent quitters</i>	51				2 (4%)	51 ± 15	75% (61.0–89.3)	23% (9.3–36.5)	13% (1.5–24.1)
Malaysia *	1998				22 (1%)	33 ± 23	62% (57.5–66.1)	19% (16.2–22.6)	14% (11.6–15.7)
<i>Cigarette Smokers</i>	1773	May 2011–Feb 2012 (May-2011)	Phone	N/A (56.7%)	16 (1%)	31 ± 12	62% (57.4–67.0)	21% (17.3–24.2)	15% (12.4–17.0)
<i>Recent quitters</i>	69				3 (4%)	32 ± 14	69% (53.1–85.8)	13% (3.2–22.2)	6% (0.4–11.5)

Table 1. Cont.

Country	N	Dates of Data Collection (Survey Mid-Point ‡)	Survey Mode	Response Rate (Average Retention Rate)	N (%) Female	Age (Mean ± SD)	% Aware of E-Cigarettes (95% CI)	% Ever-Tried E-Cigarettes (95% CI)	% Currently Using E-Cigarettes (95% CI)
Mexico *,†	2129				801 (38%)	41 ± 15	34% (30.0–37.5)	4% (3.1–5.8)	-
<i>Cigarette Smokers</i>	1747	Oct 2012–Dec 2012 (Nov-2012)	Face-to-Face	80.7% (73.6%)	646 (37%)	40 ± 15	34% (30.0–37.8)	5% (3.4–6.7)	-
<i>Recent quitters</i>	70				25 (36%)	41 ± 15	48% (28.5–68.0)	3% (0.02–6.9)	-
Brazil *,†	1215				787 (65%)	49 ± 14	37% (34.0–41.0)	8% (5.7–10.1)	-
<i>Cigarette Smokers</i>	1090	Oct 2012–Feb 2013 (Jan-2013)	Phone	10.6% (41.4%)	714 (65%)	49 ± 14	37% (33.7–41.0)	8% (5.8–10.7)	-
<i>Recent quitters</i>	45				25 (56%)	46 ± 13	38% (23.7–55.4)	8% (2.4–22.9)	-
Australia *	1492				801 (54%)	47 ± 13	66% (62.7–69.1)	20% (17.1–22.9)	7% (4.7–8.5)
<i>Cigarette Smokers</i>	1093	Feb 2013–Sep 2013 (Mar-2013)	Web or Phone	45.8% (74.5%)	586 (54%)	48 ± 13	69% (65.6–73.0)	24% (20.4–27.7)	9% (6.2–11.4)
<i>Recent quitters</i>	88				46 (52%)	44 ± 13	55% (41.2–68.1)	16% (6.8–26.0)	2% (0–5.4)

Table 1. Cont.

Country	N	Dates of Data Collection (Survey Mid-Point ‡)	Survey Mode	Response Rate (Average Retention Rate)	N (%) Female	Age (Mean ± SD)	% Aware of E-Cigarettes (95% CI)	% Ever-Tried E-Cigarettes (95% CI)	% Currently Using E-Cigarettes (95% CI)
Netherlands	1849				907 (49%)	40 ± 15	88% (86.4–90.4)	19% (16.4–20.7)	3% (2.4–4.1)
<i>Cigarette Smokers</i>	1420	May 2013–Jun 2013 (May-2013)	Web	64.6% (81.3%)	686 (48%)	41 ± 15	87% (84.8–89.6)	20% (17.4–22.5)	4% (2.8–5.0)
<i>Recent quitters</i>	284				154 (54%)	38 ± 15	92% (88.9–95.5)	14% (8.9–19.5)	1% (0–1.9)

Notes: Between-country comparisons cannot be made due to differences in survey timing and sequence of questioning in the survey; *Smokers* and *Recent quitters* categories will not add to the overall total because ‘recent quitters’ have been restricted to those who reported quitting within 6 months of the survey; Response rate for each country are for Wave 1 (except for Mexico, where the Wave 2 replenishment response rate is used as a surrogate) and an average of retention rates up to the wave being analyzed are reported above. Sample characteristics (gender and age) are unweighted and all other results are weighted by the rescaled cross-sectional weights; Prevalence estimates were rounded to the nearest whole number; SD = Standard deviation; CI = Confidence interval; N/A = Not available; * Countries where the sale of e-cigarettes containing nicotine is banned or restricted; ** estimated (exact rate could not be obtained); † The question about current use of e-cigarettes was not asked in the survey; ‡ The ‘survey date mid-point’ is the month/year on which 50% of the respondents had successfully completed the survey for that wave. Dates are listed in chronological order (earliest to latest date surveyed).

The authors would like to apologize for any inconvenience caused to readers by these changes.

Reference

1. Gravely, S.; Fong, G.T.; Cummings, K.M.; Yan, M.; Quah, A.C.K.; Borland, R.; Yong, H.-H.; Hitchman, S.C.; McNeill, A.; Hammond, D.; *et al.* Awareness, trial, and current use of electronic cigarettes in 10 countries: Findings from the ITC Project. *Int. J. Environ. Res. Public Health* **2014**, *11*, 11691–11704.

© 2015 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Gravelly, S; Fong, GT; Cummings, KM; Yan, M; Quah, ACK; Borland, R; Yong, H-H; Hitchman, SC; McNeill, A; Hammond, D; Thrasher, JF; Willemsen, MC; Seo, HG; Jiang, Y; Cavalcante, T; Perez, C; Omar, M; Hummel, K

Title:

Awareness, Trial, and Current Use of Electronic Cigarettes in 10 Countries: Findings from the ITC Project (vol 11, pg 11691, 2014)

Date:

2015-05-01

Citation:

Gravelly, S., Fong, G. T., Cummings, K. M., Yan, M., Quah, A. C. K., Borland, R., Yong, H. - H., Hitchman, S. C., McNeill, A., Hammond, D., Thrasher, J. F., Willemsen, M. C., Seo, H. G., Jiang, Y., Cavalcante, T., Perez, C., Omar, M. & Hummel, K. (2015). Awareness, Trial, and Current Use of Electronic Cigarettes in 10 Countries: Findings from the ITC Project (vol 11, pg 11691, 2014). INTERNATIONAL JOURNAL OF ENVIRONMENTAL RESEARCH AND PUBLIC HEALTH, 12 (5), pp.4631-4637. <https://doi.org/10.3390/ijerph120504631>.

Persistent Link:

<http://hdl.handle.net/11343/259079>

File Description:

Published version

License:

CC BY