

Complete Genome Sequence of *Leptospira interrogans* Serovar Bratislava, Strain PigK151

David P. Alt,^a Jennifer H. Wilson-Welder,^a Darrell O. Bayles,^a Caroline Cameron,^b Ben Adler,^c Dieter M. Bulach,^c Torsten Seemann,^d Michael J. Lehane,^e Lee R. Haines,^e Alistair C. Darby,^f Neil Hall,^f Alan D. Radford,^g Richard L. Zuerner^{a*}

National Animal Disease Center, Infectious Bacterial Diseases Research Unit, Ames, Iowa, USA^a; Department of Biochemistry and Microbiology, University of Victoria, Victoria, British Columbia, Canada^b; Departments of Microbiology and Biochemistry and Molecular Biology, Australian Research Council Centre of Excellence in Structural and Functional Microbial Genomics, Monash University, Clayton, Victoria, Australia^c; Victorian Bioinformatics Consortium, Monash University, Clayton, Victoria, Australia^d; Department of Vector Biology, Liverpool School of Tropical Medicine, Liverpool, United Kingdom^e; Institute of Integrative Biology, University of Liverpool, Liverpool, United Kingdom^f; Institute of Infection and Global Health, University of Liverpool, Liverpool, United Kingdom^g

* Present address: Richard L. Zuerner (retired), Aitkin, Minnesota, USA.

***Leptospira interrogans* serovar Bratislava infection occurs in multiple domestic and wildlife species and is associated with poor reproductive performance in swine and horses. We present the complete genome assembly of strain PigK151 comprising two chromosomes, CI (4.457 Mbp) and CII (358 kbp).**

Received 19 May 2015 Accepted 20 May 2015 Published 25 June 2015

Citation Alt DP, Wilson-Welder JH, Bayles DO, Cameron C, Adler B, Bulach DM, Seemann T, Lehane MJ, Haines LR, Darby AC, Hall N, Radford AD, Zuerner RL. 2015. Complete genome sequence of *Leptospira interrogans* serovar Bratislava, strain PigK151. *Genome Announc* 3(3):e00678-15. doi:10.1128/genomeA.00678-15.

Copyright © 2015 Alt et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Jennifer H. Wilson-Welder, jennifer.wilson-welder@ars.usda.gov.

Worldwide, leptospirosis is one of the most widespread zoonoses. The genus *Leptospira* contains pathogens serologically classified into over 250 serovars, intermediate pathogens, and saprophytes with genetic classification into 21 different species (1–3).

L. interrogans serovar Bratislava has been isolated from both free-living and domestic species (4), swine with reproductive problems (5–7), dogs with renal disease (8), and horses (4, 9). In horses, poor reproductive performance has been linked with seroreactivity (10).

Current complete genome sequences have been published for three strains of *L. interrogans*, all from human clinical cases, representing serovars Lai and Copenhageni. Here, we report the genome sequence of *L. interrogans* serovar Bratislava strain PigK151 comprising two chromosomes, CI (4.457 Mbp) and CII (358 kbp), obtained from a sow's kidney in Iowa in 1989.

High-quality genomic DNA was isolated from 5×10^{10} to 2×10^{11} pelleted *L. interrogans* following Marmur (11) with modifications. The pellet was suspended in 5 ml of 50 mM Tris-EDTA and frozen at -70°C for 1 h. Room-temperature lysozyme solution was added with mixing; the cells were thawed and then placed on ice for 45 min. One milliliter of lysis buffer (950 μl 0.5% SDS, 50 mM Tris pH 7.5, 10 mM EDTA; and 50 μl of 20 mg/ml proteinase K, 50 mM Tris pH 8.0, 1 mM CaCl_2) was added and mixed at 50°C for 1 h. Three rounds of extraction using Phase Lock Gel (5Prime) treatment with RNase-It (Agilent Technologies, Inc.) and a final chloroform extraction were performed.

Sequencing was performed using both Illumina HiSeq and Roche FLX-Titanium chemistry. Libraries were prepared according to manufacturers' directions. A fully closed genome consisting of two chromosomes was assembled using MIRA version 3.4.0 (12) coupled with the Roche gsAssembler version 2.8. MIRA as-

semblies were hybrid assemblies comprising Roche FLX sequencing reads and Illumina paired-end reads providing $40\times$ and $55\times$ genome coverage, respectively. Roche gsAssembler assemblies used only Roche sequencing data obtained from GS FLX shotgun, GS FLXplus, and GS FLX Titanium large-insert (2.7 kb and 3.7 kb) mate pair sequencing reads. Illumina sequencing data consisted of 2×54 -bp paired-end reads. Genome sequencing gaps were closed using either Gap Resolution version 1.2.2 (U.S. Department of Energy, Joint Genome Institute; <http://jgi.doe.gov/data-and-tools>) on the gsAssembler or by primer design followed by PCR amplification and Sanger sequencing. Editing was performed using GAP5 (13), from the Staden Package. The chromosomes were given a final polish by using SEQuel (14) to map Illumina reads back at approximately $180\times$ total coverage to correct any remaining base calling and pyrosequencing errors in the assembly.

Annotation was performed using the NCBI Prokaryotic Genome Annotation Pipeline. The genome comprises two chromosomes: CI (4.457 Mbp) contains 3,486 coding sequences, 37 tRNAs covering all 20 amino acids, a single copy of the 5S rRNA, and two copies of both 16S and 23S rRNA; and CII (358 kbp) contains 287 coding sequences. Availability of additional complete genomic sequence data will enable better molecular comparisons and enhance the understanding of leptospiral pathogenicity.

Nucleotide sequence accession numbers. The annotated assembly is available in GenBank under the accession numbers CP011410 (CI) and CP011411 (CII).

ACKNOWLEDGMENTS

Portions of this project were supported by a grant from the Royal College of Veterinary Surgeons (RCVS) Golden Jubilee Trust Fund (GJ08 546).

Roche 454 GS FLX titanium sequencing was performed at the Centre for Genomic Research at the University of Liverpool and at the National

Animal Disease Center. Illumina sequencing was performed at the Iowa State University DNA Facility.

We thank Rick Hornsby for his excellent technical support in this project.

Mention of trade names or commercial products in this article is solely for providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture.

REFERENCES

1. Lehmann JS, Matthias MA, Vinetz JM, Fouts DE. 2014. Leptospiral pathogenomics. *Pathogens* 3:280–308. doi:10.3390/pathogens3020280.
2. Smythe L, Adler B, Hartskeerl RA, Galloway RL, Turenne CY, Levett PN, International Committee on Systematics of Prokaryotes Subcommittee on the Taxonomy of *Leptospiraceae*. 2013. Classification of *Leptospira* genomospecies 1, 3, 4 and 5 as *Leptospira alstonii* sp. nov., *Leptospira vanthieli* sp. nov., *Leptospira terpstrae* sp. nov. and *Leptospira yanagawae* sp. nov., respectively. *Int J Syst Evol Microbiol* 63:1859–1862. <http://dx.doi.org/10.1099/ijs.0.047324-0>.
3. Saito M, Villanueva SY, Kawamura Y, Iida K, Tomida J, Kanemaru T, Kohno E, Miyahara S, Umeda A, Amako K, Gloriani NG, Yoshida S. 2013. *Leptospira idonii* sp. nov., isolated from environmental water. *Int J Syst Evol Microbiol* 63:2457–2562. <http://dx.doi.org/10.1099/ijs.0.047233-0>.
4. Hathaway SC, Little TW, Stevens AE, Ellis WA, Morgan J. 1983. Serovar identification of leptospires of the australis serogroup isolated from free-living and domestic species in the United Kingdom. *Res Vet Sci* 35:64–68.
5. Bolin CA, Cassells JA. 1990. Isolation of *Leptospira interrogans* serovar Bratislava from stillborn and weak pigs in Iowa. *J Am Vet Med Assoc* 196:1601–1604.
6. Ellis WA, McParland PJ, Bryson DG, Cassells JA. 1986. Prevalence of *Leptospira* infection in aborted pigs in Northern Ireland. *Vet Rec* 118:63–67. <http://dx.doi.org/10.1136/vr.118.3.63>.
7. Ellis WA, McParland PJ, Bryson DG, Thiermann AB, Montgomery J. 1986. Isolation of leptospires from the genital tract and kidneys of aborted sows. *Vet Rec* 118:294–295. <http://dx.doi.org/10.1136/vr.118.11.294>.
8. Nielsen JN, Cochran GK, Cassells JA, Hanson LE. 1991. *Leptospira interrogans* serovar Bratislava infection in two dogs. *J Am Vet Med Assoc* 199:351–352.
9. Rocha T, Ellis WA, Montgomery J, Gilmore C, Regalla J, Brem S. 2004. Microbiological and serological study of leptospirosis in horses at slaughter: first isolations. *Res Vet Sci* 76:199–202. <http://dx.doi.org/10.1016/j.rvsc.2003.12.003>.
10. Pinna A, Martins G, Hamond C, Medeiros MA, de Souza GN, Lilenbaum W. 2014. Potential differences between *Leptospira* serovars, host-adapted (Bratislava) and incidental (Copenhageni), in determining reproductive disorders in embryo transfer recipient mares in Brazil. *Vet Rec* 174:531. <http://dx.doi.org/10.1136/vr.101444>.
11. Marmor J. 1961. A procedure for the isolation of deoxyribonucleic acid from micro-organisms. *J Mol Biol* 3:208–218. [http://dx.doi.org/10.1016/S0022-2836\(61\)80047-8](http://dx.doi.org/10.1016/S0022-2836(61)80047-8).
12. Chevreur B, Wetter T, Suhai S. 1999. Genome sequence assembly using trace signals and additional sequence information, p 45–56. *In* Computer science and biology: Proceedings of the German Conference on Bioinformatics, GCB, Hannover, Germany.
13. Bonfield JK, Whitwham A. 2010. Gap5—editing the billion fragment sequence assembly. *Bioinformatics* 26:1699–1703. <http://dx.doi.org/10.1093/bioinformatics/btq268>.
14. Ronen R, Boucher C, Chitsaz H, Pevzner P. 2012. SEQuel: improving the accuracy of genome assemblies. *Bioinformatics* 28:i188–i196. <http://dx.doi.org/10.1093/bioinformatics/bts219>.



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Alt, DP; Wilson-Welder, JH; Bayles, DO; Cameron, C; Adler, B; Bulach, DM; Seemann, T; Lehane, MJ; Haines, LR; Darby, AC; Hall, N; Radford, AD; Zuerner, RL

Title:

Complete Genome Sequence of *Leptospira interrogans* Serovar Bratislava, Strain PigK151

Date:

2015-05-01

Citation:

Alt, D. P., Wilson-Welder, J. H., Bayles, D. O., Cameron, C., Adler, B., Bulach, D. M., Seemann, T., Lehane, M. J., Haines, L. R., Darby, A. C., Hall, N., Radford, A. D. & Zuerner, R. L. (2015). Complete Genome Sequence of *Leptospira interrogans* Serovar Bratislava, Strain PigK151. MICROBIOLOGY RESOURCE ANNOUNCEMENTS, 3 (3), <https://doi.org/10.1128/genomeA.00678-15>.

Persistent Link:

<http://hdl.handle.net/11343/259298>

File Description:

Published version

License:

CC BY