

Udayasiri Dilshan (Orcid ID: 0000-0002-3943-7862)
MacCallum Caroline (Orcid ID: 0000-0002-3258-9900)
Da Silva Nigel (Orcid ID: 0000-0003-3522-7419)

2

Title Page

Title

Impact of hospital geographic remoteness on short-term outcomes after colorectal cancer resection using state-wide administrative data.

Short running head

Outcomes post resection CRC in Victoria

Authors

Dilshan K. Udayasiri

First author

ORCID: 0000-0002-3943-7862

Degrees: M.B.B.S., F.R.A.C.S.

Academic title: Dr

Main appointment:

Colorectal research fellow

Institution/department:

Colorectal Surgical Unit, The Royal Melbourne Hospital, Victoria, Australia

Department of General Surgical Specialties, The Royal Melbourne Hospital, Victoria, Australia

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1111/ans.15992](https://doi.org/10.1111/ans.15992)

Department of Surgery, The University of Melbourne, Victoria, Australia

Caroline MacCallum

ORCID: 0000-0002-3258-9900

Degrees: M.S., M.B.B.S.

Academic title: Dr

Institution/department:

Colorectal Surgical Unit, The Royal Melbourne Hospital, Victoria, Australia

Department of General Surgical Specialties, The Royal Melbourne Hospital, Victoria, Australia

Department of Surgery, The University of Melbourne, Victoria, Australia

Nigel Da Silva

ORCID: 0000-0003-3522-7419

Degrees: M.S., M.B.B.S., F.R.A.C.S.

Institution/department:

Colorectal Surgical Unit, The Royal Melbourne Hospital, Victoria, Australia

Department of General Surgical Specialties, The Royal Melbourne Hospital, Victoria, Australia

Department of Surgery, The University of Melbourne, Victoria, Australia

Anita Skandarajah*

ORCID 0000-0001-7702-4850

Degrees: M.B.B.S., M.D., F.R.A.C.S., F.A.C.S

Academic title: Dr

Main Appointment:

Breast and Endocrine Surgeon

Co Supervisor for research

Institution/department:

Department of General Surgical Specialties, The Royal Melbourne Hospital, Victoria, Australia

Department of Surgery, The University of Melbourne, Victoria, Australia

Ian P. Hayes*

ORCID 0000000243682510

Degrees: M.B.B.S., M.S., M.Epi., F.R.A.C.S., F.R.C.S. (Gen-Surg)

Academic title: Clinical Associate Professor

Main Appointment:

Chief Supervisor of Research

Head of Colorectal Surgical Unit

Colorectal Surgeon

Institution/department:

Colorectal Surgical Unit, The Royal Melbourne Hospital, Victoria, Australia

Department of General Surgical Specialties, The Royal Melbourne Hospital, Victoria, Australia

Department of Surgery, The University of Melbourne, Victoria, Australia

*** Co-senior authors**

Author responsible for correspondence about the manuscript

Dr Dilshan Udayasiri

9 Cook Street, Ivanhoe, Victoria, Australia, postcode 3079

Telephone: +61431059868

Email: dilshan.udayasiri@gmail.com

Sources of support

Condon Colonic Disorder Scholarship, Medicine, Dentistry and Health Sciences, The University of Melbourne.

Health Technology Assessment Scholarship, The Royal Australian College of Surgeons

Word count abstract: 200

Word count body of text: 2780

Word Count Bibliography: 1037

Word Count tables: 500

Total Word Count: 4517

Number of tables: 2

Number of supplementary Figures: 2

[Disclaimers](#)

There are no conflicts of interest in this article.

Abstract

Purpose

This study aimed to use validated coding algorithms, applied to a central repository of administrative data (AD), to report on short-term outcomes following resection for colorectal cancer (CRC) comparing regional to metropolitan Victorian hospitals.

Methodology

A retrospective cohort study using prospectively-gathered AD.

The primary outcome was prolonged length of stay (LOS).

Secondary outcomes were: inpatient mortality, return to theatre (RTT), discharge destination and need for mechanical ventilation/intensive care unit support.

Outcomes were adjusted for potential confounders via multivariable logistic regression analysis.

Results

This study of 18470 patients found strong evidence for lower odds of prolonged LOS (OR 0.53, 95% CI 0.48 – 0.58, $p < 0.001$) and inpatient mortality (OR 0.67, 95%CI 0.49 – 0.91, $p = 0.01$) in inner regional hospital compared with metropolitan hospitals. For outer regional hospitals, there was strong evidence of decreased odds of prolonged LOS (OR 0.64, 95%CI 0.52 – 0.77, $p < 0.001$) and RTT (OR 0.67, 95%CI 0.47 – 0.95, $p = 0.03$).

Conclusion

This is the largest and most detailed study concerning short term outcomes following CRC resection in Victorian public hospitals.

Inner and outer regional centres had similar or better short-term outcomes than metropolitan hospitals after CRC resection.

AD with validated algorithms serves as a large accurate database to report on CRC outcomes.

Keywords: Colorectal neoplasm, surgery, administrative data, short-term outcomes

Introduction

Background

Of the studies that examine state-wide outcomes for colorectal cancer (CRC) resection in Australia¹⁻¹⁷ relatively few provide detailed data. This study uses validated algorithms of administrative data (AD) coding to access whole-of-state data and is one of the largest, detailed studies on CRC surgical outcomes in Victoria.

Centralisation of health care for CRC surgery has been advocated due to operative challenges and requirements for specialised supports¹⁸⁻²⁰. It might be expected that metropolitan hospitals with higher case and complexity volume could have better outcomes than their regional counterparts, but this has not been proven in Victoria. To study this issue requires detailed data with broad coverage to adjust for multiple confounding variables.

No multi-institutional database is perfect. Clinical databases with prospectively-gathered, detailed clinical data are the most accurate of the surgical databases; the Bi-National Colorectal Cancer Audit (BCCA) and BIOGRID are the largest colorectal surgery databases in Australia and New Zealand. However, BCCA only captures 15-20% of CRC resections, with mostly metropolitan hospital data²¹. BIOGRID only has contributions from 3 public sites in Victoria¹³.

Hospital AD, whilst not as accurate as clinical data, has access to all recorded inpatient AD for the state²². AD are collected on all patients admitted to Australian Hospitals. Patient demographics, co-morbidities, type of operation, post-operative complications, histopathology and some staging information are recorded. Trained coders review clinical notes and then assign alphanumeric codes to these data based on the International Classification of Diseases Tenth Revision, Australian

Modification (ICD-10-AM). These codes, developed for the purpose of billing, are not focused on reporting data in a clinically relevant fashion. We have previously shown that the use of algorithms of code combinations can increase the accuracy of AD for clinical research²²⁻²⁴. These algorithms can be applied to the central repository of AD in Victoria, the Victorian Admitted Episode Dataset (VAED), to create a validated alternative database with broad coverage that is more clinically relevant than unmodified AD.

Objectives

This study's aim was to use validated coding algorithms applied to the VAED to report on short-term outcomes following CRC surgery comparing regional to metropolitan hospitals.

The primary outcome was prolonged length of stay (LOS).

Secondary outcomes were: inpatient mortality, return to theatre (RTT), discharge destination.

A further important objective was to highlight the potential of AD to provide adequately detailed information for clinical outcomes research in CRC at a population level.

Methods

Study Design

This is a retrospective cohort study using prospectively-gathered AD from VAED.

This study was performed in accordance with the STROBE statement²⁵.

Ethics approval for this study was granted by the Human Research and Ethics Committee, National Health and Medical Research Council of Australia (reference number HREC/52415/MH-2019).

Participants

The algorithms created from a previous study were applied to the VAED to facilitate case finding and improve accuracy²³. These algorithms were based on comparing clinical notes with AD and using this information to construct combinations of administrative codes to improve the clinical relevance of AD. Right colon was defined as extending to and including the splenic flexure, left colon comprised the descending colon and sigmoid colon and rectum included recto-sigmoid and rectum. The algorithms combined the code for 'tumour location' with the code for 'type of operation' to eradicate illogical combinations thereby increasing accuracy²⁶. (An example of an illogical combination is a tumour location code in the ascending colon but with an operative code of a high anterior resection). When compared to clinical notes, the accuracy of validated algorithms of AD was 93%, 89% and 88% for right, left and rectum respectively. To create our study population, these algorithms were applied to the VAED to gather all patients undergoing resection for CRC in Victoria from 2008 to 2018. Data from VAED is de-identified for patient and hospital name. However, data can be requested by categories such as hospital location, case-volume etc. Patients less than 10 years old were excluded.

Variables

Exposure and Control

The principal exposure of interest was geographic location of hospital. Patients having major resection for CRC (adenocarcinoma) in regional public hospitals were compared with similar patients from metropolitan public hospitals (control) in the state of Victoria, Australia.

To define the geographical location of hospitals, the Accessibility/Remoteness Index of Australia (ARIA+) was used. ARIA+ is a nationally recognised index for defining geographical remoteness based

on accessibility to services. It divides Australia into five categories (S1)²⁶. All Victorian hospitals reside in major cities, inner regional and outer regional zones.

The control group was metropolitan hospitals and was compared separately to inner regional and outer regional hospitals.

Potential Confounders

Patient demographics, comorbidities ((Charlson Comorbidity Index score²⁷ (CCI) and American Society of Anaesthesiologists (ASA) score)), mode of surgery (open/laparoscopic approach) and pathological details were recorded. Although AD does not record TNM stage, it does report on the presence of lymph node metastases (92% accuracy) and distant metastases (88% accuracy)²³.

Hospital case volume (high volume ≥ 50 cases per year, low volume < 50 cases per year) and Colorectal Surgical Society of Australia and New Zealand (CSSANZ) status were recorded. Victorian CSSANZ hospitals have been previously defined²⁶. Although Peter MacCallum Cancer Centre is not a CSSANZ hospital, it was included in this group because of its case load, surgeon experience and research.

ASA was defined categorically: low for ASA 1-2, high for ASA 3-5. To minimise data restriction in the final multivariable analysis, a further category was assigned 'grade not recorded' if an anaesthetic code had been used without an ASA grade

CCI score was recorded as 0, 1 or 2, with ≥ 2 conferring higher mortality. The diagnostic categories used to calculate CCI were those validated for use in the ICD-10-AM²⁷. This categorisation has been used previously^{5,15,27,28}.

Surgical mode of access was defined as laparoscopic or open. Laparoscopic was defined by the use of a unique code for a laparoscopic CRC operation or an open code combined with the code for 'diagnostic laparoscopy' or 'laparoscopic division of adhesions.' The overall accuracy of coding for laparoscopic operations was 76% in our institution.

The composite variable HDU+ was a composite variable signifying patients that required more than simple ward care. It represents all patients requiring any period of postoperative mechanical ventilation and/or a monitored bed including ICU, CCU or HDU. HDU+ was also dichotomised as requiring these measures or not.

Outcomes

The primary outcome of interest was prolonged LOS (>14 days). LOS was defined as the discharge date minus the admission date. This outcome was chosen because it was likely to capture most significant types of post-operative morbidity. Secondary outcomes were inpatient mortality, RTT, discharge destination.

Inpatient mortality was defined as death during the index admission.

RTT was defined as an unplanned operation following the initial CRC resection. The algorithm used to extract this information was described previously²³.

Discharge destination was defined as home or other.

Statistical Analysis

Data was analysed using Stata version 15.0 (College Station, Texas, USA). Baseline data were presented as absolute numbers with percentages (Table1).

A univariable logistic regression analysis assessing the odds of the primary outcome (prolonged LOS) was performed for each baseline variable. Multivariable logistic regression analysis was then performed retaining geographical location as the principal exposure of interest. Co-variables from the univariable analysis which were associated with $\geq 10\%$ change in odds ratio were included in the multivariable logistic regression analysis. The co-variables, CSSANZ-status and Case-volume were not included in the multivariable analysis because CSSANZ hospitals were all metropolitan and there were no high-volume hospitals in the outer regional zone.

A similar analysis was performed for the secondary outcomes.

The sample size was not calculated for this study as the database was a fixed available sample.

A priori, it was determined that rectal cancer, high ASA score and modality of surgery (laparoscopic versus open) were potential effect modifiers. Testing for effect modification was performed on the primary outcome by using likelihood ratio testing to compare the original regression model with a further regression model including the variable of concern as an indicator variable. Effect modification was not tested for the secondary outcomes to minimise the likelihood of type 2 error.

Patients with missing data were identified and excluded from the analysis.

Results

Data extracted from the VAED included 75,460 patients totalling 1,623,753 admissions. Of the 75,460 patients, 60,819 had major colorectal resections (not necessarily for cancer). After applying the algorithms described above to improve the accuracy of the database, 33,275 patients were found who had major colorectal resections for CRC stratified by tumour position. Patients aged less than 10 years were also excluded (5 patients) as were patients operated on in private hospitals

leaving 18479. Finally, 9 duplicates were excluded leaving 18470 patients for analysis: 13511 metropolitan patients, 4202 inner regional patients and 766 outer regional patients (S2). Baseline variables were similar across the cohorts (Table 1). This dataset had remarkably little missing data; only 1% of ASA scores and 2% of discharge destination had missing data. The distribution of missing data was similar across cohorts.

Length of Stay

LOS was prolonged in 23% of metropolitan hospitals, 18% inner regional hospitals and 23% outer regional hospitals. The median LOS was: 8 days (interquartile range (IQR) 6 – 14 days) in metropolitan hospitals, 8 days (IQR 6 – 13 days) in inner regional hospitals and 9 days (IQR 7 – 14 days) in outer regional hospitals.

Univariable logistic regression analysis showed that inner regional hospitals had 26% lower odds of prolonged length of stay than metropolitan hospitals (OR 0.74, 95% CI 0.68 – 0.81, $p < 0.001$). There was no evidence of a difference in incidence of prolonged length of stay in outer regional hospitals compared with metropolitan hospitals. (OR 0.98, 95% CI 0.83 – 1.17, $p = 0.86$). Multivariable regression, adjusted for the effects of age, gender, year, ASA score, CCI score, position of tumour, mode of access, admission type and distant metastasis, decreased the odds of prolonged LOS in inner regional hospitals further to 47% (OR 0.53, 95% CI 0.48 – 0.58, $p < 0.001$). After adjusting for the same variables outer regional hospitals had 36% lower odds of prolong LOS which was now significant (OR 0.64, 95%CI 0.52 – 0.77. $p < 0.001$ (Table 2).

No evidence of effect modification was found. This suggests that the subgroups of patients having high ASA score, rectal surgery or laparoscopic surgery in regional centres did not have different odds of prolonged LOS compared with metropolitan centres.

Inpatient Mortality

Inpatient mortality was 1.9% in metropolitan hospitals, 2.1% in inner regional hospitals and 2.2% in outer regional hospitals.

Univariable logistic regression analysis showed that there was no difference in the odds of inpatient mortality in inner regional or outer regional hospitals when compared to metropolitan hospitals ((OR 1.06, 95% CI 0.83 – 1.36, p=0.62) and (OR 1.14, 95% CI 0.69 – 1.87, p=0.61) respectively).

Multivariable regression, adjusted for the effects of age, year, gender, ASA status, CCI, position of tumour, mode of access, admission type, lymph node metastases, distant metastasis and HDU+ showed that the odds of inpatient mortality were 30% less in inner regional hospitals compared with metropolitan hospitals and this was statistically significant (OR 0.67, 95%CI 0.49 – 0.91, p=0.01).

Adjusting for the same variables resulted in outer regional patients having no difference in odds of inpatient mortality when compared to metropolitan hospitals (OR 0.64, 95%CI 0.38 – 1.07, p=0.08).

Return to theatre

Only 5.6% of metropolitan patients had a RTT compared to 6.2% of inner regional patients and 4.7% of outer regional patients.

Univariable logistic regression analysis showed no difference in odds of RTT for inner regional or outer regional hospitals when compared to metropolitan hospitals ((OR 1.11, 95% CI 0.96 – 1.29, p=0.15) and (OR 0.83, 95% CI 0.59 – 1.17, p=0.28) respectively).

Multivariable regression, adjusted for the effects of gender, ASA, CCI, position of tumour, type of admission, distant metastases and HDU+ showed that the odds of RTT were no different in inner regional hospitals compared to metropolitan hospitals (OR 0.94, 95% CI 0.81 – 1.09, p=0.42).

However, adjusting for the same variables resulted in outer regional hospitals having 33% lower odds of RTT, which was statistically significant (OR 0.67, 95%CI 0.47 – 0.95, $p=0.03$).

Discharge Destination

13% of metropolitan patients were not discharged home compared to 9% of inner regional patients and 20% of outer regional patients.

Univariable logistic regression analysis showed that, compared to metropolitan patients, inner regional patients had 55% higher odds of not being discharged home (OR 1.55, 95% CI 1.41 – 1.70, $p<0.001$). Outer regional patients had 65% higher odds of not being discharged home (OR 1.65, 95% CI 1.37 – 1.99, $p<0.001$).

After adjusting for the effects of age, gender, year, ASA score, CCI score, position of tumour, mode of access, admission type, distant metastasis and HDU+ there was no difference in discharge destination between inner regional or outer regional hospitals and metropolitan hospitals ((OR 1.11, 95%CI 1.00 – 1.23, $p=0.051$) and (OR 1.08, 95%CI 0.88 – 1.33, $p=0.47$) respectively).

CSSANZ status and hospital volume were not included in the multivariable analysis as they were not applicable across all cohorts.

Compared with non-CSSANZ hospitals, CSSANZ hospitals had higher odds of prolonged LOS (OR 1.18, 95%CI 1.10 – 1.26, $p<0.001$), decreased odds of inpatient mortality (OR 0.74, 95%CI 0.60 – 0.91, $p=0.004$), no difference in odds of RTT (OR 0.97, 95%CI 0.86 – 1.10, $p=0.67$) and decreased odds of a discharge destination other than home (OR 0.63, 95%CI 0.58 – 0.69, $p<0.001$).

Compared with low volume hospitals, high volume hospitals had higher odds of prolonged LOS (OR 1.22, 95%CI 1.12 – 1.33, $p<0.001$), no difference in odds of inpatient mortality (OR 1.05, 95%CI 0.82

– 1.35, $p=0.70$), or odds of RTT (OR 1.12, 95%CI 0.96 – 1.30, $p=0.15$) but decreased odds of discharge destination other than home (OR 0.60, 95%CI 0.58 – 0.66, $p<0.001$).

Discussion

This is the largest and most detailed study concerning short term outcomes of CRC resection involving all Victorian public hospitals. Rather than present the results simply as an audit, we have chosen to subject the data to hypothesis testing to maximise their utility. With 14 exposure variables and 4 outcome variables there were many ways in which this study could have been configured. We felt that an important initial issue was to determine if there were major differences in short-term outcomes by geographical location of hospital. This would determine if future studies utilising this dataset could use the whole state's data unmodified. A study of long-term survival analysis will follow.

Overall this study shows that short-term outcomes of CRC resection in regional hospitals are comparable to and sometimes better than metropolitan hospitals. The odds of all adverse outcomes tend to be lower in regional hospitals. Strong evidence was found for lower odds of prolonged LOS and inpatient mortality in inner regional hospital compared with metropolitan hospitals. For outer regional hospitals, there was strong evidence for a decrease in the odds of prolonged length of stay and RTT. CSSANZ training centres performed worse than other hospitals in terms of LOS but better for inpatient mortality and discharge to home. High volume hospitals compared with low volume hospitals performed worse regarding LOS but better for discharge to home. Overall other co-variables had expected results with patients who were older, more comorbid, emergency cases and those requiring high dependency having worse outcomes.

The methodology of this study does not account for allocation bias in determining which patients chose to have surgery at their regional hospital, patterns of referral to metropolitan hospitals and case selection by regional surgeons. However, we have used the technique of multivariable regression analysis to adjust for co-variables which may confound outcomes. The results of this study support the current practices in surgical treatment of CRC across the state of Victoria. This study used AD rather than the clinical data to report on surgical colorectal cancer outcomes. AD has been used in the past in Australia in this setting^{1,6-10,14,29,30} however these studies have used unmodified ICD-10-AM codes. Unmodified coding data does not always match well with data from clinical notes. We have used combinations of codes which we knew to be highly accurate having compared them to clinical notes²³. With these algorithms, AD approaches the accuracy of clinical databases but with the benefit of much higher numbers and state-wide generalisability^{11-13,31,32}.

AD cannot record all complications. The most notable example in AD is the absence of a code for anastomotic leak. However, many leaks would have been captured indirectly in our primary and secondary outcomes. It should also be noted that AD does not report on outpatient neoadjuvant treatments.

We excluded private hospital data from this analysis because all our previous validation of AD was performed using public hospital data.

The results of this study are similar to other Australian studies examining LOS but using unmodified AD. A study performed in New South Wales, Australia found comparable incidence of prolonged LOS and inpatient mortality when comparing rural to metropolitan patients¹⁶. They also found that high CCI, age and emergency admission were associated with prolonged LOS. A South Australian study

found increased odds of 30-day mortality in non-metropolitan patients⁵ whereas a Queensland study did not¹⁷.

Of the international studies exploring the effects of geographical location on LOS, one of the largest studies was performed in England³³. This showed increased odds of prolonged LOS in the Central and Southern parts of England when compared to the North. An Irish study showed no difference in prolonged LOS after colorectal resection when case volume and socioeconomic status were used as markers of access to healthcare³⁴.

Conclusion

After adjustment for confounding variables, Inner and outer regional hospitals have similar or better short-term outcomes after resection for colorectal cancer than metropolitan hospitals in the state of Victoria.

AD with validated algorithms, utilising pre-existing data and requiring minimal additional resources, has demonstrated fitness for purpose as a large and detailed database to report on CRC outcomes.

Acknowledgements

We would like to thank the Centre of Victoria Data Linkage (CVDL) for data curation and data linkage.

References

1. Faragher IG, Hong MK, Stupart D, Watters DA, Yeung J. Complete state-wide outcomes in elective colon cancer surgery. *ANZ journal of surgery*. 2018;88(11):1174-1177.
2. Yu XQ, O'Connell D L, Gibberd RW, Armstrong BK. A population-based study from New South Wales, Australia 1996-2001: area variation in survival from colorectal cancer. *European journal of cancer (Oxford, England : 1990)*. 2005;41(17):2715-2721.
3. Coleman MP, Quaresma M, Berrino F, et al. Cancer survival in five continents: a worldwide population-based study (CONCORD). *The Lancet Oncology*. 2008;9(8):730-756.
4. Marwan K, Staples MP, Thursfield V, Bell SW. The rate of abdominoperineal resections for rectal cancer in the state of Victoria, Australia: a population-based study. *Diseases of the colon and rectum*. 2010;53(12):1645-1651.
5. Beckmann K, Moore J, Wattchow D, Young G, Roder D. Short-term outcomes after surgical resection for colorectal cancer in South Australia. *Journal of evaluation in clinical practice*. 2017;23(2):316-324.
6. Semmens JB, Platell C, Threlfall TJ, Holman CD. A population-based study of the incidence, mortality and outcomes in patients following surgery for colorectal cancer in Western Australia. *The Australian and New Zealand journal of surgery*. 2000;70(1):11-18.
7. Hall SE, Holman CD, Platell C, Sheiner H, Threlfall T, Semmens J. Colorectal cancer surgical care and survival: do private health insurance, socioeconomic and locational status make a difference? *ANZ journal of surgery*. 2005;75(11):929-935.
8. Beckmann KR, Bennett A, Young GP, Roder DM. Treatment patterns among colorectal cancer patients in South Australia: a demonstration of the utility of population-based data linkage. *Journal of evaluation in clinical practice*. 2014;20(4):467-477.
9. Dobbins TA, Young JM, Solomon MJ. Uptake and outcomes of laparoscopically assisted resection for colon and rectal cancer in Australia: a population-based study. *Diseases of the colon and rectum*. 2014;57(4):415-422.
10. Dobbins TA, Badgery-Parker T, Currow DC, Young JM. Assessing measures of comorbidity and functional status for risk adjustment to compare hospital performance for colorectal cancer surgery: a retrospective data-linkage study. *BMC medical informatics and decision making*. 2015;15:55.
11. Teloken PE, Spilsbury K, Platell C. Analysis of mortality in colorectal surgery in the Bi-National Colorectal Cancer Audit. *ANZ journal of surgery*. 2016;86(6):454-458.
12. Warriar SK, Kong JC, Guerra GR, et al. Risk Factors Associated With Circumferential Resection Margin Positivity in Rectal Cancer: A Binational Registry Study. *Diseases of the colon and rectum*. 2018;61(4):433-440.
13. Sammour T, Hayes IP, Jones IT, Steel MC, Faragher I, Gibbs P. Impact of anastomotic leak on recurrence and survival after colorectal cancer surgery: a BioGrid Australia analysis. *ANZ journal of surgery*. 2018;88(1-2):E6-e10.
14. Luke CG, Koczwara B, Moore JE, et al. Treatment and survival from colorectal cancer: the experience of patients at South Australian teaching hospitals between 1980 and 2002. *Clinical oncology (Royal College of Radiologists (Great Britain))*. 2005;17(5):372-381.

15. Jorgensen ML, Young JM, Dobbins TA, Solomon MJ. Assessment of abdominoperineal resection rate as a surrogate marker of hospital quality in rectal cancer surgery. *The British journal of surgery*. 2013;100(12):1655-1663.
16. Jorgensen ML, Young JM, Dobbins TA, Solomon MJ. Predictors of variation in colorectal cancer care and outcomes in New South Wales: a population-based health data linkage study. *The Medical journal of Australia*. 2014;200(7):403-407.
17. Theile DE, Philpot S, Blake M, Harrington J, Youl PH. Outcomes following colorectal cancer surgery: Results from a population-based study in Queensland, Australia, using quality indicators. *Journal of evaluation in clinical practice*. 2019;25(5):834-842.
18. Aquina CT, Probst CP, Becerra AZ, et al. High volume improves outcomes: The argument for centralization of rectal cancer surgery. *Surgery*. 2016;159(3):736-748.
19. Archampong D, Borowski D, Wille-Jorgensen P, Iversen LH. Workload and surgeon's specialty for outcome after colorectal cancer surgery. *The Cochrane database of systematic reviews*. 2012(3):Cd005391.
20. Khani MH, Smedh K. Centralization of rectal cancer surgery improves long-term survival. *Colorectal Dis*. 2010;12(9):874-879.
21. Lee CHA, Kong JCH, Heriot AG, Warriar S, Zalberg J, Sitzler P. Short-term outcome of emergency colorectal cancer surgery: results from Bi-National Colorectal Cancer Audit. *International Journal of Colorectal Disease*. 2019;34(1):63-69.
22. Hong MKY, Skandarajah AR, Hayes IP. Administrative data: what surgeons should know about big data. *ANZ J Surg*. 2017;87(9):650-651.
23. MacCallum C, Skandarajah A, Gibbs P, Hayes I. The Value of Clinical Colorectal Cancer Registries in Colorectal Cancer Research: A Systematic Review. *JAMA surgery*. 2018;153(9):841-849.
24. Hong MKY, Skandarajah AR, Higgins RD, Faiz OD, Hayes IP. International Variation in Emergency Operation Rates for Acute Diverticulitis: Insights into Healthcare Value. *World J Surg*. 2017;41(8):2121-2127.
25. von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet*. 2007;370(9596):1453-1457.
26. MacCallum C, Da Silva N, Skandarajah A, Hayes I. Study of colorectal cancer resection patterns across the state of Victoria using validated administrative data algorithms. *ANZ J Surg*. n/a(n/a).
27. Sundararajan V, Henderson T, Perry C, Muggivan A, Quan H, Ghali WA. New ICD-10 version of the Charlson comorbidity index predicted in-hospital mortality. *Journal of clinical epidemiology*. 2004;57(12):1288-1294.
28. Field K, Shapiro J, Wong HL, et al. Treatment and outcomes of metastatic colorectal cancer in Australia: defining differences between public and private practice. *Internal medicine journal*. 2015;45(3):267-274.
29. Weir K, Supramaniam R, Gibberd A, Dillon A, Armstrong BK, O'Connell DL. Comparing colorectal cancer treatment and survival for Aboriginal and non-Aboriginal people in New South Wales. *The Medical journal of Australia*. 2016;204(4):156.
30. Thompson BS, Coory MD, Lumley JW. National trends in the uptake of laparoscopic resection for colorectal cancer, 2000–2008. *Medical Journal of Australia*. 2011;194(9):443-447.

31. Sammour T, Jones IT, Gibbs P, et al. Comparing oncological outcomes of laparoscopic versus open surgery for colon cancer: Analysis of a large prospective clinical database. *Journal of surgical oncology*. 2015;111(7):891-898.
32. Lee CHA, Kong JCH, Heriot AG, Warriar S, Zalcborg J, Sitzler P. Short-term outcome of emergency colorectal cancer surgery: results from Bi-National Colorectal Cancer Audit. *International journal of colorectal disease*. 2019;34(1):63-69.
33. Pucciarelli S, Zorzi M, Gennaro N, et al. In-hospital mortality, 30-day readmission, and length of hospital stay after surgery for primary colorectal cancer: A national population-based study. *European journal of surgical oncology : the journal of the European Society of Surgical Oncology and the British Association of Surgical Oncology*. 2017;43(7):1312-1323.
34. Kelly M, Sharp L, Dwane F, Kelleher T, Comber H. Factors predicting hospital length-of-stay and readmission after colorectal resection: a population-based study of elective and emergency admissions. *BMC health services research*. 2012;12:77.

Supplementary Figure Legends

S1: ARIA+ remoteness classification.

S2: Study population selection flow diagram

S3: Algorithms to improve cohort accuracy

		Metropolitan n=13511 (73%)	Inner Regional n=4202(23%)	Outer Regional n=766(4%)
Age, deciles (years) (%total)	10-19	11 (0.1%)	1 (0.0%)	0 (0.0%)
	20s	102 (0.8%)	11 (0.3%)	2 (0.3%)
	30s	346 (2.6%)	51 (1.2%)	9 (1.2%)
	40s	908 (6.7%)	186 (4.5%)	25 (3.3%)
	50s	2095 (15.5%)	546 (13.0%)	93 (12.1%)
	60s	3429 (25.4%)	973 (23.2%)	182 (23.8%)
	70s	4118 (30.5%)	1448 (34.5%)	261 (34.1%)
	80s	2492 (18.4%)	975 (23.3%)	194 (25.3%)
	90+	10 (0.1%)	0 (0.0%)	0 (0.0%)
Sex M:F (Male%)		7581:5930 (56%)	2261:1932 (54%)	403:363 (53%)
ASA score	Grade not recorded	2390 (17.7%)	645 (15.4%)	172 (22.5%)
	1	700 (5.2%)	225 (5.4%)	47 (6.1%)
	2	4682 (34.7%)	1391 (33.2%)	244 (31.9%)
	3	4679 (34.6%)	1470 (35.1%)	260 (33.9%)
	4	896 (6.6%)	374 (8.9%)	35 (4.6%)
	5	36 (0.3%)	25 (0.6%)	2 (0.3%)
	6	0 (0.0%)	0 (0.0%)	1 (0.1%)
	Missing	128 (1.0%)	62 (1.5%)	5 (0.7%)
CCI score	0	12380 (91.6%)	3819 (91.1%)	702 (91.6%)
	1	1020 (7.6%)	337 (8.0%)	60 (7.8%)
	2	63 (0.5%)	27 (0.6%)	3 (0.4%)
	3	20 (0.2%)	8 (0.2%)	1 (0.1%)
	4	18 (0.1%)	1 (0.0%)	0 (0.0%)
	5	1 (0.0%)	1 (0.0%)	0 (0.0%)
	6	9 (0.1%)	0 (0.0%)	0 (0.0%)
Year*	2007	765	284	49
	2008	1,056	386	74
	2009	1,118	345	67
	2010	1,157	374	54
	2011	1,159	356	66
	2012	1,102	365	73
	2013	1,119	355	59
	2014	1,101	346	70
	2015	1,172	343	61
	2016	1,117	322	59
	2017	1,207	325	67
	2018	1,240	342	56
2019	198	50	11	

Admission type	Elective	11343 (84%)	3465 (83%)	590 (77%)
	Emergency	2168 (16%)	728 (17%)	176 (23%)
Side of operation	Right	5740 (42.5%)	2121 (50.6%)	411 (53.7%)
	Left	3209 (23.8%)	902 (21.5%)	184 (24.0%)
	Rectum	4562 (33.8%)	1170 (27.9%)	171 (22.3%)
Mode of access	Open	7784 (57.6%)	3160 (75.4%)	700 (91.4%)
	Laparoscopic	5727 (42.4%)	1033 (24.6%)	66 (8.6%)
Lymph node metastases	Present	4041 (29.9%)	1323 (31.6%)	206 (26.9%)
	Absent	9470 (70.1%)	2870 (68.5%)	560 (73.1%)
Distant metastases	Present	1887 (14.0%)	485 (11.6%)	97 (12.7%)
	Absent	11624 (86.0%)	3708 (88.4%)	669 (87.3%)

Table 1: Baseline characteristics

*2017 & 2019 are included in this table as patients have fallen within this timeframe due to patient dates being moved forward and backwards by fixed amounts by VAED to protect confidentiality.

		Univariable						Multivariable				
		No. total	prolonged LOS	OR	95% CI		p-value	n=	OR	95%CI		p-value
ARIA	Inner v Metro	4193 v 13511	754 v 3088	0.74	0.68	0.81	<0.001	18273	0.53	0.48	0.58	<0.001
	Outer v Metro	766 v 13511	173 v 3088	0.98	0.83	1.17	0.862	18273	0.64	0.52	0.77	<0.001
CSSANZ Hospital	Yes v No	9449 v 9021	2183 v 1832	1.18	1.10	1.26	<0.001					
Case Volume	≥50 cases/year vs <50 cases/year	14199 v 4271	3195 v 820	1.22	1.12	1.33	<0.001					
Age	>70 v <70	9498 v 8972	2440 v 1575	1.62	1.51	1.74	<0.001	18273	1.34	1.24	1.46	<0.001
Sex	Male v Female	10245 v 8225	2385 v 1630	1.23	1.14	1.32	<0.001	18273	1.16	1.08	1.26	<0.001
Year	2008-2013 v 2014 - 2018	10383 v 8087	2425 v 1590	1.25	1.16	1.34	<0.001	18273	1.13	1.05	1.23	0.002
ASA score	High vs Low	7777 v 7289	2292 v 861	3.12	2.86	3.40	<0.001	18273	2.07	1.88	2.28	<0.001
	NR vs Low	3207 v 7289	829 v 861	2.60	2.34	2.89	<0.001	18273	2.17	1.93	2.42	<0.001
CCI score	1 vs 0	1417 v 16901	310 v 3644	1.02	0.89	1.16	0.781	18273	0.95	0.83	1.10	0.522
	≥2 vs 0	152 v 16901	61 v 3644	2.44	1.76	3.38	<0.001	18273	1.64	1.14	2.35	0.007
Position	Left vs Right	4295 v 8272	877 v 1661	1.02	0.93	1.12	0.653	18273	1.02	0.92	1.12	0.754
	Rectum vs Right	5903 v 8272	1477 v 1661	1.33	1.23	1.44	<0.001	18273	1.63	1.48	1.79	<0.001
Mode of access	Laparoscopic v Open	6826 v 11644	859 v 3156	0.39	0.36	0.42	<0.001	18273	0.56	0.51	0.62	<0.001
Admission type	Emergency/Elective	3072 v 15398	1266 v 2749	3.23	2.97	3.50	<0.001	18273	2.73	2.48	3.00	<0.001
Lymph node metastases	Positive v Negative	5570 v 12900	1228 v 2787	1.03	0.95	1.11	0.504					
Distant metastases	Positive v Negative	2469 v 16001	753 v 3262	1.71	1.56	1.88	<0.001	18273	1.27	1.14	1.41	<0.001
HDU+	Yes v No	6661 v 11809	2389 v 1626	3.50	3.26	3.77	<0.001	18273	2.80	2.58	3.03	<0.001

Table 2: Univariable and multivariable logistic regression analysis – Prolonged length of stay



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Udayasiri, DK; MacCallum, C; Da Silva, N; Skandarajah, A; Hayes, IP

Title:

Impact of hospital geographic remoteness on short-term outcomes after colorectal cancer resection using state-wide administrative data

Date:

2020-05-26

Citation:

Udayasiri, D. K., MacCallum, C., Da Silva, N., Skandarajah, A. & Hayes, I. P. (2020). Impact of hospital geographic remoteness on short-term outcomes after colorectal cancer resection using state-wide administrative data. ANZ JOURNAL OF SURGERY, 90 (7-8), pp.1328-1334. <https://doi.org/10.1111/ans.15992>.

Persistent Link:

<http://hdl.handle.net/11343/275785>

File Description:

Accepted version