

Article type: Original Article

Title: Diversity and stability of coral endolithic microbial communities at a naturally high $p\text{CO}_2$ reef.

Vanessa R. Marcelino¹, Kathleen M. Morrow^{2,3}, Madeleine J.H. van Oppen^{1,3}, David G. Bourne^{3,4} and Heroen Verbruggen¹

¹ School of Biosciences, University of Melbourne, Victoria 3010, Australia

² Department of Molecular, Cellular and Biomedical Sciences, University of New Hampshire, Durham, New Hampshire 03824, United States

³ Australian Institute of Marine Science, Townsville MC, Queensland 4810, Australia

⁴ College of Science and Engineering, James Cook University, Townsville, Queensland 4811, Australia

Keywords: Microbiome, limestone-boring, ocean acidification, *Ostreobium*, coral skeletons, multi-marker metabarcoding.

Author for correspondence: Vanessa Rossetto Marcelino

School of Biosciences (bldg. 122, room 101), University of Melbourne, VIC 3010, Australia

Telephone: (+61) 03 83448931. Email: vrmarcelino@gmail.com

Running title: Endolithic microbes and ocean acidification

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as [doi: 10.1111/mec.14268](https://doi.org/10.1111/mec.14268)

This article is protected by copyright. All rights reserved

Abstract:

The health and functioning of reef-building corals is dependent on a balanced association with prokaryotic and eukaryotic microbes. The coral skeleton harbours numerous endolithic microbes, but their diversity, ecological roles and responses to environmental stress, including ocean acidification, are not well characterized. This study tests whether pH affects the diversity and structure of prokaryotic and eukaryotic algal communities associated with skeletons of *Porites* spp. using targeted amplicon (16S rRNA gene, UPA and *tufA*) sequencing. We found that the composition of endolithic communities in the massive coral *Porites* spp. inhabiting a naturally high $p\text{CO}_2$ reef (avg. $p\text{CO}_2$ 811 μatm) is not significantly different from corals inhabiting reference sites (avg. $p\text{CO}_2$ 357 μatm), suggesting that these microbiomes are less disturbed by ocean acidification than previously thought. Possible explanations may be that the endolithic microhabitat is highly homeostatic, or that the endolithic microorganisms are well adapted to a wide pH range. Some of the microbial taxa identified include nitrogen-fixing bacteria (*Rhizobiales* and cyanobacteria), algicidal bacteria in the phylum *Bacteroidetes*, symbiotic bacteria in the family *Endozoicomonaceae*, and endolithic green algae, considered the major microbial agent of reef bioerosion. Additionally, we test whether host species has an effect on the endolithic community structure. We show that the endolithic community of massive *Porites* spp. is substantially different and more diverse than that found in skeletons of the branching species *Seriatopora hystrix* and *Pocillopora damicornis*. This study reveals highly diverse and structured microbial communities in *Porites* spp. skeletons that are possibly resilient to ocean acidification.

Introduction

Ocean acidification (OA) is predicted to threaten the persistence of coral reefs by affecting the balance between constructive forces (calcification and growth of reef builders) and destructive forces (bioerosion and carbonate dissolution) (Tribollet 2008a; Andersson & Gledhill 2013). Acidification lowers the saturation state of calcium carbonate (CaCO_3), making it more difficult for calcifying organisms, such as stony corals, to build their skeletons (Orr *et al.* 2005; Hofmann *et al.* 2010). OA has been shown to slow down calcification and cause structural deformities in juvenile corals (Crook *et al.* 2013; Foster *et al.* 2016). Some studies, however, indicate that corals are able to regulate the pH at the tissue-skeleton interface, where calcification takes place, mitigating the potential consequences of OA on the calcification process (McCulloch *et al.* 2012; Venn *et al.* 2013; Georgiou *et al.* 2015). Rates of biological dissolution of CaCO_3 (bioerosion) tend to increase under low pH conditions, mostly due to an increase in biomass of the boring organisms living inside coral skeletons (Manzello *et al.* 2008; Tribollet *et al.* 2009; Crook *et al.* 2013; Fang *et al.* 2013; Reyes-Nivia *et al.* 2013; Enochs *et al.* 2016),

potentially resulting in a shift from a net reef accretion condition to one of net erosion (Andersson & Gledhill 2013).

The skeletons of live and dead corals harbour bacteria, fungi, sponges and an abundant population of limestone-boring algae, all having important roles in the reef's CaCO_3 budget (Le Campion-Alsumard *et al.* 1995; Tribollet 2008a; Verbruggen & Tribollet 2011). For example, the green alga *Ostreobium* can be responsible for 70–90% of carbonate dissolution within dead corals, eroding as much as 1 kg of reef carbonate per m^2 per year (Tribollet 2008b; Grange *et al.* 2015). Green algal biomass in live coral skeletons exceeds *Symbiodinium* biomass in coral tissues by about 16 times (Odum & Odum 1955), making the limestone attractive to grazers and further increasing bioerosion (Chazottes *et al.* 1995; Clements *et al.* 2016). However, endolithic algae also protect corals from high light stress (Yamazaki *et al.* 2008) and provide vital nutrients to corals, potentially extending the time they can survive without *Symbiodinium* during bleaching events (Schlichter *et al.* 1995; Fine & Loya 2002). Endolithic algae have exceptionally high levels of cryptic diversity (Marcelino & Verbruggen 2016; Sauvage *et al.* 2016; Del Campo *et al.* 2017), and although it is known that their biomass increases substantially upon acidification and warming (Tribollet *et al.* 2009; Reyes-Nivia *et al.* 2013), it is not known which of the cryptic species increase in relative abundance.

The endolithic community, along with the coral host and its other symbionts, constitutes the coral holobiont (Rohwer *et al.* 2002). The responses of the coral microbiome, including both prokaryotic and eukaryotic members, to acidification has gained attention as we continue to uncover vital roles played by microorganisms in holobiont health and resilience (Bourne *et al.* 2009; Sharp & Ritchie 2012; Krediet *et al.* 2013; Blackall *et al.* 2015; Bourne *et al.* 2016). Because the ocean pH naturally changes throughout seasons, along depth gradients, with productivity and other biological factors, marine microorganisms may have the physiological plasticity required to cope with the predicted levels of ocean acidification over the next 100 years (Joint *et al.* 2011). This notion is supported by several studies showing stable coral prokaryotic community when shifted from ambient to high CO_2 partial pressure ($p\text{CO}_2$) and therefore reduced seawater pH conditions (Meron *et al.* 2012; Webster *et al.* 2016; Zhou *et al.* 2016). However other studies have demonstrated that a reduced seawater pH can lead to the loss of *Symbiodinium* (coral bleaching) and trigger shifts from a healthy microbiome composition to a microbial community typically associated with diseased corals (Anthony *et al.* 2008; Vega Thurber *et al.* 2009; Meron *et al.* 2011; Webster *et al.* 2013; Morrow *et al.* 2015). These different responses of a coral's microbiome to reduced seawater pH may reflect differences in resilience across coral species to acidification or different experimental setups used in the various studies.

Reefs at the Milne Bay Province of Papua New Guinea (PNG) are in close proximity to volcanic seeps (expelling ~ 99% pure CO_2) and constitute a good model system to study the impacts of acidification *in situ* on the microbial community associated with corals. Both coral species composition and the prokaryotic microbial community associated with coral tissue and mucus differ between high $p\text{CO}_2$ seep sites and nearby reference sites with ambient $p\text{CO}_2$ (Fabricius *et al.* 2011; Morrow *et al.* 2015). However, little is known about the coral endolithic communities and how these may change

under various seawater pH conditions. Previous studies have screened 16S rRNA gene clone libraries and demonstrated contrasting results, with significant effects of OA community composition within the skeleton in an experimental system (Meron *et al.* 2011) but no significant changes in corals transplanted to a natural CO₂ seep site (Meron *et al.* 2012). One limitation with the 16S rRNA gene marker is that it underestimates the diversity of eukaryotic algae (Marcelino & Verbruggen 2016), and as a consequence, the major microbial agents of bioerosion have been overlooked.

Here we use high-throughput amplicon sequencing to investigate the effects of ocean acidification on the diversity and structure of endolithic microbial communities of corals inhabiting a high pCO₂ site in PNG. Our goals are to: 1) test whether the community composition of prokaryotes and photosynthetic eukaryotes (assessed with the 16S rRNA gene, UPA and *tufA* markers) within the skeletons of massive colonies of *Porites* spp. differs between high pCO₂ sites and nearby reference sites where pCO₂ is not affected by the volcanic seeps; 2) compare the endolithic communities of *Porites* spp. with those of two branching coral species (*Seriatopora hystrix* and *Pocillopora damicornis*) to investigate whether the microbiome in coral skeletons varies among host species; and 3) describe the endolithic community diversity found in corals of Papua New Guinea and discuss the potential functional roles of this microbiome under ocean acidification.

Methods

Field sites and sampling

Samples of massive *Porites* spp. (n = 24, six per site and month) were collected in April and November 2014 at two high pCO₂ (seep) and reference sites within the D'Entrecasteaux Islands, Milne Bay Province, Papua New Guinea. High pCO₂ samples were collected at Illi Illi (Upa-U) Seep (09.82425S, 150.81789E) and Dobu Seep (09.73646S, 150.86894E), and at nearby reference sites (ambient pCO₂) not exposed to elevated pCO₂ conditions (Illi Illi Reference, 09.82806S, 150.82028E and Dobu Reference, 09.75211S, 150.85410E) (Fabricius *et al.* 2011; Uthicke *et al.* 2013). High pCO₂ and reference sites were ~500 m and ~3 km apart from one another at Illi Illi and Dobu respectively. Samples of the branching corals *Seriatopora hystrix* (n = 3 at each site) and *Pocillopora damicornis* (n = 3 at each site) were only collected in April 2014 at Illi Illi seep and reference sites (same as above). Seawater carbonate chemistry varies in response to bubble activity and water motion at the seep sites; thus, at Illi Illi seep, corals experience a pH range (defined here as the 5th and 95th percentiles) of 7.28–8.01 (avg. pCO₂ 624 µatm) and at Dobu seep a pH range of 7.08–7.99 (avg. pCO₂ 998 µatm). At Illi Illi reference site the pH ranges from 7.91–8.09 (avg. pCO₂ 346 µatm) and at Dobu reference site the pH ranges from 7.91–8.10 (avg. pCO₂ 368 µatm) (Fabricius *et al.* 2014), which is within the range of future predictions for the year 2100 (Moss *et al.* 2010).

Coral fragments were collected using bone cutters or a hammer and chisel and placed into individual sections within a plastic tackle box, which allowed for water flow whilst underwater. After

returning to the boat, samples were immediately placed into flowing seawater sourced directly from the collection site. Large pieces of *Porites* spp. were chipped into smaller fragments, rinsed thoroughly with sterile 0.02 µm-filtered seawater and placed in 50 mL Falcon tubes with RNAlater (Ambion). Samples were kept in a cooler with ice until returned to the laboratories at the Australian Institute of Marine Science (AIMS) where they were processed.

Fragments were removed from RNAlater and soaked in 0.2 µm filtered calcium and magnesium free seawater for ~10 minutes at room temperature (CMFSW; 0.45M NaCl, 10mM KCl, 7mM Na₂SO₄, 0.5mM NaHCO₃ and milli-Q water; (Esteves *et al.* 2016)) to aid in the initial removal of tissues from the skeleton. Tissues were removed into the CMFSW using an air gun fitted with a sterile tip. Skeletons with tissues removed were placed back into the original RNAlater collection buffer and stored at -80°C until shipment to the University of Melbourne where DNA was isolated from the endolithic community (see below).

DNA isolation, library preparation and sequencing

Total DNA was isolated from coral skeletons using the Wizard Genomic DNA Purification Kit (Promega). The manufacturer's protocol for plant DNA was followed, with the exception of an extended 3 hr incubation step with the first extraction buffer to allow the DNA to leak out from the limestone into the solution. Amplified DNA products for library preparation were obtained with a two-step process described by Marcelino and Verbruggen (2016). During the first PCR step, three metabarcoding markers were amplified: the 16S rRNA gene (Klindworth *et al.* 2013); the universal plastid amplicon (UPA), which is a fragment of the 23S rRNA gene (Presting 2006; Sherwood & Presting 2007) and the elongation factor Tu (*tufA*), which targets green algae (Ulvophyceae) (Fama *et al.* 2002; Marcelino & Verbruggen 2016). During the second PCR step, barcodes and Illumina adapters were attached to both 3' and 5' ends of the amplicons. One negative control was performed with each amplification (6 in total, one per marker and per amplification step) and sequenced with the library, even though no DNA was detected in any negative control during quantification. Two mock 'blank' extractions were also performed along with the samples DNA isolation, processed through the amplification process and sequenced to further control for possible contamination. DNA isolation and PCR preparation were carried out inside a dedicated dead-air box (PCR workstation) sterilized with UV light for 15 min prior to each use. Libraries were quantified using the Quant-It PicoGreen reagent (Invitrogen) and pooled with other samples of another project. The library was sequenced using the Illumina MiSeq platform (2×300 bp paired end reads) at the Centre for Translational Pathology, University of Melbourne. Further details about the primers and library preparation are provided in Supplementary materials.

Data processing

The MiSeq run yielded one file containing all amplicons per sample, which were demultiplexed based on the primer sequences. The 3' ends of reads were trimmed to improve

consensus quality; forward and reverse reads were merged using FLASH (Magoc & Salzberg 2011) and sequences having average quality scores smaller than 35 or lengths shorter than a threshold (350 bp for 16S rRNA gene, 320 bp for UPA and 400 bp for the *tufA*) were filtered out using PRINSEQ (Schmieder & Edwards 2011). To verify that the data lost during quality control did not affect the results, we also performed the analyses with a less stringent quality filtering (Supplementary materials). Sequences were clustered into Operational Taxonomic Units (OTUs) using UPARSE (Edgar 2013). A similarity threshold of 98% was set for the *tufA* marker, a threshold near species level for this marker (Sauvage *et al.* 2016). For the other markers, the default threshold of 97% was used. The 16S rRNA gene OTUs were aligned with PyNAST (Caporaso *et al.* 2010a) while the UPA and *tufA* datasets were aligned with MAFFT (Katoh *et al.* 2002). A taxonomy affinity was assigned to the OTUs using the Naïve Bayesian Classifier (RDP) implemented in QIIME v.1.9.1 (Wang *et al.* 2007; Caporaso *et al.* 2010b). The Greengenes v.13.8 dataset (DeSantis *et al.* 2006) was used to classify the 16S rRNA gene sequences, and custom-made reference datasets (described and available in Marcelino & Verbruggen 2017) were used for *tufA* and UPA. The resulting OTU table went through a filtering process to remove OTUs found in the negative controls (3 OTUs in the 16S rRNA gene, 5 OTUs in the UPA and 2 OTUs in the *tufA* datasets) and rare OTUs (i.e. OTUs with less than 5 reads across all samples and OTUs from samples where they are present with 2 or less reads). OTUs were also filtered based on their taxonomic classification to focus on the taxonomic groups that each marker best characterizes: chloroplast sequences were excluded from the 16S rRNA gene dataset and bacterial sequences were excluded from the *tufA* dataset. Further details about the data processing pipeline are provided in the Supplementary materials.

Statistical analysis

There were no significant differences related to the time of collection (Supplementary materials), therefore all *Porites* spp. samples (n = 24) were used to investigate the effects of $p\text{CO}_2$ in endolithic communities associated with this coral genus. Rarefaction curves of the number of observed OTUs per number of reads were constructed by randomly subsampling the reads in QIIME, allowing to set a threshold for each marker where the curve reaches saturation (i.e. a plateau in the rarefaction curve), which was 2,200 reads in the 16S rRNA gene, 1,400 in the *tufA* and 7,000 in the UPA dataset (Supplementary figure 1). Samples with sequencing depths lower than these thresholds were excluded, resulting in 20 samples in the 16S rRNA gene, and 22 samples in the UPA and *tufA* datasets (Supplementary Table S1). Alpha diversity indices (Chao1 and observed OTUs) were calculated using QIIME (Caporaso *et al.* 2010b). The relative abundance of individual OTUs and taxonomic groups (i.e. OTUs grouped at phylum level for bacteria and genus level for algae) between sites were tested for significant differences with a Kruskal–Wallis test (for OTUs) and ANOVA (for taxon groups) using QIIME (Caporaso *et al.* 2010b).

To further investigate the distribution of green algal lineages, a maximum likelihood phylogeny was built with the green algal *tufA* OTUs together with reference sequences (from GenBank) using a GTR+gamma model of sequence evolution in RAxML v.8.2.6 (Stamatakis 2006).

OTUs present in less than 3 samples were excluded, their relative abundances were normalized with cumulative sum scaling (Paulson *et al.* 2013) and visualized alongside a phylogenetic tree using the R package phytools (Revell 2012).

Potential differences in community composition between high $p\text{CO}_2$ and reference sites (beta-diversity) were investigated with a combination of statistical methods. Principal coordinate analysis (PCoA) on weighted UniFrac distance matrices was performed using QIIME (Caporaso *et al.* 2010b; Lozupone *et al.* 2011) and the results visualized using the ggplot2 package in R (Wickham 2009). A multivariate generalized linear model (MGLM) was used to investigate potential differences in community composition between high $p\text{CO}_2$ and reference sites. The MGLM was computed using the mvabund R package (Wang *et al.* 2012), considering a negative binomial distribution. The null hypothesis of no difference among sites was statistically tested with analysis of deviance using 999 bootstrap interactions (R scripts provided in Supplementary materials). To verify that the results are not a consequence of PCR bias, PCoA and MGLM analyses were also performed with a distance matrix based on Sørensen similarity, which is a presence/absence index (Supplementary materials).

The number of samples of the branching species (*S. hystrix* and *P. damicornis*) did not allow statistical analyses to test for differences between high $p\text{CO}_2$ and reference sites, although it did permit a comparison of the endolithic communities associated with the different coral hosts (Supplementary Table 1). To investigate the community structure related to host species, a rarefaction threshold of 707 reads for the 16S rRNA gene, 713 for the *tufA* and 3257 for the UPA marker was used, allowing the inclusion of a larger number of samples in the analysis. Samples with sequencing depths lower than these thresholds were excluded, resulting in 35 samples in the 16S rRNA gene and UPA datasets and 27 in the *tufA* datasets (Supplementary Table S1). Alpha diversity, Kruskal–Wallis test (for OTUs), ANOVA (for taxon groups), PCoA and MGLM were performed on this dataset as previously described, but here, samples from different $p\text{CO}_2$ conditions from conspecific host species were combined in order to investigate the community structure purely associated with coral host species.

Results

Effects of $p\text{CO}_2$ conditions on the Porites spp. endolithic microbiome

A total of 6,584,274 sequence reads were recovered for the samples analysed here, 4,405,336 belonging to *Porites* spp. samples (ca. 25% of these reads belonged to a marker that is not analysed in this manuscript). After stringent pre and post OTU-clustering and filtering, a total of 119,367 (16S rRNA gene), 109,948 (*tufA*) and 393,816 (UPA) reads remained in the analysed *Porites* spp. dataset. A less stringent quality filtering which resulted in inclusion of more sequence reads did not alter the community patterns (see Supplementary materials). The alpha diversity statistics, including Chao1 and observed OTUs, indicated that the species richness of the endolithic communities associated with *Porites* spp. was not significantly different between high $p\text{CO}_2$ and reference sites (Table 1). Although

the relative abundance of some microbial taxa differed between high $p\text{CO}_2$ and reference sites (Figure 1, see also Supplementary figure 2 for a sample-based representation), the differences were not statistically significant (Bonferroni-corrected p -values = 1), neither at the OTU level (Supplementary table S2) nor at higher taxonomic levels (Supplementary table S3). Accordingly, principal coordinate analysis and MGLM did not reveal any significant pattern between sites with all three markers (Figure 2).

We further investigated whether any of the different phylogenetic lineages in the endolithic algal communities differed in abundance at high $p\text{CO}_2$ and reference sites. A phylogenetic heatmap of relative abundances (Supplementary figure 3) indicated that phylogenetic relatedness among green algae is not correlated with different abundances in high $p\text{CO}_2$ or reference sites. Notably, three algal OTUs were present in either high $p\text{CO}_2$ or reference sites, but not in both (Supplementary figure 3).

Taxonomic profiling of the Porites spp. endolithic community

The microbial community observed in the skeletons of *Porites* spp. was highly diverse and variable among samples within $p\text{CO}_2$ conditions. Prokaryotic members of the microbiome (observed in the 16S rRNA gene dataset) accounted for most of the species diversity (Table 1), and no bacterial OTUs were present in all *Porites* spp. samples. The most abundant phylum recovered was *Proteobacteria*, followed by *Bacteroidetes* and Archaea (Figure 1A, Supplementary figure 2). The relative abundance of the nitrogen-fixing order *Rhizobiales* (Alphaproteobacteria) in all *Porites* spp. samples was $9.1\% \pm 4\%$ standard deviation (hereafter $\pm$ only), and the phylum of green sulphur bacteria, *Chlorobi*, was $0.4\% \pm 1\%$. Members of the *Bacteroidetes* were twice as abundant at high $p\text{CO}_2$ sites ($12.1\% \pm 10\%$ in reference versus $23.9\% \pm 17\%$ in high $p\text{CO}_2$ sites), mostly due to a higher abundance within the classes *Cytophagia* ($4.8\% \pm 7\%$ versus $10.4\% \pm 9\%$), *Flavobacteria* ($2.2\% \pm 2\%$ versus $3.7\% \pm 3\%$) and *Saprospirae* ($2.4\% \pm 5\%$ versus $9.4\% \pm 15\%$). We also observed a lower abundance of the Archaeal class *Parvarchaea* in the high $p\text{CO}_2$ site ($1.6\% \pm 0.9\%$ versus $11.6\% \pm 12\%$ in high $p\text{CO}_2$ and reference sites, respectively). A representation of the relative abundances of bacterial phyla on a sample-by-sample basis can be found in Supplementary figure 2. These differences were, however, not statistically significant based on ANOVA and Kruskal-Wallis tests (Bonferroni-corrected p -values = 1, Supplementary tables S2 and S3).

The *tufA* dataset (Figure 1B) suggested that the algal community was dominated ($64.7\% \pm 36\%$) by lineages of the *Ostreobiaceae* (Chlorophyta, Bryopsidales). Although no *tufA* OTUs were found to be omnipresent in *Porites* spp., siphonous green algae (order Bryopsidales) were present in all samples analysed here, indicating that they are ubiquitous members of the coral skeleton core microbiome. *Ostreobium* clade #1 showed the highest relative abundance ($33\% \pm 40\%$), followed by *Ostreobium* clades #4, #3 and #2. While the relative abundance of clade #1 was similar between sites, the relative abundance of the other *Ostreobium* clades varied substantially, but not statistically significant, between reference and high $p\text{CO}_2$ sites. A high abundance of an unclassified group of OTUs belonging to the green algal order Bryopsidales was also observed, particularly in the reference site (Figure 1B).

The prevalence of green algal lineages in the skeletons of *Porites* spp. was also suggested by the UPA dataset (Figure 1C), which shows that $86.8\% \pm 15\%$ of the reads belong to the green algal order Bryopsidales. There are no UPA reference sequences for *Ostreobium* clades #1 and #2, therefore possible sequences of these clades might have been classified as clades #3, #4 or “unclassified Bryopsidales” by the RDP classifier, which may explain the differences in the abundances of *Ostreobium* clades between the *tufA* and UPA datasets. One UPA OTU was found across all *Porites* spp. samples (OTU_3). This OTU was only classified at the kingdom level as a eukaryote (Supplementary table S2) and showed little similarity to known red algae and Stramenopiles species (blastn e-values $\leq 4e-79$ but Identity $\leq 79\%$). Differences in the relative abundances between reference and high pCO_2 sites were minimal (Figure 1C, Supplementary figure 2).

Endolithic communities across different coral host species

The prokaryotic endolithic communities of *Seriatopora hystrix* and *Pocillopora damicornis* were significantly less rich than those found in *Porites* spp., as indicated by Chao1 and observed OTUs indices (p -value = 0.003, Table 2). A significant difference was detected in the relative abundances of certain OTUs belonging to the *Endozoicimonaceae* family between coral species, with the highest abundance in *P. damicornis* (Kruskal-Wallis, p -values < 0.0002 , Supplementary table S4). The *Porites* spp. samples had a higher relative abundance of an OTU related to the order *Rhizobiales* (genus *Aifella*), and *P. damicornis* showed a significantly (p -value = 0.03) higher abundance of an OTU related to the phylum *Bacteroidetes* (order *Cytophagales*; Supplementary table S4). At higher taxonomic levels, the relative abundance of the phyla *Planctomycetes*, *Bacteroidetes*, and the bacterial phylum OD1 were significantly different among coral hosts (ANOVA, p -values = 0.007 and 0.01 respectively, Figure 3A, Supplementary figure 2, Supplementary table S5). Principal coordinate analysis (PCoA) showed evident differences in the prokaryotic microbiome associated with different coral hosts: *Porites* spp. samples clustered together, clearly separated from the two branching species (Figure 4A). The MGLM analysis confirmed a significant difference between the prokaryotic communities of different coral hosts (Figure 4A).

The alpha diversity of green algae (i.e. Chao 1 and Observed OTUs within the *tufA* dataset) was significantly different between *S. hystrix* and *Porites* spp., but no significant differences were observed within the taxonomically broader spectrum of eukaryotic algae amplified with the UPA marker (Table 2). The Kruskal–Wallis test suggested no significant difference in the relative abundances of particular OTUs between host species (neither within the *tufA* nor within the UPA dataset), at least when corrected (Bonferroni) p -values were taken into consideration (Supplementary table S4). After rarefying the sequences, the *tufA* dataset was reduced to a single *P. damicornis* sample (Supplementary table S1), therefore we could not test for differences in alpha diversity or relative abundances in the *P. damicornis* community compared to other host corals. We observed a significantly different relative abundance of *Ostreobium* spp. (order Bryopsidales) among coral hosts within the UPA dataset (ANOVA, p -value = 0.000002), but not in the *tufA* dataset (p -values > 0.05 , Figure 3B and 3C, Supplementary figure 2, Supplementary table S5). *Seriatopora hystrix* had a high

and variable ($83\% \pm 57\%$) relative abundance of endolithic lineages related to the macroalga *Halimeda* spp. within the *tufA* dataset, while this group constituted a minimal fraction ($0.04\% \pm 0.2\%$) of the endolithic community of *Porites* spp. (Figure 3B). No pattern was observed within the PCoA plot of the community composition using the *tufA* marker (Figure 4B). The UPA marker, which comprised more samples of the branching species, showed that *Porites* spp. samples cluster together and away from *S. hystrix* and *P. damicornis*, although two outliers belonging to branching samples cluster with *Porites* spp. (Figure 4C). The MGLM analysis suggested no significant differences between the algal communities of different coral hosts (Figure 4B and C), except when a presence-absence distance matrix was used (Supplementary materials, Supplementary figure 4).

Discussion

Our results show that the prokaryotic and eukaryotic microbiome in the skeletons of *Porites* spp. are highly diverse but indistinguishable between corals inhabiting naturally high $p\text{CO}_2$ reefs and ambient conditions. Ocean acidification is predicted to affect the coral reef CaCO_3 budget and its biological associations (Meron *et al.* 2011; Andersson & Gledhill 2013; Morrow *et al.* 2015), and depending on the experiment, endolithic communities were shown to either exacerbate or buffer the effects of these environmental changes (Fine & Loya 2002; Tribollet *et al.* 2009; Reyes-Nivia *et al.* 2013). Our results suggest that the composition of endolithic communities, at least in *Porites* spp., is virtually unaffected by the surrounding high $p\text{CO}_2$ water from a natural volcanic seep, and therefore less likely to be disturbed by OA than we previously thought. Although homogeneous between high $p\text{CO}_2$ and reference sites, we show that the endolithic community is highly diverse and structured among coral host species.

A stable microbiome

The mechanisms influencing the structure of the endolithic microbiome (regardless of variable $p\text{CO}_2$ conditions), are currently unknown. We raise here two hypotheses that may explain our results. The "stable habitat" hypothesis assumes that the endolithic environment is highly homeostatic so that pH is maintained inside the skeletons regardless of external changes in the surrounding water. The "tolerant endolith" hypothesis is based on the notion that endolithic microorganisms have a wide pH tolerance range, potentially wider than the microbes associated with the tissues and mucus.

The first hypothesis is supported by the ability of some corals to up-regulate the pH at the tissue-skeleton interface, which allows them to calcify and grow even under high $p\text{CO}_2$ (McCulloch *et al.* 2012; Venn *et al.* 2013; Georgiou *et al.* 2015). The pH within coral cells remains relatively constant throughout the day (7.05-7.46 units), likely due to membrane transporters that extrude the excess of by-products of photosynthesis and respiration to maintain a stable intracellular pH (Laurent *et al.* 2013). This process may indirectly create a stable microhabitat within the coral skeleton that is protected from shifting pH in the surrounding seawater (see also Shashar *et al.* 1997). The observation that

radioactivity in seawater impacted corals' living tissue but did not reach their endolithic zone (Odum & Odum 1955) supports this notion.

One problem with the stable habitat hypothesis is that the pH in the skeletons of *Porites* (*compressa*) can vary daily from 7.7 to 8.5 pH units, mostly due to the by-products of respiration and photosynthesis of the coral and *Symbiodinium* that are exported to the skeleton (Shashar & Stambler 1992). This daily variation is well above the projections of OA for the near future, which predicts a pH drop of 0.4 units by 2100, and up to 0.7 units by 2300 (Raven *et al.* 2005; Hoegh-Guldberg *et al.* 2007). A reasonable counterargument is that the direction of the movement (pH reduction due to OA) may be more important than the daily variation within skeletons – for example, microorganisms living under 7.7 to 8.5 pH units not necessarily withstand a seawater pH shift from 7.8 to 7.6 pH units. The resilience of at least some microbes might also be related to their boring mechanism, which involves a sophisticated control of intracellular pH (associated with calcium pumps and protons counter-transport) in endolithic cyanobacteria (Garcia-Pichel 2006; Garcia-Pichel *et al.* 2010). Therefore our second hypothesis that organisms exposed to daily pH fluctuations within the skeleton are adapted to cope with a wide range of $p\text{CO}_2$ conditions may be more accurate. Experimental work and genomic data of endolithic organisms will help to test the tolerant endolith and the stable habitat hypotheses. For example, specialization to the low light experienced in the endolithic habitat has been observed in the plastid genome of *Ostreobium quekettii* (Marcelino *et al.* 2016) and the presumed pH tolerance may also be reflected in the genomes of endolithic organisms.

The lack of discernible differences in the endolithic community composition between high $p\text{CO}_2$ and reference corals observed here is in agreement with the results of an experiment where *Balanophyllia europaea* and *Cladocora caespitosa* corals were transplanted to a naturally high $p\text{CO}_2$ area (Meron *et al.* 2012). In an aquarium-based experiment conducted over a shorter time, the bacterial community composition present in the tissue, skeleton and mucus of *Acropora eurystroma* were found to be affected by high $p\text{CO}_2$, but further analysis using clone libraries suggested that only the prokaryotic communities of the mucus and tissue, not the skeleton, were affected by low pH (Meron *et al.* 2011). These different observations might be associated with the different time spans and experimental setups of the two studies, and it is likely that the microbial community associated with different coral taxa have different responses to acidification. The resilience of endolithic algae to acidification has also been observed: the net photosynthesis and respiration of algae growing at the surface of dead coral blocks was severely impacted upon exposure to high $p\text{CO}_2$ treatments, while the endolithic flora was unaffected (Tribollet *et al.* 2006). Studies have demonstrated that endolithic algae actually benefit from low pH and tend to increase in biomass under high $p\text{CO}_2$ conditions (Tribollet *et al.* 2009; Reyes-Nivia *et al.* 2013; Enochs *et al.* 2016).

The observation that high $p\text{CO}_2$ did not impact the endolithic community of *Porites* spp. does not necessarily imply that coral holobionts are immune to ocean acidification. It is possible that the methods used here are not sufficiently powerful to detect the effects of high $p\text{CO}_2$ on endolithic microbial communities. However, the fact that we detected significant differences among coral hosts even though the sampling size for the branching corals was smaller, indicates that our methods and

sampling design are adequate and it is unlikely that differences among high $p\text{CO}_2$ and reference sites were present but went undetected. It is possible though that high $p\text{CO}_2$ impacts the endolithic communities of other coral species that were not examined. It is interesting to note that massive *Porites* spp. dominate the reef near volcanic seeps while the presence of branching species (e.g. *Acropora* spp.) was largely reduced (Fabricius *et al.* 2011). Our analyses are restricted to the volcanic seeps of Milne Bay, which have relatively small areas under high $p\text{CO}_2$ and are surrounded by ambient seawater. Further studies at additional sites impacted by high $p\text{CO}_2$ and across a wider range of coral species is required to evaluate the results in our study across the broader ecological context of effects of OA on coral microbiomes.

Diversity and potential functional roles of the endolithic microbiome

Bacteria related to *Endozoicomonas* spp. (class Gammaproteobacteria) are predicted to have a key role in the coral holobiont. These bacteria have been shown to be endosymbionts, forming aggregations within coral tissues (Neave *et al.* 2017), potentially contributing to nutrient cycling and structuring of the microbiome through the production of quorum-sensing signalling metabolites and antimicrobial compounds (Meyer *et al.* 2014; Morrow *et al.* 2015 and references therein). The relative abundance of *Endozoicomonaceae* within coral tissues appears to be sensitive to high $p\text{CO}_2$ (Morrow *et al.* 2015; Webster *et al.* 2016), but in the skeletons of *Porites* spp. analysed here, they did not differ significantly between samples from different $p\text{CO}_2$ conditions (Supplementary Table S2). We observed a significantly higher relative abundance of two *Endozoicomonaceae* OTUs in the skeletons of *P. damicornis* when compared to the other two coral species, possibly reflecting stable associations of *Endozoicomonaceae* species with this coral host (see Neave *et al.* 2017). Although some of the sequences retrieved here may derive from other parts of the coral, other studies have also detected members of *Endozoicomonaceae* in the endolithic community (Williams *et al.* 2015; Marcelino & Verbruggen 2016; see also Ainsworth *et al.* 2015).

Bacteria in the phylum *Bacteroidetes* are often associated with coral disease and have been shown to increase in relative abundance under reduced pH (Vega Thurber *et al.* 2009). The average relative abundance of this group doubled in endolithic communities from high $p\text{CO}_2$ sites, but this difference was not significant likely due to the high level of variation in community composition among colonies within sites. This increase was mostly due to a higher abundance of the classes *Saprospirae*, *Flavobacteria* and *Cytophagia*, which contain most of the known marine algicide bacteria (Furusawa *et al.* 2003; Mayali & Azam 2004; Zozaya-Valdes *et al.* 2015). It is plausible that a higher relative abundance of these microorganisms is associated with an increase in endolithic algal biomass under high $p\text{CO}_2$ (see Reyes-Nivia *et al.* 2013, Johnson *et al.* 2017). Rather than compromising coral health, these bacteria may control excessive endolithic algal growth and may help to maintain a stable community composition under ocean acidification.

Microorganisms involved in nitrogen cycling may be fundamental to coral resilience to ocean acidification and climate change (Rädecker *et al.* 2014; Santos *et al.* 2014; Radecker *et al.* 2015). We found a diverse community of nitrogen fixing (diazotrophic) microorganisms inhabiting coral

skeletons. The majority (in terms of relative abundances) belonged to the order *Rhizobiales*, a group that appears to form stable symbiotic associations with corals (Lema *et al.* 2014). Green sulphur (also diazotrophic) bacteria in the phylum *Chlorobi*, previously documented as prevalent members of the endolithic community in the coral *Isopora* (Yang *et al.* 2016), were found at low relative abundances in the samples analysed here and in a previous study (Marcelino & Verbruggen 2016). Cyanobacterial OTUs captured with the UPA marker, while not abundant, were very diverse and mostly unclassified at lower taxonomic levels. Cyanobacteria have been shown to fix nitrogen in coral tissues (Lesser *et al.* 2004; Radecker *et al.* 2015) and can be responsible for a large fraction of the nitrogen fixation observed in their skeletons (Crossland & Barnes 1976; Davey *et al.* 2007).

Endolithic algal biomass has been shown to increase under high $p\text{CO}_2$, as phototrophic organisms benefit from the increased availability of carbon dioxide for photosynthesis (Tribollet *et al.* 2009; Reyes-Nivia *et al.* 2013). Indeed, we observed a higher relative abundance of all *Ostreobium* clades in *Porites* spp. samples from high $p\text{CO}_2$ sites, but the variability among replicates (i.e. *Porites* spp. samples within sites) was also high, making it difficult to draw conclusions about whether *Ostreobium* spp. are competitively superior to other endolithic algal lineages under OA. Whether the increase in algal biomass is a threat to corals under OA depends on whether the associated bioerosion levels will exceed reef accretion (calcification). Besides increasing bioerosion, excessive endolithic algal growth can penetrate the coral living tissue, possibly increasing their susceptibility to infections (Peters 1984; Fine *et al.* 2006). An increase in endolithic algae may also be beneficial to the coral by providing them with vital nutrients, which is especially important during coral bleaching events (Schlichter *et al.* 1995; Fine & Loya 2002).

The possibility that the endolithic microbiome contributes to the resilience of corals under future OA conditions deserves further attention. Massive *Porites* spp. may be considered more competitive under OA than branching species based on their prevalence at naturally high $p\text{CO}_2$ sites (Fabricius *et al.* 2011). We visibly observed that our massive *Porites* spp. samples had higher colonisation with endolithic algae compared to the branching species. The photosynthetic activity of *Symbiodinium* plays an important role in maintaining pH homeostasis within corals (Gibbin *et al.* 2014), and it is possible that endolithic algae provide a similar service within the skeleton. The biomass of endolithic algae may exceed that of *Symbiodinium* by 16-fold (Odum & Odum 1955) and can contribute significantly to the buffering capacity of the holobiont (see Yamazaki *et al.* 2008; Reyes-Nivia *et al.* 2014). It is noteworthy that several functionally important microorganisms (e.g. *Endozoicomonaceae* and *Bacteroidetes*) often found in coral tissues and mucus also occur in coral skeletons (Sweet *et al.* 2010; Ainsworth *et al.* 2015; Willians *et al.* 2015; Marcelino & Verbruggen 2016; this study). It is possible that some of these microorganisms were initially associated with polyp tissues, but remained after removing the tissues from the skeleton with pressurized air, especially in *Porites* spp. which is a perforate coral with tissues that penetrate the skeleton, or that they penetrated the coral skeleton when the samples were placed in the storage buffer. It is noteworthy however that multiple studies have found tissue-associated microbes within coral skeletons, and that the majority of OTUs commonly found in healthy corals have also been found in bare coral skeleton, but not in seawater or in a diseased coral tissue (Fernando *et al.* 2015). It is possible therefore that the coral

skeleton serves as a reservoir for the microbiome and provides a source of beneficial bacteria to coral tissues, analogous to the human appendix which functions as a safe house for symbiotic microbes that repopulate the intestine following acute illness (Randal Bollinger *et al.* 2007). Acute environmental stress can disrupt symbiotic relationships among hosts and symbionts (see Hawkins *et al.* 2013), and a stable endolithic community may assist in the recovery of the coral microbiome after environmental (and/or physiological) conditions stabilize.

Different host species harbour distinct endolithic communities

The endolithic communities of the branching corals *Seriatopora hystrix*, *Pocillopora damicornis* and the massive *Porites* spp. contain significantly different relative abundances of functionally important members of the microbiome (including species of *Endozoicimonaceae* and *Bacteroidetes*) and appear to separate based on morphology or taxonomy (as both branching species belong to the family *Pocilloporidae*). First, the two branching species harbour a reduced diversity of bacteria and algae. The low relative abundance of *Ostreobium* spp. in the endolithic communities of branching species is surprising, considering the generally ubiquitous nature of this alga in coral skeletons (Odum & Odum 1955; Tribollet 2008a; Gutner-Hoch & Fine 2011). Instead of *Ostreobium* spp., the coral *S. hystrix* has a high relative abundance of OTUs related to a macroalga (*Halimeda* spp.), which has only recently been reported in coral skeletons. It is possible that *Halimeda* spp. occur in the coral skeleton in the form of rhizoids that have penetrated the limestone, or most likely, as an unknown microscopic and endolithic life stage of two *Halimeda* species (*H. discoidea* and *H. micronesica*) that are commonly present in metabarcoding studies of endolithic communities (Marcelino & Verbruggen 2016; Sauvage *et al.* 2016; this study).

The observed differences in endolithic community composition among coral hosts may be a result of specialization to particular host traits or reflect co-evolution between coral hosts and endolithic species. The living tissue of *Porites (lobata)* is about five times thicker, penetrates the skeleton and contains a higher density of *Symbiodinium* than the living tissue of *P. damicornis* and *S. hystrix* (Yost *et al.* 2013). Tissue thickness would influence the amount of light that penetrates and reflects within the inner parts of the skeleton and may influence the composition of the endolithic community. Branching coral species also tend to grow faster than massive corals (Gates & Ainsworth 2011), and the branch tips collected in this study may have a younger population of endoliths in comparison to more mature sections of the colony base (a pattern reported in Pica *et al.* 2016). Future studies would benefit from examining the microbiome associated with different areas of the colony and possible specialization to skeletal features. Alternatively (and not mutually exclusively), endolithic lineages might form stable community assemblies that have co-evolved with the coral host, or the host species have some control over the composition of the endolithic community by selecting beneficial taxa. Mutualistic relationships between corals and their endolithic associates have been suggested in several studies (Odum & Odum 1955; Schlichter *et al.* 1995; Schlichter *et al.* 1997; Fine & Loya 2002; Försterra & Häussermann 2008; Titlyanov *et al.* 2009), and future research would benefit from characterizing possible co-evolutionary processes among coral species and endolithic microorganisms.

530

531 Conclusions

532 This study reports a diverse microbiome within the skeletons of *Porites* spp., and
533 demonstrates that little discernible patterns exist in this microbiome across ambient and naturally high
534 $p\text{CO}_2$ environments. We show that the endolithic community shares several functionally important
535 microbes with the coral tissue layer. Environmental stress can induce corals to lose their symbiotic
536 microorganisms, and a diverse endolithic microbial community might serve as a reservoir to recolonise
537 the microbiome in the coral tissue after the re-establishment of their physiological equilibrium. We
538 found functionally important members in the endolithic community, including members in the
539 *Endozoicimonaceae* and *Bacteroidetes*, forming distinct associations with the different host coral
540 families, an observation consistent with the endolithic reservoir proposition. The diversity and
541 community structure observed in this study form the baseline for future studies aiming to investigate
542 the roles of endolithic microorganisms in enabling corals to endure climate change.

543

544 Acknowledgements

545 This work was supported by the Australian Biological Resources Study (RFL213-08), the Australian
546 Research Council (FT110100585, DP150100705), the Holsworth Wildlife Research Endowment, the
547 Albert Shimmings Fund and the University of Melbourne (ECR grant to HV and scholarship to VRM),
548 Melbourne Bioinformatics (project UOM0007) and the Nectar Research Cloud (National Collaborative
549 Research Infrastructure Strategy). We are thankful to Emmanuelle Botté for coral collections in Papua
550 New Guinea and Sara Bell for laboratory assistance in processing samples. The expedition to conduct
551 the fieldwork and collections was funded by the Australian Institute of Marine Science. We thank the
552 reviewers (including from Axios) for their constructive feedback.

553

554 References

- 555 Ainsworth T, Krause L, Bridge T, Torda G, Raina J-B, *et al.* (2015) The coral core microbiome
556 identifies rare bacterial taxa as ubiquitous endosymbionts. *ISME Journal*, **9**, 2261-2274.
- 557 Andersson AJ, Gledhill D (2013) Ocean acidification and coral reefs: effects on breakdown,
558 dissolution, and net ecosystem calcification. *Annual Review of Marine Science*, **5**, 321-348.
- 559 Anthony KRN, Kline DI, Diaz-Pulido G, Dove S, Hoegh-Guldberg O (2008) Ocean acidification
560 causes bleaching and productivity loss in coral reef builders. *Proceedings of the National*
561 *Academy of Sciences, USA*, **105**, 17442-17446.
- 562 Blackall LL, Wilson B, van Oppen MJ (2015) Coral-the world's most diverse symbiotic ecosystem.
563 *Molecular Ecology*, **24**, 5330-5347.
- 564 Bourne DG, Garren M, Work TM, Rosenberg E, Smith GW, Harvell CD (2009) Microbial disease and
565 the coral holobiont. *Trends in Microbiology*, **17**, 554-562.

566 Bourne DG, Morrow KM, Webster NS (2016) Insights into the coral microbiome: underpinning the
567 health and resilience of reef ecosystems. *Annual Review of Microbiology*, **70**, 317-340.

568 Caporaso JG, Bittinger K, Bushman FD, DeSantis TZ, Andersen GL, Knight R (2010a) PyNAST: a
569 flexible tool for aligning sequences to a template alignment. *Bioinformatics*, **26**, 266-267.

570 Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, *et al.* (2010b) QIIME allows
571 analysis of high-throughput community sequencing data. *Nature Methods*, **7**, 335-336.

572 Chazottes V, Campion-Alsumard TL, Peyrot-Clausade M (1995) Bioerosion rates on coral reefs:
573 interactions between macroborers, microborers and grazers (Moorea, French Polynesia).
574 *Palaeogeography, Palaeoclimatology, Palaeoecology*, **113**, 189-198.

575 Clements KD, German DP, Piché J, Tribollet A, Choat JH (2016) Integrating ecological roles and
576 trophic diversification on coral reefs: multiple lines of evidence identify parrotfishes as
577 microphages. *Biological Journal of the Linnean Society*, **120**, 729-751.

578 Crook ED, Cohen AL, Rebolledo-Vieyra M, Hernandez L, Paytan A (2013) Reduced calcification and
579 lack of acclimatization by coral colonies growing in areas of persistent natural acidification.
580 *Proceedings of the National Academy of Sciences, USA*, **110**, 11044-11049.

581 Crossland CJ, Barnes DJ (1976) Acetylene reduction by coral skeletons. *Limnology and*
582 *Oceanography*, **21**, 153-156.

583 Davey M, Holmes G, Johnstone R (2007) High rates of nitrogen fixation (acetylene reduction) on coral
584 skeletons following bleaching mortality. *Coral Reefs*, **27**, 227-236.

585 Del Campo J, Pombert JF, Slapeta J, Larkum A, Keeling PJ (2017) The 'other' coral symbiont:
586 *Ostreobium* diversity and distribution. *ISME Journal*, **11**, 296-299.

587 DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, *et al.* (2006) Greengenes, a chimera-
588 checked 16S rRNA gene database and workbench compatible with ARB. *Applied and*
589 *Environmental Microbiology*, **72**, 5069-5072.

590 Edgar RC (2013) UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nature*
591 *Methods*, **10**, 996-998.

592 Enochs IC, Manzello DP, Tribollet A, Valentino L, Kolodziej G, *et al.* (2016) Elevated colonization of
593 microborers at a volcanically acidified coral reef. *PLoS One*, **11**, e0159818.

594 Esteves AI, Amer N, Nguyen M, Thomas T (2016) Sample processing impacts the viability and
595 cultivability of the sponge microbiome. *Frontiers in Microbiology* **7**, 499.

596 Fabricius KE, De'ath G, Noonan S, Uthicke S (2014) Ecological effects of ocean acidification and
597 habitat complexity on reef-associated macroinvertebrate communities. *Proceedings of the*
598 *Royal Society B*, **281**, 20132479.

599 Fabricius KE, Langdon C, Uthicke S, Humphrey C, Noonan S, *et al.* (2011) Losers and winners in
600 coral reefs acclimatized to elevated carbon dioxide concentrations. *Nature Climate Change*, **1**,
601 165-169.

602 Fama P, Wysor B, Kooistra WHCF, Zuccarello GC (2002) Molecular phylogeny of the genus
603 *Caulerpa* (Caulerpales, Chlorophyta) inferred from chloroplast *tufA* gene. *Journal of*
604 *Phycology*, **38**, 1040-1050.

605 Fang JK, Mello-Athayde MA, Schonberg CH, Kline DI, Hoegh-Guldberg O, Dove S (2013) Sponge
606 biomass and bioerosion rates increase under ocean warming and acidification. *Global Change*
607 *Biology*, **19**, 3581-3591.

608 Fernando SC, Wang J, Sparling K, Garcia GD, Francini-Filho RB, *et al.* (2015) Microbiota of the
609 major South Atlantic reef building coral *Mussismilia*. *Microbial Ecology*, **69**, 267-280.

610 Fine M, Loya Y (2002) Endolithic algae: an alternative source of photoassimilates during coral
611 bleaching. *Proceedings of the Royal Society B*, **269**, 1205-1210.

612 Fine M, Roff G, Ainsworth TD, Hoegh-Guldberg O (2006) Phototrophic microendoliths bloom during
613 coral “white syndrome”. *Coral Reefs*, **25**, 577-581.

614 Försterra G, Häussermann V (2008) Unusual symbiotic relationships between microendolithic
615 phototrophic organisms and azooxanthellate cold-water corals from Chilean fjords. *Marine*
616 *Ecology Progress Series*, **370**, 121-125.

617 Foster T, Falter JL, McCulloch MT, Clode PL (2016) Ocean acidification causes structural deformities
618 in juvenile coral skeletons. *Science Advances*, **2**, e1501130-e1501130.

619 Furusawa G, Yoshikawa T, Yasuda A, Sakata T (2003) Algicidal activity and gliding motility of
620 *Saprospira* sp. SS98-5. *Canadian Journal of Microbiology / Revue Canadienne de*
621 *Microbiologie*, **49**, 92-100.

622 Garcia-Pichel F (2006) Plausible mechanisms for the boring on carbonates by microbial phototrophs.
623 *Sedimentary Geology*, **185**, 205-213.

624 Garcia-Pichel F, Ramirez-Reinat E, Gao Q (2010) Microbial excavation of solid carbonates powered
625 by P-type ATPase-mediated transcellular Ca_2^+ transport. *Proceedings of the National*
626 *Academy of Sciences, USA*, **107**, 21749-21754.

627 Gates RD, Ainsworth TD (2011) The nature and taxonomic composition of coral symbiomes as drivers
628 of performance limits in scleractinian corals. *Journal of Experimental Marine Biology and*
629 *Ecology*, **408**, 94-101.

630 Georgiou L, Falter J, Trotter J, Kline DI, Holcomb M, *et al.* (2015) pH homeostasis during coral
631 calcification in a free ocean CO_2 enrichment (FOCE) experiment, Heron Island reef flat, Great
632 Barrier Reef. *Proceedings of the National Academy of Sciences, USA*, **112**, 13219-13224.

- 633 Gibbin EM, Putnam HM, Davy SK, Gates RD (2014) Intracellular pH and its response to CO₂-driven
634 seawater acidification in symbiotic versus non-symbiotic coral cells. *Journal of Experimental*
635 *Biology*, **217**, 1963-1969.
- 636 Grange JS, Rybarczyk H, Tribollet A (2015) The three steps of the carbonate biogenic dissolution
637 process by microborers in coral reefs (New Caledonia). *Environmental Science and Pollution*
638 *Research*, **22**, 13625-13637.
- 639 Gutner-Hoch E, Fine M (2011) Genotypic diversity and distribution of *Ostreobium quekettii* within
640 scleractinian corals. *Coral Reefs* **30**, 643-650.
- 641 Hawkins TD, Bradley BJ, Davy SK (2013) Nitric oxide mediates coral bleaching through an apoptotic-
642 like cell death pathway: evidence from a model sea anemone-dinoflagellate symbiosis. *FASEB*
643 *Journal*, **27**, 4790-4798.
- 644 Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, *et al.* (2007) Coral reefs under
645 rapid climate change and ocean acidification. *Science*, **318**, 1737-1742.
- 646 Hofmann GE, Barry JP, Edmunds PJ, Gates RD, Hutchins DA, Klinger T, Sewell MA (2010) The
647 effect of ocean acidification on calcifying organisms in marine ecosystems: an organism to-
648 ecosystem perspective. *Annual Review of Ecology, Evolution and Systematics*, **41**, 127-147.
- 649 Johnson MD, Comeau S, Lantz CA, Smith JE (2017) Complex and interactive effects of ocean
650 acidification and temperature on epilithic and endolithic coral-reef turf algal assemblages.
651 *Coral Reefs*, Online first - doi:10.1007/s00338-017-1597-2
- 652 Joint I, Doney SC, Karl DM (2011) Will ocean acidification affect marine microbes? *ISME Journal*, **5**,
653 1-7.
- 654 Katoh K, Misawa K, Kuma K-i, Miyata T (2002) MAFFT: a novel method for rapid multiple sequence
655 alignment based on fast Fourier transform. *Nucleic Acids Research*, **30**, 3059-3066.
- 656 Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, Glockner FO (2013) Evaluation of
657 general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-
658 based diversity studies. *Nucleic Acids Research*, **41**, e1.
- 659 Krediet CJ, Ritchie KB, Paul VJ, Teplitski M (2013) Coral-associated micro-organisms and their roles
660 in promoting coral health and thwarting diseases. *Proceedings of the Royal Society B*, **280**,
661 20122328.
- 662 Laurent J, Tambutte S, Tambutte E, Allemand D, Venn A (2013) The influence of photosynthesis on
663 host intracellular pH in scleractinian corals. *Journal of Experimental Biology*, **216**, 1398-
664 1404.
- 665 Le Campion-Alsumard T, Golubic S, Hutchings P (1995) Microbial endoliths in skeletons of live and
666 dead corals: *Porites lobata* (Moorea, French Polynesia). *Marine Ecology Progress Series*, **117**,
667 149-157.

- 668 Lema KA, Willis BL, Bourne DG (2014) Amplicon pyrosequencing reveals spatial and temporal
669 consistency in diazotroph assemblages of the *Acropora millepora* microbiome. *Environmental*
670 *Microbiology*, **16**, 3345-3359.
- 671 Lesser MP, Mazel CH, Gorbunov MY, Falkowski PG (2004) Discovery of symbiotic nitrogen-fixing
672 cyanobacteria in corals. *Science*, **305**, 997-1000.
- 673 Lozupone C, Lladser ME, Knights D, Stombaugh J, Knight R (2011) UniFrac: an effective distance
674 metric for microbial community comparison. *ISME Journal*, **5**, 169-172.
- 675 Magoc T, Salzberg SL (2011) FLASH: fast length adjustment of short reads to improve genome
676 assemblies. *Bioinformatics*, **27**, 2957-2963.
- 677 Manzello DP, Kleypas JA, Budd DA, Eakin CM, Glynn PW, Langdon C (2008) Poorly cemented coral
678 reefs of the eastern tropical Pacific: possible insights into reef development in a high-CO₂
679 world. *Proceedings of the National Academy of Sciences, USA*, **105**, 10450-10455.
- 680 Marcelino VR, Cremen MC, Jackson CJ, Larkum AA, Verbruggen H (2016) Evolutionary dynamics of
681 chloroplast genomes in low light: a case study of the endolithic green alga *Ostreobium*
682 *quekettii*. *Genome Biology and Evolution*, **8**, 2939-2951.
- 683 Marcelino VR, Verbruggen H (2016) Multi-marker metabarcoding of coral skeletons reveals a rich
684 microbiome and diverse evolutionary origins of endolithic algae. *Scientific Reports*, **6**, 31508.
- 685 Marcelino VR, Verbruggen H (2017) Reference datasets of *tufA* and UPA markers to identify algae in
686 metabarcoding surveys. *Data in Brief*, **11**, 273-276.
- 687 Mayali X, Azam F (2004) Algicidal bacteria in the sea and their impact on algal blooms. *The Journal*
688 *of Eukaryotic Microbiology*, **51**, 139-144.
- 689 McCulloch M, Falter J, Trotter J, Montagna P (2012) Coral resilience to ocean acidification and global
690 warming through pH up-regulation. *Nature Climate Change*, **2**, 623-627.
- 691 Meron D, Atias E, Iasur Kruh L, Elifantz H, Minz D, Fine M, Banin E (2011) The impact of reduced
692 pH on the microbial community of the coral *Acropora eurytoma*. *ISME Journal*, **5**, 51-60.
- 693 Meron D, Rodolfo-Metalpa R, Cunning R, Baker AC, Fine M, Banin E (2012) Changes in coral
694 microbial communities in response to a natural pH gradient. *ISME Journal*, **6**, 1775-1785.
- 695 Meyer JL, Paul VJ, Teplitski M (2014) Community shifts in the surface microbiomes of the coral
696 *Porites astreoides* with unusual lesions. *PLoS One*, **9**, e100316.
- 697 Morrow KM, Bourne DG, Humphrey C, Botte ES, Laffy P, *et al.* (2015) Natural volcanic CO₂ seeps
698 reveal future trajectories for host-microbial associations in corals and sponges. *ISME Journal*,
699 **9**, 894-908.
- 700 Moss RH, Edmonds JA, Hibbard KA, Manning MR, Rose SK, *et al.* (2010) The next generation of
701 scenarios for climate change research and assessment. *Nature*, **463**, 747-756.

- Neave MJ, Rachmawati R, Xun L, Michell CT, Bourne DG, Apprill A, Voolstra CR (2017) Differential specificity between closely related corals and abundant *Endozoicomonas* endosymbionts across global scales. *ISME Journal*, **11**, 186-200.
- Odum HT, Odum EP (1955) Trophic structure and productivity of a windward coral reef community on Eniwetok Atoll. *Ecological Monographs*, **25**, 291-320.
- Orr JC, Fabry VJ, Aumont O, Bopp L, Doney SC, *et al.* (2005) Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature*, **437**, 681-686.
- Paulson JN, Stine OC, Bravo HC, Pop M (2013) Robust methods for differential abundance analysis in marker gene surveys. *Nature Methods*, **10**, 1200-1202.
- Peters EC (1984) A survey of cellular reactions to environmental stress and disease in Caribbean scleractinian corals. *Helgoländer Meeresuntersuchungen*, **37**, 113-137.
- Pica D, Tribollet A, Golubic S, Bo M, Di Camillo CG, Bavestrello G, Puce S (2016) Microboring organisms in living stylasterid corals (Cnidaria, Hydrozoa). *Marine Biology Research* **12**, 573-582.
- Presting GG (2006) Identification of conserved regions in the plastid genome: implications for DNA barcoding and biological function. *Canadian Journal of Botany-Revue Canadienne De Botanique*, **84**, 1434-1443.
- Rädecker N, Meyer FW, Bednarz VN, Cardini U, Wild C (2014) Ocean acidification rapidly reduces dinitrogen fixation associated with the hermatypic coral *Seriatopora hystrix*. *Marine Ecology Progress Series*, **511**, 297-302.
- Radecker N, Pogoreutz C, Voolstra CR, Wiedenmann J, Wild C (2015) Nitrogen cycling in corals: the key to understanding holobiont functioning? *Trends in Microbiology*, **23**, 490-497.
- Randal Bollinger R, Barbas AS, Bush EL, Lin SS, Parker W (2007) Biofilms in the large bowel suggest an apparent function of the human vermiform appendix. *Journal of Theoretical Biology*, **249**, 826-831.
- Raven J, Caldeira K, Elderfield H, Hoegh-Guldberg O, Liss P, Riebesell, U., *et al.* (2005) *Ocean acidification due to increasing atmospheric carbon dioxide* The Royal Society. http://www.us-ocb.org/publications/Royal_Soc_OA.pdf
- Revell LJ (2012) phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution*, **3**, 217-223.
- Reyes-Nivia C, Diaz-Pulido G, Dove S (2014) Relative roles of endolithic algae and carbonate chemistry variability in the skeletal dissolution of crustose coralline algae. *Biogeosciences Discussions*, **11**, 2993-3021.
- Reyes-Nivia C, Diaz-Pulido G, Kline D, Guldberg OH, Dove S (2013) Ocean acidification and warming scenarios increase microbioerosion of coral skeletons. *Global Change Biology*, **19**, 1919-1929.

- 738 Rohwer F, Seguritan V, Azam F, Knowlton N (2002) Diversity and distribution of coral-associated
739 bacteria. *Marine Ecology Progress Series*, **243**, 1-10.
- 740 Santos HF, Carmo FL, Duarte G, Dini-Andreote F, Castro CB, *et al.* (2014) Climate change affects key
741 nitrogen-fixing bacterial populations on coral reefs. *ISME Journal*, **8**, 2272-2279.
- 742 Sauvage T, Schmidt WE, Suda S, Fredericq S (2016) A metabarcoding framework for facilitated
743 survey of endolithic phototrophs with *tufA*. *BMC Ecology*, **16**, 8.
- 744 Schlichter D, Kampmann H, Conrady S (1997) Trophic potential and photoecology of endolithic algae
745 living within coral skeletons. *Marine Ecology*, **18**, 299-317.
- 746 Schlichter D, Zscharnack B, Krisch H (1995) Transfer of photoassimilates from endolithic algae to
747 coral tissue. *Naturwissenschaften*, **82**, 561-564.
- 748 Schmieder R, Edwards R (2011) Quality control and preprocessing of metagenomic datasets.
749 *Bioinformatics*, **27**, 863-864.
- 750 Sharp KH, Ritchie KIMB (2012) Multi-partner interactions in corals in the face of climate change.
751 *Marine Biological Laboratory*, 66-77.
- 752 Shashar N, Banaszak AT, Lesser MP, Amrami D, Gan R (1997) Coral endolithic algae : life in a
753 protected environment. *Pacific Science*, **51**, 167-173.
- 754 Shashar N, Stambler N (1992) Endolithic algae within corals - life in an extreme environment. *Journal*
755 *of Experimental Marine Biology and Ecology*, **163**, 277-286.
- 756 Sherwood AR, Presting GG (2007) Universal primers amplify a 23S rDNA plastid marker in
757 eukaryotic algae and cyanobacteria. *Journal of Phycology*, **43**, 605-608.
- 758 Stamatakis A (2006) RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with
759 thousands of taxa and mixed models. *Bioinformatics*, **22**, 2688-2690.
- 760 Sweet MJ, Croquer A, Bythell JC (2010) Bacterial assemblages differ between compartments within
761 the coral holobiont. *Coral Reefs*, **30**, 39-52.
- 762 Titlyanov EA, Kiyashko SI, Titlyanova TV, Yakovleva IM (2009) $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in tissues of reef
763 building corals and the endolithic alga *Ostreobium quekettii* under their symbiotic and
764 separate existence. *Journal of Coral Reef Studies*, 169-175.
- 765 Tribollet A (2008a) The boring microflora in modern coral reef ecosystems: a review of its roles. In:
766 *Current developments in bioerosion* (eds. Wisshak M, Tapanila L), pp. 67-94. Springer,
767 Berlin, Heidelberg.
- 768 Tribollet A (2008b) Dissolution of dead corals by euendolithic microorganisms across the northern
769 Great Barrier Reef (Australia). *Microbial Ecology*, **55**, 569-580.
- 770 Tribollet A, Godinot C, Atkinson M, Langdon C (2009) Effects of elevated $p\text{CO}_2$ on dissolution of
771 coral carbonates by microbial euendoliths. *Global Biogeochemical Cycles*, **23**. GB3008.

772 Tribollet T, Atkinson MJ, Langdon C (2006) Effects of elevated $p\text{CO}_2$ on epilithic and endolithic
773 metabolism of reef carbonates. *Global Change Biology*, **12**, 2200-2208.

774 Uthicke S, Momigliano P, Fabricius KE (2013) High risk of extinction of benthic foraminifera in this
775 century due to ocean acidification. *Scientific Reports*, **3**, 1769.

776 Vega Thurber R, Willner-Hall D, Rodriguez-Mueller B, Desnues C, Edwards RA, *et al.* (2009)
777 Metagenomic analysis of stressed coral holobionts. *Environmental Microbiology*, **11**, 2148-
778 2163.

779 Venn AA, Tambutte E, Holcomb M, Laurent J, Allemand D, Tambutte S (2013) Impact of seawater
780 acidification on pH at the tissue-skeleton interface and calcification in reef corals.
781 *Proceedings of the National Academy of Sciences, USA*, **110**, 1634-1639.

782 Verbruggen H, Tribollet A (2011) Boring algae. *Current Biology*, **21**, R876-877.

783 Wang Q, Garrity GM, Tiedje JM, Cole JR (2007) Naïve Bayesian Classifier for rapid assignment of
784 rRNA sequences into the new bacterial taxonomy. *Applied and Environmental Microbiology*,
785 **73**, 5261-5267.

786 Wang Y, Naumann U, Wright ST, Warton DI (2012) mvabund- an R package for model-based analysis
787 of multivariate abundance data. *Methods in Ecology and Evolution*, **3**, 471-474.

788 Webster NS, Negri AP, Botte ES, Laffy PW, Flores F, *et al.* (2016) Host-associated coral reef microbes
789 respond to the cumulative pressures of ocean warming and ocean acidification. *Scientific*
790 *Reports*, **6**, 19324.

791 Webster NS, Negri AP, Flores F, Humphrey C, Soo R, *et al.* (2013) Near-future ocean acidification
792 causes differences in microbial associations within diverse coral reef taxa. *Environmental*
793 *Microbiology Reports*, **5**, 243-251.

794 Wickham H (2009) *ggplot2: Elegant Graphics for Data Analysis* Springer-Verlag New York.

795 Williams AD, Brown BE, Putschim L, Sweet MJ (2015) Age-related shifts in bacterial diversity in a
796 reef coral. *PLoS One* **10**, e0144902.

797 Yamazaki SS, Nakamura T, Yamasaki H (2008) Photoprotective role of endolithic algae colonized in
798 coral skeleton for the host photosynthesis. In: *Photosynthesis. Energy from the Sun* eds. Allen
799 JF, Gantt E, Golbeck JH, Osmond B), pp. 1391-1395. Springer Netherlands, Dordrecht.

800 Yang S-H, Lee STM, Huang C-R, Tseng C-H, Chiang P-W, *et al.* (2016) Prevalence of potential
801 nitrogen-fixing, green sulfur bacteria in the skeleton of reef-building coral *Isopora palifera*.
802 *Limnology and Oceanography*, **61**, 1078-1086.

803 Yost DM, Wang LH, Fan TY, Chen CS, Lee RW, Sogin E, Gates RD (2013) Diversity in skeletal
804 architecture influences biological heterogeneity and *Symbiodinium* habitat in corals. *Zoology*,
805 **116**, 262-269.

Zhou G, Yuan T, Cai L, Zhang W, Tian R, *et al.* (2016) Changes in microbial communities, photosynthesis and calcification of the coral *Acropora gemmifera* in response to ocean acidification. *Scientific Reports*, **6**, 35971.

Zozaya-Valdes E, Egan S, Thomas T (2015) A comprehensive analysis of the microbial communities of healthy and diseased marine macroalgae and the detection of known and potential bacterial pathogens. *Frontiers in Microbiology*, **6**, 146.

Data Accessibility

DNA sequences have been deposited in NCBI's Sequence Read Archive (SRA) under the accession IDs SAMN07251731 – SAMN07251770.

Author Contributions

KMM and DGB conceived and conducted the sampling. VRM performed DNA extractions, library preparation, analyses and drafted the manuscript. MvO, KMM, DGB and HV reviewed the analyses and contributed to writing.

Tables:

Table 1: Diversity indices based on the microbiome of *Porites* spp. skeletons from reference and high $p\text{CO}_2$ sites and standard deviations ($\pm$). Chao1 and Number of Observed OTUs between high $p\text{CO}_2$ and reference sites were compared with a two-sample t-test, p -values ≤ 0.05 suggest significant differences between sites, but were not observed. N = number of samples after rarefaction. Seqs = rarefaction threshold. OTUs = number of OTUs retrieved in each dataset after quality filtering.

	N	Seqs	OTUs	Chao1			Obs. OTUS		
				Reference	high $p\text{CO}_2$	p -value	Reference	high $p\text{CO}_2$	p -value
16S	20	2200	890	141.2 $\pm$ 52.4	148.5 $\pm$ 45.8	0.79	133.6 $\pm$ 49.9	140.7 $\pm$ 44.8	0.77
<i>tufA</i>	22	1400	59	7.8 $\pm$ 3.3	7.4 $\pm$ 2.8	0.78	7.7 $\pm$ 3.2	7.3 $\pm$ 2.8	0.74
UPA	22	7000	164	21.6 $\pm$ 7.1	24.4 $\pm$ 5.4	0.33	20.6 $\pm$ 6.7	23.2 $\pm$ 5.1	0.33

16S rRNA gene Chao1							
Group1	Group2	Group1 mean	Group1 std	Group2 mean	Group2 std	t stat	p-value
<i>S. hystrix</i>	<i>P. damicornis</i>	29.087	29.494	21.514	15.705	0.507	1.000
<i>S. hystrix</i>	<i>Porites</i> spp.	29.087	29.494	125.296	49.889	-4.363	0.003*
<i>P. damicornis</i>	<i>Porites</i> spp.	21.514	15.705	125.296	49.889	-4.854	0.003*
16S rRNA gene observed OTUs							
Group1	Group2	Group1 mean	Group1 std	Group2 mean	Group2 std	t stat	p-value
<i>S. hystrix</i>	<i>P. damicornis</i>	21.767	19.758	19.083	12.331	0.258	1.000
<i>S. hystrix</i>	<i>Porites</i> spp.	21.767	19.758	101.274	37.282	-4.865	0.003*
<i>P. damicornis</i>	<i>Porites</i> spp.	19.083	12.331	101.274	37.282	-5.138	0.003*
tufA Chao1							
Group1	Group2	Group1 mean	Group1 std	Group2 mean	Group2 std	t stat	p-value
<i>S. hystrix</i>	<i>P. damicornis</i>	2.033	0.858	3.450	0.000	1.168	1.000
<i>S. hystrix</i>	<i>Porites</i> spp.	2.033	0.858	9.606	3.849	-3.264	0.006*
<i>P. damicornis</i> ¹	<i>Porites</i> spp.	3.450	0.000	9.606	3.849	-1.531	0.309 ¹
tufA observed OTUs							
Group1	Group2	Group1 mean	Group1 std	Group2 mean	Group2 std	t stat	p-value
<i>S. hystrix</i>	<i>P. damicornis</i>	1.933	0.736	3.000	0.000	1.024	1.000
<i>S. hystrix</i>	<i>Porites</i> spp.	1.933	0.736	9.330	3.623	-3.388	0.003*
<i>P. damicornis</i> ¹	<i>Porites</i> spp.	3.000	0.000	9.330	3.623	-1.673	0.192 ¹
UPA Chao 1							
Group1	Group2	Group1 mean	Group1 std	Group2 mean	Group2 std	t stat	p-value
<i>S. hystrix</i>	<i>P. damicornis</i>	19.353	14.158	13.386	16.038	0.624	1.000
<i>S. hystrix</i>	<i>Porites</i> spp.	19.353	14.158	21.732	6.187	-0.591	1.000
<i>P. damicornis</i>	<i>Porites</i> spp.	13.386	16.038	21.732	6.187	-1.921	0.162
UPA observed OTUs							
Group1	Group2	Group1 mean	Group1 std	Group2 mean	Group2 std	t stat	p-value
<i>S. hystrix</i>	<i>P. damicornis</i>	18.100	12.882	12.567	14.816	0.630	1.000
<i>S. hystrix</i>	<i>Porites</i> spp.	18.100	12.882	19.539	5.572	-0.394	1.000
<i>P. damicornis</i>	<i>Porites</i> spp.	12.567	14.816	19.539	5.572	-1.754	0.327

Table 2: Comparison of endolithic community diversity indices between *Porites* spp., *Seriatopora hystrix* and *Pocillopora damicornis* corals. Chao1 and Number of observed OTUs were compared with a two-sample t-test, *p*-values ≤ 0.05 indicate significant differences between host species and are marked with an asterisk. Std = standard deviations.

1 - The *tufA* dataset has only one *P. damicornis* sample (after rarefaction), therefore the significance cannot be calculated. See Supplementary table S1 for number of samples.






Figure 1: Relative abundances of the most common microbial taxa in coral skeletons of *Porites* spp. from high $p\text{CO}_2$ and reference sites.

Results were based on all samples from each site, averaging the relative abundances at the taxonomic level displayed in the legend.

A) biodiversity survey targeting prokaryotes based on the 16S rRNA gene; B) survey of the eukaryotic green algal members of the microbiome based on the *tufA* marker; C) biodiversity survey using the Universal Plastid Amplicon.




Figure 2: Principal Coordinate Analysis of microbial communities present in limestone skeletons of *Porites* spp. from high $p\text{CO}_2$ and reference sites. The analyses were based on weighted UniFrac distance matrices calculated with OTU-level abundances for each metabarcoding marker: A) prokaryotic 16S rRNA gene marker; B) eukaryotic green algae *tufA* marker; C) Universal Plastid Amplicon marker. The results of the MGLM analysis (Deviance and P value) are shown.






Figure 3: Relative abundances of the most common microbial taxa in coral skeletons of *Pocillopora damicornis*, *Seriatopora hystrix* and *Porites* spp. Results were based on all samples from each coral species, averaging the relative abundances at the taxonomic level displayed in the legend. A) Biodiversity survey targeting prokaryotes based on the 16S rRNA gene; B) biodiversity survey targeting eukaryotic members of the microbiome based on the *tufA* marker; C) biodiversity survey using the Universal Plastid Amplicon




Figure 4. Principal Coordinate Analysis of microbial communities present in limestone skeletons of three coral host species collected in high pCO_2 and reference sites. The analyses were based on weighted UniFrac distance matrices calculated with OTU-level abundances for each metabarcoding marker: A) prokaryotic 16S rRNA gene marker; B) eukaryotic green algae *tufA* marker; C) Universal Plastid Amplicon marker. The results of the MGLM analysis (Deviance and P value) are shown.