



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Hauser, CE;Southwell, D;Lahoz-Monfort, JJ;Rumpff, L;Benshemesh, J;Burnard, T;van Hesperen, R;Wright, J;Wintle, B;Bode, M

Title:

Adaptive management informs conservation and monitoring of Australia's threatened malleefowl

Date:

2019-05-01

Citation:

Hauser, C. E., Southwell, D., Lahoz-Monfort, J. J., Rumpff, L., Benshemesh, J., Burnard, T., van Hesperen, R., Wright, J., Wintle, B. & Bode, M. (2019). Adaptive management informs conservation and monitoring of Australia's threatened malleefowl. *Biological Conservation*, 233, pp.31-40. <https://doi.org/10.1016/j.biocon.2019.02.015>.

Persistent Link:

<https://hdl.handle.net/11343/348173>

# Adaptive management informs conservation and monitoring of Australia's threatened malleefowl

Cindy E. Hauser<sup>1\*</sup>, Darren Southwell<sup>1</sup>, José J. Lahoz-Monfort<sup>1</sup>, Libby Rumpff<sup>1</sup>, Joe Benshemesh<sup>2,3</sup>, Tim Burnard<sup>4</sup>, Rosanna van Hesp<sup>1</sup>, John Wright<sup>5</sup>, Brendan Wintle<sup>1</sup>, & Michael Bode<sup>1,6</sup>

<sup>1</sup>School of BioSciences, University of Melbourne, Parkville Vic 3010 Australia

<sup>2</sup>Victorian Malleefowl Recovery Group, Vic Australia

<sup>3</sup>La Trobe University, Bundoora Vic Australia

<sup>4</sup>National Malleefowl Recovery Team, Vic Australia

<sup>5</sup>Parks Victoria, Melbourne, Vic Australia

<sup>6</sup>School of Mathematical Sciences, Queensland University of Technology, Brisbane, Qld Australia

\*corresponding author. Email: [chauser@unimelb.edu.au](mailto:chauser@unimelb.edu.au). Phone: +61-3-8344 3201. Postal address as listed above.

## Abstract

Monitoring is an essential component of adaptive management, and a carefully designed program is needed to ensure high-quality data and inferences over realistic time scales. Co-operation among agencies and incorporating citizen science may help enhance learning whilst reducing the financial costs of monitoring. We seek to realise this potential while conserving the Australian malleefowl (*Leipoa ocellata*). An established network of citizen scientists provide low-cost, sustainable annual monitoring data, yet the most effective actions for conserving malleefowl remain highly uncertain. The continent-wide species' distribution presents significant challenges, including multiple environmental strata to sample and numerous management jurisdictions. We outline an adaptive management framework that aims to unify malleefowl conservation priorities nationally, and target monitoring efforts. We elicited a model structure for the drivers of, and threats to, malleefowl persistence in a workshop with land managers and advocates. We parameterised 80 uncertain interactions within this structure using novel ensemble modelling techniques and identified the effectiveness of predator control as a critical uncertainty affecting malleefowl persistence. We developed a classical, spatially replicated experimental design to test whether malleefowl breed more frequently where predators are suppressed. The proposed monitoring design will rely on the contributions of several dozen land managers and 200-300 citizen scientists annually. We have developed a broad stakeholder base, a proactive communication strategy, and an agile approach to accessing resources to foster resilience and longevity in the monitoring program. If malleefowl conservation successfully adapts in response to monitoring outcomes, it will become one of the largest adaptive management programs on the planet.

**Keywords:** citizen science, *Leipoa ocellata*; feral predator; structured decision making; stakeholder engagement; ensemble modelling; statistical power

## 41 **1 Introduction**

42 A key challenge within the science of conservation biology is assessing the relative importance of the  
43 various threats to species and ecosystems, and consequently identifying the actions that will most  
44 effectively conserve those species and ecosystems. Approaches such as structured decision-making  
45 (Gregory et al. 2012) and management strategy evaluation (Bunnefeld et al. 2011) offer a logical way  
46 of deconstructing and assessing this problem to arrive at a preferred course of action. While these  
47 approaches can accommodate a range of knowledge formats, such as field measurements, system  
48 models, stakeholder preferences and expert opinion, it is also prudent to specify what inevitably  
49 remains uncertain. Adaptive management is a specific form of structured decision-making that  
50 allows the most effective actions to be identified in the presence of uncertainty (Walters 1986, Lyons  
51 et al. 2008).

52 Adaptive management was initially applied within the context of natural resource management  
53 (Walters 1986), where uncertainty in population growth and carrying capacity was accounted for in  
54 the management of fisheries (e.g. Smith & Walters 1981). Waterfowl hunting across North America  
55 has been guided by adaptive management for three decades (Johnson et al. 1997), in the face of  
56 uncertainty around individual survival rates and the strength of density dependence in recruitment.  
57 Applications of adaptive management to conservation have taken longer to develop (Runge 2011).  
58 These have included predator control (Innes et al. 1999, Parkes et al. 2006, Whitehead et al. 2008),  
59 weed suppression (Gannon et al. 2013), habitat management (Aldridge et al. 2004, Nicol et al. 2015),  
60 fire management (Moore et al. 2011), and reintroductions (Armstrong et al. 2007).

61 Embarking on adaptive management is particularly relevant if uncertainty is present, if learning (i.e.  
62 reducing uncertainty) is feasible on a management-relevant time-scale, and where there are  
63 opportunities to adjust management as a response to learning (Doremus 2010). Quantitative  
64 methods for planning adaptive management are well developed (Chadès et al. 2017). These  
65 methods require the specification of management objectives and performance measures,  
66 characterisation of uncertainty as multiple alternative models of how actions could influence  
67 performance, and a monitoring program design (Lyons et al. 2008). Past adaptive management  
68 programs have typically identified a small set of uncertainties (often just one or two) that crucially  
69 affect the optimal choice of action (e.g. Walters & Hilborn 1976, Johnson et al. 1997, although see  
70 Armstrong et al. 2007 for an exception). Value-of-information analysis can be used to evaluate the  
71 benefits of resolving uncertainty in terms of the management objective (Walters 1986). Quantitative  
72 analysis can guide how actions should be adjusted as a response to learning (often using control  
73 theory and other optimisation approaches; Walters 1986), although socio-political factors are also  
74 important (Walters 2007).

75 The broad philosophy of adaptive management is participatory and collaborative, extending to the  
76 person(s) with the authority to make decisions and take actions and the many other stakeholders  
77 concerned with the consequences of those actions (Allen & Gunderson 2011, Hopkinson et al. 2017).  
78 It is important that their range of perspectives are represented in the management objectives  
79 (Susskind et al. 2012). Stakeholders, researchers and other system experts may propose a diverse  
80 range of actions and system models for consideration. Leaders or champions, facilitators,  
81 researchers and system experts without a stake in management decisions are needed to direct the  
82 process and ensure all voices are heard (Allen & Gunderson 2011, Susskind et al. 2012). Similarly,  
83 'bridging organisations' can be crucial for facilitating communication and holding participants  
84 accountable during planning, action and monitoring (Allen et al. 2011, Hopkinson et al. 2017).  
85 Engaging and sustaining such a diverse team is difficult, and a common barrier to successful adaptive  
86 management (Allen & Gunderson 2011, Susskind et al. 2012)

87 Monitoring provides essential feedback within adaptive management (Lyons et al. 2008), and a lack  
88 of sustained funding for monitoring is a widespread barrier to programs' success (Walters 2007,  
89 Westgate et al. 2013). Recruiting citizen scientists has been proposed as a low-cost and participatory  
90 monitoring approach, and it requires commitment from program managers to engage and motivate  
91 participants (Aceves-Bueno et al. 2015). Citizen science programs must be rigorous and carefully  
92 targeted to meet the requirements of adaptive management.

93 In this study, we outline our approach to designing a large-scale adaptive management program for  
94 the threatened malleefowl (*Leipoa ocellata*). Actions that influence malleefowl persistence are taken  
95 by a range of land managers including government agencies, mining companies, Traditional Owners,  
96 farmers and other private landholders and leaseholders. Furthermore, the various levels of  
97 government have varying authority over different tenures and legislation designed to protect species  
98 and habitats (see Appendix A for a summary). A National Malleefowl Recovery Team (henceforth the  
99 'recovery team') is responsible for providing recommendations to these land managers. Their task is  
100 hampered by significant uncertainty about which threats and conservation actions most strongly  
101 influence malleefowl persistence (Benshemesh et al. 2007, Bode & Brennan 2011, Walsh et al.  
102 2012). The recovery team therefore called for a unified adaptive management framework that  
103 integrates the interests and capabilities of varying stakeholders across jurisdictions, and prioritises  
104 conservation actions in the presence of uncertainty. The recovery team already supported co-  
105 ordinated citizen science monitoring of malleefowl breeding mounds as a potential foundation for  
106 program monitoring (Benshemesh et al. 2018). A key challenge to overcome was that prominent  
107 methods within adaptive management were not suitable for eliciting and analysing the multiple  
108 threats and uncertainties present in the mallee system.

109 In this paper, we describe the extension and adjustment of adaptive management practices to  
110 develop a unified malleefowl conservation plan. In collaboration with the recovery team, we acted  
111 as a 'bridging organisation' that engaged and co-ordinated regular communication among land  
112 managers, technical experts, and the existing citizen science program. We introduced ensemble  
113 modelling as a technique for representing and analysing multi-dimensional uncertainty. We  
114 addressed the challenge of sustaining monitoring by building on the existing citizen science program.  
115 The effectiveness of our proposed monitoring design was evaluated using a general power analysis  
116 structure that was tailored to management priorities.

## 117 **2 An adaptive management framework for malleefowl conservation**

### 118 **2.1 Management context**

119 The malleefowl is an iconic megapode that spans the Australian continent (3.3 million km<sup>2</sup>, BirdLife  
120 International 2010; Figure 1). It is nationally listed as vulnerable, and endangered in the eastern  
121 states of Australia. Malleefowl have experienced substantial range contraction from their pre-  
122 European distribution, following large-scale habitat clearance for agriculture (Benshemesh 2007).  
123 Although clearance has slowed considerably in recent decades, it is uncertain if populations continue  
124 to decline (Benshemesh et al. 2007). In addition to historic habitat clearance, malleefowl persistence  
125 may be threatened by predation from introduced and native species and changing fire regimes  
126 (Benshemesh 2007). There is significant uncertainty about which threats and conservation actions  
127 most strongly influence malleefowl persistence (Benshemesh et al. 2007, Bode & Brennan 2011,  
128 Walsh et al. 2012).

129 An extensive malleefowl mound monitoring program has developed over the past three decades,  
130 growing from a handful of sites in the early 1990s to more than 150 sites in 2016 (Figure 1b). Initially  
131 monitoring was often performed by government agencies and consultants. Recruitment of citizen

132 scientists occurred gradually in most areas as locals interested in the species were casually invited to  
133 participate in monitoring, and in some cases undertook monitoring of their local site. As word of the  
134 program spread through volunteer and friendship networks, the citizen scientist component grew  
135 and the program was adapted accordingly to ensure data accuracy. As funding became scarce,  
136 citizen scientists increasingly played a pivotal role in the monitoring program. Their involvement has  
137 led to substantial cost savings and reduced the program's reliance on continuous funding  
138 (Benshemesh et al. 2018).

139 Historic malleefowl monitoring has focused on annual observations of nesting activity at known  
140 breeding mounds, because this is generally regarded as the most feasible and accurate method of  
141 measuring trends in the species' conservation trajectory. Adult malleefowl breed most years except  
142 following winter drought (Booth & Seymour 1984). Current and past malleefowl breeding mounds  
143 are identified via ground searches or remote sensing techniques, such as photogrammetry  
144 (Thompson et al. 2015) and LiDAR (Saffer & Peake 2014). Malleefowl pairs occasionally abandon old  
145 mounds to create new ones, and so the recovery team recommends that these searches are  
146 repeated every 10 years to identify new mounds. Known mounds are visited annually by citizen  
147 scientists to observe signs of breeding and predator activity (evidenced by tracks or scats).

148 More recently, motion-triggered cameras have also been piloted as a means of monitoring  
149 malleefowl, their predators, and competitors (Benshemesh 2013). Citizen scientists retrieved data  
150 cards from camera traps and sorted through the resulting tens of thousands of photographs.  
151 Benshemesh (2013) found that volunteers were readily available, accurate, timely and enthusiastic  
152 for these tasks.

153 While the recovery team had already developed recommendations and established monitoring  
154 practices with the co-operation of diverse land managers and citizen scientists (Benshemesh et al.  
155 2018), there was a need to prioritise multiple threats and conservation actions to effectively protect  
156 malleefowl populations. In the project outlined in this paper, we addressed most stages of the  
157 adaptive management cycle (Figure 2) and established a plan for co-ordinated malleefowl  
158 conservation among land managers. First, we engaged malleefowl advocates and land managers in  
159 framing the conservation problem (2.2). Using their conceptual models of cause and effect, we  
160 characterised and prioritised uncertainty across multiple dimensions (2.3). We designed a  
161 monitoring program targeting the highest priority uncertainty, taking advantage of spatial  
162 replication (2.4). Conservation action and monitoring rely on the co-operation of many land  
163 managers, so we explicitly addressed the challenges of leadership and long-term engagement,  
164 communication, and sustained funding (2.5).

## 165 *2.2 Problem framing*

166 We convened a workshop and used structured decision-making (Gregory et al. 2012) to frame the  
167 malleefowl conservation challenge, understand the current state of knowledge, and identify relevant  
168 issues for future prioritisation. The 24 attendees included: community advocates and citizen  
169 scientists; state government staff responsible for research integration, for environmental  
170 management planning and for park management; federal government staff responsible for  
171 threatened species support; and university researchers with expertise in structured decision-making,  
172 adaptive management, and/or mallee ecosystems.

173 There was broad agreement among workshop participants that their fundamental objective was to  
174 foster the long-term persistence of a self-sustaining malleefowl population. The spatial extent of the  
175 population was left unspecified, acknowledging that there has been substantial historic range  
176 contraction and there may be future range shifts as a response to climate change. Three indicators

177 were nominated as the quantitative means for assessing this fundamental objective: adult  
178 abundance, juvenile abundance, and population occupancy/range. Given the population fluctuations  
179 that malleefowl typically exhibit as a response to fluctuating resources, participants noted that high  
180 values for these indicators need not be achieved every year in order to fulfil the fundamental  
181 objective.

182 In smaller independent teams, participants developed lists and cause-effect diagrams of the drivers  
183 and threats that influence the three indicators. These, in combination with potential management  
184 actions, formed the foundation of our system models. While numerous influences were collated,  
185 four were broadly agreed-upon as drivers of malleefowl population dynamics: predation, rainfall,  
186 grazing and fire. Predation by foxes, cats, dingoes, dogs, raptors and goannas were predicted to  
187 directly reduce adult and juvenile malleefowl abundance. Rainfall, grazing and fire potentially  
188 interact to affect habitat structure, which provides food, nesting materials, and cover from  
189 predators. Herbivores (rabbits, goats, sheep and kangaroos) may compete directly with malleefowl  
190 for food, and also deplete the understorey vegetation that is thought to protect malleefowl from  
191 predation and provide nesting materials. Fire was expected to cause direct mortality and  
192 dramatically alter habitat structure, reducing food availability and exposing malleefowl to predators.  
193 Malleefowl have rarely been observed to breed in recently (< 15 years) burned areas and thus  
194 broad-scale fires could affect the population's range.

195 Actions (which can be grouped into strategies or alternatives; Figure 2) were proposed to address  
196 threats to malleefowl persistence, including reduction of grazing pressure (e.g. by closing water  
197 points or erecting fencing strategically), reducing the density of grazers and predators, strategic fire  
198 management, influencing land change and protection, malleefowl translocation, habitat  
199 revegetation, supplementary feeding, and erecting road signs. Participants contributed knowledge  
200 and located documents that could help characterise the cause-and-effect relationships that affect  
201 malleefowl persistence. The relative importance of the identified threats, and our capacity to  
202 address them, was not resolved in the literature and so there was substantial uncertainty to  
203 characterise through an adaptive management approach.

### 204 *2.3 Characterising and prioritising uncertainties*

205 The initial workshop (2.2) revealed a complex array of interacting environmental and biotic factors  
206 and potential conservation actions influencing malleefowl persistence. We focused on one  
207 participant group's causal ecosystem model (Figure 3a), which included 80 interactions between 14  
208 ecosystem components (Bode et al. 2017). While each interaction was captured qualitatively, it was  
209 not feasible to specify the interaction strengths quantitatively. Faced with a model with this  
210 complexity: (1) participants were unlikely to know about all possible interactive responses, (2)  
211 representing uncertainty and differing opinions among stakeholders was difficult, and (3) eliciting all  
212 interactions would have placed an excessive burden on participants.

213 The quantitative strength of each interaction in the ecosystem, i.e. 80 parameters, formed the set of  
214 uncertainties for the adaptive management problem. We used a single ecosystem model structure,  
215 prescribing an equation for each of the 14 ecosystem components (e.g. malleefowl density, fox  
216 density, rainfall quantity, see Figure 3a for full list) that included density dependence and responses  
217 to every other model component via an interaction parameter (Appendix B). Assuming that we knew  
218 only the direction of the interaction (positive, negative or zero; indicated in Figure 3a) from  
219 workshop participants, we generated  $10^9$  different plausible parameterisations of the ecosystem  
220 model (Bode et al. 2017). This finite, discrete ensemble of model parameterisations formed a more

221 tractable expression of uncertainty than a set of 80 interaction parameters with continuous  
222 probability distributions (Chadès et al. 2017).

223 We subsequently refined the set of plausible parameterisations by excluding models that  
224 contradicted what was known about and had been observed in the system. First, if the long-term  
225 equilibrium of a given parameterisation included the extinction of any ecosystem components, it  
226 was removed from the set, given that all model components are currently observed in the system.  
227 Second, we engaged malleefowl experts via a second workshop and elicited more detailed,  
228 quantitative predictions of how the ecosystem would respond to perturbations in each component.  
229 Participants described the range of possible component trajectories over 5 years following a  
230 perturbation, e.g. changes in fox population density as a response to 3 years in which herbivores  
231 occurred at double their long-term average density. We selected the 5% ( $\sim 10^5$ ) of the remaining  
232 ecosystem parameterisations that aligned most consistently with participants' responses (Bode et al.  
233 2017).

234 Using this reduced ensemble capturing plausible ecosystem dynamics, we predicted the  
235 consequences of candidate management actions (Recovery plan interventions; Benshemesh 2007)  
236 addressing six threats: disease and inbreeding; fire intensity and severity; predation by introduced  
237 cats; predation by introduced foxes; competition by herbivores; and habitat loss and fragmentation  
238 (see Appendix B for more details). For each action and each parameterisation in the ensemble, we  
239 simulated all components for 5 years and recorded the final malleefowl abundance. We also  
240 simulated each parameterisation in the absence of management action, and thus calculated the  
241 predicted change in malleefowl abundance arising from each management action. Five of the six  
242 actions had the potential to positively or negatively affect malleefowl abundance; addressing disease  
243 and inbreeding was very likely to positively affect malleefowl abundance (Figure 3b). Actions  
244 addressing fox and cat predation included possibilities for the largest malleefowl population  
245 increases, while addressing fox predation and fire intensity included possibilities for the most rapid  
246 population declines. Other studies have similarly highlighted uncertainty surrounding the roles of fox  
247 predation and control in malleefowl population dynamics (Benshemesh et al. 2007, Bode & Brennan  
248 2011, Walshe et al. 2012). Thus, understanding fox predation and control emerged as a top research  
249 priority, with cat predation and fire management also warranting consideration.

#### 250 *2.4 Monitoring design*

251 As we reviewed and expanded the malleefowl monitoring program in this project, we focused on our  
252 capacity to learn more about and act to address the threats of fox and cat predation. Learning about  
253 the effect of predator control on malleefowl persistence could be accelerated further by spatially  
254 replicating management across the species' range (Walters 1986). We investigated a control-impact  
255 design by identifying clusters of unmanaged ('control') and predator-managed ('impact') sites with  
256 similar characteristics, such as habitat quality and rainfall patterns. Each site within the cluster was  
257 expected to experience similar temporal fluctuations in environmental conditions, thus constraining  
258 random spatial variation, and enhancing our ability to estimate the effect of predator management.

259 Even with the advantage of spatial replication, multiple years of data are required from each cluster  
260 to effectively distinguish local population dynamics from any global responses to predator  
261 management. The monitoring program does not have consistent long-term funding and staffing to  
262 guarantee that uncertainty will be resolved by a control-impact experiment (Hopkinson et al. 2017),  
263 so it was important that the benefits of implementing adaptive management were assessed over  
264 realistic time frames and resource constraints (Walters 2007, Doremus 2010, Allen et al. 2011). We

265 developed a statistical power analysis to evaluate the capacity of the malleefowl monitoring  
266 program to resolve uncertainty around the benefits of fox (and potentially feral cat) management.

#### 267 *2.4.1 Statistical power analysis for a predator control experiment*

268 The power analysis estimated the probability that a spatially-replicated control-impact experiment  
269 could detect a pre-set range of improvements in malleefowl mound activity arising from predator  
270 management (Figure 4). The analysis used historic mound activity data to estimate temporal and  
271 spatial fluctuations in the number of active mounds at monitoring sites, and simulated data  
272 collection from multiple control-impact two-site clusters occupied by malleefowl (see Appendix C).  
273 An experiment of five years duration comprising 36 or more sites (in 18 two-site clusters) had an  
274 89% probability of detecting a 22% increase in mound activity. Mound activity is expected to be  
275 proportional to adult malleefowl abundance (and mediated by rainfall). Thus, the ensemble  
276 modelling suggests effect sizes of 0-100% are possible (i.e. no effect up to a doubling of malleefowl  
277 abundance; Figure 3b). Population viability analysis has previously suggested that smaller effect sizes  
278 of 0-35% are plausible (Bode & Brennan 2011). After reviewing the power analysis, the recovery  
279 team aspired to engage 40 sites for control-impact monitoring over five years.

280 We are now in the process of identifying suitable control and impact sites for targeted monitoring.  
281 Land managers are advising us of which neighbouring malleefowl sites have comparable habitat and  
282 contrasting predator management. In some cases the paired control-impact design will be extended  
283 to a cluster of three or more sites grouped together and assigned the same spatial random effect  
284 (Appendix C). While grouping near neighbours strengthens our inference in the face of spatial  
285 variation, it conflicts with our need to ensure that the malleefowl populations respond to the action  
286 implemented in their site and not actions in neighbouring sites (e.g., through dispersal of malleefowl  
287 or predators). Thus, we will set buffers of predator control around monitored impact sites, and  
288 minimum distances between control and impact sites.

289 A five-year control-impact experiment might not detect an effect of predator control on malleefowl  
290 mound activity, even if it exists. This could occur if mound activity were below the nominated  
291 average across most sites, if the effect of predator control were weaker than estimated, or if the  
292 spatial or temporal variation were larger than estimated. Finally, the nature of the power analysis is  
293 such that, even under ideal conditions, we accept some probability of failing to detect a significant  
294 effect of management even though it is present (i.e. type II error). This probability is estimated to be  
295 less than 0.1% if malleefowl mound activity doubles in the presence of predator control (effect size  
296 101%, Figure 3), but is as high as 76% if predator control only induces a 10% increase in mound  
297 activity.

298 Alternatively, the implemented predator control may not increase malleefowl mound activity. This  
299 could arise if: (a) the implemented predator control does not successfully reduce predator activity,  
300 or (b) reduced predator activity does not increase malleefowl mound activity. Distinguishing  
301 between these cases would be beneficial for adaptive management. If the implemented actions do  
302 not reduce predator activity, then alternative predator control methods may still be worth exploring  
303 for the potential benefit of malleefowl and other native prey. However, if malleefowl breeding  
304 activity is found not to improve under reduced predator activity, then predator control can be  
305 abandoned as a malleefowl management action. We proposed supplementary monitoring at the  
306 experimental sites to test whether hypothesis (a) is true.

#### 307 *2.4.2 Supplementary monitoring of predator activity*

308 We aimed to distinguish whether predator management at 'impact' sites reduces predator activity  
309 below the predator activity observed at predator-unmanaged 'control' sites. We proposed installing

310 an array of solar-powered camera traps at the sites included in the predator control experiment (van  
311 Hespen et al. in press) to estimate predator activity as a count of fox or cat photos from each  
312 camera. We performed a second power analysis to estimate the number of camera traps that would  
313 be required at each site to distinguish control-impact differences in predator activity from random  
314 variation.

315 To parameterise our analysis, we used data from 16 camera traps collected by Benshemesh (2013)  
316 at Wandown Nature Reserve, a 20 km<sup>2</sup> patch of remnant mallee vegetation not subject to fox  
317 control and surrounded by agricultural land. From these data we estimated spatial and temporal  
318 variation in fox photo counts. The model structure closely resembled the power analysis of mound  
319 activity (Appendix C), although the spatial and temporal scales had higher resolution (in terms of  
320 between-camera variation and monthly variation, respectively). The power analysis estimated the  
321 difference in mean photographic trapping rate between a 'control' and an 'impact' site, and we  
322 deemed the difference to be statistically significant at the customary  $\alpha = 0.05$  significance level (van  
323 Hespen et al. in press). If predator control efforts reduce fox activity by 75% from an unmanaged  
324 density of 2 foxes/km<sup>2</sup>, we estimated that 6 cameras per site would be sufficient to achieve a 96%  
325 probability of detecting the difference in fox densities over 12 months (Figure 5a).

326 Stakeholders queried whether more expensive cameras with broader detection zones would  
327 generate higher quality data to support this experiment. The power analysis indicated that it is more  
328 important to capture the high estimated spatial variation in photo counts than to increase the  
329 number of fox detections from each camera. Consequently, a fixed budget would be more  
330 effectively spent on a larger number of cheaper cameras with reduced detection zones (Figure 4b).

### 331 *2.5 Strategies for collaboration and commitment*

332 Effective monitoring and conservation of malleefowl relies on continued co-operation among  
333 stakeholders and land managers. We evaluate our co-operative approach using Aceves-Bueno et  
334 al.'s (2015) three stakeholder criteria for using citizen science within adaptive management. They  
335 propose that: (1) stakeholders must be identified and engaged, (2) managers must provide  
336 appropriate motivation and incentives to participants, and (3) decision-makers must be accountable  
337 to stakeholders.

338 First, we have identified and engaged community stakeholders. The recovery team communicates,  
339 shares knowledge and trains volunteers continuously. Researchers join them annually to conduct  
340 workshops for participants and potential participants in the predator control experiment. The first of  
341 these workshops focused on a project design that was feasible and would allow for meaningful  
342 inference, and guided the monitoring design outlined in this article. Subsequent workshops have  
343 been used to maintain the momentum of the project, by assessing progress, sharing knowledge and  
344 solving logistical challenges collaboratively (Allen et al. 2011). As sites submit data in the future,  
345 workshops will expand to presentation of analyses, reflecting on what has been learned, and  
346 potentially developing adjustments to management and monitoring protocols.

347 Second, there are incentives and motivation for participation. For land managers, there is an  
348 opportunity to adopt a rigorous national standard of data collection. In the medium term, managers  
349 will receive trend analyses that exceed their local capacity and in the long term, they expect to learn  
350 more about the role of predator control in malleefowl conservation, drawn from the co-operation of  
351 many participants. The value of this information should not be underestimated. Many land  
352 managers have expressed frustration at not having resources to test the efficacy of their predator  
353 control programs. Furthermore, they often have other external motivations to monitor and report

354 their predator control and malleefowl conservation activities (e.g. state directives, reports to NGO  
355 sponsors); we worked with managers to develop protocols that align with these other motivators.

356 Citizen scientists' motivations may be less tangible, primarily focused on individuals' connections to  
357 nature and fellow malleefowl enthusiasts. They may rely on leadership from one of numerous  
358 community 'champions' (Garnett et al. 2018). Malleefowl are much loved in the regional  
359 communities where they persist, perhaps because their maintenance of 1m tall nest mounds has  
360 earned them a reputation as one of the hardest working birds in the world (Department of the  
361 Environment and Energy 2006). The program offers flexibility in involvement at several levels: the  
362 difficult physical work of monitoring in remote locations with harsh climates (earning the catch  
363 phrase 'bushwalking with a purpose'), organising equipment and volunteers, managing and verifying  
364 data, and classifying the images captured by camera traps. Ensuring ongoing citizen science  
365 involvement is a major challenge, and it is important that citizen scientists see that their  
366 contributions are valued and influential. To this end, the recovery team engages volunteers by co-  
367 hosting community training days (up to 11 sessions across regional Australia in one year, attracting  
368 over 250 participants); providing online access to the national database; disseminating newsletters  
369 covering news, personal stories and research findings from the program; and holding a National  
370 Malleefowl Forum every 3-4 years.

371 The third stakeholder criterion is that decision-makers must be accountable to stakeholders (Aceves-  
372 Bueno et al. 2015). The network of agencies with the authority to decide on malleefowl conservation  
373 actions is diffuse. Within the predator control experiment, none of them have been called on to  
374 change their actions. Malleefowl monitoring is within the remit of participants, and we aimed to  
375 build a culture of accountability within the annual workshops. In this way the recovery team has  
376 acted as a 'bridging organisation', despite its lack of regulatory authority or funding to incentivise  
377 particular management actions (Allen et al. 2011).

378 Many other studies cite a need for strong leadership or a 'champion' to drive successful adaptive  
379 management (Gregory et al. 2006, Allen & Gunderson. 2011, Walters 2007). The malleefowl project  
380 has these leaders. First, there is a leading malleefowl ecological expert with a long-term  
381 commitment to the species' conservation and excellent communication skills. Second, the recovery  
382 team created paid National Co-ordinator and Engagement Officer roles with explicit responsibilities  
383 for progressing adaptive management and monitoring. The co-ordinator and engagement officer are  
384 facilitators: organised, enthusiastic and persistent, and not expected to drive the scientific research  
385 design (Walters 2007). These leaders are supported by a project team with a shared vision and  
386 purpose, initially negotiated and articulated in a joint co-funding proposal, with varied and  
387 complementary skills and knowledge to contribute.

388 Sustaining funding is a common challenge in adaptive management (Allen et al. 2011, Walters 2007)  
389 that applies to the malleefowl program. Characteristic of long-term research projects, the recovery  
390 team has drawn funding from a variety of sources. The initial research plan was launched via an  
391 Australian Research Council Linkage grant with supplementary funding from one state government  
392 agency and one environmental offsetting program, mediated via a citizen scientists' group.  
393 Researcher salaries and annual workshop expenses have been secured through universities, large  
394 environmental management, and threatened species research centres. The recovery team has  
395 financially supported the stakeholder co-ordinators and database managers. Members of the  
396 recovery team have facilitated small grant applications to NGOs seeking the equipment and training  
397 needed at individual sites participating in the predator control experiment. Future data collection  
398 relies on the in-kind support of numerous professional agencies and community groups. This wide-  
399 ranging strategy demonstrates that the recovery team can be proactive and flexible in its resource

400 acquisition. While this can be daunting, we hope that the approach will prove robust, freeing the  
401 program from reliance on the long-term financial commitment of any one agency.

### 402 **3 Discussion**

403 We have established the minimum conditions needed to justify adaptive management for  
404 malleefowl conservation: uncertainty is present, learning (i.e. reducing uncertainty) is feasible on a  
405 management-relevant time-scale, and there are opportunities to adjust management as a response  
406 to learning (Doremus 2010). We have developed an adaptive management framework and  
407 communication strategy that facilitate and formalise collaboration among the numerous agencies  
408 and community groups concerned with malleefowl conservation. The adaptive management  
409 framework has steered their monitoring efforts towards a better understanding of the uncertain  
410 effects of predator control. Nevertheless, the ensemble model structure and regularly scheduled  
411 workshops offer some flexibility for future management choices and learning priorities. If malleefowl  
412 conservation successfully adapts in response to the new monitoring data we expect to collect, it will  
413 become one of the largest adaptive management programs on the planet, both in terms of  
414 geographic extent and the number of jurisdictions and citizen scientists involved.

415 Our project was well aligned with more general frameworks for adaptive management in  
416 conservation (Runge 2011; Figure 2). We posed and structured the malleefowl conservation  
417 challenge clearly in collaboration with experts and stakeholders (2.2). Our approach to evaluating  
418 consequences and uncertainties was new to the adaptive management context (2.3), and we  
419 designed a rigorous, feasible monitoring program (2.4). However, some components of the  
420 framework were less thoroughly addressed, and we discuss these here.

#### 421 *3.1 Evaluating consequences and critical uncertainties to identify the preferred alternative*

422 We introduced ensemble modelling into the adaptive management space, as a means of  
423 representing and prioritising a large number of uncertainties (i.e. 80 interactions between mallee  
424 system components; Figure 3a). The consequences of six candidate management actions (Appendix  
425 A) were predicted for malleefowl abundance, including all associated uncertainty. However, we did  
426 not construct an objective function or use any optimisation methods to derive a preferred  
427 conservation action. We identified critical uncertainties informally by inspecting the range of  
428 plausible malleefowl responses to each candidate action, and prioritising actions with broad  
429 uncertainty and high possible benefits. A more formal value-of-information analysis using an  
430 objective function could estimate the expected benefits of resolving uncertainty. However, there  
431 were barriers to using these more traditional analyses. Any combination of the six modelled  
432 conservation actions could theoretically be pursued simultaneously, but it was unclear which  
433 combinations of agencies were responsible for implementing actions and not all agencies could  
434 afford all possible action combinations (as it was, some could only be involved as control sites). We  
435 focused instead on how targeted monitoring could help resolve the highest priority uncertainty  
436 identified by the literature and the ensemble modelling: the effect of fox and cat predation on  
437 malleefowl persistence.

#### 438 *3.2 Implementing action*

439 We have collaborated intensively with land managers to identify suitable control and impact sites for  
440 the project. Under the proposed experimental design, participants continue to implement their  
441 current predator control action. Predator and malleefowl populations are expected to already occur  
442 at their equilibrium densities with respect to their assigned management action. The experiment  
443 therefore poses no extra risk of negative outcomes than status quo management. However, stronger  
444 statistical inference would arise from random allocation of actions to sites (e.g. Lyons et al. 2008). As

445 it stands, there may be systematic reasons why predator control has been selected by managers *a*  
446 *priori* for some sites and not others. These reasons are confounded with malleefowl's response to  
447 predator control in the proposed design.

### 448 *3.3 Monitoring*

449 Our experimental monitoring design calls on numerous agencies to standardise and share their  
450 monitoring data, much of it collected by citizen scientists. The proposed program sets a foundation  
451 for fulfilling the four monitoring criteria set by Aceves-Bueno et al. (2015) to successfully utilise  
452 citizen science within adaptive management. First, the historic mound monitoring program and  
453 camera trap pilot program have demonstrated that monitoring can be achieved. Monitoring in the  
454 first predator control experiment clusters is underway, and there are strategies in place to continue  
455 engaging participants (2.5). Second, the monitoring design is relevant and rigorous. Power analyses  
456 of malleefowl mound monitoring and camera trapping of predators have helped us build evidence  
457 that an experiment can generate meaningful inference regarding an important uncertainty over  
458 feasible temporal and spatial scales (40 sites in 20 control-impact pairs/clusters monitored for five  
459 years each; Figure 3) with realistic resourcing (e.g. Figure 4).

460 Third, we have strived for cost-effectiveness, although we have not performed a value-of-  
461 information analysis that could directly weigh the benefits of the experimental design against its  
462 costs. Using remote sensing to locate new mounds and cameras for monitoring predators are  
463 notable financial costs that exceed status quo management. Camera installation was not a  
464 mandatory requirement for participation, and the recovery team has worked with site managers to  
465 secure funding where assistance was needed. The monitoring program has been structured to  
466 'piggy-back' on existing initiatives wherever possible (Westgate et al. 2013), harnessing volunteer  
467 efforts and spreading costs among many participants (Walters 2007).

468 Fourth, monitoring will accommodate multiple temporal and spatial scales. It is not expected that all  
469 site clusters will commence monitoring in the same year. Rather than planning an experiment that  
470 will conclude in five years, we expect staggered involvement of various site clusters, each having a  
471 goal of at least five years' participation. Learning the range-wide malleefowl response to predator  
472 control over five-to-ten years is the overarching goal of the experiment, but data and feedback can  
473 also be examined at finer resolutions. Malleefowl mound and predator activity data will be collated  
474 at each site annually (with predator activity measured as monthly photo counts), and the recovery  
475 team can deliver annual cluster-specific reports.

### 476 *3.4 Updating and learning*

477 If participants collect and submit the proposed data to the recovery team, we anticipate  
478 opportunities to update knowledge and share learning at a number of scales. These range from  
479 refining our definition of predator control, to opportunistically learning about uncertainties beyond  
480 predation, and reprioritising the critical uncertainties.

481 Our power analyses treat predator control as a binary action with a common response across all  
482 managed sites, but this is unlikely to be the case for at least three reasons. First, predator control is  
483 applied differently across Australia, involving baiting at varying frequencies and densities, trapping  
484 and/or fencing. Second, foxes and cats occur at different densities and in different ratios across  
485 malleefowl's range and they respond differently to the same control action. Third, malleefowl  
486 populations may respond differently to the same reduction in predator density, for example  
487 depending on the shelter from predators provided by a site's vegetation structure. All three  
488 phenomena could be assessed through the planned camera trapping to estimate predator (fox or  
489 cat) activity. After data have been collected, the method of predator control may be included as a

490 more nuanced covariate in analyses, and other variables that mediate the response of malleefowl to  
491 predator control could also be included as interaction terms. Thus, the power analysis was designed  
492 to provide only a rough guide to the realised statistical power.

493 An upcoming technical challenge will be updating the ensemble models (Bode et al. 2017) over a  
494 relevant time frame in response to the predator control experiment findings and other malleefowl  
495 monitoring data. Further refinement of the ensemble model set should consider the reliability of the  
496 new data (e.g. via sample size or confidence/credible intervals). The model set represents  
497 uncertainty in 80 parameters with a common model structure, and we expect that this could be  
498 updated numerically using Bayesian methods (Chadès et al. 2017). More research is needed to  
499 evaluate tools for representing this kind of multi-dimensional uncertainty within adaptive  
500 management, including ensemble modelling and Bayesian belief networks (Nyberg et al. 2006,  
501 Rumpff et al. 2011).

502 Regardless of whether reducing fox and cat predation is feasible and beneficial to malleefowl  
503 persistence, fire and grazing management may warrant exploration (Figure 2b). The targeted  
504 monitoring and power analysis design described in this article could be transferred and tailored to  
505 these issues. However, we anticipate slow ecosystem responses and land manager commitments far  
506 beyond the five years nominated in the predator control experiment. Furthermore, making strategic  
507 changes to fire and grazing management may be even more difficult than altering predator control.  
508 Scenario planning may be a preferable alternative to adaptive management in these cases (Allen et  
509 al. 2011).

510 There will be some capacity for opportunistic learning about the role of fire and grazing  
511 management within the current design. Our understanding of the mallee system can potentially be  
512 updated with more than just malleefowl mound activity and predator photo counts. Identifying  
513 other species among the photographs collected can also provide supplementary information,  
514 including malleefowl, predators such as cats and dogs, and potential competitors such as kangaroos,  
515 goats and rabbits. Successful fox control has the potential to influence the abundance of these other  
516 malleefowl predators and competitors (e.g. Figure 3a), and thus indirectly affect malleefowl  
517 persistence. We expect to collect other supplementary data, such as rainfall (which affects food  
518 availability) and fire history (which affects mortality, food availability and cover from predators).  
519 These could all potentially be used as covariates in a model of mound activity (based on the  
520 equations in Appendix C) and also to refine the ensemble model set. However, learning is expected  
521 to be slow and erratic in the absence of a targeted monitoring design.

522 Successful adaptive management makes space for uncertainty, complexity, reflection and critical  
523 discussion (Allen & Curtis 2005, Gregory et al. 2006, Scarlett 2013). Over the five-to-ten years  
524 needed to undertake the predator control experiment, funding, policy, personnel and agency  
525 priorities are all likely to undergo changes. Extreme natural events may occur. It is possible, even  
526 probable, that the data generated in the predator control experiment will include surprises,  
527 generating new doubts and questions. The philosophy of adaptive management welcomes these  
528 events as opportunities for learning, generating new hypotheses and collaboration. An effective  
529 recovery team will consciously create a culture of openness and constructive debate to address  
530 these upheavals, and allow for revisions of the program strategy and design (Allen & Gunderson  
531 2011).

### 532 *3.5 Can adaptive management of malleefowl succeed?*

533 If the uncertainties in malleefowl conservation are effectively resolved, it remains to be seen  
534 whether management can be shifted accordingly. Conservation actions such as fox and cat

535 suppression, strategic fire management, reducing grazing pressure, and reducing land clearing have  
536 implications for other valued native species, ecological communities and economic interests. The  
537 recovery team and surrounding community of malleefowl advocates must engage with broader  
538 processes of decision-making, identifying trade-offs within the malleefowl-specific actions and  
539 among malleefowl and other land use priorities. A rigorous monitoring program will form an  
540 important foundation of evidence as these trade-offs are investigated. By focusing first on the  
541 management actions already taken by participants and the monitoring established with citizen  
542 scientists, the program can build co-operation and credibility, and prepare for future changes.

#### 543 **Acknowledgements**

544 This research project has been funded by an Australian Research Council Linkage grant  
545 (LP120100490) in partnership with Parks Victoria, the Victorian Malleefowl Recovery Group and  
546 Iluka Resources Ltd, and received additional support from the Australian Government (Dept of the  
547 Environment and Energy) through the National Environmental Research Program Environmental  
548 Decisions Hub and the National Environmental Science Program Threatened Species Recovery Hub.  
549 We are grateful for the ongoing support of the National Malleefowl Recovery Team, the knowledge  
550 shared by experts in our elicitation workshops and the enthusiasm of the broader malleefowl  
551 conservation community.

#### 552 **4 References**

- 553 Aceves-Bueno, E., Adeleye, A.S., Bradley, D., Brandt, W.T., Callery, P., Feraud, M., Garner, K.L.,  
554 Gentry, R., Huang, Y., McCullough, I., Pearlman, I., Sutherland, S.A., Wilkinson, W., Yang, Y., Zink, T.,  
555 Anderson, S.E. & Tague, C. (2015) Citizen science as an approach for overcoming insufficient  
556 monitoring and inadequate stakeholder buy-in in adaptive management: criteria and evidence.  
557 *Ecosystems* 18: 493-506.
- 558 Allen, C. & Curtis, A. (2005) Nipped in the bud: why regional scale adaptive management is not  
559 blooming. *Environmental Management* 36: 414-425.
- 560 Allen, C.R., Fontaine, J.J., Pope, K.L. & Garmestani, A.S. (2011) Adaptive management for a turbulent  
561 future. *Journal of Environmental Management* 92: 1339–1345.
- 562 Allen, C.R. & Gunderson, L.H. (2011) Pathology and failure in the design and implementation of  
563 adaptive management. *Journal of Environmental Management* 92: 1379–1384.
- 564 Armstrong, D.P., Castro, I. & Griffiths, R. (2007) Using adaptive management to determine  
565 requirements of re-introduced populations: the case of the New Zealand hihi. *Journal of Applied*  
566 *Ecology* 44, 953-962.
- 567 Benshemesh, J. (2004) Monitoring malleefowl: options, problems and solutions. In *Proceedings of*  
568 *the National Malleefowl Forum 2004*. Victorian Malleefowl Recovery Group, Mildura, Victoria,  
569 Australia, pp 128-134. <http://www.malleefowlvictoria.org.au/2004Forum/Benshemesh2.pdf>
- 570 Benshemesh, J. (2007) National Recovery Plan for Malleefowl. Department for Environment and  
571 Heritage, South Australia. [https://www.environment.gov.au/system/files/resources/dd346674-  
572 08ab-403d-8c11-5b88e8247e8f/files/malleefowl.pdf](https://www.environment.gov.au/system/files/resources/dd346674-08ab-403d-8c11-5b88e8247e8f/files/malleefowl.pdf)
- 573 Benshemesh, J., Barker, R. & MacFarlane, R. (2007) Trend analysis of Malleefowl monitoring data.  
574 Population Assessment and Conservation Action Project, Bundoora, Victoria.  
575 <http://malleefowlvictoria.org.au/documents/nationalProject/milestone1-3.pdf>

576 Benshemesh, J. (2013) Trial of motion sensitive cameras for monitoring a range of animals in  
577 Malleefowl monitoring sites. *Report to VMRG and Malleefowl management committee*, 25pp.  
578 <http://www.nationalmalleefowl.com.au/uploads/file/camera%20trap%20report%202013%20ALL%20w%20appendices.pdf>  
579

580 Benshemesh, J., Southwell, D., Lahoz-Monfort, J.J., Hauser, C., Rumpff, L., Bode, M., Burnard, T.,  
581 Wright, J. & Wintle, B.A. (2018) The national Malleefowl monitoring effort: citizen scientists,  
582 databases and adaptive management. In *Monitoring Threatened Species and Ecological*  
583 *Communities*, eds Legge, Lindenmayer, Robinson, Scheele, Southwell & Wintle. CSIRO Publishing,  
584 Melbourne, pp 387-396.

585 Birdlife International (2010) Species factsheet: *Leipoa ocellata*. Accessed from www.birdlife.org on  
586 16/10/2017, <http://datazone.birdlife.org/species/factsheet/malleefowl-leipoa-ocellata>.

587 Bode, M., Baker, C., Benshemesh, J., Burnard, T., Rumpff, E., Hauser, C.E., Lahoz-Monfort, J.J. &  
588 Wintle, B.A. (2017) Revealing beliefs: using ensemble ecosystem modeling to extrapolate expert  
589 beliefs to novel ecological scenarios. *Methods in Ecology & Evolution* 8: 1012-1021.

590 Bode, M. & Brennan, K.E.C. (2011) Using population viability analysis to guide research and  
591 conservation actions for Australia's threatened malleefowl *Leipoa ocellata*. *Oryx* 45: 513-521.

592 Booth, D.T. & Seymour, R.S. (1984) Effect of adding water to Malleefowl mounds during a drought.  
593 *Emu* 84: 116-118.

594 Bunnefeld, N., Hoshino, E. & Milner-Gulland, E.J. (2011) Management strategy evaluation: a  
595 powerful tool for conservation? *Trends in Ecology and Evolution* 26: 441-447. Chadès I., Nicol S., Rout  
596 T.M., Peron M., Dujardin Y., Pichancourt J.-B., Hastings A. & Hauser C.E. (2017) Optimization  
597 methods to solve adaptive management problems. *Theoretical Ecology* 10: 1-20.

598 Department of the Environment and Energy (2006) Malleefowl (*Leipoa ocellata*) Threatened Species  
599 Day fact sheet. Accessed from [www.environment.gov.au](http://www.environment.gov.au) on 4/12/2018,  
600 <https://www.environment.gov.au/resource/malleefowl-leipoa-ocellata>.

601 Doremus, H. (2010) Adaptive management as an information problem. *North Carolina Law Review*  
602 89: 1455-1498.

603 Gannon, J.J., Shaffer, T.L. & Moore, C.T. (2013) Native prairie adaptive management: a multi region  
604 adaptive approach to invasive plan management on Fish and Wildlife Service owned native prairies.  
605 U.S. Geological Survey, Reston, Virginia. <https://pubs.usgs.gov/of/2013/1279/pdf/of2013-1279.pdf>

606 Garnett, S., Latch, P., Lindenmayer, D. & Woinarski, J.C.Z. (2018) *Recovering Australian Threatened*  
607 *Species: a Book of Hope*. CSIRO Publishing, Clayton Victoria.

608 Gregory, R., Ohlson, D. & Arvai, J. (2006) Deconstructing adaptive management: criteria for  
609 applications to environmental management. *Ecological Applications* 16:2411–25.

610 Gregory, R., Failing, L., Harstone, M., Long, G., McDaniels, R. & Ohlson, D. (2012) *Structured Decision*  
611 *Making: A Practical Guide to Environmental Management Choices*. Wiley-Blackwell, Chichester.

612 Hopkinson, P., Huber, A., Saah, D. & Battles, J.J. (2017) A word to the wise: advice for scientists  
613 engaged in collaborative adaptive management. *Environmental Management* 59: 752-761.

614 Innes, J., Hay, R., Flux, I., Bradfield, P., Speed, H. & Jansen, P. (1999) Successful recovery of Island  
615 kokako *Callaeas cinereal wilsoni* populations, by adaptive management. *Biological Conservation* 87:  
616 201-214.

617 Johnson, F.A., Moore, C.T., Kendall, W.L., Dubovsky, J.A., Caithamer, D.F., Kelley, J.R. & Williams, B.K.  
618 (1997) Uncertainty and the management of mallard harvests. *Journal of Wildlife Management* 61:  
619 202–216.

620 Lyons, J.E., Runge, M.C., Laskowski, H.P. & Kendall, W.L. (2008) Monitoring in the context of  
621 structured decision-making and adaptive management. *Journal of Wildlife Management* 72: 1683–  
622 1692.

623 Nicol, S., Fuller, R.A., Iwamura, T. & Chadès, I. (2015) Adapting environmental management to  
624 uncertain but inevitable change. *Proceedings of the Royal Society B* 282: 20142984.  
625 <http://dx.doi.org/10.1098/rspb.2014.2984>

626 Nyberg, J.B., Marcot, B.G. & Sulyma, R. (2006) Using Bayesian belief networks in adaptive  
627 management. *Canadian Journal of Forest Research* 36: 3104-3116.

628 Parkes, J.P., Robley, A., Forsyth, D.M. & Choquenot, D. (2006) Adaptive management experiments in  
629 vertebrate pest control in New Zealand and Australia. *Wildlife Society Bulletin* 34: 229-236.

630 Rumpff, L., Duncan, D.H., Vesk, P.A., Keith, D.A. & Wintle, B.A. (2011) State-and-transition modelling  
631 for adaptive management of native woodlands. *Biological Conservation* 144: 1224-1236.

632 Runge, M.C. (2011) An introduction to adaptive management for threatened and endangered  
633 species. *Journal of Fish and Wildlife Management* 2: 220-233.

634 Saffer, V. & Peake, T. (2014) The use of LiDAR to determine the presence of Malleefowl mounds. In  
635 *Proceedings of the 5th National Malleefowl Forum*, eds Bannerman & Davies. Pintak Pty Ltd. Dubbo,  
636 New South Wales, pp. 140-150.

637 Scarlett, L. (2013) Collaborative adaptive management: challenges and opportunities. *Ecology and*  
638 *Society* 18(3): 26.

639 Smith, A.D.M. & Walters, C.J. (1981) Adaptive management of stock-recruitment systems. *Canadian*  
640 *Journal of Fisheries and Aquatic Sciences* 38: 690-703.

641 Susskind, L., Camacho, A.E. & Schenk, T. (2012) A critical assessment of collaborative adaptive  
642 management in practice. *Journal of Applied Ecology* 49: 47-51.

643 Thompson, S., Thompson, G., Sackmann, J., Spark, J. & Brown, T. (2015) Using high-definition aerial  
644 photography to search in 3D for malleefowl mounds is a cost-effective alternative to ground  
645 searches. *Pacific Conservation Biology* 21: 208-213.

646 van Hespen, R., Hauser, C.E., Benshemesh, J., Rumpff, L. & Lahoz-Monfort, J.J. (in press) Designing a  
647 camera trap monitoring program to measure efficacy of invasive predator management. *Wildlife*  
648 *Research*.

649 Walsh, J.C., Wilson, K.A., Benshemesh, J. & Possingham, H.P. (2012) Unexpected outcomes of  
650 invasive predator control: the importance of evaluating conservation management actions. *Animal*  
651 *Conservation* 15: 319-328.

652 Walters, C. (1986) *Adaptive management of renewable resources*. The Blackburn Press, Caldwell  
653 New Jersey.

654 Walters, C.J. (2007) Is adaptive management helping to solve fisheries problems? *AMBIO* 36: 304–7.

655 Walters, C.J. & Hilborn, R. (1976) Adaptive control of fishing systems. *Journal of the Fisheries*  
656 *Research Board of Canada* 33: 145-159.


- 657 Westgate, M.J., Likens, G.E. & Lindenmayer, D.B. (2013) Adaptive management of biological systems:  
658 a review. *Biological Conservation* 158: 128–39.
- 659 Whitehead, A.L., Edge, K.-A., Smart, A.F., Hill, G.S. & Willans, M.J. (2008) Large scale predator control  
660 improves the productivity of a rare New Zealand riverine duck. *Biological Conservation* 141: 2784-  
661 2794.

662 (a)




663

664 (b)




665

666 *Figure 1. (a) An adult malleefowl in the wild, photographed by Graeme Tonkin. (b) The presumed*  
667 *original distribution of malleefowl across Australia, based on historical records (dark shading), with*  
668 *locations of long-term monitoring sites (black circles), and locations of candidate sites where*  
669 *monitoring data will support the predator control experiment (white triangles).*




670  
 671 *Figure 2. A schematic of adaptive management, originally published by Runge (2011), including the*  
 672 *contributions that this paper makes to adaptive malleefowl conservation. We developed the problem*  
 673 *framing, conservation objectives and alternative management options with stakeholders (2.2);*  
 674 *developed a network model and identified critical uncertainties (2.3); designed a monitoring*  
 675 *program to address the critical uncertainty (2.4); and formulated strategies for collaboration that*  
 676 *support conservation action and monitoring (2.5). We have not yet formally identified preferred*  
 677 *conservation actions, nor developed a plan for updating the network model.*

(a)



(b)



678

679


680

681

682


683

Figure 3. (a) The malleefowl ecosystem model structure developed for ensemble modelling. Arrows indicate the direction of influence, with the sign label indicating the direction of influence (see Appendix A for associated equations); absent connections between components indicate that there is no direct influence. (b) 95% credible intervals on predicted malleefowl population change as a response to a Recovery plan intervention (respectively: Dis = disease and inbreeding, Fire = fire intensity and severity, Cat = predation by cats, Fox = predation by foxes, Graz = competition from grazing herbivores, Hab = habitat loss and fragmentation). Diagrams adapted from Bode et al. (2017).



685  
686  
687  
688  
689  
690  
691  
692

Figure 4. Statistical power to detect a proportional change in malleefowl mound activity, over 5 years of monitoring, given the total number of unmanaged and managed sites included in the study and assuming a type I error rate of 0.05. Lines indicate the power to reject a null hypothesis of no change based on mound activity effect sizes ranging from a 10% to a 101% (i.e. two-fold) increase. See Appendix for details of model construction.



693  
694  
695  
696  
697  
698  
699

Figure 5. Power to detect a difference in predator activity (baiting effect) between a pair of control and impact sites, (a) given the number of cameras allocated to each site over a 12-month period, and (b) given the camera budget at each site for two alternative camera models. Images adapted from van Hespén et al. (in press).

700 **Supplementary Material**

701 **Appendix A: Summary of roles and responsibilities for malleefowl conservation**

702

703 *Table A1. A list of agencies involved in the conservation of malleefowl and/or other land management that is likely to influence malleefowl persistence. State*  
 704 *abbreviations are: Western Australia (WA), South Australia (SA), Victoria (Vic) and New South Wales (NSW). Other common abbreviations are: Natural*  
 705 *Resource Management (NRM), Catchment Management Authority (CMA), Local Land Service (LLS), and Indigenous Protected Area (IPA). While substantial*  
 706 *efforts have been made to engage broadly, this may not form a comprehensive account of all agencies concerned with malleefowl conservation.*

<i>Role</i>	<i>Agency names</i>	<i>Responsibilities</i>
Federal government	Department of the Environment and Energy (DOTEE)	Listing of threatened species under the Environmental Protection and Biodiversity Conservation (EPBC) Act 1999 Establishing a recovery team for listed species
Species recovery team (National Malleefowl Recovery Team)	National Malleefowl Recovery Team (NMRT, the 'recovery team' in this manuscript)	Developing a recovery plan Advising land managers of preferred management actions to conserve target species
State government	WA Department of Biodiversity Conservation and Attractions (DBCA); SA Department of Environment and Water (DEW), Parks Victoria (PV), Vic Department of Environment, Land, Water and Planning (DELWP), NSW Office of Environment and Heritage (OEH), NSW National Parks and Wildlife Service (NPWS)	Predator control in state and national parks Control of pest species, e.g. rabbits, goats Weed management Habitat restoration (revegetation) Fire management on public and private land Land clearing permits across all land types Implementing the Flora & Fauna Guarantee Act 1986 (FFG, Vic), Threatened Species Conservation Act (NSW) Malleefowl mound monitoring across all land types
Local area coordinators of land management	WA: South West Catchment Council, Northern Agriculture Catchment Council, Wheatbelt NRM, South Coast NRM, Rangelands NRM. SA: Eyre Peninsula NRM, Northern and Yorke NRM, Arid NRM, Alinytjara Wilurara NRM, South East NRM, SA Murray Darling Basin NRM. Vic: Mallee CMA, Wimmera CMA, North Central CMA NSW: Western LLS, Riverina LLS, Central West LLS	Implement actions from the recovery plan utilising federal funding under the National Landcare Partnerships program (17 projects with primary or secondary benefits for malleefowl worth AU\$24.8m over 2013-2023) Land clearing permits on private land (NSW)

Traditional Owners	Aboriginal Corporations: Maralinga Tjarutja (SA)*, Anangu-Pitjantjatjara- Yankunytjatjara (SA)*, Pila Nguru (WA), Ngaanyatjarra (WA) Indigenous Protected Area: Ninghan (WA), Mawonga (NSW), Walalkara (SA)*, Watarru (SA)* Other: Rick Farley Reserve (NSW)	Predator control on the IPA Malleefowl mound monitoring on Indigenous managed lands Camera monitoring predators on Indigenous managed lands
Non-government organisations (NGOs)	Bush Heritage Australia, Australian Wildlife Conservancy, Great Victorian Desert Biodiversity Trust, Gondwana Link, Gunduwa Regional Conservation Association	Predator control on NGO land Fire management on NGO land aligned with state government directives Malleefowl mound monitoring on NGO land Camera monitoring predators on NGO land
Community volunteer groups	Victorian Malleefowl Recovery Group (VMRG), WA Malleefowl Recovery Group, National Malleefowl Recovery Group, Kalgoorlie Field Naturalists, Dubbo Field Naturalists, Mid Murray Field Naturalists	Malleefowl mound monitoring as permitted across all tenures
Mining companies	Wiluna Uranium mine, St Ives mining, Mount Gibson Iron Limited Extension Hill Operations, Cliffs Resources, Cameco Yeelirrie	Predator control on leased land Malleefowl mound monitoring on leased land Camera monitoring predators on NGO land
Universities	Federation University, Melbourne University, La Trobe University	Research Predator control on leased land Malleefowl mound monitoring on leased land Camera monitoring predators on NGO or leased land??
Private property	Many individuals	Predator control on private land Malleefowl mound monitoring on private land Camera monitoring predators on private land Weed control and habitat restoration on private land

707 \* Aboriginal people of the Great Victoria Desert in SA and WA.

708

709 **Appendix B: Ensemble model structure and analysis**

710 The ecosystem model (Figure 3a) followed the Lotka-Volterra structure:

711 
$$\frac{dN_i}{dt} = r_i N_i + \sum_{j=1}^C a_{ij} N_i N_j$$

712 for ecosystem components  $i = 1, 2, \dots, C$ .  $N_i(t)$  was the quantity of component  $i$  at time  $t$  years (the  
713 abundance or density of a population, volume of rainfall, etc.), and  $C = 14$  was the number of  
714 components in the ecosystem. The intrinsic growth rate of component  $i$  was  $r_i$ , and the per-unit  
715 effect of component  $j$  on each unit of component  $i$  was  $a_{ij}$  (Bode et al. 2017).

716  
717 All ecosystem component quantities  $N_i(t)$  were scaled to lie between 0 and 1; plausible values for  
718 the interaction parameters  $a_{ij}$  fell between -1 and 1. In the first instance, we used workshop  
719 participants' knowledge to restrict parameters to  $[-1, 0)$  when the interaction between components  
720 was thought to be negative (red lines in Figure 3a), 0 when there was no interaction between  
721 components, and  $(0, 1]$  when the interaction between components was thought to be positive (blue  
722 lines in Figure 3a). We assigned uniform prior distributions over the above ranges to describe  
723 parameter uncertainty in component interactions. We assigned inverse uniform distributions to the  
724 intrinsic growth rates, i.e.  $1/(r_i + 1) \sim U(0, 1)$ . We used Latin hypercube sampling to generate  $10^9$   
725 plausible parameterisations of the ecosystem model.

726

727 Following the refinement of the model set described in 2.2 we developed six candidate actions or  
728 interventions, with each one designed to address a threat to malleefowl that had been identified in  
729 the National Recovery Plan (Benshemesh 2007):

- 730 1. *Disease and inbreeding*: the specific management action was unclear, but the result of  
731 successful management was predicted to be a 10% increase in malleefowl population  
732 growth.
- 733 2. *Fire intensity and severity*: the management action and component response was a 50%  
734 increase in the area burned by fire.
- 735 3. *Predation by introduced cats*: baiting that targeted cats was predicted to reduce cat  
736 populations by 85%.
- 737 4. *Predation by introduced foxes*: baiting that targeted foxes was predicted to reduce fox  
738 populations by 95%.
- 739 5. *Competition by herbivores*: Mustering feral goats was predicted to reduce their population  
740 by 30%, and baiting rabbits was predicted to reduce their population by 30%.
- 741 6. *Habitat loss and fragmentation*: Active restoration was predicted to increase seedling and  
742 vegetation components by 15%.

743

744 **Appendix C: Power analysis of control-impact monitoring on malleefowl mound activity**

745 In developing an initial power analysis, we assume clusters of two sites each with similar  
746 characteristics (e.g. climate, vegetation), that can be used as a ‘control’ site (no predator  
747 management) and an ‘impact’ site (intense predator control). To ensure that predator activity around  
748 the monitored mounds reflects the chosen management, we have recommended that predator  
749 management be uniformly applied to a 10000 ha area surrounding the mounds as a ‘buffer’. To limit  
750 the possibility that predators travel between unmanaged and managed sites, we also recommend that  
751 sites be placed at least 8km apart (and often more, depending on the location of tracks).

752 For each site, the response data is the *number of active mounds* in a given year. We assume a common  
753 average mound activity with superimposed temporal and spatial fluctuations. The paired nature of  
754 the experiment (and consequent model structure) helps disentangle the long-term (average) effect of  
755 predator management on malleefowl breeding, from the spatial and temporal fluctuations, as:

- 756 1. some sites consistently show higher nesting activity than others, due to e.g. more suitable  
757 habitat for breeding. Pairing together similar sites, that are ‘consistently high’ or ‘consistently  
758 low’, helps to isolate the effect of predator management from the effects of breeding activity  
759 due to local habitat quality.
- 760 2. some malleefowl breeding seasons will be ‘better’ or ‘worse’ than average across the entire  
761 range. This influences how we view the impact of predator management, e.g. we are less likely  
762 to attribute a breeding activity increase to predator management if it occurs in both control  
763 and impact sites.

764 The purpose of this power analysis is to determine – in advance of data collection – the probability  
765 that a control-impact experiment would correctly recognise the benefits of predator management,  
766 assuming it has a given effect size.

767 **Model specification**

768 For each site  $s$  and year  $t$ , the number of active mounds (random variable  $a$ ) can be described with a  
769 Poisson distribution with year- and site-specific mean  $\lambda_{s,t}$ :

770 
$$a_{s,t} \sim \text{Pois}(\lambda_{s,t}).$$

771 We model the mean number of active mounds  $\lambda$  as a log regression with time and cluster-specific  
772 variation as:

773 
$$\log(\lambda(s, t)) = \beta_0 + \varepsilon_S(s) + \varepsilon_T(t) + b \cdot c(s)$$

774 where  $\beta_0$  is the average log-rate (intercept),  $\varepsilon_S$  is a site-level random term (spatial variation),  $\varepsilon_T$  is a  
775 year-specific random term (temporal variation) and  $b$  is the effect of predator management (the  
776 indicator covariate  $c(s)$  is 1 for ‘impact’ sites and 0 otherwise). The random effects are described as  
777 normally distributed with zero mean and variances:

778 
$$\varepsilon_S \sim N(0, \sigma_S^2)$$

779 
$$\varepsilon_T \sim N(0, \sigma_T^2)$$

780 Note that the site random effect  $\varepsilon_S(s)$  is shared by each cluster of control-impact sites.

781 **Simulation parameters**

782 We analysed historical mound activity data in the state of Victoria from the National Malleefowl  
783 Monitoring Database (<http://database.malleefowlvictoria.org.au/Start.aspx>) to obtain values to  
784 parameterise realistic simulation of mound activity data. The analysis was based on the model above  
785 (without the predator management term  $b \cdot c(s)$ ). In particular, we estimated:

- 786 • long-term **average mound activity**:  $\lambda=3.07$  active mounds per site (equivalent to  $\beta_0=1.123$ ).
- 787 • spatial variation in mound activity across sites:  $\sigma_s=1.172$
- 788 • temporal variation (sample variance) in mound activity, from year-to-year:  $\sigma_T=0.411$

789 We investigated the amount of monitoring effort that would be needed to draw scientifically rigorous  
 790 conclusions under several different predator management responses (i.e. different ‘effect sizes’), from  
 791 no effect ( $b = 0$ ; average active mounds  $\lambda = 3.07$  at both sites) to doubling the average number of  
 792 active mounds ( $b = 0.7$ ; from  $\lambda = 3.07$  mounds at a control site to  $\lambda = 6.19$  mounds at an impact site;  
 793 Table A1).

#### 794 **Power analysis simulations**

795 We performed a power analysis on the above defined control-impact experiment, for scenarios  
 796 defined by the number of sites  $S$ , number of years  $T$  and an effect size  $b$  using the following steps:

- 797 1) Simulate a data set of mound activity observations (active/inactive) using the model above  
 798 (assumed as the reference “truth”)
- 799 2) Analyse the simulated data set using the same model structure
- 800 3) Determine whether the estimation of the existing effect of predator control ( $\hat{b}$ ) is found to be  
 801 statistically significant at the customary  $\alpha = 0.05$  significance level (i.e. whether the 95%  
 802 Credible Interval does not include the value zero)<sup>1</sup>.
- 803 4) Repeat steps 1 to 3 for 200 iterations
- 804 5) Estimate the statistical power  $G$  for each scenario as the percentage of simulations in that  
 805 scenario in which the assumed effect of predator management is detected.

806 The simulation study was conducted using the software R (R Core Team 2015, version 3.2.0). To fit  
 807 the model specified above to each data set in step 2, we performed Bayesian Markov chain Monte  
 808 Carlo sampling in the rjags package (Plummer 2016) and monitored the effect of predator control ( $b$ ,  
 809 step 3). Note that for each site cluster, the proportional increase in activity between control and  
 810 impact sites depends on the number of active mounds per site, because  $\lambda$  includes site- and time-  
 811 specific random effects. Therefore the percentage increases in mound activity in Table A1 are  
 812 accurate only at the landscape level, and the parameter  $b$  is the true underlying difference that is  
 813 used directly in the model.

814 *Table C1. Conversions of effect size  $b$  to mean number of active mounds  $\lambda$  and percentage increase in mound*  
 815 *activity between unmanaged and managed sites. The % increase compared to no management is only valid for*  
 816 *the mean of  $\lambda = 3.07$ , and is shown here to provide an intuitive idea of the effect size.*

effect size ( $b$ ), log scale	0.0	0.1	0.2	0.3	0.4	0.5	0.6	0.7
corresponding average $\lambda$	3.07	3.40	3.75	4.15	4.59	5.07	5.60	6.19
% increase in managed site above unmanaged site	---	10.5%	22.1%	35.0%	49.2%	64.9%	82.2%	101.4%

817

#### 818 **References**

- 819 Plummer, M. (2016) rjags: Bayesian graphical models using MCMC. R package version 4-6.  
 820 <https://CRAN.R-project.org/package=rjags>
- 821 R Core Team (2015) R: a language and environment for statistical computing. Version 3.2.0. R  
 822 Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>

<sup>1</sup> The 5% significance level implies that there is a 5% chance of declaring that an effect exists when in fact it does not; a trade-off exists between this error and statistical power, so that lowering that probability of falsely detecting an effect implies a lower power to detect a true effect.