



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Schwab, RHM;Amin, N;Flanagan, DJ;Johanson, TM;Pheese, TJ;Vincan, E

Title:

Wnt is necessary for mesenchymal to epithelial transition in colorectal cancer cells

Date:

2018-03-01

Citation:

Schwab, R. H. M., Amin, N., Flanagan, D. J., Johanson, T. M., Pheese, T. J. & Vincan, E. (2018). Wnt is necessary for mesenchymal to epithelial transition in colorectal cancer cells. *Developmental Dynamics*, 247 (3), pp.521-530. <https://doi.org/10.1002/dvdy.24527>.

Persistent Link:

<https://hdl.handle.net/11343/293101>

Wnt is necessary for mesenchymal to epithelial transition in colorectal cancer cells.

Renate H. M. Schwab¹, Nancy Amin¹, Dustin J. Flanagan¹, Timothy M. Johanson¹, Toby J. Phesse^{1,2^}, Elizabeth Vincan^{1,3^*}

¹Molecular Oncology Laboratory, University of Melbourne and the Victorian Infectious Diseases Reference Laboratory, Doherty Institute of Infection and Immunity, 792 Elizabeth Street, Melbourne, Victoria 3000, Australia

²European Cancer Stem Cell Research Institute, School of Biosciences, Cardiff University, Hadyn Ellis Building, Maindy Road, Cardiff CF24 4HQ, Wales, UK

³School of Biomedical Sciences, Curtin University, Perth, Western Australia 6845, Australia

[^]These authors contributed equally

**To whom correspondence should be addressed*

Elizabeth Vincan, PhD

Molecular Oncology Laboratory, University of Melbourne and the Victorian Infectious Diseases Reference Laboratory,

Doherty Institute for Infection and Immunity,

792 Elizabeth Street,

Melbourne,

VIC 3000

Phone: +613 9342 9348

Fax: +613 9342 9696

Email: evincan@unimelb.edu.au

Funding: Project grants from the Australian National Health & Medical Research Council (NHMRC) and Melbourne Health to EV and TJP. Fellowships from the Cancer Council of Victoria to DJF and from Cardiff University/Capital Medical University to TJP.

Key words: epithelial to mesenchymal transition, EMT, mesenchymal to epithelial transition, MET, metastasis, Wnt signaling, Frizzled7, Wnt2B, Wnt3A, colorectal cancer

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version record](#). Please cite this article as [doi:10.1002/dvdy.24527](https://doi.org/10.1002/dvdy.24527).

Abstract

Background: Metastasis underlies most colorectal cancer mortality. Cancer cells spread through the body as single cells or small clusters of cells that have an invasive, mesenchymal, non-proliferative phenotype. At the secondary site, they revert to a proliferative ‘tumor constructing’ epithelial phenotype to rebuild a tumor. We previously developed a unique *in vitro* 3D model, called LIM1863-*Mph*, which faithfully recapitulates these reversible transitions that underpin colorectal cancer metastasis. Wnt signaling plays a key role in these transitions and is initiated by the coupling of extracellular Wnt to Frizzled (FZD). Using the LIM1863-*Mph* model system we demonstrated that the Wnt receptor FZD7 is necessary for mesenchymal to epithelial transition (MET). Here we investigate the role of Wnt in MET. **Results:** Wnt secretion is dependent on palmitoylation by Porcupine (PORC). A PORC inhibitor (IWP2) that prevents Wnt secretion, blocked the epithelial transition of mesenchymal LIM1863-*Mph* cells. Wnt gene array analysis identified several Wnts that are upregulated in epithelial compared to mesenchymal LIM1863-*Mph* cells, suggesting these ligands in MET. Wnt2B was the most abundant differentially expressed Wnt gene. Indeed, recombinant Wnt2B could overcome the IWP2-mediated block in epithelial transition of mesenchymal LIM1863-*Mph* cells.

Conclusion: Wnt2B co-operates with Frizzled7 to mediate MET in colorectal cancer.

Introduction

Colorectal cancer (CRC) is a major health problem worldwide and, similar to other forms of cancer, it is much harder to treat once it has spread to other organs in the body. A necessary step in the genesis of most CRCs is deregulation of the β -catenin dependent Wnt signaling pathway (Clevers and Nusse, 2012), which occurs through genetic mutation of downstream pathway components, primarily the tumor suppressor gene *Adenomatous Polyposis Coli (APC)* (Fearon, 2011). At a molecular level, the Wnt/ β -catenin pathway regulates the cytoplasmic and nuclear levels of β -catenin via a cytoplasmic destruction complex that includes APC, Axin and GSK3 β . Mutations of *APC* in CRC disrupts this regulation and β -catenin accumulates in the cytoplasm, translocates to the nucleus and leads to the constitutive activation of TCF/LEF/ β -catenin transcription of target genes (Korinek et al., 1997; van de Wetering et al., 2002). However, the level of nuclear β -catenin is variable in CRC despite activating mutations in the downstream components of the pathway (Brabletz et al., 1998; Brabletz et al., 2001), indicating additional regulation of the Wnt/ β -catenin pathway by the tumor environment (Vincan and Barker, 2008).

The majority of CRCs are well differentiated forming organized tubular structures, yet they still metastasize, and therefore differentiation status is often a poor prognostic factor for this cancer. Both primary and secondary (metastases) well-differentiated tumors have clearly defined localized invasive areas where the tubular structures of the CRC is reorganized yielding tumor cell sheets and isolated tumor cells interspersed with the stroma (Brabletz et al., 1998; Brabletz et al., 2001). Histological detection of invasive areas in the tumor predicts poor outcome, independent of tumor staging (Wong et al., 2003; Ueno et al., 2014). Tumor cells engaged in tubular structures are well differentiated with membrane β -catenin (polarized epithelial cells) that stain with Ki-67. In contrast, cell division (Ki-67 staining) is decreased or undetected in de-differentiated (migratory, invasive, mesenchymal) tumor cells with intense cytoplasmic and nuclear β -catenin at invasive areas (Brabletz et al., 2001). Based on these phenotypic changes, de-differentiation at the invasive front is referred to as an epithelial to mesenchymal transition (EMT) (Kirchner and Brabletz, 2000). Once in the secondary organs, the reverse transition occurs (MET) which re-instates the epithelial, proliferative phenotype. Notably, the pathology of primary and secondary tumors for most CRCs is the same (Thiery, 2002; Brabletz et al., 2005a). Transitions between epithelial and mesenchymal states are thus dynamic and reversible. Presumably, this reversal of de-differentiated

disseminated cells to an epithelial phenotype is necessary to re-instate cell division so that the tumor structure can be re-built at the secondary site (Thiery, 2002; Vincan, 2004).

During metastasis cancer cells disseminate through the body as single cells or small clusters of cells and can remain 'dormant' for many years (MacDonald et al., 2002; Naumov et al., 2002). Moreover, disseminated cancer cells (mesenchymal, non-proliferating) evade conventional anti-cancer therapies, which in general target proliferating cells. The ability to eliminate these 'dormant' cancer cells is critical for curative treatment. Initiation of tumor growth at the secondary site is the rate-limiting step in metastasis (Chambers et al., 2002). Consequently, circumventing release from dormancy offers a further avenue for therapeutic intervention. However, the molecular mechanisms controlling early events of release from 'dormancy' are still largely unknown due to the complexity this presents for analysis *in vivo*. To overcome this limitation, we have established a unique *in vitro* morphogenesis culture system using the human colorectal cancer cell line LIM1863 (Whitehead et al., 1987). In this model system, LIM1863 cells growing as a two-dimensional (2D) monolayer undergo MET to form 3D highly organized, multicellular structures that resemble enclosed carcinoma tubules (termed organoids). We demonstrated cyclic transition from 2D to 3D and refer to this adapted cell line as LIM1863-*Mph* (morphogenetic) (Vincan et al., 2007a; Vincan et al., 2007b; Vincan et al., 2008). The dynamic phenotype transitions between 2D and 3D faithfully recapitulate the pathology of invasive (deconstructing tubular structures) and central (constructing tubular structures) areas of carcinoma tissues (Vincan et al., 2007a). We recorded nuclear β -catenin in the monolayer cells (low or undetectable Ki-67 staining), then a transient, short, sharp increase in nuclear β -catenin at the initiation of MET, just as the cells lift off the tissue culture plastic and become Ki-67 positive. This intense nuclear β -catenin rapidly subsides to undetectable levels as the organoids form. In contrast, Ki-67 staining is retained as all the organoid cells proliferate. Using RNAi-mediated knock-down in LIM1863-*Mph*, we demonstrated that the Wnt receptor Frizzled7 (FZD7) is necessary for MET (Vincan et al., 2007b). FZD7-mediated Wnt/ β -catenin signaling is known to orchestrate tissue "constructing" morphogenetic events during embryonic development (e.g. epithelial organization of somites (Linker et al., 2005)) and our findings indicate a similar role for FZD7 in the context of CRC (Vincan et al., 2007c).

However, the Wnt ligands that co-operate with FZD7 during the initiation of MET are not known. Here we show that Wnt is required for MET in the LIM1863-*Mph* model as

inhibiting Wnt secretion blocks MET. Secretion of Wnt is dependent on porcupine (PORC) (van den Heuvel et al., 1993), an enzyme in the endoplasmic reticulum that post translationally palmitoylates Wnts at a highly conserved serine residue (Lum and Clevers, 2012; Madan et al., 2016). Inhibition of PORC with IWP2 (Chen et al., 2009a; Chen et al., 2009b) blocked MET. We identify the Wnts that are differentially expressed by mesenchymal and epithelial LIM1863-*Mph* cells using an array platform; and demonstrate that recombinant Wnt2B or the archetypical β -catenin activating Wnt, Wnt3A, can partially rescue MET when Wnt secretion is blocked by IWP2.

Results

Wnt expression and signaling in LIM1863-*Mph* cells

The LIM1863-*Mph* cells undergo cyclic transitions between monolayer and free-floating organoid clusters. The organoids are spheres of highly polarized epithelial cells around a central lumen (Vincan et al., 2007a; Vincan et al., 2007b), shown diagrammatically in Fig. 1A. Having identified a role for FZD7 in orchestrating MET in the LIM1863-*Mph* model (Vincan et al., 2007b), and given that the expression of FZD7 does not change during these transitions (data not shown), the next important step was to identify which Wnt genes were differentially expressed by monolayer and organoid cells, and thus could potentially cooperate with FZD7 in this process. To answer this question, we employed the Human Wnt Signaling Pathway Array (Qiagen). Total RNA was extracted from LIM1863-*Mph* monolayer and organoid cells, cDNA synthesized and assayed using the cDNA array. Relative expression was determined using HMBS as reference (Fig. 1B). Wnt2B, Wnt3 and Wnt10A were markedly up-regulated in the organoids compared to the monolayer cells; expression of Wnt4 and Wnt11 were relatively unchanged, while the other Wnt genes were expressed at very low levels. Looking at expression levels of Wnt/ β -catenin target genes (Fig. 1C), these were similar in both cell types with less than two-fold changes in expression. The differences in Wnt2B, Wnt3 (and Wnt3A albeit expressed at very low levels) and Wnt target gene expression identified by the array analyses were validated by independent qRT-PCR on a separate set of cell preparations (Fig. 2A & 2B). Wnt2B was the most abundant differentially expressed Wnt and was thus investigated further. Wnt2B expression in the LIM1863-*Mph* organoids was confirmed by immunofluorescence and confocal microscopy, in which we

observed Wnt2B localized in puncta apical to the nuclei of LIM1863-*Mph* organoid cells (Fig. 2C). As shown previously (Vincan et al., 2007a; Vincan et al., 2007b), *laminin- γ 2*, a marker for the invasive front (Hlubek et al., 2001) is very strongly up-regulated (~30 fold) in the mesenchymal cells compared to the mature organoids. Similarly, the EMT marker *Claudin1* (Suh et al., 2013) is also up-regulated in the mesenchymal cells (Fig. 2B).

Wnt is necessary for MET

Wnt2B functions to promote the activity of intestinal stem cell Wnts (Farin et al., 2012), which are also the cell-of-origin in colorectal cancer (Barker et al., 2009). We have recently shown that Wnt2B co-operates and co-immunoprecipitates with FZD7 in intestinal stem cells (Flanagan et al., 2015), and therefore we focused on the function of this Wnt in MET as stem cell properties are often hijacked by cancer cells (Barker et al., 2009; Pesse et al., 2014).

First, we investigated whether Wnt is necessary for MET. Wnts are secreted growth factors and require post-translational modification by PORC to be secreted (van den Heuvel et al., 1993; Chen et al., 2009a; Chen et al., 2009b; Lum and Clevers, 2012; Madan et al., 2016). Inhibition of PORC enzyme activity has been shown to inhibit the secretion of Wnts, and therefore we used the PORC inhibitor IWP2 as a tool to inhibit Wnt function. We confirmed inhibition of Wnt function using Wnt3A-L-cells, a mouse fibroblast continuous cell line that was stably transfected with a Wnt3A construct and secretes Wnt3A into the supernatant (Willert et al., 2003). We collected conditioned medium from parental L-cells, and Wnt3A-L-cells with or without the PORC inhibitor IWP2 (5 μ M). These conditioned media were then added to parental L-cells. We showed by immunoblot that Wnt3a conditioned medium induced the accumulation of β -catenin and that this activity was lost when the PORC inhibitor IWP2 was added during preparation of the Wnt3A-conditioned medium, thus confirming that IWP2 is functioning to inhibit Wnt secretion in these cells (Fig. 3A).

Next, monolayer organoids were treated with 5 μ M IWP2 as they were just starting to re-organize to form organoids i.e. undergoing MET. Inhibition of PORC blocked MET and the monolayer patch had a “cotton wool” appearance with no budding organoids, consistent

with a mesenchymal phenotype (Vincan et al., 2007b). In contrast, numerous generating organoids were observed in the carrier-treated monolayer patch (Fig. 3B). Expression of the mesenchymal markers, *laminin- γ 2* and *Claudin1* remained high in the IWP2 treated cells and was increased approximately two fold when compared to the control DMSO treated cells, confirming their mesenchymal phenotype (Fig. 3C). The expression of Wnt target genes was relatively comparable between the IWP2 treated and control cells (Fig. 3C). This is consistent with our observation that these genes are relatively unchanged when mesenchymal and epithelial LIM1863-*Mph* cells are compared (Fig. 1C).

Wnt2B and Wnt3A function in MET

Following the demonstration that inhibiting Wnt secretion blocked MET, we next investigated if addition of recombinant Wnts could rescue/restore MET. To better visualize the changes in the morphology during the MET process, the cells were seeded into chamber slides and stained for either E-cadherin or ZO-1, to demonstrate the cell architecture. Cultures were treated with DMSO, 5 μ M IWP2 or with 5 μ M IWP2 with either Wnt2B (100 ng/ml) or the archetypical β -catenin-activating Wnt, Wnt3a (100 ng/ml) (Willert et al., 2003). The E-cadherin stained DMSO treated cultures show that organoids were re-assembling because the monolayer cell patches have “holes”, i.e. areas where the cells are lifting off (arrows) and starting to re-organize themselves into organoids (indicated by the circles in Fig. 4A). In contrast, the IWP2 treated cells remain a tight patch and the E-cadherin staining does not distinguish any organoid formation or cells lifting off the tissue culture plastic (Fig. 4B). Adding recombinant Wnt2B (Fig. 4C) or Wnt3A (Fig. 4D) restores organoid assembly with “holes” observed in the monolayer patches, indicating the cells are lifting off and re-organizing into organoids. The ZO-1 staining shows the central lumen of the organoids being assembled in the DMSO treated cultures (Fig. 5A, circles indicate one assembling organoid). These clusters of apical ZO-1 staining are not observed in the cultures treated with IWP2 (Fig. 5B). In contrast, the clusters of apical ZO-1 were clearly evident when recombinant Wnt was added to the IWP2-treated cultures (Fig. 5C & 5D). We have previously demonstrated that FZD7 is required for MET (Vincan et al., 2007b) and binds to Wnt2B (Flanagan et al., 2015). Thus, our collective data indicates that FZD7 transmits signals from Wnt2B to orchestrate MET in this model of colorectal cancer morphogenesis (Fig. 6).

Discussion

In CRC, aberrant activation of the Wnt/ β -catenin signaling pathway leads to the accumulation of β -catenin in the nucleus and thus the hyper-activation of β -catenin-TCF/LEF gene transcription. Indeed, the over-activation of the Wnt signaling pathway, caused by mutations to some of its intracellular components, is found in nearly all CRCs, both sporadic and familial (Kinzler et al., 1991). Most notable is the tumor suppressor gene *APC*, which is found to be mutated in approximately 80-90% of CRCs. However, in addition to these *APC* mutations, the over-expression of certain upstream signaling components can regulate Wnt signaling activity, irrespective of the downstream pathway-activating mutations (Caldwell et al., 2004; Suzuki et al., 2004; Vincan et al., 2005; Vincan et al., 2007b; Ueno et al., 2009). Importantly, additional Wnt signaling underlies malignant progression as the acquisition of invasive capabilities and metastasis relies on dynamic reversible processes, EMT and MET (Thiery, 2002). Thus, it is of great importance to understand how these processes are coordinated and regulated, as it may provide novel avenues to block metastasis, which is ultimately the cause of cancer death. The results presented in this study demonstrate the potential of blocking the interaction between Wnt and FZD7, as a therapeutic target to prevent MET and cancer recurrence (Vincan and Barker, 2008).

Although additional signaling by the Wnt-FZD receptor complex was initially considered unnecessary in CRC, it is now accepted that the upstream components participate in the initiation and progression of CRC. This is based on several lines of evidence. Expression of nuclear β -catenin in early adenoma is not as strong as it is in later stages (Phelps et al., 2009a; Phelps et al., 2009b). Also, naturally occurring inhibitors of Wnt-FZD interaction, the secreted Frizzled related proteins (sFRPs), are *bona fide* tumor suppressors in CRC (Caldwell et al., 2004; Suzuki et al., 2004), suggesting that the additional modulation of the β -catenin-dependent Wnt signaling pathway is via the upstream components of the pathway. Furthermore, epigenetic silencing of sFRP can occur before truncating mutations in *APC* (Suzuki et al., 2004), indicating that aberrant signaling from the Wnt-FZD complex may precede aberrant signaling due to truncation of *APC*. Furthermore, both Wnts (Kato et al., 1996; Dimitriadis et al., 2001; Kirikoshi et al., 2001; Ru et al., 2008) and FZDs (Sagara et al., 1998; Holcombe et al., 2002; Vincan, 2004; Vincan et al., 2007b; Ueno et al., 2008; He et al.,

2011) are over-expressed in CRC tissues and cell lines and can additionally modulate the β -catenin-dependent Wnt signaling pathway (Vincan et al., 2005; Vincan et al., 2007b; Ueno et al., 2009), despite mutation to *APC* or *CTNN1* (the gene that codes for β -catenin). Notably, Wnt signaling appears to be constrained in CRC cells [reviewed in (Pheesse et al., 2016)], giving way to bursts of localized increased signaling as occurs at the invasive front (Brabletz et al., 2005b) where aberrant activation of the Wnt pathway can induce EMT in tumor cells (Mariadason et al., 2001; Kim et al., 2002; Muller et al., 2002). Our collective data in the LIM1863-*Mph* model indicates that the initiation of MET is associated with a brief burst of even higher Wnt/ β -catenin signaling than that seen in the dormant mesenchymal cells at the invasive front (Brabletz et al., 2001; Vincan et al., 2007b).

Using the LIM1863-*Mph* model system, we demonstrated a requirement for FZD7 in MET that is necessary to assemble the organoids (Vincan et al., 2007b), but the question of which Wnt ligand was signaling via FZD7 remained unanswered. There are 19 mammalian Wnts and here we investigated which of these Wnt ligands interacts with FZD7 to activate MET. First, we employed a gene array and identified Wnt2B, Wnt3 and Wnt10A as differentially expressed by mesenchymal and organoid LIM1863-*Mph* cells. Next, we demonstrated a requirement for Wnt in the assembly of organoids by demonstrating that inhibition of Wnt secretion using IWP2 blocked MET. We chose to further investigate the role of Wnt2B in the LIM1863-*Mph* cells as we had identified this Wnt as a signaling partner of FZD7 in intestinal stem cells (Flanagan et al., 2015) and Wnt2B can activate β -catenin-dependent Wnt signaling (Goss et al., 2009). Indeed, we demonstrate that recombinant Wnt2B [or Wnt3A, the archetypical β -catenin-activating Wnt (Willert et al., 2003)] could restore MET when Wnt secretion was inhibited in the monolayer LIM1863-*Mph* cells. Notably, Wnt2B mediated β -catenin-dependent Wnt signaling is implicated during the morphogenesis of renal tubules (Iglesias et al., 2007), which is one of the key MET events during development. Thus a role for Wnt2B in promoting the formation of glandular structures in CRC is consistent with its role during development. Furthermore, Wnt2B was also one of the first Wnts demonstrated to be over-expressed in CRC cell lines (Katoh et al., 1996; Bafico et al., 2004); Notably, Wnt3 is also overexpressed in colorectal tumors (Voloshanenko et al., 2013). Thus the role we identify for Wnt2B, and potentially Wnt3, are consistent with over-expression reported for these Wnts in colorectal cancer.

Wnt3 produced by Paneth cells in the epithelium is essential for intestinal stem cells (Sato et al., 2009; Sato et al., 2011). In the absence of Wnt3, Wnt2B can compensate (Farin et al., 2012). More recently, it was shown, that high levels of Wnt2B are predominantly expressed in Gli1 positive sub-epithelial mesenchymal cells, which are adjacent to the crypts (Valenta et al., 2016). This localization of Wnt2B is consistent with previous reports showing Wnt2b expression in the mesenchyme of mouse intestine (Gregorieff et al., 2005). Collectively, these studies identify Wnt2B and Wnt3 as intestinal stem cell Wnts. Intriguingly, the data presented here indicates that these stem cell Wnts also function in colorectal cancer morphogenesis indicating that the stem cell Wnts, like FZD7, feature in colorectal cancer. Notably, we demonstrated that FZD7 stem cell function can be partially compensated by closely related FZD genes, FZD1 and FZD2 (Flanagan et al., 2015), however, the same compensatory mechanism does not appear to occur in cancer as knock-down of FZD7 blocks MET (Vincan et al., 2007b), and inhibiting FZD7 function with a dominant-negative receptor ectodomain has potent antitumor activity in colorectal cancer (Vincan et al., 2005) and other cancers such as liver cancer [reviewed in (Vincan and Barker, 2008)].

Colorectal cancer (CRC) is a major health problem worldwide that is exacerbated by an aging population. CRC can effectively be cured if it remains localized and is surgically resected, however cancer cells often spread to other organs with metastases underlying most CRC mortality. Clearly better strategies are needed to stop tumor recurrence, which will ultimately lead to better management and treatment of advanced CRC, and consequently, increase patient survival. Furthermore, reversible EMT and MET underpins metastasis of other solid tumors (Ocana et al., 2012), thus the mechanisms for MET reported here might also be relevant to other forms of cancer.

Experimental Procedures

Cell lines, cell culture and live cell imaging

Cells lines were either purchased from ATCC (HEK293), provided by the Ludwig Institute for Cancer Research (LIM1863) (Whitehead et al., 1987) or the Nusse laboratory (parental L-cells and Wnt3a-Lcells) (Willert et al., 2003). The adapted LIM1863-*Mph* line

has been previously described (Vincan et al., 2007a; Vincan et al., 2008). Cells were cultured in RPMI 1640 medium (GIBCO, 11875-119), supplemented with 10% heat inactivated fetal calf serum (FSC, Chemicon International, Embryo Max FCS, ES qualified ES-009-A), 20 mM HEPES, Glutamine and antibiotics (Penicillin/Streptomycin). The LIM1863-*Mph* cells were cultured in rich medium with additional supplements [1 $\mu\text{g}/\text{mL}$ hydrocortisone (Solu-Cortef for injection), $\sim 0.9\text{-}1\mu\text{g}/\text{mL}$ insulin (Insulin for injection, 100U/mL, from Pharmacy) and 10^{-4}M 1-thioglycerol (Sigma, M6145)]. The cells were incubated at 37°C in a humidified atmosphere containing 10% CO_2 .

To plate the LIM1863-*Mph* so that they preferentially form monolayer patches, adherent LIM1863-*Mph* cells were detached with Trypsin/EDTA solution [0.05% (w/v) trypsin 1:250, 0.02% (w/v) EDTA, pH7.0] and seeded at 5×10^3 cells/ cm^2 tissue culture well surface area as previously described (Vincan et al., 2008).

Differential interference contrast (DIC) images of LIM1863-*Mph* cultures were taken on a Nikon Ti-E microscope using either a $4\times$ Plan Fluor NA 0.13 objective or a $10\times$ Plan Fluor NA 0.3 objective. A focal stack of images was collected $10\mu\text{m}$ apart and processed through the “Best Focus” function of MetaMorph v7.7.7 (Molecular Devices) to generate the final image (Vincan et al., 2007b).

Detection of β -catenin by immunoblot

L-cells were seeded in replicate wells treated with conditioned medium (CM) harvested from the L-cell parental line, or the Wnt3a-producing cell line (Willert et al., 2003) with or without the Porcupine inhibitor IWP2 (Chen et al., 2009a; Chen et al., 2009b). After treatment with the conditioned media, the L-cells were washed with warm calcium and magnesium free phosphate buffered saline (PBS^-) and then scraped from the well in to ~ 1 ml of PBS^- . The cells were then pelleted at 1200 rpm for 2min and resuspended in $50\mu\text{l}$ of PBS^- , followed immediately by the addition of hot ($\geq 95^{\circ}\text{C}$) 2x Laemmli Sample Buffer (Bio-Rad Laboratories, 161-0737) with 0.5M β -mercaptoethanol and then incubated at $\geq 95^{\circ}\text{C}$ for 5 min. The whole cell lysate protein samples were separated by SDS-PAGE (sodium dodecyl sulfate polyacrylamide gel electrophoresis) and the separated proteins transferred to a nitrocellulose membrane (Hybond C-Extra, Amersham, RPN303E). After transfer, the membrane was blocked with 5% (w/v) skim milk powder in Tris buffered saline (TBS) with 0.1% Tween-20 (TBS-T) for 1hr at room temperature, rinsed and incubated overnight in

primary antibody (mouse anti- β -catenin, BD Transduction Labs, 610154) at 4°C. Following the overnight incubation, the primary antibody was washed off with TBS-T and the bound antibody detected with anti-mouse-HRP (DAKO, P0161) and Enhanced Chemiluminescence (ECL) [Clarity Western ECL Substrates; Luminol/enhancer reagent and Peroxide reagent (Bio-Rad, 1705060)].

Immunofluorescence staining

To stain LIM1863-*Mph* during MET, detached monolayer cells were seeded at $5 \times 10^3/\text{cm}^2$ in 2-chamber slides (Lab-Tek, 177429) as described (Vincan et al., 2008) and given the indicated treatments. Following treatment, the growth medium was removed and the cells were washed twice in PBS⁺⁺ (PBS with Mg⁺⁺/Ca⁺⁺) and fixed *in situ* with methanol at -20°C for 20 min. The cells were incubated for 1hr in 5% FCS in 0.2% PBST to block nonspecific binding of the antibodies. The primary antibody was diluted in 5% FCS in 0.05% PBST (PBS + 0.05% Triton X-100), and left overnight at 4°C. After washing, the cells were incubated in the secondary antibody for 1hr at room temperature and then DAPI (4'-6-Diamidino-2-phenylindole) was added to stain the nuclei. The cells were mounted in 70% glycerol for confocal microscopy. The primary antibodies used were mouse anti-E-cadherin (BD Transduction Labs, 610181) and mouse anti-ZO-1 (Invitrogen, 339100). Secondary antibody was Alexa anti-mouse 568. Detection of the staining for the chamber slides was captured on a Leica LAS AF SP5 microscope using a 20× HCX Plan Apo NA 0.7 IMM objective.

Free-floating organoids were fixed in 4.0% paraformaldehyde in PBS⁼ and permeabilized with 0.2% PBST (PBS⁼ with 0.2% Triton X-100). Primary antibody was E-cadherin and rabbit anti-Wnt2B (abcam, ab50575). Secondary antibodies were Alexa anti-mouse 594 and Alexa anti-rabbit 488. Staining of the free-floating organoids was captured using the Zeiss LSM Meta confocal.

Z-stacks of images was collected and processed in ImageJ v1.43u, then finalized in Adobe Photoshop CS4 v11.0.2 to generate the final image. All images from an experiment were processed in the same way.

RNA extraction, cDNA synthesis and quantitative PCR

Gene expression by qRT-PCR (quantitative reverse transcriptase polymerase chain reaction) was performed as we previously described (Vincan et al., 2005). Briefly, following treatment, the cells were harvested in TRIzol (Invitrogen 15596-018) and total RNA was purified and DNase treated using RNeasy mini-prep columns (Qiagen 74104). cDNA was synthesized from 4 μ g of total RNA using anchored oligodT primers (Promega) and Moloney Murine Leukemia Virus Reverse Transcriptase (M-MLV RT, Promega M1705). Quantitative PCR on the cDNA samples was performed using SYBR green (Invitrogen 4309155) and ABI PRISM 7500 Sequence Detector (Applied Biosystems). The housekeeping gene hydroxymethyl-bilane synthase (HMBS) (Vandesompele et al., 2002) was used to calculate the fold change ($2^{\Delta\Delta CT}$ method) as described (Vincan et al., 2005). Primer sequences are available on request.

For the array analysis, cDNA (four separate monolayer and organoid populations) was synthesized as above and then assayed on the Human Wnt Signaling Pathway Array (Qiagen, PAHS-043YA). HMBS expression was determined by qPCR as above and used to calculate the fold change ($2^{\Delta\Delta CT}$ method) as described (Vincan et al., 2005).

Statistical analysis

Data are expressed as mean \pm SEM, where mean represents number of experiments (three or more). Statistical tests used are Student's t test or Mann-Whitney with Prism6 (GraphPad software) where *P* values of ≤ 0.05 were considered significant.

References

- Bafico A, Liu GZ, Goldin L, Harris V, Aaronson SA. 2004. An autocrine mechanism for constitutive Wnt pathway activation in human cancer cells. *Cancer Cell* 6:497-506.
- Barker N, Ridgway RA, van Es JH, van de Wetering M, Begthel H, van den Born M, Danenberg E, Clarke AR, Sansom OJ, Clevers H. 2009. Crypt stem cells as the cells-of-origin of intestinal cancer. *Nature* 457:608-611.
- Brabletz T, Hlubek F, Spaderna S, Schmalhofer O, Hiendlmeyer E, Jung A, Kirchner T. 2005a. Invasion and metastasis in colorectal cancer: epithelial-mesenchymal transition, mesenchymal-epithelial transition, stem cells and beta-catenin. *Cells Tissues Organs* 179:56-65.
- Brabletz T, Jung A, Hermann K, Gunther K, Hohenberger W, Kirchner T. 1998. Nuclear overexpression of the oncoprotein beta-catenin in colorectal cancer is localized predominantly at the invasion front. *Pathol Res Pract* 194:701-704.

- Brabletz T, Jung A, Reu S, Porzner M, Hlubek F, Kunz-Schughart LA, Knuechel R, Kirchner T. 2001. Variable beta-catenin expression in colorectal cancers indicates tumor progression driven by the tumor environment. *Proc Natl Acad Sci U S A* 98:10356-10361.
- Brabletz T, Jung A, Spaderna S, Hlubek F, Kirchner T. 2005b. Opinion: migrating cancer stem cells - an integrated concept of malignant tumour progression. *Nat Rev Cancer* 5:744-749.
- Caldwell GM, Jones C, Gensberg K, Jan S, Hardy RG, Byrd P, Chughtai S, Wallis Y, Matthews GM, Morton DG. 2004. The Wnt antagonist sFRP1 in colorectal tumorigenesis. *Cancer Res* 64:883-888.
- Chambers AF, Groom AC, MacDonald IC. 2002. Dissemination and growth of cancer cells in metastatic sites. *Nat Rev Cancer* 2:563-572.
- Chen B, Dodge ME, Tang W, Lu J, Ma Z, Fan CW, Wei S, Hao W, Kilgore J, Williams NS, Roth MG, Amatruda JF, Chen C, Lum L. 2009a. Small molecule-mediated disruption of Wnt-dependent signaling in tissue regeneration and cancer. *Nat Chem Biol* 5:100-107.
- Chen S, Borowiak M, Fox JL, Maehr R, Osafune K, Davidow L, Lam K, Peng LF, Schreiber SL, Rubin LL, Melton D. 2009b. A small molecule that directs differentiation of human ESCs into the pancreatic lineage. *Nat Chem Biol* 5:258-265.
- Clevers H, Nusse R. 2012. Wnt/beta-catenin signaling and disease. *Cell* 149:1192-1205.
- Dimitriadis A, Vincan E, Mohammed IM, Roczo N, Phillips WA, Baidur-Hudson S. 2001. Expression of Wnt genes in human colon cancers. *Cancer Lett* 166:185-191.
- Farin HF, Van Es JH, Clevers H. 2012. Redundant sources of wnt regulate intestinal stem cells and promote formation of paneth cells. *Gastroenterology* 143:1518-1529 e1517.
- Fearon ER. 2011. Molecular genetics of colorectal cancer. *Annu Rev Pathol* 6:479-507.
- Flanagan DJ, Pheesse TJ, Barker N, Schwab RH, Amin N, Malaterre J, Stange DE, Nowell CJ, Currie SA, Saw JT, Beuchert E, Ramsay RG, Sansom OJ, Ernst M, Clevers H, Vincan E. 2015. Frizzled7 functions as a Wnt receptor in intestinal epithelial Lgr5(+) stem cells. *Stem Cell Reports* 4:759-767.
- Goss AM, Tian Y, Tsukiyama T, Cohen ED, Zhou D, Lu MM, Yamaguchi TP, Morrisey EE. 2009. Wnt2/2b and beta-catenin signaling are necessary and sufficient to specify lung progenitors in the foregut. *Dev Cell* 17:290-298.
- Gregorieff A, Pinto D, Begthel H, Destree O, Kielman M, Clevers H. 2005. Expression pattern of Wnt signaling components in the adult intestine. *Gastroenterology* 129:626-638.
- He W, Wong SC, Ma B, Ng SS, Lam MY, Chan CM, Au TC, Chan JK, Chan AT. 2011. The expression of frizzled-3 receptor in colorectal cancer and colorectal adenoma. *Journal of Clinical Oncology* 29.
- Hlubek F, Jung A, Kotzor N, Kirchner T, Brabletz T. 2001. Expression of the invasion factor laminin gamma2 in colorectal carcinomas is regulated by beta-catenin. *Cancer Res* 61:8089-8093.
- Holcombe RF, Marsh JL, Waterman ML, Lin F, Milovanovic T, Truong T. 2002. Expression of Wnt ligands and Frizzled receptors in colonic mucosa and in colon carcinoma. *Mol Pathol* 55:220-226.
- Iglesias DM, Hueber PA, Chu L, Campbell R, Patenaude AM, Dziarmaga AJ, Quinlan J, Mohamed O, Dufort D, Goodyer PR. 2007. Canonical WNT signaling during kidney development. *American Journal of Physiology-Renal Physiology* 293:F494-F500.
- Kato M, Hirai M, Sugimura T, Terada M. 1996. Cloning, expression and chromosomal localization of Wnt-13, a novel member of the Wnt gene family. *Oncogene* 13:873-876.
- Kim K, Lu ZF, Hay ED. 2002. Direct evidence for a role of beta-catenin/LEF-1 signaling pathway in induction of EMT. *Cell Biology International* 26:463-476.
- Kinzler KW, Nilbert MC, Su LK, Vogelstein B, Bryan TM, Levy DB, Smith KJ, Preisinger AC, Hedge P, McKechnie D, et al. 1991. Identification of FAP locus genes from chromosome 5q21. *Science* 253:661-665.
- Kirchner T, Brabletz T. 2000. Patterning and nuclear beta-catenin expression in the colonic adenoma-carcinoma sequence. Analogies with embryonic gastrulation. *Am J Pathol* 157:1113-1121.

- Kirikoshi H, Sekihara H, Katoh M. 2001. WNT10A and WNT6, clustered in human chromosome 2q35 region with head-to-tail manner, are strongly coexpressed in SW480 cells. *Biochemical and Biophysical Research Communications* 283:798-805.
- Korinek V, Barker N, Morin PJ, van Wichen D, de Weger R, Kinzler KW, Vogelstein B, Clevers H. 1997. Constitutive transcriptional activation by a beta-catenin-Tcf complex in APC^{-/-} colon carcinoma. *Science* 275:1784-1787.
- Linker C, Lesbros C, Gros J, Burrus LW, Rawls A, Marcelle C. 2005. {beta}-Catenin-dependent Wnt signalling controls the epithelial organisation of somites through the activation of paraxis. *Development* 132:3895-3905.
- Lum L, Clevers H. 2012. Cell biology. The unusual case of Porcupine. *Science* 337:922-923.
- MacDonald IC, Groom AC, Chambers AF. 2002. Cancer spread and micrometastasis development: quantitative approaches for in vivo models. *Bioessays* 24:885-893.
- Madan B, Ke Z, Harmston N, Ho SY, Frois AO, Alam J, Jeyaraj DA, Pendharkar V, Ghosh K, Virshup IH, Manoharan V, Ong EH, Sangthongpitag K, Hill J, Petretto E, Keller TH, Lee MA, Matter A, Virshup DM. 2016. Wnt addiction of genetically defined cancers reversed by PORCN inhibition. *Oncogene* 35:2197-2207.
- Mariadason JM, Bordonaro M, Aslam F, Shi L, Kuraguchi M, Velcich A, Augenlicht LH. 2001. Down-regulation of beta-catenin TCF signaling is linked to colonic epithelial cell differentiation. *Cancer Res* 61:3465-3471.
- Muller T, Bain G, Wang X, Papkoff J. 2002. Regulation of epithelial cell migration and tumor formation by beta-catenin signaling. *Experimental Cell Research* 280:119-133.
- Naumov GN, MacDonald IC, Weinmeister PM, Kerkvliet N, Nadkarni KV, Wilson SM, Morris VL, Groom AC, Chambers AF. 2002. Persistence of solitary mammary carcinoma cells in a secondary site: a possible contributor to dormancy. *Cancer Res* 62:2162-2168.
- Ocana OH, Corcoles R, Fabra A, Moreno-Bueno G, Acloque H, Vega S, Barrallo-Gimeno A, Cano A, Nieto MA. 2012. Metastatic Colonization Requires the Repression of the Epithelial-Mesenchymal Transition Inducer Prrx1. *Cancer Cell*.
- Phelps RA, Broadbent TJ, Stafforini DM, Jones DA. 2009a. New perspectives on APC control of cell fate and proliferation in colorectal cancer. *Cell Cycle* 8:2549-2556.
- Phelps RA, Chidester S, Dehghanizadeh S, Phelps J, Sandoval IT, Rai K, Broadbent T, Sarkar S, Burt RW, Jones DA. 2009b. A two-step model for colon adenoma initiation and progression caused by APC loss. *Cell* 137:623-634.
- Pheesse T, Flanagan D, Vincan E. 2016. Frizzled7: A Promising Achilles' Heel for Targeting the Wnt Receptor Complex to Treat Cancer. *Cancers (Basel)* 8.
- Pheesse TJ, Buchert M, Stuart E, Flanagan DJ, Faux M, Afshar-Sterle S, Walker F, Zhang HH, Nowell CJ, Jorissen R, Tan CW, Hirokawa Y, Eissmann MF, Poh AR, Malaterre J, Pearson HB, Kirsch DG, Provero P, Poli V, Ramsay RG, Sieber O, Burgess AW, Huszar D, Vincan E, Ernst M. 2014. Partial inhibition of gp130-Jak-Stat3 signaling prevents Wnt-beta-catenin-mediated intestinal tumor growth and regeneration. *Sci Signal* 7:ra92.
- Ru MX, Hang ESU, Balraj P, Lim P, Jamal R. 2008. Overexpression of WNT2 and TSG101 genes in colorectal carcinoma. *Tropical Biomedicine* 25:46-57.
- Sagara N, Toda G, Hirai M, Terada M, Katoh M. 1998. Molecular cloning, differential expression, and chromosomal localization of human frizzled-1, frizzled-2, and frizzled-7. *Biochem Biophys Res Commun* 252:117-122.
- Sato T, van Es JH, Snippert HJ, Stange DE, Vries RG, van den Born M, Barker N, Shroyer NF, van de Wetering M, Clevers H. 2011. Paneth cells constitute the niche for Lgr5 stem cells in intestinal crypts. *Nature* 469:415-418.
- Sato T, Vries RG, Snippert HJ, van de Wetering M, Barker N, Stange DE, van Es JH, Abo A, Kujala P, Peters PJ, Clevers H. 2009. Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. *Nature* 459:262-265.

- Suh Y, Yoon CH, Kim RK, Lim EJ, Oh YS, Hwang SG, An S, Yoon G, Gye MC, Yi JM, Kim MJ, Lee SJ. 2013. Claudin-1 induces epithelial-mesenchymal transition through activation of the c-Abl-ERK signaling pathway in human liver cells. *Oncogene* 32:4873-4882.
- Suzuki H, Watkins DN, Jair KW, Schuebel KE, Markowitz SD, Dong Chen W, Pretlow TP, Yang B, Akiyama Y, Van Engeland M, Toyota M, Tokino T, Hinoda Y, Imai K, Herman JG, Baylin SB. 2004. Epigenetic inactivation of SFRP genes allows constitutive WNT signaling in colorectal cancer. *Nat Genet* 36:417-422.
- Thiery JP. 2002. Epithelial-mesenchymal transitions in tumour progression. *Nat Rev Cancer* 2:442-454.
- Ueno H, Shinto E, Kajiwaru Y, Fukazawa S, Shimazaki H, Yamamoto J, Hase K. 2014. Prognostic impact of histological categorisation of epithelial-mesenchymal transition in colorectal cancer. *Br J Cancer* 111:2082-2090.
- Ueno K, Hazama S, Mitomori S, Nishioka M, Suehiro Y, Hirata H, Oka M, Imai K, Dahiya R, Hinoda Y. 2009. Down-regulation of frizzled-7 expression decreases survival, invasion and metastatic capabilities of colon cancer cells. *Br J Cancer* 101:1374-1381.
- Ueno K, Hiura M, Suehiro Y, Hazama S, Hirata H, Oka M, Imai K, Dahiya R, Hinoda Y. 2008. Frizzled-7 as a potential therapeutic target in colorectal cancer. *Neoplasia* 10:697-705.
- Valenta T, Degirmenci B, Moor AE, Herr P, Zimmerli D, Moor MB, Hausmann G, Cantu C, Aguet M, Basler K. 2016. Wnt Ligands Secreted by Subepithelial Mesenchymal Cells Are Essential for the Survival of Intestinal Stem Cells and Gut Homeostasis. *Cell Reports* 15:911-918.
- van de Wetering M, Sancho E, Verweij C, de Lau W, Oving I, Hurlstone A, van der Horn K, Battle E, Coudreuse D, Haramis AP, Tjon-Pon-Fong M, Moerer P, van den Born M, Soete G, Pals S, Eilers M, Medema R, Clevers H. 2002. The beta-catenin/TCF-4 complex imposes a crypt progenitor phenotype on colorectal cancer cells. *Cell* 111:241-250.
- van den Heuvel M, Harryman-Samos C, Klingensmith J, Perrimon N, Nusse R. 1993. Mutations in the segment polarity genes wingless and porcupine impair secretion of the wingless protein. *EMBO J* 12:5293-5302.
- Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F. 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* 3:RESEARCH0034.
- Vincan E. 2004. Frizzled/WNT signalling: the insidious promoter of tumour growth and progression. *Front Biosci* 9:1023-1034.
- Vincan E, Barker N. 2008. The upstream components of the Wnt signalling pathway in the dynamic EMT and MET associated with colorectal cancer progression. *Clin Exp Metastasis* 25:657-663.
- Vincan E, Brabletz T, Faux MC, Ramsay RG. 2007a. A human three-dimensional cell line model allows the study of dynamic and reversible epithelial-mesenchymal and mesenchymal-epithelial transition that underpins colorectal carcinogenesis. *Cells Tissues Organs* 185:20-28.
- Vincan E, Darcy PK, Farrelly CA, Faux MC, Brabletz T, Ramsay RG. 2007b. Frizzled-7 dictates three-dimensional organization of colorectal cancer cell carcinoids. *Oncogene* 26:2340-2352.
- Vincan E, Darcy PK, Smyth MJ, Thompson EW, Thomas RJ, Phillips WA, Ramsay RG. 2005. Frizzled-7 receptor ectodomain expression in a colon cancer cell line induces morphological change and attenuates tumor growth. *Differentiation* 73:142-153.
- Vincan E, Swain RK, Brabletz T, Steinbeisser H. 2007c. Frizzled7 dictates embryonic morphogenesis: implications for colorectal cancer progression. *Front Biosci* 12:4558-4567.
- Vincan E, Whitehead RH, Faux MC. 2008. Analysis of Wnt/FZD-mediated signalling in a cell line model of colorectal cancer morphogenesis. *Methods Mol Biol* 468:263-273.
- Voloshanenko O, Erdmann G, Dubash TD, Augustin I, Metzger M, Moffa G, Hundsrucker C, Kerr G, Sandmann T, Anchang B, Demir K, Boehm C, Leible S, Ball CR, Glimm H, Spang R, Boutros M. 2013. Wnt secretion is required to maintain high levels of Wnt activity in colon cancer cells. *Nature Communications* 4.

Whitehead RH, Jones JK, Gabriel A, Lukies RE. 1987. A new colon carcinoma cell line (LIM1863) that grows as organoids with spontaneous differentiation into crypt-like structures in vitro. *Cancer Res* 47:2683-2689.

Willert K, Brown JD, Danenberg E, Duncan AW, Weissman IL, Reya T, Yates JR, 3rd, Nusse R. 2003.

Wnt proteins are lipid-modified and can act as stem cell growth factors. *Nature* 423:448-452.

Wong SC, Lo ES, Chan AK, Lee KC, Hsiao WL. 2003. Nuclear beta catenin as a potential prognostic and diagnostic marker in patients with colorectal cancer from Hong Kong. *Mol Pathol* 56:347-352.

Acknowledgments

Many thanks to Gavin Mitchell for assistance with the artwork and Damian Neate for technical assistance. Funding is gratefully acknowledged from the following: National Health and Medical Research Council of Australia (NHMRC, 566679 and APP1099302), Melbourne Health project grants (605030 and PG-002), and Early career researcher grant (GIA-033) awarded to DJF, Cancer Council of Victoria project grants (CCV, APP1020716) awarded to EV/TJP, CCV Fellowship awarded to DJF and Cardiff University Research Fellowship awarded to TJP.

Accepted Article

Figure Legends

Fig. 1 A: Representative DIC images of LIM1863-*Mph* cells as mesenchymal monolayer (MONO, EMT), during organoid assembly (MET) and as a cluster of free-floating organoids (ORG) (scale bar is 100 μ). Schematic of a LIM1863-*Mph* organoid cross-section showing polarised epithelial cells around a central lumen. **B:** Wnt gene expression (qRT-PCR, array analysis) in monolayer (Mono) and organoid (ORG) LIM1863-*Mph* cells (mean \pm SEM, n=4 independent cell preparations; *p<0.05). **C:** Wnt target gene expression (qRT-PCR, array analysis) in monolayer (Mono) and organoid (ORG) LIM1863-*Mph* cells (mean \pm SEM, n=4 independent cell preparations; *p<0.05).

Fig. 2. A: Wnt gene expression (Sybr green qRT-PCR) in LIM1863-*Mph* monolayer (EMT), MET and organoid (ORG) (mean \pm SEM, n=4 independent cell preparations; *p<0.05). **B:** Gene expression (Sybr green qRT-PCR) in LIM1863-*Mph* monolayer (EMT), MET and organoid (ORG) (mean \pm SEM, n=4 independent cell preparations; *p<0.05). Laminin γ 2 (LAM γ 2), Claudin1 (CLDN1), c-MYC (MYC), Cyclin D1 (CCND1) and CD44. **C:** Confocal immunofluorescence microscopy for Wnt2B (green) and E-cadherin (red) localisation (nuclei are blue, DAPI stained). The organoid diagram depicts the plane shown in cross section.

Fig. 3. A: Immunoblot to detect β -catenin in lysates prepared from parental L-cells treated with conditioned medium (CM) collected from parental L-cells (L-cell CM), Wnt3A CM prepared in the absence (Wnt3A-CM) or presence of the porcupine inhibitor IWP2 (Wnt3A/IWP2 CM). The arrow indicates β -catenin. **B:** DIC images of control (DMSO) and IWP2 treated LIM1863-*Mph* cells. The circles indicate examples of organoids forming (scale bar is 100 μ). **C:** Gene expression (Sybr green qRT-PCR) in LIM1863-*Mph* monolayer cells treated with DMSO or IWP2 (5 μ M) (mean \pm SEM, n=3 independent cell preparations; *p<0.05). Laminin γ 2 (LAM γ 2), Claudin1 (CLDN1), c-Myc (MYC), Cyclin D1 (CLND1) and CD44.

Fig. 4. Confocal immunofluorescence microscopy for E-cadherin on LIM1863-*Mph* monolayer cultures treated with carrier (**A**), IWP2 (**B**), IWP2 and Wnt2b (**C**), or IWP2 and Wnt3a (**D**). The arrows and circles indicate organoids forming; the “holes” in the monolayer indicate that cells have lifted off the tissue culture plastic to initiate organoid formation (i.e. MET, seen in **A**, **C** and **D**); the square in the merge indicates the area enlarged in the image on the right (scale bars are 50 μ).

Fig. 5. Confocal immunofluorescence microscopy for ZO-1 on LIM1863-*Mph* monolayer cultures treated with carrier (**A**), IWP2 (**B**), IWP2 and Wnt2b (**C**), or IWP2 and Wnt3a (**D**). The circles indicate organoids forming as indicated by the ring of apical ZO-1 staining, the square in the merge indicates the area enlarged in the image on the right. Shown are Z-stacks at the centre of organoids i.e. a cross-section (scale bars are 50 μ).

Fig. 6. Schematic showing the cell morphology transitions in the LIM1863-*Mph* model and the role of FZD7 and Wnt2B. The shading in the nuclei indicates intensity of β -catenin (reference Vincan et. al., 2007b).




Fig.1 A: Representative DIC images of LIM1863-Mph cells as mesenchymal monolayer (MONO, EMT), during organoid assembly (MET) and as a cluster of free-floating organoids (ORG) (scale bar is 100 μ). Schematic of a LIM1863-Mph organoid cross-section showing polarised epithelial cells around a central lumen. B: Wnt gene expression (qRT-PCR, array analysis) in monolayer (Mono) and organoid (ORG) LIM1863-Mph cells (mean \pm SEM, n=4 independent cell preparations; *p<0.05). C: Wnt target gene expression (qRT-PCR, array analysis) in monolayer (Mono) and organoid (ORG) LIM1863-Mph cells (mean \pm SEM, n=4 independent cell preparations; *p<0.05).

55x72mm (300 x 300 DPI)




Fig.2. A: Wnt gene expression (Sybr green qRT-PCR) in LIM1863-Mph monolayer (EMT), MET and organoid (ORG) (mean \pm SEM, n=4 independent cell preparations; *p<0.05). B: Gene expression (Sybr green qRT-PCR) in LIM1863-Mph monolayer (EMT), MET and organoid (ORG) (mean \pm SEM, n=4 independent cell preparations; *p<0.05). Laminin Y2 (LAMY2), Claudin1 (CLDN1), c-MYC (MYC), Cyclin D1 (CCND1) and CD44. C: Confocal immunofluorescence microscopy for Wnt2B (green) and E-cadherin (red) localisation (nuclei are blue, DAPI stained). The organoid diagram depicts the plane shown in cross section.

55x72mm (300 x 300 DPI)




Fig. 3. A: Immunoblot to detect β -catenin in lysates prepared from parental L-cells treated with conditioned medium (CM) collected from parental L-cells (L-cell CM), Wnt3A CM prepared in the absence (Wnt3A-CM) or presence of the porcupine inhibitor IWP2 (Wnt3A/IWP2 CM). The arrow indicates β -catenin. B: DIC images of control (DMSO) and IWP2 treated LIM1863-Mph cells. The circles indicate examples of organoids forming (scale bar is 100 μ). C: Gene expression (Sybr green qRT-PCR) in LIM1863-Mph monolayer cells treated with DMSO or IWP2 (5 μ M) (mean \pm SEM, n=3 independent cell preparations; *p<0.05). Laminin 2Y (LAMY2), Claudin1 (CLDN1), c-Myc (MYC), Cyclin D1 (CLND1) and CD44.

27x67mm (300 x 300 DPI)




Fig. 4. Confocal immunofluorescence microscopy for E-cadherin on LIM1863-Mph monolayer cultures treated with carrier (A), IWP2 (B), IWP2 and Wnt2b (C), or IWP2 and Wnt3a (D). The arrows and circles indicate organoids forming; the “holes” in the monolayer indicate that cells have lifted off the tissue culture plastic to initiate organoid formation (i.e. MET, seen in A, C and D); the square in the merge indicates the area enlarged in the image on the right (scale bars are 50 μ).

112x105mm (300 x 300 DPI)

AC




Fig. 5. Confocal immunofluorescence microscopy for ZO-1 on LIM1863-Mph monolayer cultures treated with carrier (A), IWP2 (B), IWP2 and Wnt2b (C), or IWP2 and Wnt3a (D). The circles indicate organoids forming as indicated by the ring of apical ZO-1 staining, the square in the merge indicates the area enlarged in the image on the right. Shown are Z-stacks at the centre of organoids i.e. a cross-section (scale bars are 50 μ).

112x104mm (300 x 300 DPI)

AC





Fig. 6. Schematic showing the cell morphology transitions in the LIM1863-Mph model and the role of FZD7 and Wnt2B. The shading in the nuclei indicates intensity of β -catenin (reference Vincan et. al., 2007b).

55x21mm (300 x 300 DPI)

Accepted



Wnt/FZD in tumor morphogenesis

162x86mm (300 x 300 DPI)

Accepted

Graphical Abstract (49 words)

The LIM1863-*Mph* cells are a unique tissue culture model of the transitions between mesenchymal (tumour invasive front) and epithelial (tumour centre) colorectal cancer cells. Here we show that endogenous Wnt secretion is necessary for epithelial transition of mesenchymal LIM1863-*Mph* cells and identify a role for Wnt2B in this process.

Accepted Article