

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Machado, PP;Steele, EM;Levy, RB;da Costa Louzada, ML;Rangan, A;Woods, J;Gill, T;Scrinis, G;Monteiro, CA

Title:

Ultra-processed food consumption and obesity in the Australian adult population

Date:

2020-12-01

Citation:

Machado, P. P., Steele, E. M., Levy, R. B., da Costa Louzada, M. L., Rangan, A., Woods, J., Gill, T., Scrinis, G. & Monteiro, C. A. (2020). Ultra-processed food consumption and obesity in the Australian adult population. *Nutrition and Diabetes*, 10 (1), <https://doi.org/10.1038/s41387-020-00141-0>.

Persistent Link:

<https://hdl.handle.net/11343/272235>

License:

[CC BY](#)

ARTICLE

Open Access

Ultra-processed food consumption and obesity in the Australian adult population

Priscila Pereira Machado^{1,2}, Eurídice Martinez Steele², Renata Bertazzi Levy^{2,3}, Maria Laura da Costa Louzada^{2,4}, Anna Rangan⁵, Julie Woods¹, Timothy Gill⁶, Gyorgy Scrinis⁷ and Carlos Augusto Monteiro^{2,4}

Abstract

Background: Rapid simultaneous increases in ultra-processed food sales and obesity prevalence have been observed worldwide, including in Australia. Consumption of ultra-processed foods by the Australian population was previously shown to be systematically associated with increased risk of intakes of nutrients outside levels recommended for the prevention of obesity. This study aims to explore the association between ultra-processed food consumption and obesity among the Australian adult population and stratifying by age group, sex and physical activity level.

Methods: A cross-sectional analysis of anthropometric and dietary data from 7411 Australians aged ≥ 20 years from the National Nutrition and Physical Activity Survey 2011–2012 was performed. Food consumption was evaluated through 24-h recall. The NOVA system was used to identify ultra-processed foods, i.e. industrial formulations manufactured from substances derived from foods and typically added of flavours, colours and other cosmetic additives, such as soft drinks, confectionery, sweet or savoury packaged snacks, microwaveable frozen meals and fast food dishes. Measured weight, height and waist circumference (WC) data were used to calculate the body mass index (BMI) and diagnosis of obesity and abdominal obesity. Regression models were used to evaluate the association of dietary share of ultra-processed foods (quintiles) and obesity indicators, adjusting for socio-demographic variables, physical activity and smoking.

Results: Significant (P -trend ≤ 0.001) direct dose–response associations between the dietary share of ultra-processed foods and indicators of obesity were found after adjustment. In the multivariable regression analysis, those in the highest quintile of ultra-processed food consumption had significantly higher BMI (0.97 kg/m^2 ; 95% CI 0.42, 1.51) and WC (1.92 cm; 95% CI 0.57, 3.27) and higher odds of having obesity (OR = 1.61; 95% CI 1.27, 2.04) and abdominal obesity (OR = 1.38; 95% CI 1.10, 1.72) compared with those in the lowest quintile of consumption. Subgroup analyses showed that the trend towards positive associations for all obesity indicators remained in all age groups, sex and physical activity level.

Conclusion: The findings add to the growing evidence that ultra-processed food consumption is associated with obesity and support the potential role of ultra-processed foods in contributing to obesity in Australia.

Introduction

Non-communicable diseases (NCDs) are estimated to account for 89% of all deaths in Australia¹, and high body mass index (BMI) remains the second greatest risk factor driving most death and disability in the country¹. In the past 20 years, Australian prevalence of obesity has risen dramatically—19% in 1995, 27% in 2012 and 31% in 2018²—and currently has the fifth highest rate of obesity

Correspondence: Carlos Augusto Monteiro (carlosam@usp.br)

¹Institute for Physical Activity and Nutrition, School of Exercise and Nutrition Sciences, Deakin University, Geelong, VIC 3220, Australia

²Center for Epidemiological Research in Nutrition and Health, University of Sao Paulo, Av. Dr. Arnaldo, 715, Sao Paulo 01246-904, Brazil

Full list of author information is available at the end of the article

© The Author(s) 2020



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

among the Organisation for Economic Co-operation and Development countries³.

The increase in global obesity rates appears to be a consequence of changes in global food systems^{4,5}, leading to the displacement of dietary patterns based on traditional meals by those that are increasingly made up of ultra-processed foods⁶. A growing body of evidence from cross-sectional and longitudinal studies conducted worldwide has shown that ultra-processed food consumption is consistently associated with weight gain and obesity^{7–14}.

Ultra-processed foods are defined by the NOVA food classification system as industrial formulations manufactured from substances derived from foods (e.g. modified starch, maltodextrin, hydrogenated oils, protein isolates) and typically added of flavours, colours and other cosmetic additives¹⁵. The poor nutrient profile of these foods (high in salt or added sugar and unhealthy fats and low in dietary fibre, micronutrients and phytochemicals) and the processing itself (altered physical and structural characteristics, removal of water and use of flavours, flavour enhancers, colours and other cosmetic additives) make them intrinsically nutritionally unbalanced, hyperpalatable and habit-forming. They dispense the necessity of culinary preparation and are ubiquitous, which make them convenient and accessible. Their manufacture using low-cost ingredients and the aggressive marketing of these products amplify their market advantages over unprocessed or minimally processed foods and freshly prepared meals. All those factors contribute to the replacement of traditional dietary patterns by others based on ultra-processed foods and also encourage the excessive consumption of energy^{15,16}, conditions potentially related to the increased risk of obesity¹⁶.

A recent randomised controlled trial (RCT) showed that, compared to a diet with no ultra-processed foods, a diet with >80% of ultra-processed foods caused an increase in energy intake of near 500 kcal per day and that, in 2 weeks, participants exposed to the ultra-processed diet gained 0.9 kg while participants exposed to the non-ultra-processed diet lost 0.9 kg¹⁷. Interestingly, this was despite both diets being designed to be matched for energy, macronutrients, fibre and sodium content, suggesting that mechanisms other than nutrient profile might explain observed differences.

Ultra-processed food sales are increasing globally, including in Australia^{13,18}. A previous study based on the Australian 2011–2012 National Nutrition and Physical Activity survey found that the increased dietary share of ultra-processed foods was systematically associated with intakes of nutrients outside levels recommended for the prevention of obesity and other NCDs¹⁹. This same survey simultaneously collected food intake and anthropometric data and thus allows testing of the hypothesis that

increases in the dietary share of ultra-processed foods are associated with increases in the risk of obesity among the adult population, regardless of age group, sex and physical activity level.

Methods

Data source and collection

The data source is the National Nutrition and Physical Activity Survey (NNPAS), a household survey that collects information about the Australian population's health, including anthropometric, food consumption and physical activity data. This survey recruited a random sample of the Australian population obtained by using a complex, stratified, multistage probability cluster sampling design based on the selection of strata, households and people within households. The NNPAS was conducted between May 2011 and June 2012, covering 9519 households where 12,153 Australians were interviewed. Information was obtained about one adult and, where possible, one child aged 2–17 years in each selected household²⁰.

Food consumption

Data on food consumption were collected based on two non-consecutive 24-h dietary recalls using an electronic survey, which guides the interviewer towards preventing the interviewee from forgetting consumption items frequently omitted by those interviewed (USDA Automated Multiple-Pass Method). The first recall was applied through a face-to-face interview ($n = 12,153$) while the second recall ($n = 7735$) was applied via a telephone interview conducted ≥ 8 days after the first interview²⁰.

Energy was estimated based on the Australian Food and Nutrient Database (AUSNUT 2011–2013), which contains information for 5740 foods and beverages consumed during the survey²¹.

Mixed dishes composed of two or more food items were disaggregated using the AUSNUT 2011–2013 Food Recipe File (45% of food codes). Reported single food items and the underlying ingredients of mixed dishes were classified according to the NOVA classification system into the following four groups (and subgroups within these groups): (a) unprocessed or minimally processed foods, e.g. cereals, legumes, vegetables, fruits, milk, meat; (b) processed culinary ingredients, e.g. table salt, table sugar, honey, vegetable oils, butter; (c) processed foods, e.g. canned vegetables in brine, salted or sugared nuts, canned fish, freshly made breads and cheeses; and (d) ultra-processed foods, e.g. mass-produced packaged breads, carbonated soft drinks, confectionery, cookies, breakfast 'cereals', flavoured yoghurts, reconstituted meat products, ready to heat meals and packaged instant soups and noodles¹⁵. More information regarding the application of the NOVA system to AUSNUT 2011–13 and access to the coding can be found elsewhere⁽²⁰⁾.

Obesity indicators

Weight, height and waist circumference (WC) measurements were obtained and registered in the surveys by interviewers using digital scales, vertical stadiometers and a metal tape measure, respectively, following standard measurement techniques²⁰. BMI (weight (kg)/height (m)²) and WC (cm) were used as indicators of adiposity. Obesity was defined as BMI ≥ 30 kg/m²²², and abdominal obesity as WC ≥ 88 cm for women and ≥ 102 cm for men²³.

Covariates

Demographic covariates of interest include age, sex, educational attainment, socio-economic status (assessed with the Socio-Economic Index of Disadvantage for Areas (SEIFA), a ranking based on the relative socio-economic advantage and disadvantage of the location of the household), zones (urbanity of the household location based on the Australian Standard Geographical Classification) and country of birth. Physical activity, based on total minutes undertaken in physical activity for fitness, recreation, sport or (for) transport in last week, and current smoking status were also accounted in the analysis²⁰.

Inclusion and exclusion criteria

The analytical sample was restricted to adults aged 20–85 years. Individuals were included in the analyses if they had complete data for BMI and WC. A total of 9238 participants were in the appropriate age span and eligible to be included in the analyses. Of these, pregnant and lactating women, participants who reported implausible energy intakes (<1st or >99th percentile of energy intake) and individuals with missing data for outcomes were excluded. All individuals presented information for the exposure. Thus the final sample of this study was 7411 (Fig. 1).

Data analysis

The first 24-h recall was used for the analyses. The population was first stratified into quintiles of the dietary share of ultra-processed foods (percentage of total energy intake), with the lowest consumers belonging to the first quintile and the highest consumers to the fifth. Thereafter, the characteristics of participants (demographics, physical activity, smoking status and total energy intake) according to quintiles of ultra-processed food consumption were assessed. Differences in those characteristics across the dietary share of ultra-processed foods were evaluated by Pearson's χ^2 test of independence (categorical variables) and unadjusted linear regression models (treating quintile of ultra-processed food consumption as an ordinal variable).

Linear and logistic regression analyses were performed to assess the association between the dietary contribution of ultra-processed foods (quintiles) and obesity indicators,

i.e. BMI (as a continuous variable and categorised to identify obesity) and WC (continuous and categorised to identify abdominal obesity). For all outcomes, we ran an unadjusted model and thereafter a multivariable model adjusted for sociodemographic variables, physical activity and smoking status. Multivariable adjusted subgroups analysis using ultra-processed food consumption as continuous were performed for age group (20–39, 40–59, ≥ 60 years), sex (male, female) and physical activity level (active, inactive).

Additional multivariable adjusted analysis to account for potential effect of reverse causality in the relationship between ultra-processed food consumption (quintiles) and obesity indicators were performed excluding 2 801 individuals with extreme BMI values, following 'special diets' (i.e. on a diet to lose weight and/or for health reasons) at the time of the survey, or who reported a diagnosis of diabetes, heart disease or kidney disease, which could be associated with long term dietary behaviour change (Fig. 1).


All multivariable regression models were adjusted for sex (male/female), age (continuous), years of education (completed ≤ 9 years including never attended, completed 10–12 years with no graduate degree, completed 12 years with graduate degree), income (SEIFA—quintiles), zones (major cities of Australia, inner regional and other, which includes outer regional, remote and very remote Australia), country of birth (Australia or English country/other), level of physical activity (inactive/active, classified as active when physical activity last week met the 150 min recommended guidelines) and smoking status (never smoked, former smoker and current smoker). The fit of the model was verified by residual distribution plots, which should follow a normal distribution.

Weighted analyses were performed using Stata survey module (version 14) to consider the effect of complex sampling procedures adopted in the NNPAS 2011–2012 and in order to allow extrapolation of results for the Australian population (Stata Corp., College Station, United States).

This study was a secondary analysis using de-identified data from the ABS Basic Confidentialised Unit Record Files, and permission to use the data was obtained. Ethics approval for the survey was granted by the Australian Government Department of Health and Ageing Departmental Ethics Committee in 2011²⁰.

Results

Table 1 describes characteristics of the overall Australian adult population and of strata of this population that correspond to quintiles of the dietary share of ultra-processed foods. Ultra-processed foods represented 38.9% of total energy intake among Australian adults, ranging from 12.7% (range 0–21.7%) in the lowest quintile of



ultra-processed food consumption to 74.2% (range 62.1–100%) in the highest quintile. Compared with participants in the lowest quintile, individuals in the highest quintile of ultra-processed food consumption were younger (Q5 = 40.8 vs. Q1 = 48.4 years, $P < 0.001$), more likely to belong to the poorest SEIFA quintile (22.8 vs. 15.5%, $P < 0.001$), be Australian or from English country (88.0 vs. 70.3%, $P < 0.001$), inactive (53.5 vs. 41.8%, $P < 0.001$), current smoker (26.4 vs. 17.0%, $P < 0.001$) and have higher total energy intake (8 951.8 vs. 8 055.9 kJ, $P < 0.001$) and less likely to be higher educated (17.7 vs. 29.9%, $P < 0.001$) and to live in major cities (68.6 vs. 75.2%, $P = 0.002$) (Table 1).

The mean BMI and WC in the Australian adult population were 27.4 kg/m² and 92.8 cm, respectively (Table 2), whereas the prevalence of obesity and abdominal obesity was 26.5% and 40.2%, respectively (Table 3). Crude and multivariable models showed that the dietary share of ultra-processed foods was significantly associated with higher BMI and WC (Table 2) and greater prevalence of both obesity and abdominal obesity among Australian adults (Table 3) (P -trend ≤ 0.001 for all outcomes). Significant direct dose–response associations between the

dietary share of ultra-processed foods and BMI (Table 2) and obesity (Table 3) were found after adjusting for sociodemographic variables, physical activity and smoking. In the multivariable regression analyses, we observed that those in the highest quintile of ultra-processed food consumption had mean BMI 0.97 kg/m² (95% confidence interval (CI) 0.42; 1.51) and WC 1.92 cm (95% CI 0.57; 3.27) higher compared with those in the lowest quintile of consumption (Table 2). The adjusted odds ratios (ORs) of having obesity and abdominal obesity were, respectively, 1.61 (95% CI 1.27; 2.04) and 1.38 (95% CI 1.10; 1.72) in the top quintile of ultra-processed food consumption in regard to the lowest (Table 3).

Positive associations for all obesity indicators were also observed in stratified analysis across all age groups (though did not reach statistical significance with WC and abdominal obesity among the youngest), sex and physical activity level (Table 4). The association of ultra-processed food consumption on BMI and WC was stronger among people aged ≥ 40 years, female and inactive. The association of ultra-processed food consumption on obesity was stronger among people aged ≥ 60 years, male and inactive and on abdominal obesity

Table 1 Characteristics of the population according to dietary share of ultra-processed foods.

	Dietary share of ultra-processed foods (quintiles) ^a						P value*
	All	Q1	Q2	Q3	Q4	Q5	
<i>Age group (%)</i>							<0.001
20–39 years	38.5	31.8	22.7	25.5	40.9	56.3	
40–59 years	36.4	41.7	38.7	38.0	32.3	28.0	
≥60 years	25.1	26.5	27.6	26.5	26.8	15.7	
<i>Sex (%)</i>							0.493
Male	51.7	50.4	50.7	51.7	53.5	52.9	
Female	48.3	49.6	49.3	48.3	46.5	47.1	
<i>Years of education (%)</i>							<0.001
≤9 years	13.5	13.7	13.3	13.8	14.4	12.1	
10–12 years	61.8	56.4	59.3	62.4	64.1	70.2	
10–12 years with graduate degree	24.7	29.9	27.4	23.8	21.5	17.7	
<i>SEIFA (%)</i>							<0.001
Quintile 1—greater disadvantage	17.7	15.5	16.2	16.6	19.6	22.8	
Quintile 2	19.9	18.7	20.4	19.9	19.9	20.7	
Quintile 3	21.1	19.7	18.9	23.4	23.0	21.4	
Quintile 4	19.1	19.8	21.6	18.8	17.3	16.3	
Quintile 5—greater advantage	22.2	26.3	22.8	21.4	20.2	18.8	
<i>Zones (%)</i>							0.002
Major cities	71.5	75.2	72.9	72.9	66.4	68.6	
Inner regional	19.3	16.5	18.6	18.0	22.5	22.5	
Other	9.2	8.3	8.5	9.1	11.1	8.9	
<i>Country of birth (%)</i>							<0.001
Australia or English country	79.8	70.3	77.1	81.7	86.4	88.0	
Other	20.2	30.0	22.8	18.2	13.5	12.0	
<i>Physical activity level (%)^b</i>							<0.001
Inactive	48.0	41.8	47.1	47.5	53.1	53.5	
Active	52.0	58.2	52.9	52.5	46.9	46.5	
<i>Smoking status (%)</i>							<0.001
Never smoked	49.8	50.4	52.1	51.3	48.2	45.7	
Former smoker	31.8	32.6	33.4	31.8	32.2	27.9	
Current smoker	18.4	17.0	14.5	16.9	19.6	26.4	
<i>Total energy intake (kJ)^c</i>	8421.9	8055.9	8388.7	8376.2	8523.1	8951.8	<0.001

Australian population aged ≥20 years (NNPAS 2011–2012), *n* = 7411.

*P value for continuous variables is estimated through unadjusted linear regression, treating quintile of ultra-processed food consumption as an ordinal variable, and Pearson's χ^2 for categorical variables.

^aPercentage of energy intake from ultra-processed foods. Mean (range): All = 38.9 (0–100); Q1 = 12.7 (0–21.7); Q2 = 28.4 (21.7–34.6); Q3 = 40.3 (34.6–46.6); Q4 = 54.0 (46.6–62.1); Q5 = 74.2 (62.1–100).

^bActive whether physical activity last week met 150 min recommended guidelines.

^c1 kcal = 4.186 kJ.

was stronger among people aged ≥40 years, male and inactive. However, these differences were not statistically significant (Table 4).

Additional analyses considering the potential effect of reverse causality showed an increase in the magnitude of the associations in the fifth quintile of ultra-processed

Table 2 Association of dietary share of ultra-processed foods (% of total energy) with BMI and WC among Australians aged ≥20 years (NNPAS 2011–2012), *n* = 7411.

Quintiles of the dietary contribution of ultra-processed foods (% of total dietary energy) ^a	BMI (kg/m ²)					WC (cm)				
	Mean	Mean difference	(95% CI)	Mean difference, adjusted ^b	(95% CI)	Mean	Mean difference	(95% CI)	Mean difference, adjusted ^b	(95% CI)
Q1 (lowest)	26.7	0.00	Ref.	0.00	Ref.	91.1	0.00	Ref.	0.00	Ref.
Q2	27.3	0.66	(0.21; 1.12)	0.52	(0.07; 0.95)	92.9	1.73	(0.48; 2.97)	1.26	(0.19; 2.33)
Q3	27.6	0.86	(0.38; 1.32)	0.66	(0.20; 1.11)	93.2	2.00	(0.71; 3.29)	1.42	(0.30; 2.54)
Q4	27.9	1.26	(0.77; 1.75)	0.96	(0.47; 1.45)	94.8	3.63	(2.31; 4.94)	2.66	(1.46; 3.87)
Q5 (highest)	27.7	1.06*	(0.50; 1.61)	0.97*	(0.42; 1.51)	92.9	1.77*	(0.29; 3.26)	1.92*	(0.57; 3.27)
Total	27.4	–	–	–	–	92.8	–	–	–	–

BMI body mass index, WC waist circumference, CI confidence interval, Ref. reference group.

**P*-trend ≤ 0.001.

^aSee previous table.

^bAdjusted for sex, age, educational attainment, income, zones, country of birth, level of physical activity and smoking status.

Table 3 Association of dietary share of ultra-processed foods (% of total energy) with obesity and abdominal obesity among Australians aged ≥20 years (NNPAS 2011–2012), *n* = 7411.

Quintiles of the dietary contribution of ultra-processed foods (% of total dietary energy) ^a	Obesity (BMI ≥ 30 kg/m ²)					Abdominal obesity ^b				
	Percent	OR	(95% CI)	OR, adjusted ^c	(95% CI)	Percent	OR	(95% CI)	OR, adjusted ^c	(95% CI)
Q1 (lowest)	20.7	1.00	Ref.	1.00	Ref.	35.1	1.00	Ref.	1.00	Ref.
Q2	26.3	1.36	(1.11; 1.67)	1.29	(1.05; 1.59)	41.0	1.29	(1.07; 1.55)	1.24	(1.02; 1.51)
Q3	27.2	1.43	(1.16; 1.76)	1.33	(1.07; 1.64)	39.7	1.22	(1.01; 1.47)	1.16	(0.95; 1.42)
Q4	29.9	1.62	(1.30; 2.00)	1.44	(1.15; 1.80)	46.2	1.59	(1.31; 1.92)	1.53	(1.24; 1.88)
Q5 (highest)	30.9	1.71*	(1.36; 2.14)	1.61*	(1.27; 2.04)	40.5	1.26*	(1.03; 1.55)	1.38*	(1.10; 1.72)
Total	26.5	–	–	–	–	40.2	–	–	–	–

BMI body mass index, OR odds ratio, CI confidence interval, Ref. reference group.

**P*-trend ≤ 0.001.

^aSee previous table.

^bDefined as waist circumference ≥88 cm for women and ≥102 cm for men.

^cAdjusted for sex, age, educational attainment, income, zones, country of birth, level of physical activity and smoking status.

food consumption (regarding the first) for all obesity indicators in comparison to the multivariable models performed in the full analytical sample. Significant direct dose–response associations between the dietary share of ultra-processed foods and WC and abdominal obesity were observed in these analyses (Table S1).

Discussion

In this nationally representative cross-sectional study, the association of ultra-processed food consumption with obesity among Australian adults was investigated. It was found that higher consumption of ultra-processed foods was significantly associated with greater BMI and WC and greater odds of having obesity and abdominal obesity.

Trend towards positive associations of ultra-processed food consumption and obesity indicators were observed in both men and women and across age groups (but not significantly associated among the youngest age groups) and levels of physical activity.

Australians whose diets were based on ultra-processed foods (>62% of total energy intake) had 0.97 units higher BMI, 1.92 cm greater WC and were 61 and 38% more likely of having obesity and abdominal obesity, respectively, than individuals whose diets were not based on ultra-processed foods (<22% of energy intake). The findings of the present study are supported by existing literature showing a causal relationship between ultra-processed food consumption and weight gain¹⁷. In an

Table 4 Association of dietary share of ultra-processed foods^a with indicators of adiposity by age, sex and physical activity level.

	Percentage of energy intake from ultra-processed foods	BMI (kg/m ²)		WC (cm)		Obesity (BMI ≥ 30 kg/m ²)		Abdominal obesity ^c	
		Mean	Mean difference ^b (95% CI)	Mean	Mean difference ^b (95% CI)	Percent	OR ^b (95% CI)	Percent	OR ^b (95% CI)
Age, years									
20–39	43.4	26.1	0.15 (0.02 to 0.28)	88.2	0.27 (−0.06 to 0.60)	18.5	1.09 (1.02 to 1.16)	24.8	1.05 (0.99 to 1.11)
40–59	36.2	28.1	0.20 (0.08 to 0.32)	94.6	0.59 (0.29 to 0.89)	30.9	1.08 (1.02 to 1.14)	44.6	1.08 (1.03 to 1.14)
≥60	36.2	28.5	0.22 (0.05 to 0.39)	97.6	0.37 (0.02 to 0.77)	32.5	1.11 (1.04 to 1.18)	57.3	1.08 (1.01 to 1.14)
Sex									
Male	39.3	27.7	0.11 (0.02 to 0.21)	97.6	0.31 (0.05 to 0.57)	26.2	1.09 (1.03 to 1.14)	35.4	1.08 (1.03 to 1.13)
Female	38.5	27.1	0.23 (0.10 to 0.35)	87.8	0.47 (0.17 to 0.77)	26.9	1.07 (1.02 to 1.12)	45.2	1.05 (1.01 to 1.09)
Physical activity level									
Inactive	40.7	28.0	0.21 (0.08 to 0.32)	94.9	0.49 (0.20 to 0.78)	31.1	1.09 (1.04 to 1.14)	47.6	1.08 (1.03 to 1.13)
Active	37.3	26.9	0.14 (0.03 to 0.24)	91.0	0.29 (0.02 to 0.56)	22.3	1.07 (1.02 to 1.13)	33.3	1.05 (1.01 to 1.10)

Australian population aged ≥20 years (NNPAS 2011–2012), *n* = 7411.

BMI body mass index, WC waist circumference, OR odds ratio, CI confidence interval.

^aFor an increase of 10% of the proportion of ultra-processed food intake in the diet.^bAdjusted for sex, age, educational attainment, income, zones, country of birth, level of physical activity and smoking status.^cDefined as waist circumference ≥88 cm for women and ≥102 cm for men.

RCT conducted by Hall et al., at the end of 2 weeks participants gained, on average, 0.9 ± 0.3 kg during the ultra-processed diet and lost, on average, 0.9 ± 0.3 kg during the diet based on non-ultra-processed foods.

Also supporting our findings, consumption of ultra-processed foods was found to be associated with 9-year incidence of overweight or obesity in a prospective cohort of Spanish middle-aged adult university graduates⁹, incidence of obesity or of higher weight gain among Brazilian¹⁴ and U.K. adults²⁴ and in cross-sectional studies involving nationally representative sample of adults in the U.S.¹⁰, Canada¹² and Brazil⁸. An ecological study including 19 European countries found a significant positive association between national household availability of ultra-processed foods and national prevalence of obesity among adults¹⁰. Similarly, a study across 80 high- and middle-income countries found a positive association of annual changes in sales per capita of ultra-processed products with adult BMI trajectories¹³.

The mechanisms underlying the association between ultra-processed food consumption and obesity are not fully established and may result from a combination of the obesogenic nutritional profile of these foods²⁵, non-nutritional mechanisms related to the processing itself^{26–28} and the displacement of nutritious unprocessed and minimally processed foods and fresh meals prepared with these foods²⁹.

Population-based studies conducted in several countries have shown that the energy share of ultra-processed foods impacted negatively on the intake of nutrients linked to obesity, such as free or added sugars, total fats, dietary energy density and fibre^{19,30–34}. In Australia, the risk of having diets that do not comply with dietary goals recommended for the prevention of obesity increased linearly across quintiles of dietary share of ultra-processed foods, attaining the astonishing 3.9 higher risk of excessive free sugar intake among the highest consumers of ultra-processed foods¹⁹.

Processing techniques applied in the manufacture of ultra-processed foods, such as the partial or total withdrawal of water, the deconstruction of the original food matrix structure and the use of high amounts of sugar, salt, fats, and cosmetic additives, which enhance oro-sensory properties and energy density of these foods, may increase eating rate (grams consumed per minute) and override endogenous satiety and appetite signalling, thereby resulting in greater overall intake^{17,27,35,36}. Ultra-processed beverages may have an even stronger effect by adding to total energy intake without displacing energy from solid foods, as well as affecting subsequent meals due to incomplete compensatory reduction in energy intake³⁷. Recent evidence indicates that part of this mechanism may be explained by alteration in the gut microbiota^{26,38,39}. In fact, the role of food processing on the gut system has recently been emphasised with evolving evidence showing

the effect of the gut microbiota on energy homeostasis and lipid accumulation of the host, and consequent weight gain and obesity^{26,39,40}. Besides the disruption of the gut–brain satiety signalling, diets based on ultra-processed foods may induce microbiota dysbiosis, gut inflammation and decrease gut barrier function due to the presence of non-caloric artificial sweeteners, emulsifiers, advanced glycation end products and the low content of micro-nutrients and phytosterols in these foods^{26,38,39}.

Ultra-processed foods, such as soft drinks, ready meals, confectionaries and biscuits, are frequently designed to be convenient and able to be consumed anywhere, as snacks rather than as regular meals^{6,16,29}. They are accessible²⁵, affordable⁴¹, aggressively marketed^{42–44} and their portion sizes are increasing over time⁴⁵. These characteristics may also contribute to the displacement of freshly prepared meals and stimulate overconsumption of energy¹⁶.

The consumption of ultra-processed foods is increasing worldwide, already comprising the majority of calories consumed in high-income countries, such as the U.K.³³, the U.S.³² and Canada³¹. Besides weight gain and obesity, the consumption of ultra-processed foods has been associated with increased risk of all-cause mortality^{46–48}, hypertension and cardiovascular diseases^{49,50}, metabolic syndrome⁵¹, cancer⁵², diabetes⁵³, depression⁵⁴ and gastrointestinal disorders⁵⁵.

Given the adverse outcomes related to ultra-processed food consumption, dietary advice and policy actions could be aimed at decreasing consumption of these foods, while promoting the availability, accessibility and affordability of unprocessed and minimally processed foods¹⁶. Furthermore, the role of food processing, in particular of ultra-processed foods in contributing to obesity in Australia, should be considered in the current discussion for an overarching strategy to tackle obesity in the country⁵⁶.

This study has several strengths. To the best of our knowledge, this is the first study to analyse the association of ultra-processed food consumption with obesity in Australia. We used the most up-to-date, individual-level dietary survey data taken from a nationally representative sample of Australian adults, increasing generalisability. The analyses were based on the NOVA food classification system¹⁵, which has been recognised by UN agencies as a relevant approach for linking dietary intake, obesity and NCDs^{16,57,58}. The NOVA system was applied into disaggregated food codes in the Australian food composition database, which enabled determining food processing level based on standardised, objective and clear criteria, reducing the chance of misclassification. The availability of sociodemographic, physical activity and smoking data allowed adjustment for several confounders, to test consistency among population groups and provided novel evidence on group-specific associations. Although BMI is considered a useful tool to assess body mass at the

population level, the inclusion of a second indicator of adiposity (WC) strengthens our findings and adds valuable information on body fat distribution⁵⁹.

Nevertheless, there are also limitations to the interpretation of the findings. First, this is a cross-sectional study, and thus temporality and causality cannot be established. However, the results are biologically plausible and consistent with the randomised controlled trial that has assessed the short-term impact of ultra-processed diets on energy intake and weight gain¹⁷, and with a few longitudinal studies that have assessed the association between the dietary share of ultra-processed foods and the incidence of obesity or of higher weight gain^{9,14,24}. Besides, reverse causality cannot be ruled out. In fact, the magnitude of the associations increased in the sensitivity analyses excluding individuals on 'special diets', with extreme BMI values or with diagnosis of diet-related chronic diseases that may have changed dietary behaviour (Table S1). Although a vast array of potential confounders were controlled for, residual confounding due to unmeasured confounders (e.g. parity, menopause) could explain, at least in part, the observed associations.

In addition, limitations related to the dietary assessment instrument deserve mention. Analyses were based on a single recall and may not represent usual diet, possibly biasing studied association towards the null. Analyses were based on a single rather than two recalls because the lower response rate for the second day (64%) could have introduced sampling bias. Obesity could have changed individuals' health behaviours, including diet, reducing overall intake of ultra-processed foods among obese people, hence attenuating the magnitude of the associations. Misreporting is an inherent potential bias of the 24-h recall. Some studies suggest that foods usually considered unhealthy (e.g. ultra-processed foods like confectionary, cakes, chips) are more likely to be under-reported⁶⁰. However, this may be partly mitigated by having excluded participants who reported implausible energy intakes, and if differential information bias occurred, the associations would be biased towards the null. Finally, the dietary survey and food composition database were not designed specifically to categorise foods according to characteristics of industrial processing, and so some misclassification of foods at the individual level cannot be excluded. However, standardised, objective and clear criteria were considered, plus several independent researchers reviewed the classification, and a conservative approach (assigning lower level of processing) was used in case of uncertainty.

In conclusion, these findings add to the growing evidence that ultra-processed food consumption is associated with increased risk of obesity and support the potential role of ultra-processed foods in contributing to obesity in Australia. Despite the cross-sectional nature of

the study, the results are biologically plausible and underpinned by evidence derived from experimental and cross-sectional and longitudinal studies from several high- and middle-income countries showing similar results. Importantly, this study contributes to the evolving evidence on the role of food processing in adiposity and is the first one to present an association between ultra-processed food consumption and obesity in a nationally representative sample of Australian adults.

Future studies should be extended to populations around the world, to present context-dependent magnitudes and drivers of ultra-processed food consumption and obesity. Mechanistic studies are needed to clarify underlying plausible causal pathways that explain links between food processing and adiposity. This evidence is relevant to inform policy makers and for dietary advice at the population and clinical levels.

Acknowledgements

This work is part of the PhD thesis (2019) of P.P.M. at the Graduate Program in Nutrition in Public Health, School of Public Health, University of Sao Paulo, Sao Paulo, Brazil. This work was supported by the Sao Paulo Research Foundation (FAPESP), grant #2015/14900-9, grant #2016/13168-5 and grant #2017/24601-4 (P.P.M. was a beneficiary of a doctoral scholarship and of a Research Internships Abroad fellowship). FAPESP had no role in the design, analysis or writing of this manuscript.

Author details

¹Institute for Physical Activity and Nutrition, School of Exercise and Nutrition Sciences, Deakin University, Geelong, VIC 3220, Australia. ²Center for Epidemiological Research in Nutrition and Health, University of Sao Paulo, Av. Dr. Arnaldo, 715, Sao Paulo 01246-904, Brazil. ³Departamento de Medicina Preventiva, Faculdade de Medicina, Universidade de São Paulo, Av. Dr. Arnaldo, 455, Sao Paulo 01246-903, Brazil. ⁴Departamento de Nutrição, Faculdade de Saúde Pública, Universidade de São Paulo, Av. Dr. Arnaldo, 715, Sao Paulo 01246-904, Brazil. ⁵School of Life and Environmental Sciences, Charles Perkins Centre, The University of Sydney, Camperdown, NSW 2050, Australia. ⁶Boden Institute of Obesity, Nutrition, Exercise and Eating Disorders, Charles Perkins Centre, The University of Sydney, Camperdown, NSW 2050, Australia. ⁷School of Agriculture and Food, The University of Melbourne, Parkville, Melbourne, VIC 3010, Australia

Conflict of interest

The authors declare that they have no conflict of interest.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Supplementary Information accompanies this paper at (<https://doi.org/10.1038/s41387-020-00141-0>).

Received: 1 March 2020 Revised: 16 July 2020 Accepted: 22 July 2020

Published online: 05 December 2020

References

1. AIHW. Australian Institute of Health and Welfare. Australia's health 2016. Australia's health series no. 15 2016. <https://www.aihw.gov.au/reports/australias-health/australias-health-2016/contents/summary> (2016).
2. Keating, C. et al. Prevalence of class-I, class-II and class-III obesity in Australian adults between 1995 and 2011-12. *Obes. Res. Clin. Pract.* **9**, 553–562 (2015).
3. OECD. Obesity update 2017. www.oecd.org/health/obesity-update.htm (2017).

4. Swinburn, B. A. et al. The global syndemic of obesity, undernutrition, and climate change: the Lancet Commission report. *Lancet* **393**, 791–846 (2019).
5. GBD 2015 Risk Factors Collaborators. Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* **388**, 1659–1724 (2016).
6. Monteiro, C. A., Moubarac, J. C., Cannon, G., Ng, S. W. & Popkin, B. Ultra-processed products are becoming dominant in the global food system. *Obes. Rev.* **14**(Suppl 2), 21–28 (2013).
7. Canella, D. S. et al. Ultra-processed food products and obesity in Brazilian households (2008–2009). *PLoS ONE* **9**, e92752 (2014).
8. Louzada, M. L. et al. Consumption of ultra-processed foods and obesity in Brazilian adolescents and adults. *Prev. Med.* **81**, 9–15 (2015).
9. Mendonça, R. D. et al. Ultraprocessed food consumption and risk of overweight and obesity: the University of Navarra Follow-Up (SUN) cohort study. *Am. J. Clin. Nutr.* **104**, 1433–1440 (2016).
10. Juul, F., Martinez-Steele, E., Parekh, N., Monteiro, C. A. & Chang, V. W. Ultra-processed food consumption and excess weight among US adults. *Br. J. Nutr.* **120**, 90–100 (2018).
11. Monteiro, C. A. et al. Household availability of ultra-processed foods and obesity in nineteen European countries. *Public Health Nutr.* **21**, 18–26 (2018).
12. Nardocci, M. et al. Consumption of ultra-processed foods and obesity in Canada. *Can. J. Public Health* **110**, 4–14 (2018).
13. Vandevijvere, S. et al. Global trends in ultraprocessed food and drink product sales and their association with adult body mass index trajectories. *Obes. Rev.* **20**(Suppl 2), 10–19 (2019).
14. Canhada, S. L. et al. Ultra-processed foods, incident overweight and obesity, and longitudinal changes in weight and waist circumference: the Brazilian Longitudinal Study of Adult Health (ELSA-Brasil). *Public Health Nutr.* **23**, 1076–1086 (2020).
15. Monteiro, C. A. et al. Ultra-processed foods: what they are and how to identify them. *Public Health Nutr.* **22**, 936–941 (2019).
16. Monteiro, C. A., Cannon, G., Lawrence, M., Costa Louzada, M. L. & Pereira Machado, P. *Ultra-Processed Foods, Diet Quality, and Health using the NOVA Classification System* (Food and Agriculture Organization of the United Nations, Rome, 2019).
17. Hall, K. D. et al. Ultra-processed diets cause excess calorie intake and weight gain: an inpatient randomized controlled trial of ad libitum food intake. *Cell Metab.* **30**, 67.e3–77.e3 (2019).
18. Sievert, K., Lawrence, M., Naika, A. & Baker, P. Processed foods and nutrition transition in the Pacific: regional trends, patterns and food system drivers. *Nutrients* **11**, 1328 (2019).
19. Machado, P. P. et al. Ultra-processed foods and recommended intake levels of nutrients linked to non-communicable diseases in Australia: evidence from a nationally representative cross-sectional study. *BMJ Open* **9**, e029544 (2019).
20. Australian Bureau of Statistics. Australian Health Survey: users' guide, 2011–13. <http://www.abs.gov.au/ausstats/abs@nsf/Lookup/4363.0.55.001Chapter2002011-13> (2013).
21. Food Standards Australia New Zealand. AUSNUT 2011–2013 - Food Composition Database. <http://www.foodstandards.gov.au/science/monitoringnutrients/ausnut/pages/default.aspx> (2014).
22. World Health Organization. *Obesity and Overweight* (WHO, 2018).
23. Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults-the Evidence Report. National Institutes of Health. *Obes. Res.* **6**(Suppl 2), 51S–209S (1998).
24. Rauber, F., Chang, K., Vámos, E. P. et al. Ultra-processed food consumption and risk of obesity: a prospective cohort study of UK Biobank. *Eur. J. Nutr.* (2020). <https://doi.org/10.1007/s00394-020-02367-1>.
25. Spiteri, S. A., Olstad, D. L. & Woods, J. L. Nutritional quality of new food products released into the Australian retail food market in 2015 - is the food industry part of the solution? *BMC Public Health* **18**, 222 (2018).
26. Miclotte, L. & Van de Wiele, T. Food processing, gut microbiota and the globesity problem. *Crit. Rev. Food Sci. Nutr.* **60**, 1769–1782 (2019).
27. Fardet, A., Méjean, C., Labouret, H., Andreeva, V. A. & Feron, G. The degree of processing of foods which are most widely consumed by the French elderly population is associated with satiety and glycemic potentials and nutrient profiles. *Food Funct.* **8**, 651–658 (2017).
28. Ludwig, D. S. Technology, diet, and the burden of chronic disease. *JAMA* **305**, 1352–1353 (2011).
29. Monteiro, C. A. et al. The UN decade of nutrition, the NOVA food classification and the trouble with ultra-processing. *Public Health Nutr.* **21**, 5–17 (2018).
30. Parra, D. C. et al. The association between ultra-processed food consumption and the nutrient profile of the Colombian diet in 2005. *Salud Pública Méx.* **61**, 147–154 (2019).
31. Moubarac, J. C., Batal, M., Louzada, M. L., Martinez Steele, E. & Monteiro, C. A. Consumption of ultra-processed foods predicts diet quality in Canada. *Appetite* **108**, 512–520 (2017).
32. Martinez Steele, E., Popkin, B. M., Swinburn, B. & Monteiro, C. A. The share of ultra-processed foods and the overall nutritional quality of diets in the US: evidence from a nationally representative cross-sectional study. *Popul. Health Metr.* **15**, 6 (2017).
33. Rauber, F. et al. Ultra-processed food consumption and chronic non-communicable diseases-related dietary nutrient profile in the UK (2008–2014). *Nutrients* **10**, 587 (2018).
34. Louzada, M. L. D. C. et al. The share of ultra-processed foods determines the overall nutritional quality of diets in Brazil. *Public Health Nutr.* **21**, 94–102 (2018).
35. de Graaf, C. & Kok, F. J. Slow food, fast food and the control of food intake. *Nat. Rev. Endocrinol.* **6**, 290–293 (2010).
36. Forde, C. G., Mars, M. & de Graaf, K. Ultra-processing or oral processing? A role for energy density and eating rate in moderating energy intake from processed foods. *Curr. Dev. Nutr.* **4**, nzaa019 (2020).
37. DellaValle, D. M., Roe, L. S. & Rolls, B. J. Does the consumption of caloric and non-caloric beverages with a meal affect energy intake? *Appetite* **44**, 187–193 (2005).
38. Small, D. M. & DiFeliceantonio, A. G. Processed foods and food reward. *Science* **363**, 346–347 (2019).
39. Zinocker, M. K. & Lindseth, I. A. The Western diet-microbiome-host interaction and its role in metabolic disease. *Nutrients* **10**, 365 (2018).
40. Hamilton, M. K. & Raybould, H. E. Bugs, guts and brains, and the regulation of food intake and body weight. *Int. J. Obes. Suppl.* **6**(Suppl 1), S8–S14 (2016).
41. Moubarac, J. C. et al. International differences in cost and consumption of ready-to-consume food and drink products: United Kingdom and Brazil, 2008–2009. *Glob. Public Health* **8**, 845–856 (2013).
42. Zorbas, C. et al. The frequency and magnitude of price-promoted beverages available for sale in Australian supermarkets. *Aust. NZ J. Public Health* **43**, 346–351 (2019).
43. Kelly, B. et al. Global benchmarking of children's exposure to television advertising of unhealthy foods and beverages across 22 countries. *Obes. Rev.* **20**(Suppl 2), 116–128 (2019).
44. Pulker, C. E., Scott, J. A. & Pollard, C. M. Ultra-processed family foods in Australia: nutrition claims, health claims and marketing techniques. *Public Health Nutr.* **21**, 38–48 (2018).
45. Zheng, M. et al. Typical food portion sizes consumed by Australian adults: results from the 2011–12 Australian National Nutrition and Physical Activity Survey. *Sci. Rep.* **6**, 19596 (2016).
46. Kim, H., Hu, E. A. & Rebholz, C. M. Ultra-processed food intake and mortality in the USA: results from the Third National Health and Nutrition Examination Survey (NHANES III, 1988–1994). *Public Health Nutr.* **22**, 1777–1785 (2019).
47. Schnabel, L. et al. Association between ultraprocessed food consumption and risk of mortality among middle-aged adults in France. *JAMA Intern. Med.* **179**, 490–498 (2019).
48. Rico-Campa, A. et al. Association between consumption of ultra-processed foods and all cause mortality: SUN prospective cohort study. *BMJ* **365**, 11949 (2019).
49. Srour, B. et al. Ultra-processed food intake and risk of cardiovascular disease: prospective cohort study (NutriNet-Sante). *BMJ* **365**, 11451 (2019).
50. Mendonça, R. D. et al. Ultra-Processed food consumption and the incidence of hypertension in a Mediterranean cohort: the Seguimiento Universidad de Navarra Project. *Am. J. Hypertens.* **30**, 358–366 (2017).
51. Martinez Steele, E., Juul, F., Neri, D., Rauber, F. & Monteiro, C. A. Dietary share of ultra-processed foods and metabolic syndrome in the US adult population. *Prev. Med.* **125**, 40–48 (2019).
52. Fiolet, T. et al. Consumption of ultra-processed foods and cancer risk: results from NutriNet-Santé prospective cohort. *BMJ* **360**, k322 (2018).
53. Srour, B. et al. Ultraprocessed food consumption and risk of type 2 diabetes among participants of the NutriNet-Sante prospective cohort. *JAMA Intern. Med.* **180**, 283–291 (2019).
54. Adjibade, M. et al. Prospective association between ultra-processed food consumption and incident depressive symptoms in the French NutriNet-Sante cohort. *BMC Med.* **17**, 78 (2019).

55. Schnabel, L. et al. Association between ultra-processed food consumption and functional gastrointestinal disorders: results from the French NutriNet-Santé cohort. *Am. J. Gastroenterol.* **113**, 1217–1228 (2018).
56. Australian Health Department. *Overweight and Obesity* (AHD, Canberra, 2019).
57. Food and Agriculture Organization of the United Nations & Food Climate Research Network. *Plates, Pyramids, Planet. Developments in National Healthy and Sustainable Dietary Guidelines: A State of Play Assessment* (FAO/University of Oxford, 2016).
58. Pan American Health Organization. *Ultra-Processed Products in Latin America: Trends, Impact on Obesity, Policy Implications* (Pan American Health Organization/World Health Organization, Washington, DC, 2015).
59. Stevens, J., Katz, E. G. & Huxley, R. R. Associations between gender, age and waist circumference. *Eur. J. Clin. Nutr.* **64**, 6–15 (2010).
60. Lafay, L. et al. Does energy intake underreporting involve all kinds of food or only specific food items? Results from the Fleurbaix Laventie Ville Santé (FLVS) study. *Int. J. Obes. Relat. Metab. Disord.* **24**, 1500–1506 (2000).