



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Feng, QY;Griffiths, F;Parsons, N;Gunn, J

Title:

An exploratory statistical approach to depression pattern identification

Date:

2013-02-15

Citation:

Feng, Q. Y., Griffiths, F., Parsons, N. & Gunn, J. (2013). An exploratory statistical approach to depression pattern identification. *Physica A Statistical Mechanics and Its Applications*, 392 (4), pp.889-901. <https://doi.org/10.1016/j.physa.2012.10.025>.

Publication Status:

Accepted manuscript

Persistent Link:

<https://hdl.handle.net/11343/41849>

1 **An Exploratory Statistical Approach to Depression**
2 **Pattern Identification**

3
4 Qing Yi Feng,¹ Frances Griffiths,^{2*} Nick Parsons,² Jane Gunn³

5
6
7 ¹Complexity Science Doctoral Training Centre, University of Warwick, Coventry,
8 United Kingdom

9
10 ²Warwick Medical School, University of Warwick, Coventry, United Kingdom

11
12 ³General Practice and Primary Health Care Academic Centre, The University of
13 Melbourne, Melbourne, Australia

14
15
16
17
18 ***Corresponding author:**

19 Frances Griffiths, Warwick Medical School, University of Warwick, Coventry CV4 7AL,
20 United Kingdom

21 Email: F.E.Griffiths@warwick.ac.uk

22 Fax: 44-24-76528375 Tel: 44-24-76522534

1 Nomenclatures

- \mathbf{x} information set, each set contains n information
- $d\mathbf{x}$ unit of the ensemble
- J information flux
- \bar{J} averaged information flux
- t time
- f constraint
- $S_{\bar{J}}$ information entropy
- c integral constant in Eq. (6-8)
- x a piece of information

2 *Greek Symbols*

- ρ probability density
- α constant in Eq. (5-8)
- β coefficient in Eq. (5-8)
- η constant in Eq. (8)
- γ coefficient in Eq. (8)
- σ coefficient in Eq. (10)
- μ constant in Eq. (10)
- Φ potential function
- λ damping coefficient
- ξ order parameter
- ζ constant in Eq. (13)

1 *Subscripts*

i, j, k, l number to indicate the depressed individual

n total number of information of the depressed individual

1 Abstract

2 Depression is a complex phenomenon thought to be due to the interaction of
3 biological, psychological and social factors. Currently depression assessment uses
4 self-reported depressive symptoms but this is limited in the degree to which it can
5 characterise the different expressions of depression emerging from the complex causal
6 pathways that are thought to underlie depression. In this study, we considered the
7 depressed individual as a subsystem of an open complex system, proposed Generalized
8 Information Entropy (GIE) to represent the general characteristics of information entropy
9 of the system, and then implemented Maximum Entropy Estimates to derive equations for
10 depression patterns. We also introduced a numerical simulation method to process the
11 depression related data obtained by the Diamond Cohort Study which has been underway
12 in Australia since 2005 involving 789 people. Unlike traditional assessment, we obtained
13 a unique value for each depressed individual which gives an overall assessment of the
14 depression pattern. Our work provides a novel way to visualise and quantitatively
15 measure the depression pattern of the depressed individual which could be used for
16 pattern categorisation. This may have potential for tailoring health interventions to
17 depressed individuals to maximize health benefit.

18 *Keywords:* Depression; Pattern Identification; Generalized Information Entropy;
19 Maximum Entropy Estimates; Quantitative analysis; Visual Representation

1 **1 Introduction**

2 Depression is a common emotional disorder [1]. For example, in the U.S. depressive
3 disorders affect approximately 18.8 million adults, or about 9.5 percent of the population
4 age 18 and older, in a given year [2]. According to Australian Government statistics,
5 everyone will at some time in their life be affected by depression – their own or someone
6 else's [3]. The WHO report on mental illness released in 2001 suggests that depression
7 will be the second largest cause of disability burden after heart disease by 2020 [1].

8 Depression remains a problem for primary care practice and research [4]. Studies
9 over the past decade have consistently found low rates of detection in primary care
10 settings, where most depressed individuals go if they seek care [5]. Until recently there
11 has been little research evidence for how to identify those at risk of chronicity and relapse
12 [6]. However, a number of cohort studies [7-9] are now identifying such risk factors in
13 order to develop evidence on how to tailor medical interventions [10], with the aim of
14 preventing chronicity and relapse.

15 In this paper we describe an exploratory study using an analysis technique from
16 complexity science that has potential to characterise patients individually and so
17 eventually contribute to identifying which treatments are most likely to benefit which
18 patients. We conceptualise depression as a pattern of contributing characteristic factors
19 for each individual at each time point for which data is available. In this exploratory
20 paper we describe analysis for a small number of cases of people with depression. The
21 next step would be to apply this analysis approach to a larger number of cases and link
22 the pattern of depression for each individual with appropriate outcomes. This may then

1 pave the way to undertaking research to evaluate interventions for depression where there
2 is close matching and tailoring of interventions to each individual's depression pattern.

3 Complexity sciences has provided prospects of achieving the above aim [11], since
4 depressed individuals can be considered as complex systems adjusting, adapting and
5 sometimes transforming with their environments [10]. Patterns of a given system can be
6 investigated by various simulation methods using nonlinear interactions among elements,
7 such as self-organized criticality (SOC), cellular automata (CA), agent-based models,
8 swarm systems etc [12-14]. These methods provide an understanding of the emergence of
9 patterns by assuming rules of interaction. For some simple or well-studied systems the
10 assumptions are reasonable, but if we take depressed individuals as systems, it is much
11 more difficult to set up regular rules: as with all people, those living with depression are
12 constantly interacting with their context and interacting with other people. They are also
13 changing with their context over time – the things they have done or experienced
14 influences how they are themselves and their relationships with other people. Thus the
15 pattern of the depressed individual emerges through a dynamic play of interactions
16 including feedback within the body as well as relationships with the physical and social
17 environment [15-16].

18 Given the difficulties of rule-based simulations, we examined data from depressed
19 individuals, which included assessments of their experiences, wellbeing and context that
20 are known to be strongly related to depression and which may have a causative
21 relationship in themselves or in combination. Using this data we aimed to identify
22 patterns of depression based on the data about each depressed individual. Assessments of
23 depression currently in use mostly ask about depression symptoms alone. Depressive

1 symptoms are an important aspect of patterns of depression and treatment of symptoms
2 can be beneficial. However our aim was to describe patterns of depression based on a
3 wider selection of associated factors. In our study, since we are dealing with information
4 rather than assumed rules, we used statistical analysis to develop a new approach to
5 depression pattern identification.

6 **2 Methods**

7 *2.1 Statistical modelling for depression pattern*

8 From the perspective of complexity science, the individual is considered as a “case”
9 in the sociological rather than the medical sense, with health and illness as just one aspect
10 of the individual [10]. Traditional epidemiology and clinical trials use statistical
11 approaches where there is typically an underlying assumption that the individual cases, in
12 our example, the people with depressive symptoms, are independent of one another [17].
13 This is a reasonable assumption to make for a great deal of health related research even
14 though it could be argued, for example, that there cannot be complete independence of
15 individuals if they live in the same locality or share the same cultural values.

16 We propose that depressed individuals can be considered as complex systems. Since
17 we cannot isolate depressed individuals from their sociocultural context, we can consider
18 the individuals as subsystems of a larger system, capture all the interactions, and then
19 apply statistical analysis to integrate their information. This analysis approach assumes
20 that individuals living within a locality are not completely independent of each other.

21 What might the system in this study be like? It is not a closed system described by
22 the law of thermodynamics, in which there is no order or pattern existing when

1 information entropy goes to the maximum. Classic dissipative systems using the concept
2 of negentropy, first introduced to conservative systems by Schrödinger to derive ordered
3 patterns [18-19], still separate the research objects from their environments by boundaries
4 and utilize two information entropy terms to describe the system. This makes it difficult
5 to identify patterns in systems that have more complex interactions. Therefore, the system
6 we need must be an open complex system connected to its environment by information
7 flux which should reflect both internal interactions and external connections of the
8 system.

9 Suppose we have collected information related to depression from several people,
10 and these people form, in some sense, an open complex system. In the Diamond Cohort
11 Study which we present here, the participants were recruited from 30 general practices in
12 rural and urban areas of the State of Victoria, Australia. Although the individual cohort
13 members are unlikely to interact directly with each other, we suggest they can be
14 assumed to be part of the same complex system as they experience a similar geophysical
15 environment, share similar social structures/organisations, economy and culture, have
16 access to the same broadcast media and access the same health care system. The complex
17 system can be specified if these people had direct interactions with each other but the
18 system can also be generalized if there exist only indirect interactions among them
19 through living within the same State of Australia. The observed data of the depressed
20 individual is represented by the information set $\mathbf{x}_k(x_{k1}, x_{k2}, \dots, x_{kn})$, where the subscript k
21 indicates the depressed individual.

22 As in classical statistical mechanics [20], we consider that all possible information
23 microstates of the system comprise of a continuous range – an ensemble, and then

1 $d\mathbf{x}_k = dx_{k1} dx_{k2} \dots dx_{kn}$ is a unit of the ensemble. The probability for the state of all units $d\mathbf{x}$ at
 2 time t is $\rho(\mathbf{x}, t) d\mathbf{x}$, here $\rho(\mathbf{x}, t)$ is the distribution function of the ensemble, which satisfies
 3 the normalization condition $\int \rho(\mathbf{x}, t) d\mathbf{x} = 1$. The parameter J combines all possible
 4 information microstates [21-23]:

$$5 \quad J = \eta + \sum_i \gamma_i \mathbf{x}_i + \sum_{ij} \gamma_{ij} \mathbf{x}_i \mathbf{x}_j + \sum_{ijk} \gamma_{ijk} \mathbf{x}_i \mathbf{x}_j \mathbf{x}_k + \sum_{ijkl} \gamma_{ijkl} \mathbf{x}_i \mathbf{x}_j \mathbf{x}_k \mathbf{x}_l + \dots \quad (1)$$

6 Here η is a constant and γ are coefficients which vary with different interactions within or
 7 among \mathbf{x}_k . J is the information flux existing within the unit $d\mathbf{x}$ at time t , the averaged flux
 8 function over all possible information microstates is then:

$$9 \quad \bar{J} = \int \rho(\mathbf{x}, t) J(\mathbf{x}) d\mathbf{x} \quad (2)$$

10 Here \bar{J} represents the general characteristics of information entropy of the system and
 11 we call it the Generalized Information Entropy (GIE).

12 Without boundaries, to fully define a system we need constraints. From the
 13 perspective of GIE, constraints \mathbf{f} including all kinds of conservations of the system can be
 14 generated by certain interactions among information sets \mathbf{x} which can be sufficiently
 15 transformed into one to four order momentum quadratures:

$$16 \quad \langle \mathbf{x}_i \rangle = f_1, \langle \mathbf{x}_i \mathbf{x}_j \rangle = f_2, \langle \mathbf{x}_i \mathbf{x}_j \mathbf{x}_k \rangle = f_3, \langle \mathbf{x}_i \mathbf{x}_j \mathbf{x}_k \mathbf{x}_l \rangle = f_4 \quad (3)$$

17 Normally constraints are expressed by many partial differential equations (PDEs) with
 18 temporal-spatial boundary conditions. However for our system, the constraints are
 19 usually unmeasurable, for example people share the same culture; in this scenario, how
 20 do we formulate the PDEs? Even if the system has measurable constraints, suppose these
 21 people are neighbours and they have regular interactions with each other, given that the
 22 interactions are complex, then the PDEs will be mainly nonlinear making numerical

1 solutions difficult to find [23-24]. The treatments for constraints we use are also different
 2 from the popular simulation methods mentioned above which set interaction rules to
 3 correspond with the non-linear PDEs. Here we believe the constraints can be investigated
 4 from the information sets \mathbf{x} – if these people share the same culture or live in the same
 5 community they must have something in common, which is reflected by their information
 6 set interactions.

7 Given defined information entropy \bar{J} and constraints, we can implement
 8 Maximum-Entropy Estimates, or in other words, use the Maximum Entropy Principle to
 9 derive equations for depression patterns, by this means depression patterns emerge when
 10 the GIE of the whole system goes to a maximum:

$$11 \quad \begin{cases} S_{\bar{J}} = \max \bar{J} = \int \rho(\mathbf{x}, t) J(\mathbf{x}) d\mathbf{x} \\ \langle \mathbf{x}_i \rangle = f_1, \langle \mathbf{x}_i \mathbf{x}_j \rangle = f_2, \langle \mathbf{x}_i \mathbf{x}_j \mathbf{x}_k \rangle = f_3, \langle \mathbf{x}_i \mathbf{x}_j \mathbf{x}_k \mathbf{x}_l \rangle = f_4 \end{cases} \quad (4)$$

12 Equation (4) is actually a self-organized optimization model, but unlike other rule-setting
 13 models (e.g. Generic Algorithm, Ant Colony Optimization) the optimization here is based
 14 on the input information.

15 By using a Lagrange multiplier Eq. (4) yields:

$$16 \quad J(\rho) + \rho J'(\rho) - \alpha - \sum_i \beta_i \mathbf{x}_i - \sum_{ij} \beta_{ij} \mathbf{x}_i \mathbf{x}_j - \sum_{ijk} \beta_{ijk} \mathbf{x}_i \mathbf{x}_j \mathbf{x}_k - \sum_{ijkl} \beta_{ijkl} \mathbf{x}_i \mathbf{x}_j \mathbf{x}_k \mathbf{x}_l - \dots = 0 \quad (5)$$

17 Here α is a constant and β are coefficients which also reflect the interactions. Then:

$$18 \quad J = \alpha + \sum_i \beta_i \mathbf{x}_i + \sum_{ij} \beta_{ij} \mathbf{x}_i \mathbf{x}_j + \sum_{ijk} \beta_{ijk} \mathbf{x}_i \mathbf{x}_j \mathbf{x}_k + \sum_{ijkl} \beta_{ijkl} \mathbf{x}_i \mathbf{x}_j \mathbf{x}_k \mathbf{x}_l + \dots + \frac{c}{\rho} \quad (6)$$

19 Here c is an integral constant. Rearranging Eq. (6) gives:

$$20 \quad \rho = \frac{c}{J - \alpha - \sum_i \beta_i \mathbf{x}_i - \sum_{ij} \beta_{ij} \mathbf{x}_i \mathbf{x}_j - \sum_{ijk} \beta_{ijk} \mathbf{x}_i \mathbf{x}_j \mathbf{x}_k - \sum_{ijkl} \beta_{ijkl} \mathbf{x}_i \mathbf{x}_j \mathbf{x}_k \mathbf{x}_l - \dots} \quad (7)$$

1 Substituting Eq. (1) for the J in Eq (7):

$$2 \quad \rho = \frac{c}{\eta - \alpha + \sum_i (\gamma_i - \beta_i) \mathbf{x}_i + \sum_{ij} (\gamma_{ij} - \beta_{ij}) \mathbf{x}_i \mathbf{x}_j + \sum_{ijk} (\gamma_{ijk} - \beta_{ijk}) \mathbf{x}_i \mathbf{x}_j \mathbf{x}_k + \sum_{ijkl} (\gamma_{ijkl} - \beta_{ijkl}) \mathbf{x}_i \mathbf{x}_j \mathbf{x}_k \mathbf{x}_l + \dots} \quad (8)$$

3 We obtained the ensemble distribution function of the system. Using the power series for
4 e^x :

$$5 \quad e^x = 1 + x + \frac{1}{2!} x^2 + \frac{1}{3!} x^3 + \dots \quad (9)$$

6 We further obtained the potential function of the system is as follows:

$$7 \quad \Phi(\sigma, \mathbf{x}) = \mu + \sum_i \sigma_i \mathbf{x}_i + \sum_{ij} \sigma_{ij} \mathbf{x}_i \mathbf{x}_j + \sum_{ijk} \sigma_{ijk} \mathbf{x}_i \mathbf{x}_j \mathbf{x}_k + \sum_{ijkl} \sigma_{ijkl} \mathbf{x}_i \mathbf{x}_j \mathbf{x}_k \mathbf{x}_l + \dots \quad (10)$$

8 The potential function regulates the asymptotic stabilization of the system. Since the
9 system here is formed by the individuals with certain depressive symptoms, Eq. (10)
10 controls the characteristics of the depression patterns of depressed individuals.

11 According to the connection of potential functions and dynamic evolutionary
12 equations, we deduced the stochastic differential equations (SDEs) of depression pattern
13 from Eq. (10) as:

$$14 \quad \frac{\partial \mathbf{x}_k}{\partial t} = \lambda_k \mathbf{x}_k + S_k(x_1, x_2, \dots, x_n) + F_k(t) \quad (11)$$

15 Here the information entropy term $S_k(x_1, x_2, \dots, x_n)$ is from the non-linear
16 interactions among information. If the interactions are linear which means all information
17 is independent, there will be no information entropy term and then the equation is exactly
18 the Langevin equation. Therefore, the SDEs deduced by our proposed modelling
19 methodology are the generalized Langevin equations, which are a modified form of the
20 classical Langevin equations. They represent the dynamical evolution of patterns in a
21 system, and the stochastic force $F(t)$, which also appears in the Langevin equations,

1 often makes the pattern jump from one state to another [25].

2 To derive a potential function related to patterns, we mortify Eq. (10) and
3 diagonalize the constant in second order:

$$4 \quad \xi_k = \sum_{i=1}^n a_{ki} x_{ki} \quad (12)$$

5 Here α are coefficients directly related the information x_{ki} of the depressed individual and
6 the logic chain between γ and α is: $\gamma \Rightarrow \alpha$. Since γ represents the possible information
7 microstates and α information interactions, the logic chain we obtain through statistical
8 modelling sets up a bridge connecting microscopic dynamics and macroscopic
9 characteristics which ensure a more reliable way to identify patterns from observable
10 information rather than assumed rules. And the potential function then becomes:

$$11 \quad \Phi(\lambda, \xi) = \zeta + \sum_k \lambda_k \xi_k^2 + \dots \quad (13)$$

12 As the main parameter in the potential function, ξ_k is an order parameter defined in
13 Synergetics by Haken [26] which represents the emerged patterns of the system. From the
14 transformation equation (12), it is clear that ξ_k is a combination of an individual's
15 information related to depression, thus the value of ξ_k can be interpreted as a depression
16 pattern value and the corresponding a_{ki} are the weights or the contribution of information
17 x_{ki} to his/her depression pattern. Thus, with n different kinds of information related to
18 depression in the observed data for each depressed individual in the complex system, individual
19 k has an information set $\mathbf{x}_k(x_{k1}, x_{k2}, \dots, x_{kn})$ which determines the characteristics of the individual's
20 depression pattern; since we know the weight a_{ki} of information variable x_{ki} by maximizing the
21 GIE of the whole system, we can then characterize the individual's depression by the combination
22 of information variable x_{ki} – the depression pattern value ξ_k .

1 The potential function equation (13) represents a physical picture of how the system
 2 reaches optimization when we implement maximum-entropy estimation based on GIE:
 3 when the system is in the process of information microstates interacting or allocating
 4 information flux, two kinds of patterns will exist: 1) inactive patterns ($\lambda_k < 0$) which cannot
 5 obtain enough information flux in the line of least resistance and will be eliminated in the
 6 optimization; 2) active patterns ($\lambda_k > 0$) which can obtain sufficient information flux and
 7 make the whole system trend towards the optimization state and will survive and emerge.

8 **2.2 Simulation Algorithms**

9 Self-Organized Feature Maps (SOMs), a kind of artificial neural network (ANN),
 10 are an unsupervised learning mechanism dependent on non-linear interactions among
 11 neurons competing for activation and a “winner-take-all” strategy: only the neuron with
 12 the highest activation stays active while all other neurons shut down, with other
 13 variations allowing more than one neuron to be active also existing. The computational
 14 principles of SOMs and the maximum-entropy estimate based on GIE are completely
 15 equivalent, introducing non-linear connections and optimization to the complex system.
 16 Therefore a SOM is an ideal simulation algorithm to analyze depression pattern
 17 quantitatively.

18 In detail, we take the pre-treated information set $\mathbf{x}_k(x_{k1}, x_{k2}, \dots, x_{kn})$ of the depressed
 19 individual k as input to a SOM network. At the beginning, the SOM network has a group
 20 of arbitrary numbers x_i^r (or random connection weights w_{ij} among artificial neurons,
 21 where w_{ij} is equivalent to the connection among information x_i represented by a_{ki} in Eq.
 22 (5)) as the original random state of information x_i . In order to implement the optimization,
 23 the SOM network calculates the mean Euclidean distance d between x_i and x_i^r , the

1 smallest distance is the victorious state with corresponding connection weight w_{ij} . SOM
2 networks use positive feedback to adjust x_i^r and repeat this process which is called
3 “training”. Training, usually several hundreds of steps when the whole network tends
4 towards stability, results in a final set of connection weights w_{ij} that reflect the
5 contribution of information x_i to the emerged depression pattern of the depressed
6 individual.

7 **2.3 Data source**

8 The data used is the cleaned data from the Diamond Cohort Study, a large
9 longitudinal study approved by The University of Melbourne’s Human Research Ethics
10 Committee, conducted in Victoria Australia, which commenced in 2005. **Figure 1**
11 summarises the data collection process: 1) 30 general practices (GPs) in Victoria were
12 recruited from a randomly selected list of 200 GPs provided by the Health Insurance
13 Commission. GPs were eligible to participate if they had seen at least 600 individuals
14 aged 18–75 years in the previous year; were able to generate a computerised list of the
15 details of these people; agreed to complete a survey; and were the only GP in their
16 practice to take part in the study. 2) Practice staff supported by research assistants
17 searched the GP’s computerized records to randomly sample 600 eligible individuals
18 from each of the 30 GPs. In total almost 18,000 individuals were sent a screening survey.
19 Individuals were eligible for the screening survey if they: were able to read English; were
20 not terminally ill; were aged 18–75 years; and did not reside in a nursing home. 3) 7,667
21 individuals (43%) returned a completed screening survey. 789 individuals with ‘probable
22 clinical depression’ who declared an interest in hearing more about the study and had
23 provided their first name and telephone number recruited into cohort. 4) Cohort

1 participants were then required to complete a baseline postal survey and follow-up postal
2 surveys were sent at 3, 6, 9 and 12 months. Computer-assisted telephone interviews were
3 undertaken with participants at baseline and 12 months. A full description of data
4 collection is detailed elsewhere [9, 27].

5 ***2.4 Data selection and pre-treatment***

6 Based on our proposed model, the more exhaustive the information available for the
7 depressed individual, the more accurately the patterns will be identified. However, in
8 practice we cannot collect complete information about a system. For example, a
9 questionnaire cannot cover all aspects of the participant, their life and context. However
10 comprehensive a questionnaire aims to be, participants do not answer all the questions.
11 We suggest that for our objective, to identify patterns of depression and how they change
12 over time, it might only be necessary to have information about each individual that is
13 known to be important in relation to depression. From the variables collected as part of
14 the Diamond Cohort study, we selected 14 factors known to be strongly related to
15 depression [6-7] as the information set \mathbf{x} , shown in **Tab.1**. Some variables were used as
16 collected from participants e.g. "employment status" but others were formed by
17 combining a number of variables, e.g. the value of "total exercise" is the sum of
18 "vigorous exercise" and "moderate exercise".

19 In order to ensure the final depression pattern values consistently quantified the state
20 of depression, we reversed the values of some variables so a higher value of the variable
21 always contributed to increasing the likelihood of depression. For example, in the original
22 data for "social relationship" 100 represented a person with excellent social relationship
23 so this was reversed so that 100 meant this person had no social relationships at all. Also

1 as a requirement for neural network simulation, we rescaled all the data to the interval [0,
2 1].

3 **3 Analysis**

4 Technically by using the methodology described above we can identify the pattern
5 of every depressed person involved in the Cohort since all the participants form a large
6 complex system. In this paper, we present exploratory analysis of information about 28
7 individuals from the cohort. These individuals were chosen for this exploratory analysis
8 as based on a mixed methods longitudinal analysis concurrently underway these cases
9 were known to have a diversity of depression trajectories over one year.

10 ***3.1 Numerical simulation of different systems***

11 By utilizing the baseline data of the chosen 28 depressed individuals we formed a
12 small complex system to identify their depression patterns. Since we chose 14 variables
13 according to Eq. (5) and normalized them into the range of [0 1], the value of ξ_k in this
14 study ranges from 0 to 14 with a higher value indicating a more severe state of depression.
15 We then implemented our proposed simulation algorithms using SOMs by choosing a
16 network of 4×7 and 500 training steps.

17 After numerical simulation, we obtained an evolutionary figure to show how ξ_k
18 values change in the SOM training process and a set of connection weights w_{ij} of
19 variables for the depressed individual. We averaged the last 100 stable w_{ij} values as the
20 contribution of information x_i to the emerged depression pattern represented by a_{ki} in Eq.
21 (5). Since every set of connection weights can be used to identify a unique pattern for the

1 depressed individual, we called this identified pattern the “depression fingerprint” of the
2 depressed individual.

3 Furthermore, from Eq. (11) we know that the shapes of “depression fingerprint” of
4 the depressed individual will change over time. Thus if we consider the depressed
5 individual as one temporal complex system then we can utilize his/her data at different
6 time points to analyze the changes of his/her depression pattern over time. Here for every
7 depressed person we had 5 time points: baseline, 3 month, 6 month, 9 month and 12
8 month.

9 *3.2 Comparison with traditional methods*

10 The 20-item Centre for Epidemiologic Studies Depression Scale (CES-D) is a
11 common measurement of depressive symptoms used in the community, and scores range
12 from 0 to 60 with high scores indicating higher depressive symptoms [28]. We averaged
13 the last 100 steady ξ_k values as the depression pattern value of the depressed individual
14 compared to his/her CES-D score shown in **Tab. 2** and the standard deviation (SD) of ξ_k
15 is also provided to show the stabilities of the emerged patterns.

16 **4 Results**

17 *4.1 Depression pattern features in baseline*

18 It can be seen in **Fig. 2** that after several steps the ξ_k values gradually approach an
19 attractor and remain within a relatively narrow range. Since ξ_k is the order parameter in
20 the system, **Fig. 2** also presents a precise picture of how the depression pattern of the
21 depressed individual emerges through information interactions and gives a visual view of

1 pattern formation dynamics.

2 Unlike the CES-D score, the depression pattern value is a unique value for each
3 depressed individual in the system and gives an overall assessment of the pattern. This
4 depression pattern value could be used for pattern categorization. For example, the
5 individuals labelled as study id 7171 and study id 8051 have the same CES-D score 54,
6 suggesting a similar level of depressive symptoms and as such they may be treated
7 medically in the same manner, while their depression pattern values of 5.45 and 7.05
8 respectively suggests that they probably have very different depression experiences which
9 may require quite different treatment approaches. Conversely; study id 4020 with
10 depression pattern value 5.47 may be classified as similar to the case with study id 7171
11 (depression pattern value 5.45) although their CES-D scores differ by 18 points.

12 In **Fig. 3** we show four depression fingerprints. The blue areas in the circles indicate
13 dysfunction, so the greater the size of the blue area the more dysfunction, the smaller the
14 blue area the less dysfunction. The fingerprints in **Fig. 3 (A)** and **(D)** are for individuals
15 with a similar CES-D score (id 4020: 36 and id 8286: 39) yet the depression pattern value
16 is different (5.47 and 7.46 respectively). The difference in pattern is also clear on viewing
17 the fingerprints. Similarly in **Fig. 3 (B)** and **(C)**, the individuals have the same CES-D
18 score (54) but different pattern values (id 7171: 5.45 and id 8051: 7.05). Again the
19 difference in depression pattern is clear on viewing the fingerprint. The shape of the
20 fingerprint represents the factors contributing to the depression that were included in the
21 analysis. In **Fig. 3 (B)** and **(C)** for example: for study id 7171, social factors (living alone,
22 poor social relationships, lack of community participation), physical factors (poor
23 physical status, lack of exercise) and financial problems (poor income management) are

1 major contributors. For study id 8051 childhood abuse and poor environmental status are
2 prominent.

3 ***4.2 Temporal changes of depression patterns***

4 **Figure 4** shows the temporal changes of depression pattern for depressed
5 individuals together with values comparing with corresponding CES-D scores over 5
6 different time points. The trend of temporal depression value is similar to the one of
7 CES-D score but with less fluctuation.

8 **5 Discussion**

9 ***5.1 Practical application of depression pattern values***

10 There is ongoing controversy and debate about using symptom based diagnostic
11 criteria for depression without taking into account function or other factors which may be
12 associated with a poor outcome [17, 29]. Clinicians usually diagnose depression based on
13 symptoms often supported by a patient-completed questionnaire about symptoms such as
14 the PHQ-9, HADS or the BDI-II [30-33]. Rather than focusing on symptoms we devised
15 a modelling methodology to draw together multiple types of information related to
16 factors strongly associated with increased risk of depression, and which may have
17 causative influence, including functional measures, perceived health status, social
18 measures and risk factors. We introduced a novel way to identify a depression pattern by
19 calculating depression pattern values ξ_k and weights a_{ki} that represents the importance of
20 a certain factor related to depression. We were also able to demonstrate that the
21 depression pattern values obtained varied over time even though some of the variables as

1 measured are unchanging (e.g. having had the experience of childhood abuse). If the data
2 were available a measure of the impact of the childhood experience might provide a
3 variable more sensitive to change. It is arguable whether the 14 variables we used here
4 are sufficient to identify depression patterns with utility for clinical practice. However,
5 these variables were selected to represent a biopsychosocial and functional model of
6 depression causation. There is some evidence that the depression pattern values changed
7 over time with a similar trend to the of CES-D scores. As would be expected, the
8 depression value that includes multiple factors, some of which do not change or change
9 slowly, fluctuates differently to symptoms.

10 ***5.2 Practical application of “depression fingerprint” and the temporal changes***

11 Assessing patients as individuals is part of the traditional art of doctoring [34], and
12 patients often want advice that is individually tailored [35]. Our analytical approach
13 suggests that it is possible to identify for each depressed person within a particular
14 complex system (for example, the community of the State of Victoria, Australia), a
15 unique “depression fingerprint”. The “depression fingerprint” provides both a numerical
16 score and a visual pattern. The fingerprint could potentially be used to tailor interventions
17 to the fingerprint pattern and track change over time for individuals. In research this
18 analysis method may have potential for identifying subgroups of depression.

19 ***5.3 Future research***

20 The results presented in this paper are only preliminary; however the methodology
21 proposed here has shown promise and has clinical relevance. Future research is needed to
22 test the potential for this approach to analysis including analysis of information from
23 large numbers of individuals, evaluation of the temporal changes of depression patterns,

1 and relating the depression patterns to depression outcomes.

2 **6 Conclusions**

3 As a common emotional disorder, depression remains a problem for primary care
4 both in practice and research. Considering different interventions might be appropriate for
5 different types of depression, so it is important to develop evidence on whether an
6 intervention is appropriate to a certain type of depression. In this paper we aimed to
7 identify depression patterns by utilizing the data of depressed individuals about how they
8 functioned and related to their context.

9 This paper suggests that the depressed individual can be considered as subsystems of
10 an open complex system. This complex system is connected with its environment by its
11 information flux which describes the status of the system both internally and externally to
12 the environment. By averaging the information flux function over all possible
13 information microstates, the Generalized Information Entropy has been defined. By
14 implementing Maximum-Entropy Estimation (used the Maximum Entropy Principle) to
15 derive equations for depression patterns and the stochastic differential equations we
16 deduced what we believed to be a modified form of the classical Langevin equations.
17 Based on the proposed model, we also developed a numerical simulation method using
18 Self-Organized Feature Maps (SOMs) to process the depression related data obtained by
19 the Diamond Cohort Study.

20 Our work devised a modelling methodology to integrate the depression information
21 of depressed individuals and introduced a novel way to identify individual depression
22 patterns by calculating depression pattern values and the weights which represent the
23 importance of a certain factor related to depression. We also developed a visual method to

- 1 show a unique “depression fingerprint” of the depressed individual and temporal changes
- 2 of depression pattern which may provide practical applications for tailoring medical
- 3 interventions for individuals with depression.

Acknowledgements

The Diamond study was initiated with pilot funding from the beyondblue Victorian Centre of Excellence and the main cohort has received project grant funding from the National Health and Medical Research Council (ID 299869, 454463, 566511 & 1002908). The one year Computer Assisted Telephone Interview was funded by a Stream 3 grant from the Australian Primary Health Care Research Institute (APHCRI). No funding body had a role in study design; the collection, analysis, and interpretation of data; the writing of the manuscript; or the decision to submit this manuscript for publication. We acknowledge the 30 dedicated GPs, their patients and practice staff for making this research possible. We thank the cohort participants for their ongoing involvement in the study. We also thank the diamond project investigators and project team staff who provided assistance in preparing the data for analysis and providing the data files. Furthermore, we thank Dr. Victoria Mason for editing the paper.

References

- [1] WHO, The World Health Report 2001 - Mental Health: New Understanding, New Hope, World Health Organization, Geneva, 2001.
- [2] NIMH, Depression Research, National Institute of Mental Health, Bethesda, 1999.
- [3] AIHW, National Health Priority Areas Mental Health: A Report Focusing on Depression, Australian Institute of Health and Welfare, Canberra, 1998.
- [4] D. Pilgrim, The survival of psychiatric diagnosis, *Social Science & Medicine* 65 (2007) 536-547.
- [5] RAND Corporation, The Societal Promise of Improving Care for Depression, Rand Corporation, Santa Monica, 2008.
- [6] G. Gilchrist, J. Gunn, Observational studies of depression in primary care: what do we know? *BMC Family Practice* 8 (2007) 1-18.
- [7] H. Herrman, D. L. Patrick, P. Diehr, et al., Longitudinal investigation of depression outcomes in primary care in six countries: the LIDO study, Functional status, health service use and treatment of people with depressive symptoms, *Psychological Medicine* 32 (2002) 889-902.
- [8] J. Gunn, Depression as a chronic and disabling experience, The DIAMOND study: a longitudinal naturalistic study of depression in primary care in Victoria, Australia, *Journal of Affective Disorders* 91 (2010) S21-S22.

- [9] J. M. Gunn, G. P. Gilchrist, P. Chondros P, et al., Who is identified when screening for depression is undertaken in general practice? Baseline findings from the Diagnosis, Management and Outcomes of Depression in Primary Care (diamond) longitudinal study, *The Medical Journal of Australia* 188 (2008) S119-S125.
- [10] F. Griffiths, J. Borkan, D. Byrne, et al., Developing evidence for how to tailor medical interventions for the individual patient, *qualitative health research* 20 (2010) 1629-1641.
- [11] M. Lima, *Visual Complexity: Mapping Patterns of Information*, Princeton Architectural Press, New York, 2011.
- [12] P. Bak, *How Nature Works: The Science of Self-Organized Criticality*, Copernicus, New York, 1996.
- [13] M. I. Rabinovich, A. B. Ezersky, P. Weidman, *The Dynamics of Patterns*, World Scientific, Singapore, 2000.
- [14] J. G. Bazan, J. F. Peters, A. Skowron, Behavioral pattern identification through rough set modelling, *Lecture Notes in Computer Science* 3642 (2005) 688-697.
- [15] C. Battersby, *The Phenomenal Woman: Feminist Metaphysics and the Patterns of Identity*, Routledge, New York, 1998.
- [16] D. Meyers, Feminist perspectives on the self, in: Zalta E N (Ed.), *The Stanford Encyclopedia of Philosophy*, spring 2010 ed., 2004, Available from: <http://plato.stanford.edu/archives/spr2010/entries/feminism-self/>.

- [17]H. Herrman, M. Maj, N. Sartorius, *Depressive Disorders*, third ed., John Wiley & Sons, Chichester, 2009.
- [18]E. Schrodinger, *What is Life? With Mind and Matter and Autobiographical Sketches*, Cambridge University Press, Cambridge, 1992.
- [19]I. Prigogine, *From Being to Becoming*, W. H. Freeman, San Francisco, 1980.
- [20]E. T. Jaynes, Information theory and statistical mechanics, *Physical Review* 106 (1957) 620-630.
- [21]L. Chai, M. Shoji, Self-organization and self-similarity in boiling systems, *Journal of Heat Transfer* 124 (2002) 507-515.
- [22]Q. Feng, L. Chai, A new statistical dynamic analysis on vegetation patterns in land ecosystems, *Physica A: Statistical Mechanics and its Applications* 387 (2008) 3583-3593.
- [23]M. C. Cross, P. Hohenberg, Pattern formation outside of equilibrium, *Reviews of Modern Physics* 65 (1993) 851-1112.
- [24]S. A. Kauffman, *Investigation*, Oxford University Press, New York, 2000.
- [25]L. Chen, L. Chai, A theoretical analysis on self-organized formation of microbial biofilms, *Physica A: Statistical Mechanics and its Applications* 370 (2006) 793-807.
- [26]H. Haken, *Information and Self-Organization*, second enlarged, Springer, Berlin, 2000.
- [27]J. M. Gunn, D. R. Ayton, K. Densley, et al., The association between chronic illness, multimorbidity and depressive symptoms in an Australian primary care

- cohort, *Social Psychiatry and Psychiatric Epidemiology* (2010) DOI: 10.1007/s00127-010-0330-z [Epub ahead of print].
- [28] L. S. Radloff, The CES-D scale: a self-report depression scale for research in the general population, *Applied Psychological Measurement* 1 (1977) 385-401.
- [29] C. Dowrick, *Beyond Depression: A New Approach to Understanding and Management*, Oxford University Press, Oxford, 2004.
- [30] K. Kroenke, R.L. Spitzer, J.B. Williams, The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 16 (2001) 606-613.
- [31] A. S. Zigmond, R.P. Snaith, The hospital anxiety and depression scale. *Acta Psychiatr Scand* 67 (1983) 361-370.
- [32] A. T. Beck, R.A. Steer, R. Ball, et al., Comparison of Beck Depression Inventories-IA and -II in psychiatric outpatients. *J Personal* 67 (1996) 588-597.
- [33] National Institute for Health and Clinical Excellence, *Common mental health disorders: Identification and pathways to care*. CG123. London: National Institute for Health and Clinical Excellence, 2011.
- <http://publications.nice.org.uk/common-mental-health-disorders-cg123/guidance#step-1-identification-and-assessment>
- [34] E. Cassell, *The Nature of Suffering and the Goals of Medicine*, Oxford University Press, Oxford, 2004.
- [35] R. Massé, F. Légaré, The limitations of a negotiation model for perimenopausal women, *Sociology of Health and Illness* 23 (2001) 44-64.

Figure Legends

Fig. 1 The flow chart of the collection of depression related data done by the Diamond Cohort.

Fig. 2 How ξ_k value changes in the SOM training process.

Fig. 3 Four depression fingerprints of depressed individuals (in every fingerprint, the numbers 1-14 stand for the 14 variables in **Tab. 1**, red numbers are the stable values of connection weights w_{ij} of variables, and blue area in every fingerprint shows dysfunction the depressed individual). **(A)** Depression fingerprint for study id 4020. **(B)** Depression fingerprint for study id 7171. **(C)** Depression fingerprint for study id 8051. **(D)** Depression fingerprint for study id 8286.

Fig. 4 The temporal changes of depression pattern for depressed individuals together with values comparing with corresponding CES-D scores over 5 different time points (the subplot figures in each figure are the shapes of depression patterns of the depressed person). **(A)** The temporal changes of depression pattern of study id 4020. **(B)** The temporal changes of depression pattern of study id 7171. **(C)** The temporal changes of depression pattern of study id 8051.

Figures




Fig. 1







Fig. 2




A



B




C




D


Fig. 3



A



B



C

Fig. 4

Table Legends

Tab. 1 14 major factors highly related to depression

Tab. 2 Depression pattern values of the depressed individual at baseline and standard deviations (SD) with CES-D score in the ascending order

Tables

Tab. 1

Variables Number*	Variables Name	Score Range and Meaning	Whether Reversed
		[1, 3]	
1	Employ Status	1 = "Employed/Student" 2 = "Not employed/Not in paid employment" 3 = "Unable to work"	No
		[0, 1]	
2	Live Alone Status	0 = "No" 1 = "Yes"	No
		[1, 5]	
3'	Income Management	1 = "Easily" 2 = "Not to bad" 3 = "Difficult some of the time" 4 = "Difficult all of the time" 5 = "Impossible"	No
		[1, 5]	
3'	Money Meets Needs	1 = "Not at all" 2 = "Slightly" 3 = "Somewhat" 4 = "To a great extent" 5 = "Completely"	Yes

		[0, 100]	
4	Environmental Score	0 = "very poor environmental status" 100 = "excellent environmental status"	Yes
		[0, 100]	
5	Social Relationship	0 = "no social relationship at all" 100 = "excellent social relationship"	Yes
		[0, 100]	
6	Psychological Score	0 = "very poor psychological status" 100 = "excellent psychological status"	Yes
		[0, 100]	
7	Physical Score	0 = "very poor physical status" 100 = "excellent physical status"	Yes
		[0, 30]	
8 ^f	Somatic Symptoms	0 = "no somatic symptoms at all" 30 = "having all the measured somatic symptoms"	Yes
	Absent Days	[0, 273.75]	
8 ^f	Because of Physical Problems	0 = "no absent day at all" 273.75 = "be absent everyday in a period of 9 months"	No
		[0, 10]	
9	Total Exercise	0 = "no exercise at all" 10 = "doing exercise every day"	Yes

		[0, 90] (Baseline survey)	
		[0, 70] (Follow-up surveys)	
10	Community Participation	0 = “no community participation at all” 90 or 70 = “having community participation very often”	Yes
		[0,13] (Baseline survey)	
	Recent Life	[0, 18] (Follow-up surveys)	
11	Positive Impacts	0 = “no positive impacts at all” 13 or 18 = “had every measured positive impacts”	Yes
		[0,13] (Baseline survey)	
	Recent Life	[0, 18] (Follow-up surveys)	
12	Negative Impacts	0 = “no negative impacts at all” 13 or 18 = “had every measured negative impacts”	No
		[0, 3]	
	Total	0 = “no childhood sexual abuse history”	
13	Childhood Sexual Abuse	3 = “had childhood sexual abuse more than twice”	No
		[0, 6]	
	Total	0 = “no childhood physical abuse history”	
14	Childhood Physical Abuse	3 = “had childhood physical abuse more than 5 times”	No

*These numbers (1-14) were also used in **Fig. 3** and **Fig. 4** presenting the same variables.

^fThe variable was used in the numerical simulations of temporal changes of depression fingerprint

ⁱThe variable was used in the numerical simulations of temporal changes of depression pattern

Tab. 2

Study id	Depression pattern value (range 0-14)	SD	CES-D score (range 0-60)
9026	2.16	0.06	1
8155	1.98	0.05	2
8274	2.34	0.10	3
16237	3.41	0.30	5
21178	2.53	0.07	8
11055	4.14	0.14	9
30026	4.04	0.13	11
11067	4.84	0.12	15
13056	4.36	0.18	15
23092	2.98	0.11	15
15135	2.78	0.09	20
8096	3.53	0.16	21
6087	2.94	0.17	23
4108	2.62	0.12	26
26041	5.17	0.15	26
11118	4.09	0.20	27
20038	4.80	0.12	28
22074	4.10	0.13	30
6057	4.71	0.14	31
11088	5.73	0.24	32
4020	5.47	0.24	36
8286	7.46	0.33	39
3062	4.90	0.13	42
32125	5.48	0.28	45
18238	5.98	0.23	47
26087	3.61	0.16	47
7171	5.45	0.15	54
8051	7.05	0.33	54

- Depressed individuals can be considered subsystems of an open complex system
- Individuals with depression can each be characterised with a unique pattern value
- Over time, changes in individual pattern values are similar to traditional measures
- Individual pattern values have potential for assessing and monitoring depression