



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Díaz-Tapia, P;Maggs, CA;Macaya, EC;Verbruggen, H

Title:

Widely distributed red algae often represent hidden introductions, complexes of cryptic species or species with strong phylogeographic structure

Date:

2018-12-01

Citation:

Díaz-Tapia, P., Maggs, C. A., Macaya, E. C. & Verbruggen, H. (2018). Widely distributed red algae often represent hidden introductions, complexes of cryptic species or species with strong phylogeographic structure. *Journal of Phycology*, 54 (6), pp.829-839. <https://doi.org/10.1111/jpy.12778>.

Persistent Link:

<https://hdl.handle.net/11343/284397>

1

2 MS. PILAR DÍAZ TAPIA (Orcid ID : 0000-0003-4680-4867)

3 DR. HEROEN VERBRUGGEN (Orcid ID : 0000-0002-6305-4749)

4

5

6 Article type : Regular Article

7

8

9 Widely distributed red algae often represent hidden introductions, complexes of cryptic  
10 species or species with strong phylogeographic structure

11

12 *Pilar Díaz-Tapia*<sup>2</sup>

13 Coastal Biology Research Group, Faculty of Sciences and Centre for Advanced  
14 Scientific Research (CICA), University of A Coruña, 15071, A Coruña. Spain

15 School of BioSciences, University of Melbourne, Victoria 3010, Australia

16 *Christine A. Maggs*

17 Portaferry Marine Laboratory, Queen's University Belfast, Belfast BT22 1PF, Northern  
18 Ireland

19 *Erasmó C. Macaya*

20 Laboratorio de Estudios Algales (ALGALAB), Departamento de Oceanografía,  
21 Facultad de Ciencias Naturales y Oceanográficas, Universidad de Concepción,  
22 Concepción, Chile

23 Millennium Nucleus Ecology and Sustainable Management of Oceanic Island (ESMOI)  
24 Centro FONDAP de Investigaciones en Dinámica de Ecosistemas Marinos de Altas  
25 Latitudes (IDEAL)

26 *Heroen Verbruggen*

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/jpy.12778-18-070](https://doi.org/10.1111/jpy.12778-18-070)

This article is protected by copyright. All rights reserved

27 School of BioSciences, University of Melbourne, Victoria 3010, Australia

28 <sup>2</sup>Author for correspondence: e-mail pdiaz@udc.es

29 **Running title:** Red algae with wide geographical distributions

30 Editorial Responsibility: K. Müller (Associate Editor)

31 ABSTRACT

32 Despite studies suggesting that most seaweeds are poor dispersers, many red algal  
33 species are reported to have circumglobal distributions. Such distributions have mostly  
34 been based on morphological identifications, but molecular data have revealed a range  
35 of issues with morphologically defined species boundaries. Consequently, the real  
36 distribution of such reportedly circumglobal species must be questioned. In this study,  
37 we analyzed molecular datasets (*rbcL* gene) of nine species in the Rhodomelaceae for  
38 which samples were available from widely spaced geographical locations. Three overall  
39 patterns were identified: 1) species showing strong phylogeographic structure (i.e.,  
40 phylogenetic similarity correlates with geographical provenance), often to the point that  
41 populations from different locations could be considered as different species  
42 (*Lophosiphonia obscura*, *Ophidocladus simpliciusculus*, *Polysiphonia villum* and  
43 *Xiphosiphonia pinnulata*); 2) species with a broad distribution that is explained, in part,  
44 by putative human-mediated transport (*Symphyocladia dendroidea* and *Polysiphonia*  
45 *devoniensis*); and 3) non-monophyletic complexes of cryptic species, most with a more  
46 restricted distribution than previously thought (*Herposiphonia tenella*, *S. dendroidea*  
47 and the *X. pennata* complex that includes the species *X. pinnulata* and *S. spinifera*).  
48 This study shows that widely distributed species are the exception in marine red algae,  
49 unless they have been spread by humans.

50

51 *Key words:* introductions, new record, phylogeography, Rhodomelaceae,  
52 Polysiphonieae, Pterosiphonieae, Herposiphonieae, *rbcL*, species boundaries, species  
53 complexes

54

55 INTRODUCTION

56 Phylogeography of marine organisms is influenced by barriers to dispersal and  
57 geographical distance, as well as by aspects of their life-history, physiology and ecology  
58 (Jackson 1974, Palumbi 1994, Riginos et al. 2011). The dispersal ability of seaweeds is  
59 generally very limited, of the order of tens of meters or less (Santelices 1990, Kinlan  
60 and Gaines 2003, Destombe et al. 2009). However, long-distance dispersal is known in  
61 brown seaweeds with buoyant structures (Fraser et al. 2009, Macaya and Zuccarello  
62 2010), which can act as rafts promoting in turn the dispersal of the epiphytic species  
63 that they host (Fraser et al. 2013, Macaya et al. 2016, López et al. 2017, 2018). Still, a  
64 large proportion of macroalgae are epilithic, so their expected dispersal ability is very  
65 limited and consequently their distribution range is expected to be relatively small.  
66 Paradoxically, many macroalgal species are reported to be very widely or even globally  
67 distributed.

68 Records are usually based only on morphological identification, which can be  
69 inaccurate due to morphological plasticity within species as well as similarity between  
70 cryptic species (e.g., Verbruggen 2014, Schneider et al. 2017). Closer investigation of  
71 material from distant regions using DNA data commonly leads to the discovery of  
72 cryptic species (e.g., Won et al. 2009, Bustamante et al. 2014, Schneider et al. 2017).  
73 Even though studies combining morphological and molecular data are increasing, DNA  
74 databases are still very limited for most algal groups and molecular data are often  
75 available only for some regions. As a consequence, the true distribution of many  
76 seaweed species should be regarded as uncertain. Few studies have reassessed the  
77 distribution of widely reported red algal species using molecular data from a broad  
78 sampling area. Complexes of look-alike species, as well as widely distributed species,  
79 have been detected (Zuccarello et al. 1999, Zuccarello et al. 2002a, Zuccarello and West  
80 2003, Won et al. 2009). Among the widely distributed species, some exhibit high  
81 genetic variability and strong phylogeographic structure that often distinguishes  
82 populations from different basins (Zuccarello et al. 2002a,b, Won et al. 2009). Other  
83 widely distributed species lack phylogeographic signal, suggesting long-distance  
84 dispersal processes by unknown mechanisms (Zuccarello et al. 2002a, Fraser et al.  
85 2013). Therefore, red algal phylogeographic patterns are highly heterogeneous and  
86 depend on evolutionary histories and dispersal abilities.

87 In addition to natural dispersal mechanisms, human-mediated vectors can transport  
88 seaweeds from native areas to other world regions and rapidly alter distribution patterns

89 (Straub et al. 2016). More than 208 red algal species have been considered as introduced  
90 or cryptogenic in one or several regions (Thomsen et al. 2016). Cryptic introductions  
91 are common in the red algae and non-native species often remain unnoticed until  
92 diversity surveys use molecular tools (McIvor et al. 2001, Zuccarello et al. 2002b, Díaz-  
93 Tapia et al. 2013b, 2017a). Considering the low dispersal ability of non-buoyant  
94 epilithic red algae, we hypothesize that the distribution of most truly cosmopolitan  
95 species can be explained by human-mediated transport – which is frequently provided  
96 as a potential explanation for wide distributions of species (Zuccarello et al. 2002a,b,  
97 Fraser et al. 2013).

98 The Rhodomelaceae, with >1,000 recognized species, is the most diverse red algal  
99 family (Guiry and Guiry 2018). It includes numerous examples of widely reported  
100 species and, as in most red algal groups, cryptic diversity is common (e.g., Zuccarello et  
101 al. 2002a, Díaz-Tapia and Bárbara 2013, Bustamante et al. 2014, Savoie and Saunders  
102 2016, Zuccarello et al. 2018). Among the red algae the Rhodomelaceae accounts for the  
103 largest number of introduced species (Williams and Smith 2007). Members of this  
104 family are often major components of algal turfs where canopy-forming brown algae  
105 with buoyant structures are rare as a consequence of the stressful conditions imposed by  
106 the presence of sediment (Airoldi 1998, Díaz-Tapia et al. 2013a). This makes the family  
107 a good candidate to test hypotheses about species distributions and phylogeographic  
108 patterns.

109 The objective of this paper is to reassess the wide reported distributions of nine turf-  
110 forming species of the family Rhodomelaceae using DNA sequences. Using molecular  
111 data from distant locations within each species' reported distribution range, we evaluate  
112 whether these are indeed widely distributed species, analyze the observed  
113 phylogeographic patterns, and consider whether these species may have been introduced  
114 into one or several regions by human activities.

115

## 116 MATERIALS AND METHODS

117 Material of *Herposiphonia tenella*, *Lophosiphonia obscura*, *Ophidocladus*  
118 *simpliciusculus*, *Polysiphonia villum*, *P. devoniensis*, *Symphyocladia spinifera*, *S.*  
119 *dendroidea*, *Xiphosiphonia pennata* and *X. pinnulata* was collected in Norway, United

120 Kingdom, France, Spain, Portugal, Italy, Brazil, Chile, Australia and South Africa  
121 during general sampling surveys of the family Rhodomelaceae (Table S1 in the  
122 Supporting Information). All these species form epilithic turfs, most of them on  
123 intertidal sand-covered rocks (Womersley 2003, Díaz-Tapia and Bárbara 2013).  
124 *Lophosiphonia obscura* was found in North Atlantic brackish water coastal lagoons or  
125 estuaries, and *Symphyocladia dendroidea* was collected in Australian and Chilean  
126 shallow subtidal turfs. Distribution maps of records for these species (Figs. 1-3, lines)  
127 were drawn up based on information available in AlgaeBase and references therein  
128 (Guiry and Guiry 2018).

129 DNA was extracted from silica gel-dried material following Saunders and McDevit  
130 (2012), using the Qiagen DNeasy Plant Mini Kit (Qiagen GmbH, Hilden, Germany) or  
131 the Promega Wizard Magnetic 96 DNA Plant System kit (Promega Corporation,  
132 Madison, USA), following the manufacturer's instructions. PCR amplification was  
133 carried out for the *rbcL* gene using primers F7/RrbcStart, F7/R893 or F57/rbcLrevNEW  
134 (Freshwater and Rueness 1994, Mamoozadeh and Freshwater 2011, Saunders and  
135 Moore 2013), as well as the newly designed primers F2  
136 (TGTCTAACTCTGTAGAACAACGGA), F8 (ACTCTGTAGAASAACGGACAMG),  
137 R1008 (AACTACTACAGTACCAGCATG), R1464  
138 (AACATTAGCTGTTGGAGTTTCYAC) and R1452  
139 (TGGAGTTTCYACRAAGTCAGCTGT). Names of these primers indicate their  
140 position in the *rbcL* gene (e.g., first base of F2 primer corresponds with the second base  
141 of the *rbcL* gene). PCR reactions were performed in a total volume of 25 µl containing  
142 1× MyTaq™ reaction buffer, 0.28 µM of forward and reverse primers, 0.125 units My  
143 Taq™ DNA Polymerase (Bioline, London, UK) and 1 µl template DNA. The PCR  
144 profile consisted of initial denaturation (93°C for 3 min), 35 cycles of denaturation  
145 (94°C for 30 s), primer annealing (45°C for 30 s), and extension (74°C for 90 s) and  
146 final extension (74°C for 5 min). The PCR products were purified and sequenced at  
147 Queen's University of Belfast on an AB3730xl DNA Analyzer (Applied Biosystems,  
148 Foster City, CA, USA) or commercially by Macrogen or the sequencing service of the  
149 University of A Coruña.

150 A total of 128 new *rbcL* sequences were generated in this study and an additional  
151 91 sequences were downloaded from GenBank (Table S1). Length of sequences ranged  
152 from 585 to 1467 (Table S1). Sequences were aligned using Muscle in Geneious 6.1.8  
153 (Kearse et al. 2012). As a first stage, we analyzed these sequences in taxon-rich datasets

154 for the tribes Herposiphoniae, Pterosiphoniae, Polysiphoniae and Streblocladiae to  
155 verify that the target species were monophyletic. Based on the resulting trees, we  
156 selected all sequences corresponding to the species (or group of closely related species)  
157 that are the focus of this paper. These datasets were analyzed species by species using  
158 the unweighted pair group method with arithmetic mean (UPGMA). For two complexes  
159 of non-sister species in our initial taxon-rich trees (*Symphyocladia dendroidea* and  
160 *Herposiphonia tenella*), we included wider species sampling considering the available  
161 data for the respective genera (Table S1). We performed maximum likelihood (ML)  
162 analyses separately for each of the two genera using RAxML 8.1.X (Stamatakis 2014).  
163 GTR-Gamma was used as the nucleotide model and branch support was estimated with  
164 1000 bootstrap replicates. Three species of *Xiphosiphonia* and *Dipterosiphonia* were  
165 selected as the respective outgroups for the *Symphyocladia* and *Herposiphonia* trees  
166 based on our phylogenomic analyses of the major lineages of the Rhodomelaceae (Díaz-  
167 Tapia et al. 2017b).

168

## 169 RESULTS

170 The taxonomy of several of the studied species is complex and details are provided in  
171 Appendix S1 in the Supporting Information.

### 172 *Ophidocladus simpliciusculus*

173 *Ophidocladus simpliciusculus* was collected in four out of the six world regions where it  
174 has been reported (Fig. 1a). The UPGMA analyses included 15 newly determined *rbcL*  
175 sequences and two downloaded from GenBank (Table S1). Sequences comprised four  
176 haplotypes (Fig. 1b): haplotype 1, seven samples from Europe (Atlantic and  
177 Mediterranean); haplotype 2, six samples from Australia; haplotype 3, one sample from  
178 South Africa; and haplotype 4, three samples from Brazil. The South African sample  
179 differed by only 0.1% (1 bp) from the Australian samples, while Brazilian samples were  
180 the most divergent (up to 0.8% and 11 bp) from samples from other regions. Our results  
181 indicate that *O. simpliciusculus* has a unique *rbcL* haplotype in each region, but it has a  
182 strong phylogeographic structure.

183

### 184 *Lophosiphonia obscura*

185 *Lophosiphonia obscura* has been reported in the Atlantic and Indo-Pacific and  
186 sequences are available from Europe and Australia (Fig. 1c). Furthermore, our dataset  
187 also included *Polysiphonia hemisphaerica* from Norway and *P. boldii* from Texas, USA  
188 which may be conspecific as suggested by the low *rbcL* divergence with *L. obscura* (see  
189 Appendix S1). We analyzed six newly determined *rbcL* sequences and two downloaded  
190 from GenBank for *Lophosiphonia obscura*, *Polysiphonia hemisphaerica* and *P. boldii*  
191 (Table S1). Four haplotypes were found (Fig. 1d): 1) four samples from Spain (Atlantic  
192 and Mediterranean) and Norway; 2) a sample from the United Kingdom; 3) a sample  
193 from USA; and 4) two samples from Australia. Atlantic samples differed by 0.1-0.2%  
194 (1-2 bp), while Australian samples were 0.7-0.9% (8-11) divergent from the Atlantic  
195 samples. Our results indicate that the lineage formed by these three taxa is moderately  
196 variable in the North Atlantic, and is clearly separated from the Australian populations.

197

#### 198 *Polysiphonia villum*

199 Molecular data were obtained from two regions where *Polysiphonia villum* (as *P.*  
200 *scopulorum* var. *villum*, see Appendix S1) had previously been reported (Fig. 1e).  
201 Furthermore, it was also sampled in Spain, the French Mediterranean and Australia,  
202 where it is here newly recorded. The 13 sequences determined for *P. villum* and the two  
203 downloaded from GenBank (Table S1) belong to three haplotypes (Fig. 1f): 1) four  
204 samples from Australia, 2) three samples from Brazil and 3) eight samples from the  
205 North Atlantic (North Carolina, Spain and France). Australian samples were 0.5-0.6%  
206 (4-7 bp) divergent from the Atlantic samples and the Brazilian sequences differed by  
207 0.2% (2 bp) from the North Atlantic samples. Thus, *P. villum* shows a clear  
208 phylogeographic structure.

209

#### 210 *Polysiphonia devoniensis*

211 Our dataset included samples from the previously recorded distribution in Atlantic  
212 Europe, as well as from the northwestern Mediterranean (Italy and France), the Adriatic  
213 Sea (Italy) and Victoria (Australia), from where *P. devoniensis* is here recorded for the  
214 first time (Fig. 2a, Table S1). Furthermore, sequences of *P. kapraunii* from North  
215 Carolina were also included in our dataset (see Appendix S1). Analyses including an

216 *rbcL* sequence of *P. kapraunii* from GenBank and 21 newly determined sequences of *P.*  
217 *devoniensis* (Table S1) showed eight haplotypes (Fig. 2b). One haplotype was found in  
218 the northwestern Mediterranean, Atlantic Spain and Australia; one occurred in the  
219 Adriatic Sea and the northwestern Mediterranean; and six haplotypes were each  
220 represented by a single sample (two from Wales, two from the Adriatic Sea, one from  
221 the northwestern Mediterranean and one from North Carolina). The North Carolina  
222 sample identified as *P. kapraunii* was 0.2-0.3% (3-4 bp) divergent from two of the  
223 European samples (PD301 and PD2430). These three samples differ from the others by  
224 sequence divergences of 1-1.4% (12-18 bp), while divergence between the other five  
225 haplotypes is 0.1-1% (1-9 bp). The lineage formed by samples assigned to *P.*  
226 *devoniensis* and *P. kapraunii* has a high genetic diversity and the distribution of  
227 haplotypes lacks geographic structure.

228

#### 229 *Symphyocladia dendroidea* complex

230 Sequences of *Symphyocladia dendroidea* are available from most of the previously  
231 known distribution (British Columbia, California, Chile, Peru, Japan and the  
232 Mediterranean). Some of these sequences were labelled as *Pterosiphonia tanakae* (see  
233 Appendix S1). Furthermore, we collected this species in a Galician marina  
234 (northwestern Spain) and in Australia (Victoria), where it is here recorded for the first  
235 time (Fig. 2c).

236 The *rbcL* data for *Symphyocladia dendroidea* reveal cryptic diversity in the  
237 Americas, as specimens from Peru and Chile and specimens from British Columbia  
238 (here referred as *S. dendroidea* 2) do not constitute a clade (Fig. S1 in the Supporting  
239 Information, Table S1). In addition to these regions, both were recorded in California.  
240 *Symphyocladia dendroidea* is resolved as sister to *S. parasitica* with high support, while  
241 *S. dendroidea* 2 is placed in a moderately supported clade together with *S. brevicaulis*  
242 and *S. baileyi* (Fig. S1). Molecular data show that *S. dendroidea* has a wide distribution  
243 in the Pacific and occurs in some European locations, while *S. dendroidea* 2 is  
244 apparently restricted to Pacific North America.

245 In total, 28 *rbcL* sequences were analyzed for *Symphyocladia dendroidea* (some  
246 sequences labelled as *S. tanakae*, see Appendix S1) including 13 newly determined and  
247 15 downloaded from GenBank (Table S1). The UPGMA dendrogram shows seven

248 haplotypes (Fig. 2d) of which five comprise samples from Pacific South America, one  
249 includes samples from Australia and Japan, and the other consists of samples from  
250 California and Europe. Maximum variability between South American haplotypes is  
251 0.6% (8 bp), and sequence divergence between them and the two other clades is 0.4-1%  
252 (3-9 bp). These levels of *rbcL* variation suggest that this entity may consist of multiple  
253 species or highly differentiated populations.

254  
255 *Xiphosiphonia pennata* complex, including *X. pinnulata* and *Symphyocladia spinifera*  
256 *Xiphosiphonia pennata* has been reported in the Atlantic and Indo-Pacific (Fig 2e) and  
257 this morphological species is a complex of at least three non-sister species. Their  
258 taxonomy has been resolved with the clarification of the identity of *X. pinnulata* and *S.*  
259 *spinifera* that have been misidentified as *X. pennata* (see Appendix S1).

260 At present, 39 *rbcL* sequences (16 newly determined and 23 downloaded from  
261 GenBank) are available for *Symphyocladia spinifera* from California, Pacific South  
262 America, Australia and Korea (Fig. 2f). The UPGMA shows 10 haplotypes of which  
263 four correspond to Korean samples, four to Peruvian samples, one to Australian samples  
264 and one to a Washingtonian sample (Fig. 2f). Sequence divergence among haplotypes is  
265 up to 0.9% (7 bp). Australian samples match the morphological concept of  
266 *Xiphosiphonia pennata*, but our molecular data reveal that none of them grouped with  
267 the European *X. pennata* but instead are mostly closely related to *S. spinifera*.  
268 Therefore, *X. pennata* should be excluded from the recorded Australian flora and  
269 replaced by *S. spinifera*. Interestingly, all the Australian samples belong to a single  
270 haplotype, which contrasts with the four haplotypes found in both Peru and Korea.

271 *Xiphosiphonia pinnulata* sequences were resolved as three haplotypes of which  
272 two were found in European samples and one in Brazilian samples (Fig. 2g). Sequence  
273 divergence among them is up to 0.7% (9 bp) and between the two European clades is up  
274 to 0.3% (3 bp). *X. pennata* was only found in the Atlantic Iberian Peninsula. Therefore,  
275 the widely reported *X. pennata* (as *Pterosiphonia pennata*) is apparently restricted to  
276 European shores. *X. pinnulata* is restricted to the Atlantic, where it has a strong  
277 phylogeographic structure with divergences that may even suggest they are separate  
278 species. *S. spinifera* is restricted to the Pacific and it has a high genetic variability  
279 between regions and within regions in Korea and South America.

280

281 *Herposiphonia tenella* complex

282 In total, 27 *rbcL* sequences were obtained for samples morphologically identified as  
283 *Herposiphonia tenella* from Europe, North America and Queensland (Australia; Fig. 3).  
284 They were analyzed together with the available *rbcL* data for the genus (15 species).  
285 The phylogeny resolved *H. tenella* in seven lineages, four from the Atlantic and three  
286 from Queensland (Fig. 4). Sequence divergence among the lineages was at least 1.9%,  
287 while divergence within them was up to 0.7%. Only two of these lineages were resolved  
288 as sisters (1.9-2.1% sequence divergence), while the others, despite morphological  
289 similarities, were more closely related to other lineages. Thus, *Herposiphonia tenella* is  
290 a large species complex that requires taxonomic revision to better understand its cryptic  
291 diversity and the distribution of the resulting new species. Its type locality is in the  
292 Mediterranean, where three of the four European lineages were collected.

293

294 DISCUSSION

295 *Species complexes*

296 In this work we detected several complexes of non-sister species (*Xiphosiphonia*  
297 *pennata*, *Symphyocladia dendroidea* and *Herposiphonia tenella*). Also, we found  
298 species-level taxa that represent monophyletic lineages containing several haplotypes  
299 that in most cases are distributed in accordance with geographic regions. They could  
300 also be classified as species complexes, as sequence divergences between haplotypes  
301 are often large (up to 1.4%), possible evidence for multiple species. Interpretations of  
302 genetic divergences when delineating species boundaries vary among authors. For  
303 example, *Melanothamnus harveyi/japonicus* and other closely related species have been  
304 interpreted as a single species with an intraspecific variability in the *rbcL* gene  $\leq 2.1\%$   
305 (McIvor et al. 2001, as *Polysiphonia*) or as a species complex in which interspecific  
306 variability in the *rbcL* gene is 0.3-0.7% (Savoie and Saunders 2015, as *Neosiphonia*).  
307 The species concept has been hotly debated, but there is a general consensus that  
308 speciation is a process that takes place when gene flow is interrupted as a consequence  
309 of isolation of populations (Coyne et al. 1988, Leliaert et al. 2014). In the present work,  
310 assessing species boundaries was not always straightforward, and we used information  
311 based on genetic divergences, species distribution and, in one lineage, interbreeding

312 experiments described by Rueness (1973). The first scenario we encountered consists of  
313 species with a variety of haplotypes found in distant locations. Genetic isolation by  
314 distance seems obvious considering our data and, in some cases, where the divergences  
315 between distant populations are relatively large ( $\leq 0.9\%$ ), one might consider them  
316 different species (*Ophidocladus simpliciusculus* from Europe vs. Brazil vs.  
317 Australia/South Africa, *Lophosiphonia obscura* from the North Atlantic vs. Australia,  
318 *Polysiphonia villum* from the Atlantic vs. Australia, and *Xiphosiphonia pinnulata* from  
319 Brazil vs. Europe). However, the low number of samples in some regions or species, as  
320 well as the lack of sampling in other regions where these species were recorded or may  
321 be still unknown precludes a definitive conclusion. Perhaps the observed large sequence  
322 divergences between the lineages within these species would be less evident with larger  
323 datasets. A second scenario is similar to the former, as it consists of species with a  
324 variety of haplotypes, but in this case several haplotypes share the same distribution (*P.*  
325 *devoniensis*, *Symphyclocladia dendroidea*, *S. spinifera*). Thus, despite *rbcL* divergences  
326 among some haplotypes ( $\leq 1.4\%$ ) being even larger than in the previous group ( $\leq$   
327  $0.9\%$ ), whether they are at present reproductively isolated and should be considered as  
328 distinct species is uncertain. Interbreeding experiments may assist to clarify if these  
329 species should be considered as distinct or not. While successful reproduction may have  
330 multiple interpretations (Leliaert et al. 2014), unsuccessful reproduction indicates  
331 reproductive incompatibility. The third scenario we found in this work is represented by  
332 *Lophosiphonia obscura* whose eastern and western Atlantic populations have low  
333 genetic distances (0.1-0.2 %) in the *rbcL* gene, and also in the more variable *cox1*  
334 marker (0.6-1.2 %, HQ412544-5 as *P. hemisphaerica* and *P. boldii*, MF094025).  
335 Despite this, crossing experiments demonstrate that isolates from Texas and from  
336 Norway fail to produce fully fertile progeny (Rueness 1973, as *P. hemisphaerica* and *P.*  
337 *boldii*). This suggests that these two populations are reproductively isolated, and that  
338 divergent selection may be acting on these populations but *rbcL* and *cox1* gene  
339 sequences do not reflect this isolation (Nosil et al. 2009). These three scenarios show  
340 different evolutionary patterns even among closely related species (e.g., *P. villum* vs. *P.*  
341 *devoniensis*). Therefore, application of genetic distances in delineating species  
342 boundaries should be evaluated on a case by case basis. While these are very interesting  
343 issues from a taxonomic perspective, they are not the focus of this paper. From a  
344 phylogeographic point of view, whether these closely related monophyletic lineages are

345 different species or not is of minor importance, because either way they share a  
346 common ancestor from which several genetic entities evolved.

347

#### 348 *Phylogeographic patterns*

349 The paradox between expected dispersal limitation (Santelices 1990, Kilan and Gaines  
350 2003) and wide reported species distributions led us to hypothesize that such widely  
351 distributed species would either have strong phylogeographic structure or were spread  
352 by humans. Our results confirmed these hypotheses and exposed a third scenario, where  
353 the morphologically defined species was in fact a complex of non-sister cryptic species.

354 Three of the species exhibit genetic variability with clear phylogeographic structure  
355 in Australia, and the North and South Atlantic (*Ophidocladus simpliciusculus*,  
356 *Lophosiphonia obscura* and *Polysiphonia villum*). This result is not unexpected  
357 considering that genetic divergence is promoted by the isolation among populations  
358 separated by large geographic distances (Palumbi 1994). However, the observed genetic  
359 divergence is relatively low ( $\leq 0.9\%$ ) considering that Australia and the North and  
360 South Atlantic have been separated since about 80 My (Jordan et al. 2016). Therefore,  
361 rather than this genetic divergence resulting from an 80 My old vicariant evolution,  
362 long-distance dispersal processes acting on a common ancestor and subsequent  
363 divergence into differentiated populations are invoked to explain the observed patterns.  
364 Mechanisms responsible for this long-distance dispersal are obscure considering that  
365 these species either occur in coastal lagoons/estuaries or on sand-covered rocks where  
366 buoyant macroalgae that can act as rafts are rare (Airoldi 1998, Díaz-Tapia et al.  
367 2013a). Molecular data have provided evidence for long-distance dispersal in other red  
368 algal species but mechanisms remain unknown (Zuccarello et al. 2002a, Fraser et al.  
369 2013). The genetic separation among geographically distant lineages may indicate that  
370 long-distance dispersal occurs at a low rate. Alternatively, density-dependent processes  
371 are involved and once a population colonizes a new region it prevents the establishment  
372 of latecomers (Waters et al. 2013). Furthermore, available data for the three species  
373 mentioned above indicate different evolutionary histories and/or dispersal paths. For  
374 instance, in *L. obscura* and *P. villum* the largest sequence divergences are between  
375 Australian and Atlantic populations, whereas in *O. simpliciusculus* the Australian

376 haplotype is relatively close to South African and European haplotypes but the  
377 divergence across the Atlantic (Brazil vs. Europe) is much larger.

378 Several species showed a diversity of haplotypes sharing the same region: the  
379 Pacific *Symphyocladia spinifera* and *S. dendroidea*, as well as the Atlantic *Polysiphonia*  
380 *devoniensis*. The origin of this diversity must be related to processes of isolation that led  
381 to genetic differentiation, followed by local dispersal events. As for *Ophidocladus*  
382 *simpliciusculus*, *Lophosiphonia obscura* and *Polysiphonia villum*, dispersal mechanisms  
383 for *S. spinifera* and *P. devoniensis* are unknown. In contrast, *S. dendroidea* has been  
384 reported growing on stranded holdfasts of the floating alga *Durvillaea antarctica*  
385 (Macaya et al. 2016, López et al. 2017, 2018), which could contribute to dispersal after  
386 genetic differentiation influenced its genetic structure.

387 The disjunct distribution of a second group of species (*Polysiphonia devoniensis*  
388 and *Symphyocladia dendroidea*) can be explained by human-mediated introduction  
389 events. The human transport of species from native (donor) to introduction (recipient)  
390 regions causes the rapid expansion of species' distribution and alters natural  
391 phylogeographic patterns (Straub et al. 2016). The discovery of *P. devoniensis* in  
392 Victoria (Australia), exhibiting a single haplotype that is also present in Europe,  
393 suggests that this species has been introduced into this country from Europe, possibly  
394 Atlantic Spain or the NW Mediterranean. *Symphyocladia dendroidea* was recorded as  
395 an introduced species in the French Mediterranean in 2005 (Boudouresque and  
396 Verlaque 2008, as *P. tanakae*) and our recent discovery of the same haplotype in a  
397 marina in Atlantic Spain probably represents a secondary introduction and suggests that  
398 the species is spreading in Europe via hull fouling. The presence of several genetically  
399 separated lineages of *S. dendroidea* in Pacific South America contrasts with the  
400 occurrence of a single haplotype in Japan, Australia and California. Japan and Australia  
401 have the same haplotype, suggesting that one or both populations could be introduced.  
402 Genetic diversity of seaweeds in the introduced regions is either similar or reduced  
403 relative to the native area (McIvor et al. 2001, Voisin et al. 2005, Provan et al. 2008,  
404 Geoffroy et al. 2016). The finding of diverse haplotypes in the introduced region is  
405 indicative of an introduction involving several haplotypes or multiple introductions,  
406 depending on the phylogeographic structure in the native area (McIvor et al. 2001,  
407 Voisin et al. 2005, Geoffroy et al. 2016). A single haplotype of both *S. dendroidea* and  
408 *P. devoniensis* has been detected in the areas where the introduction of these species is

409 certain, suggesting that their introduction is the result of a single event in which a single  
410 haplotype was involved. However, much more complex scenarios could explain the  
411 observed patterns and a better understanding of the phylogeographic patterns in native  
412 and introduced areas would be needed to elucidate the introduction dynamics.

413 The third group of species analyzed here involved species complexes of non-sister  
414 cryptic species (*Xiphosiphonia pennata* including *X. pinnulata* and *Symphycladia*  
415 *spinifera*; *S. dendroidea*; and *Herposiphonia tenella*). In both cases species found in the  
416 Atlantic and Pacific basins differ, but in addition several species were found with  
417 overlapping distributions in some regions of each basin. Therefore, the distribution of  
418 these widely reported species is much narrower than previously thought. Cryptic algal  
419 species often involve a group of morphologically similar species that are genetically  
420 differentiated, but resolve as a monophyletic group (Zuccarello et al. 2002a, Won et al.  
421 2009, Payo et al. 2013). However, examples of non-monophyletic cryptic “species”  
422 have also been documented in the red algae (Zuccarello et al. 2018). Morphological  
423 similarity among non-monophyletic groups of cryptic species can be explained by  
424 evolutionary convergence, morphological stasis or developmental constraints (Leliaert  
425 et al. 2014, Zuccarello et al. 2018). *X. pennata*, *S. spinifera* and *S. dendroidea* are  
426 placed in a tribe (Pterosiphonieae) with high morphological variation ranging from  
427 filiform to foliose species (Díaz-Tapia et al. 2017b). The body plan of both species is  
428 filiform, among the simplest observed in the tribe, and morphological stasis is a  
429 plausible explanation for their similarity. In the tribe Herposiphonieae all species are  
430 very similar in morphology, with limited differences in their body plans (Díaz-Tapia et  
431 al. 2017b) and the cryptic diversity in *Herposiphonia tenella* might result from  
432 morphological stasis or developmental constraints.

433 Understanding the processes underlying phylogeographic patterns requires the  
434 study of numerous specimens from across the entire distribution of the species. In this  
435 regard, we recognize important limitations in our work that prevent us from fully  
436 elucidating causes of the observed phylogeographic patterns, leading to some tentative  
437 conclusions about the potentially introduced status of some of the analyzed populations.  
438 However, most of the species treated here are rare in all or part of their known  
439 distribution range so improving the datasets would be very difficult. For example,  
440 *Lophosiphonia obscura*, despite being widely reported, is very rare in the regions here  
441 studied: the sample from the UK used in this study is the first one collected since 1970

442 (Maggs and Hommersand 1993). In Spain, we found it only once in the Atlantic and  
443 once in the Mediterranean, and the species is here recorded for only the third time in  
444 Australia. Our work should thus be interpreted as one of the first attempts to understand  
445 phylogeographic patterns of widely distributed red algal species. Even though their  
446 evolutionary history is not well known, our analyses provide clear examples of 1)  
447 species with wide distributions and strong phylogeographic structure that reflects the  
448 geographical distance; 2) species with a broad distribution that can be only explained by  
449 human-mediated transport; and 3) species complexes in which non-monophyletic  
450 cryptic diversity has been found . This study indicates that widely distributed species  
451 are the exception in red algae, except when they have been spread by humans.

452

### 453 **Acknowledgements**

454 We thank Francis Bunker, Fabio Rindi, Iara Chapuis, Jan Rueness and Max  
455 Hommersand for providing samples and Joana Costa, as well as other researchers  
456 included in Table S1, for assistance in the field. We thank Marisa Pasella for carrying  
457 out part of the molecular work. PDT acknowledges support by the postdoctoral program  
458 “Axudas de apoio á etapa de formación posdoutoral”, as well as the funding program  
459 “Axudas para a consolidación e estruturación de unidades de investigación competitivas  
460 do SUG”, grant GPC2015/025 (Xunta de Galicia). Field work in the Mediterranean Sea  
461 was funded by a grant awarded to PDT by the British Phycological Society. Part of the  
462 research presented here was funded through the Australian Biological Resources Study  
463 (ABRS), including participation in a Bush Blitz expedition, a Bush Blitz Strategic  
464 Taxonomy Grant (TTC216-03) and a National Taxonomy Research Grant (RFL213-  
465 08). Sampling in Western Australia, Queensland and Tasmania was made possible  
466 through funding from the Holsworth Wildlife Research Endowment. ECM  
467 acknowledges support by the Chilean Millennium Initiative (NC120030) grant.

468

### 469 **References**

470 Airoidi, L. 1998. Roles of disturbance, sediment stress, and substratum retention on  
471 spatial dominance in algal turf. *Ecology* 79:2759–70.

472 Boudouresque, C. F. & Verlaque, M. 2008. Biological pollution in the Mediterranean  
473 Sea: invasive versus introduced macrophytes. *Mar. Pollut. Bull.* 44:32-8.

474 Bustamante, D.E., Won, B.Y. & Cho, T.O. 2014. *Polysiphonia dokdoensis* sp. nov.  
475 (Rhodomelaceae, Ceramiales) based on a population previously known as *Polysiphonia*  
476 *atlantica sensu* Kim and Lee from Korea. *Bot. Mar.* 57:281-9.

477 Coyne, J.A., Orr, H.A., & Futuyma, D.J. 1988. Do we need a new species concept?  
478 *Syst. Biol.* 37:190–200.

479 Destombe, C., Godin, J., Lefebvre, C., Dehorter, O., Vernet, Ph. 2009. Differences in  
480 dispersal abilities of haploid and diploid spores of *Gracilaria verrucosa* (Gracilariales,  
481 Rhodophyta). *Bot. Mar.* 35:93-8.

482 Díaz-Tapia, P. & Bárbara, I. 2013. Seaweeds from sand-covered rocks of the Atlantic  
483 Iberian Peninsula. Part. 1. The Rhodomelaceae (Ceramiales, Rhodophyta). *Cryptog.*  
484 *Algol.* 34:325-422.

485 Díaz-Tapia, P., Bárbara, I. & Díez, I. 2013a. Multi-scale spatial variability in intertidal  
486 benthic assemblages: Differences between sand-free and sand-covered rocky habitats,  
487 *Estuar. Coast. Shelf Sci.* 133:97-108.

488 Díaz-Tapia, P., Kim, M.S., Secilla, A., Bárbara, I., Cremades, J. 2013b. Taxonomic  
489 reassessment of *Polysiphonia foetidissima* (Rhodomelaceae, Rhodophyta) and similar  
490 species, including *P. schneideri*, a newly introduced species in Europe. *Eur. J. Phycol.*  
491 48:345-62.

492 Díaz-Tapia, P., Bárbara, I., Cremades, J., Verbruggen, H. & Maggs C.A. 2017a. Three  
493 new cryptogenic species in the tribes Polysiphonieae and Streblocladieae  
494 (Rhodomelaceae, Rhodophyta). *Phycologia* 56:605-23.

495 Díaz-Tapia, P., Maggs, C. A., West, J. A. & Verbruggen, H. 2017b. Analysis of  
496 chloroplast genomes and a supermatrix inform reclassification of the Rhodomelaceae  
497 (Rhodophyta). *J. Phycol.* 53:920–37.

498 Fraser, C.I., Nikula, R., Spencer, H.G., & Waters, J.M. 2009. Kelp genes reveal effects  
499 of subantarctic sea ice during the Last Glacial Maximum. *Proc. Natl. Acad. Sci. USA*  
500 106:3249-53.

501 Fraser, C.I., Zuccarello, G.C., Spencer, H.G., Salvatore, L.C., Garcia, G.R. & Waters,  
502 J.M. 2013. Genetic affinities between trans-oceanic populations of non-buoyant  
503 macroalgae in the high latitudes of the southern hemisphere. *PLoS ONE* 8:e69138.

504 Freshwater, D.W. & Rueness, J. 1994. Phylogenetic relationships of some European  
505 *Gelidium* (Gelidiales, Rhodophyta) species based upon *rbcL* nucleotide sequence  
506 analysis. *Phycologia* 33:187-94.

507 Geoffroy, A., Destombe, C., Kim, B, Mauger, S., Raffo, M.P., Kim, M.S. & Le Gall, L.  
508 2016. Patterns of genetic diversity of the cryptogenic red alga *Polysiphonia morrowii*  
509 (Ceramiales, Rhodophyta) suggest multiple origins of the Atlantic populations. *Ecol.*  
510 *Evol.* 6:5635-47.

511 Guiry, M.D. & Guiry, G.M. 2018. *AlgaeBase*; World-Wide Electronic Publication,  
512 National University of Ireland: Galway, UK, 2018; Available online:  
513 <http://www.algaebase.org> (accessed on 6 January 2018).

514 Jackson, J.B.C. 1974. Biogeographic consequences of eurytopy and stenotopy among  
515 marine bivalves and their evolutionary significance. *Amer. Nat.* 108:541–60.

516 Jordan, S.M.R., Barraclough T.G. & Rosindell, J. 2016. Quantifying the effects of the  
517 break up of Pangaea on global terrestrial diversification with neutral theory. *Phil. Trans.*  
518 *R. Soc. B* 371:20150221.

519 Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton,  
520 S, Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P. &  
521 Drummond, A. 2012. Geneious Basic: An integrated and extendable desktop software  
522 platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–9.

523 Kinlan, B.P. & Gaines, S.D. 2003. Propagule dispersal in marine and terrestrial  
524 environments: a community perspective. *Ecology* 84:2007–20.

525 Leliaert F., Verbruggen H., Vanormelingen P., Steen, F., López-Bautista J.M.,  
526 Zuccarello, G.C. & De Clerck O. 2014. DNA based species delimitation in algae. *Eur.*  
527 *J. Phycol.* 49: 179–96.

528 López, B.A., Tellier, F., Retamal-Alarcón, J.C., Pérez-Araneda, K., Fierro, A.O.,  
529 Macaya, E.C., Tala, F. & Thiel, M. 2017. Phylogeography of two intertidal seaweeds,

530 *Gelidium lingulatum* and *G. rex* (Rhodophyta: Gelidiales), along the South East Pacific:  
531 patterns explained by rafting dispersal? *Mar. Biol.* 164:188.

532 López, B.A., Macaya E. C., Rivadeneira M., Tala F., Tellier F. & Thiel M. 2018.  
533 Epibiont communities on stranded kelp rafts of *Durvillaea Antarctica* (Fucales,  
534 Phaeophyceae) – do positive interactions facilitate range extensions? *J. Biogeogr.*  
535 45:1833-45.

536 Macaya, E.C. & Zuccarello, G.C. 2010. Genetic structure of the giant kelp *Macrocystis*  
537 *pyrifera* along the southeastern Pacific. *Mar Ecol Prog Ser* 420:103–12.

538 Macaya, E.C., López, B., Tala, F., Tellier, F. & Thiel, M. 2016. Float and raft: role of  
539 buoyant seaweeds in the phylogeography and genetic structure of non-buoyant  
540 associated flora. In Hu, Z.M. & Fraser, C.I. [Eds.] *Seaweed phylogeography*. Springer,  
541 Dordrecht, pp. 97–130.

542 Maggs, C.A. & Hommersand, M.H. 1993. *Seaweeds of the British Isles. Volume 1.*  
543 *Rhodophyta. Part 3A. Ceramiales*. HMSO, London, 444 pp.

544 Mamoozadeh, N.R. & Freshwater, D.W. 2011. Taxonomic notes on Caribbean  
545 *Neosiphonia* and *Polysiphonia* (Ceramiales, Florideophyceae): five species from  
546 Florida, USA and Mexico. *Bot. Mar.* 54:269-92.

547 McIvor, L., Maggs, C.A., Provan, J. & Stanhope, M.J. 2001. *rbcL* sequences reveal  
548 multiple cryptic introductions of the Japanese red alga *Polysiphonia harveyi*. *Mol. Ecol.*  
549 10: 911–9.

550 Nosil, P., Funk, D.J. & Ortiz-Barrientos, D. 2009. Divergent selection and  
551 heterogeneous genomic divergence. *Molec. Ecol.* 18:375–402.

552 Palumbi, S.R. 1994. Genetic divergence, reproductive isolation, and marine speciation.  
553 *Annu. Rev. Ecol Syst.* 25:547-72.

554 Payo, D.A., Leliaert, F., Verbruggen, H., D’hondt, S., Calumpong, H. P. & De Clerck,  
555 O. 2013. Extensive cryptic species diversity and fine-scale endemism in the marine red  
556 alga *Portieria* in the Philippines. *Proc. R. Soc. Lond. B Biol. Sci.* 280: 20122660.

557 Provan, J., Booth, D., Todd, N.P., Beatty, G.E. & Maggs, C.A. 2008. Tracking  
558 biological invasions in space and time: elucidating the invasive history of the green alga  
559 *Codium fragile* using old DNA. *Divers. Distributions* 14:343–54.

- 560 Riginos, C., Douglas, K. E., Jin, Y., Shanahan, D. F. & Treml, E.A. 2011. Effects of  
561 geography and life history traits on genetic differentiation in benthic marine fishes.  
562 *Ecography* 34:566-75.
- 563 Santelices, B. 1990. Patterns of reproduction, dispersal and recruitment in seaweeds.  
564 *Oceanogr. Mar. Biol. Ann. Rev.* 28:177-276.
- 565 Saunders, G.W. & McDevit, D.C. 2012. Methods for DNA Barcoding Photosynthetic  
566 Protists Emphasizing the Macroalgae and Diatoms. In Kress, W. & Erickson, D. [Eds.]  
567 *DNA Barcodes: Methods in Molecular Biology (Methods and Protocols)*. Humana  
568 Press, Totowa, NJ, USA, pp. 207-22
- 569 Saunders, G.W. & Moore, T.E. 2013. Refinements for the amplification and sequencing  
570 of red algal DNA barcode and RedToL phylogenetic markers: a summary of current  
571 primers, profiles and strategies. *Algae* 28:31-43.
- 572 Savoie, A. M. & Saunders, G. W. 2015. Evidence for the introduction of the Asian red  
573 alga *Neosiphonia japonica* and its introgression with *Neosiphonia harveyi* (Ceramiales,  
574 Rhodophyta) in the Northwest Atlantic. *Mol. Ecol.* 24:5927–37.
- 575 Savoie, A.M. & Saunders, G.W. 2016. A molecular phylogenetic and DNA barcode  
576 assessment of the tribe Pterosiphonieae (Ceramiales, Rhodophyta) emphasizing the  
577 Northeast Pacific. *Botany* 94: 917-39.
- 578 Schneider, C.W., Quach, P.K. & Lane, C.E. 2017. A case for true morphological  
579 cryptsis: Pacific *Dasya anastomosans* and Atlantic *D. cryptica* sp. nov. (Dasyaceae,  
580 Rhodophyta). *Phycologia* 56:359-68.
- 581 Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-  
582 analysis of large phylogenies. *Bioinformatics* 30:1312–13.
- 583 Straub, S.C., Thomsen, M.S. & Wernberg, T. 2016. The dynamic biogeography of the  
584 anthropocene: the speed of recent range shifts in seaweeds. In Hu, Z.-M. & Fraser, C.  
585 [Eds.] *Seaweed Phylogeography*. Springer, Amsterdam, pp. 63–93.
- 586 Thomsen M.S., Wernberg T., South P.M. & Schiel D.R. 2016. Non-native seaweeds  
587 drive changes in marine coastal communities around the world. In Hu, Z.M. & Fraser,  
588 C.I. [Eds.] *Seaweed phylogeography*. Springer, Dordrecht, pp. 147-185.

589 Verbruggen, H. 2014. Morphological complexity, plasticity, and species diagnosability  
590 in the application of old species names in DNA-based taxonomies. *J. Phycol.* 50:26-31.

591 Voisin, M., Engel, C.R. & Viard, F. 2005. Differential shuffling of native genetic  
592 diversity across introduced regions in a brown alga: Aquaculture vs. maritime traffic  
593 effects. *Proc. Natl. Acad. Sci. USA* 102:5432-7.

594 Waters, J.M., Fraser, C.I. & Hewitt, G.M. 2013. Founder takes all: density-dependent  
595 processes structure biodiversity. *Trends Ecol. Evol.* 28:78-85.

596 Williams, S.L. & Smith, J.E. 2007. A global review of the distribution, taxonomy, and  
597 impacts of introduced seaweeds. *Annu. Rev. Ecol. Evol. Syst.* 38:327-59.

598 Womersley, H.B.S. 2003. *The marine benthic flora of southern Australia - Part IIID*  
599 *Ceramiales - Delesseriaceae, Sarcomeniaceae, Rhodomelaceae*. Australian Biological  
600 Resources Study and State Herbarium of South Australia, Canberra and Adelaide, 533  
601 pp.

602 Won, B.Y., Cho, T.O. & Fredericq, S. 2009. Morphological and molecular  
603 characterization of species of the genus *Centroceras* (Ceramiales, Ceramiaceae),  
604 including two new species. *J. Phycol.* 45:227-50.

605 Zuccarello, G., West, J. & King, R. 1999. Evolutionary divergence in the *Bostrychia*  
606 *moritziana/B. radicans* complex (Rhodomelaceae, Rhodophyta): molecular and  
607 hybridization data. *Phycologia* 38:234-44.

608 Zuccarello, G.C., Sandercock, B. & West, J.A. 2002a. Diversity within red algal  
609 species: variation in world-wide samples of *Spyridia filamentosa* (Ceramiales) and  
610 *Murrayella pericladus* (Rhodomelaceae) using DNA markers and breeding studies. *Eur.*  
611 *J. Phycol.* 37:403-18.

612 Zuccarello, G. C., West, J. & Rueness, J. 2002b. Phylogeography of the cosmopolitan  
613 red alga *Caulacanthus ustulatus* (Caulacanthaceae, Gigartinales). *Phycol. Res.* 50:163-  
614 72.

615 Zuccarello, G.C. & West, J. 2003. Multiple cryptic species: molecular diversity and  
616 reproductive isolation in the *Bostrychia radicans/B. moritziana* complex  
617 (Rhodomelaceae, Rhodophyta) with focus on North American isolates. *J. Phycol.*  
618 39:948-59.

619 Zuccarello, G.C., West J. A. & Kamiya, M. 2018. Non-monophyly of *Bostrychia*  
620 *simpliciuscula* (Ceramiales, Rhodophyta): Multiple species with very similar  
621 morphologies, a revised taxonomy of cryptic species. *Phycological Res.* 66:100-7.

622

623 Figure legends

624 **Figure 1** Distribution and UPGMA unrooted distance phylogram based on *rbcL*  
625 sequences of (a, b) *Ophidocladus simpliciusculus*, (c, d) *Lophosiphonia obscura* (as  
626 *Polysiphonia hemisphaerica* and *P. boldii* in Norway and Texas, respectively, see  
627 Appendix S1), and (e, f) *P. villum*. In panels a, c and e, circles indicate the regions from  
628 which sequences are available and their colors indicate the distribution of haplotypes.  
629 Areas outlined in red are regions where the species is recorded for the first time.  
630 Coastline in black shows the reported distribution (Guiry and Guiry 2018). In panel c,  
631 black coastlines represent the recorded distribution of *Lophosiphonia obscura*, red line  
632 *P. hemisphaerica* and yellow line *P. boldii*. Scale bars: 5 mm in (a), 8 mm in (c), 6 mm  
633 in (e).

634 **Figure 2** Distribution and UPGMA unrooted distance phylogram based on *rbcL*  
635 sequences of (a, b) *Polysiphonia devoniensis* (as *P. kaprauni* in North Carolina), (c, d)  
636 *Symphyclocladia dendroidea* and (e, f, g) *S. spinifera*/*Xiphosiphonia pennata*/*X.*  
637 *pinnulata*. Symbols are as in Figure 1, and pie divisions in panels a, c and e indicate  
638 proportions of each haplotype when multiple haplotypes were present. In panel e, circles  
639 with white border correspond to *S. spinifera* and the ones with black border to *X.*  
640 *pinnulata*. Encircled areas marked with red color are regions where the species are here  
641 recorded for the first time. In panel a, black lines represent the recorded distribution of  
642 *P. devoniensis* and red line the distribution of *P. kapraunii*. In panel c, black lines  
643 represent the recorded distribution of *S. dendroidea*, blue lines the regions where it was  
644 recorded as *S. tanakae*, red line the region where molecular data demonstrated that *S.*  
645 *dendroidea 2* is present, the asterisk indicates the area where both *S. dendroidea 2* and  
646 *S. tanakae* were reported based on molecular data, and plus symbols the regions from  
647 which sequences of *S. dendroidea 2* are available. In panel e, black lines represent the  
648 recorded distribution of *X. pennata*; yellow lines the regions where molecular data  
649 showed the presence of *S. spinifera* instead *X. pennata*, red lines regions where only *X.*  
650 *pinnulata* has been recorded based on molecular data and blue line the region where

651 both *X. pinnulata* and *X. pennata* have been recorded based on molecular data. Scale  
652 bars: 6 mm in (a), 7 mm in (c), 4 mm in (e).






653 **Figure 3** Distribution of *Herposiphonia tenella*. Asterisks indicate the regions from  
654 which sequences are available. Scale bar: 1 mm.

655 **Figure 4** RAxML tree based on *rbcL* sequences of the genus *Herposiphonia*. Samples  
656 that morphologically correspond with *Herposiphonia tenella* are in bold. Bootstrap  
657 values are indicated on the nodes when > 80. BE (Belize), CA (Canada), CI (Canary  
658 Islands), FR (France), IN (India), IT (Italy), KO (Korea), NC (North Carolina), PO  
659 (Portugal), QL (Queensland), SP (Spain), WA (Western Australia), VIC (Victoria).


660

661 **Appendix S1.** Taxonomic notes.

662 **Table S1.** GenBank accession numbers of the *rbcL* sequences included in the UPGMA  
663 and phylogenetic analysis.



jpy\_12778-18-070\_f1.tif



ipy\_12778-18-070\_f2.tif



jpy\_12778-18-070\_f3.tif

