



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Dower, PM;Zhang, H

Title:

A max-plus primal space fundamental solution for a class of differential Riccati equations

Date:

2017-09-01

Citation:

Dower, P. M. & Zhang, H. (2017). A max-plus primal space fundamental solution for a class of differential Riccati equations. *Mathematics of Control Signals and Systems*, 29 (3), <https://doi.org/10.1007/s00498-017-0200-2>.

Persistent Link:

<https://hdl.handle.net/11343/283308>

A max-plus primal space fundamental solution for a class of differential Riccati equations*

Peter M. Dower · Huan Zhang

Initial submission date: December 2015.

Abstract A class of differential Riccati equations (DREs) is considered for which the evolution of any solution can be identified with the propagation of a value function of a corresponding optimal control problem arising in \mathcal{L}_2 -gain analysis. By exploiting the semigroup properties inherited from the attendant dynamic programming principle, a max-plus primal space fundamental solution semigroup of max-plus linear max-plus integral operators is developed that encapsulates all such value function propagations. Using this semigroup, a one-parameter fundamental solution semigroup of matrices is constructed for the aforementioned class of DREs. It is demonstrated that this semigroup can be used to compute particular solutions of these DREs, and to characterize finite escape times (should they exist) in a simple way.

Keywords: Max-plus methods, fundamental solutions, dynamic programming semigroups, differential Riccati equations.

1 INTRODUCTION

Differential Riccati equations (DREs) arise naturally in linear optimal control and dissipative systems theory [1–4]. A typical finite dimensional DRE applicable in the verification of the \mathcal{L}_2 -gain property for linear systems is an ordinary differential equation of the form

$$\dot{P}_t = A'P_t + P_tA + P_tBB'P_t + C'C, \quad (1)$$

in which $A \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{n \times m}$, $C \in \mathbb{R}^{p \times n}$, $n, m, p \in \mathbb{N}$, are fixed problem specific matrices. A symmetric matrix-valued map $t \mapsto P_t \in \mathbb{S}^{n \times n}$ describes a

[†]Dower and Zhang are with the Department of Electrical & Electronic Engineering at the University of Melbourne, Victoria 3010, Australia. {pdower, hzhang5}@unimelb.edu.au

*Research partially supported by the Australian Research Council and AFOSR/AOARD grants DP120101549 and FA2386-12-1-4084.

particular solution to (1), evolved forward from an initial condition

$$P_0 \in \mathbb{S}_{>M}^{n \times n} \doteq \{P \in \mathbb{S}^{n \times n} \mid P - M > 0\}, \quad M \in \mathbb{S}^{n \times n}, \quad (2)$$

residing in a subset $\mathbb{S}_{>M}^{n \times n}$ of the space of symmetric matrices $\mathbb{S}^{n \times n}$, to any time $t \in [0, t^*)$ up to some maximal horizon of existence $t^* = t^*(P_0) \in \mathbb{R}_{>0}^+ \doteq \mathbb{R}_{>0} \cup \{+\infty\}$. Related DREs arise in linear \mathcal{H}_2 - and \mathcal{H}_∞ -control and filtering, etc, see for example [2–4].

A fundamental solution for DRE (1), (2) is an object that characterizes every possible solution of the differential equation (1), as parameterized by its initial (or terminal) condition (2). One such fundamental solution is the *symplectic fundamental solution*, which is itself the solution of a (derived) Hamiltonian system of linear ordinary differential equations, see for example [1, 5, 6]. Another fundamental solution is the *max-plus dual space fundamental solution* [7–10], which is constructed by exploiting semiconvex duality [11] and max-plus linearity of the Lax-Oleinik semigroup [12] of dynamic programming evolution operators for an associated optimal control problem, see also [12–16].

In this paper, a *max-plus primal space fundamental solution* is constructed for DREs of the form (1), (2). This fundamental solution can be used to evaluate particular solutions of (1), analogously to the symplectic and max-plus dual space fundamental solutions. Its development is complementary to that of the max-plus dual space fundamental solution documented in [7, 8, 10], and parallels the corresponding recent primal space development for difference Riccati equations [9]. It is shown that this new fundamental solution provides a simple test for establishing existence of solutions of (1), (2).

In terms of organization, the symplectic fundamental solution for DRE (1), (2) is recalled in Section 2 for comparative purposes, to formalize existence of solutions, and to construct a specific particular solution to (1) of utility later. The max-plus primal space fundamental solution, and corresponding fundamental solution semigroup, is subsequently constructed in Sections 3 and 4, using the aforementioned particular solution. An illustration of its application is provided in Section 5, followed by some brief concluding remarks in Section 6. Proofs are largely delayed to the appendices.

Throughout, \mathbb{N} , \mathbb{Q} , \mathbb{R} , $\mathbb{R}_{\geq 0}$, \mathbb{R}^n , $\mathbb{R}^{n \times n}$ denote respectively the natural, rational, real, and nonnegative real numbers, n -dimensional Euclidean space, and the space of $n \times n$ matrices with real entries. \mathbb{R}^\pm denotes the analogous extended reals $\mathbb{R} \cup \{\pm\infty\}$. Similarly, $\mathbb{S}^{n \times n}$, $\mathbb{S}_{\geq 0}^{n \times n}$, $\mathbb{S}_{>0}^{n \times n}$ denote the spaces of symmetric, nonnegative symmetric, and positive definite symmetric elements of $\mathbb{R}^{n \times n}$ respectively. Further extending this notation, $\mathbb{S}_{>M}^{n \times n}$ denotes the subset of $\mathbb{S}^{n \times n}$ of matrices P satisfying $P - M \in \mathbb{S}_{>0}^{n \times n}$, etc. The transpose of $P \in \mathbb{R}^{n \times n}$ is denoted by $P' \in \mathbb{R}^{n \times n}$. The corresponding identity is denoted by $I \in \mathbb{S}^{n \times n}$. Given $U \in \mathbb{R}^{2n \times 2n}$, the two-by-two block matrix representation

$$U = \begin{bmatrix} U^{11} & U^{12} \\ U^{21} & U^{22} \end{bmatrix} \in \mathbb{R}^{2n \times 2n}, \quad (3)$$

with $U^{ij} \in \mathbb{R}^{n \times n}$, $i, j \in \{1, 2\}$, is used where convenient.

2 SYMPLECTIC FUNDAMENTAL SOLUTION

Existence of a unique solution to DRE (1), subject to (2), may be verified by application of Banach's fixed point theorem, see for example [8, Theorem 2.4]. Alternatively, it may be constructed directly as

$$P_t = Y_t X_t^{-1} \quad (4)$$

in which $X_t, Y_t \in \mathbb{R}^{n \times n}$ are defined with respect to the *symplectic fundamental solution* $\Sigma_t \in \mathbb{R}^{2n \times 2n}$ for DRE (1), (2) by

$$\begin{aligned} \begin{bmatrix} X_t \\ Y_t \end{bmatrix} &= \Sigma_t \begin{bmatrix} I \\ P_0 \end{bmatrix}, \quad t \in [0, t^*(P_0)), \\ \Sigma_t &\doteq \exp(\mathcal{H}t), \quad \mathcal{H} \doteq \begin{bmatrix} -A & -BB' \\ C'C & A' \end{bmatrix}, \end{aligned} \quad (5)$$

in which the maximal horizon of existence $t^*(P_0) \in \mathbb{R}_{>0}^+$ of the unique particular solution P_t in (4) is characterized by

$$t^*(P_0) \doteq \sup \left\{ t \in \mathbb{R}_{>0} \left| \begin{array}{l} X_s^{-1} \text{ exists } \forall s \in (0, t] \\ \text{with } X_s \text{ given by (5)} \\ \text{subject to } P_0 \in \mathbb{S}^{n \times n} \end{array} \right. \right\}, \quad (6)$$

see [5, 17, 18]. This maximal horizon of existence is either strictly positive and finite, or infinite. Where $t^*(P_0)$ is strictly positive, the solution P_t experiences a finite escape at $t = t^*(P_0)$. Otherwise, no such finite escape time exists, and P_t may be evolved to any arbitrarily large time horizon $t \in \mathbb{R}_{>0}$. For example, under the conditions of the strict bounded real lemma (e.g. [3, Theorem 2.1] or [4, Lemma 3.7.7]), $P_0 = 0 \in \mathbb{S}^{n \times n}$ implies that $t^*(P_0) = +\infty$.

By inspection, the symplectic fundamental solution Σ_t , defined by (4), (5), (6) satisfies the properties of a fundamental solution for DRE (1), (2). In particular, it can be evolved independently of any specific DRE initial condition P_0 , and can be used to recover any such particular solution via an operation involving that P_0 . It is a standard tool for the representation and computation of solutions to DREs of this form. In Section 3, it is used to construct a particular solution that is employed in the construction the max-plus primal space fundamental solution of interest here.

3 MAX-PLUS FUNDAMENTAL SOLUTION

A max-plus fundamental solution for DRE (1), (2) is developed via the following steps: 1) the standard symplectic fundamental solution (5) is identified with the solution of an auxiliary optimal control problem, 2) max-plus and semiconvex analysis concepts are applied to represent this value function in terms of a horizon indexed max-plus linear max-plus integral operator, and 3) all particular solutions of DRE (1), (2) are described in terms of the Hessian of the quadratic kernel of this max-plus linear max-plus integral operator. As will

be shown, a one-parameter family of these Hessians need only be computed once, whereupon it can be used in the evaluation of any particular solution of DRE (1), (2), on any time horizon **for which that solution exists**. The relationship between this fundamental solution and that of [7] is also summarized.

As a prelude to the aforementioned development, a brief summary of the pertinent max-plus and semiconvex analysis concepts is first provided.

3.1 Max-plus algebra and semiconvex duality

The max-plus algebra [12,16] is a commutative semifield over \mathbb{R}^- , equipped with addition and multiplication operators defined respectively by $a \oplus b \doteq \max(a, b)$ and $a \otimes b \doteq a + b$. It is an idempotent algebra, as the \oplus operation is idempotent (i.e. $a \oplus a = a$), and a semifield as additive inverses do not exist. The max-plus integral of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^-$ over a subset $\mathcal{Y} \subset \mathbb{R}^n$ of its domain is $\int_{\mathcal{Y}}^{\oplus} f(y) dy \doteq \sup_{y \in \mathcal{Y}} f(y)$. The max-plus delta function $\delta^- : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^-$ is defined for all $x, y \in \mathbb{R}^n$ by

$$\delta^-(x, y) \doteq \begin{cases} 0, & x = y, \\ -\infty, & x \neq y. \end{cases} \quad (7)$$

In developing a max-plus fundamental solution, it is useful to introduce spaces of uniformly semiconvex and semiconcave functions, defined with respect to $K \in \mathbb{S}^{n \times n}$, by

$$\begin{aligned} \mathcal{S}_+^K &\doteq \left\{ f : \mathbb{R}^n \rightarrow \mathbb{R}^- \mid \begin{array}{l} f + \frac{1}{2} \langle \cdot, K \cdot \rangle \\ \text{convex} \end{array} \right\}, \\ \mathcal{S}_-^K &\doteq \left\{ a : \mathbb{R}^n \rightarrow \mathbb{R}^- \mid \begin{array}{l} a - \frac{1}{2} \langle \cdot, K \cdot \rangle \\ \text{concave} \end{array} \right\}, \end{aligned} \quad (8)$$

respectively. Semiconvex duality is a duality between these spaces of semiconvex and semiconcave functions, that is established via the semiconvex transform [11]. The semiconvex transform is a generalization of the Legendre-Fenchel transform [19–21], in which **affine support (see for example [21, Theorem 8.13]) is weakened to quadratic support, using a quadratic *basis* function** $\varphi : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$. **Here, this basis function is defined for all $x, z \in \mathbb{R}^n$ by**

$$\varphi(x, z) \doteq \frac{1}{2}(x - z)' M (x - z) = \frac{1}{2} \begin{bmatrix} x \\ z \end{bmatrix}' \mu(M) \begin{bmatrix} x \\ z \end{bmatrix}, \quad (9)$$

in which $M \in \mathbb{S}^{n \times n}$ and $\mu(M) \in \mathbb{R}^{2n \times 2n}$, with the latter defined in terms of the former via a map $\mu : \mathbb{S}^{n \times n} \rightarrow \mathbb{S}^{2n \times 2n}$, where

$$\mu(P) \doteq \begin{bmatrix} +P & -M \\ -M & +M \end{bmatrix} \in \mathbb{S}^{2n \times 2n}, \quad (10)$$

for all $P \in \mathbb{S}^{n \times n}$.

Assumption 1 Matrix $M \in \mathbb{S}^{n \times n}$ defining the basis (9) satisfies the following properties:

- 1) $M^{-1} \in \mathbb{S}^{n \times n}$ exists;
- 2) $t^*(M) = +\infty$, cf. (6).

Standard conditions under which Assumption 1 holds are controllability and observability of (A, B) and (C, A) respectively, or via the strict bounded real lemma, see for example [3, 4]. The details are postponed to Lemma 1, see Section 3.3.

The semiconvex transform and its inverse are well-defined with respect to the basis function φ of (9) by

$$\mathcal{D}_\varphi \psi \doteq - \int_{\mathbb{R}^n}^{\oplus} \varphi(x, \cdot) \otimes (-\psi(x)) dx, \quad (11)$$

$$\mathcal{D}_\varphi^{-1} a \doteq \int_{\mathbb{R}^n}^{\oplus} \varphi(\cdot, z) \otimes a(z) dz, \quad (12)$$

for all $\psi \in \text{dom}(\mathcal{D}_\varphi) \doteq \mathcal{S}_+^{-M}$ and $a \in \text{dom}(\mathcal{D}_\varphi^{-1}) \doteq \mathcal{S}_-^{-M}$, see also [16, 7, 8, 22]. For quadratic functions, (11) and (12) define a pair of matrix operations on corresponding spaces of Hessians. In particular, with $\psi : \mathbb{R}^n \rightarrow \mathbb{R}$ defined with respect to some $P \in \mathbb{S}_{>M}^{n \times n}$ by $\psi(x) \doteq \frac{1}{2}x'Px$ for all $x \in \mathbb{R}^n$, application of (11) yields a well-defined semiconvex dual $a \in \mathcal{S}_-^{-M}$. In particular, $a(z) \doteq (\mathcal{D}_\varphi \psi)(z) = \frac{1}{2}z'\Upsilon(P)z$ for all $z \in \mathbb{R}^n$, with $\Upsilon : \mathbb{S}^{n \times n} \rightarrow \mathbb{S}^{n \times n}$ defined by

$$\begin{aligned} \Upsilon(P) &\doteq -M - M(P - M)^{-1}M, & P &\in \text{dom}(\Upsilon), \\ \text{dom}(\Upsilon) &\doteq \mathbb{S}_{>M}^{n \times n}. \end{aligned} \quad (13)$$

Similarly, the inverse semiconvex transform (12) corresponds to the inverse map Υ^{-1} , with

$$\begin{aligned} \Upsilon^{-1}(P) &\doteq M - M(P + M)^{-1}M, & P &\in \text{dom}(\Upsilon^{-1}), \\ \text{dom}(\Upsilon^{-1}) &\doteq \mathbb{S}_{<-M}^{n \times n}. \end{aligned} \quad (14)$$

Remark 1 The domains specified in (13) and (14) may be extended to $\mathbb{S}_{\geq M}^{n \times n}$ and $\mathbb{S}_{\leq -M}^{n \times n}$ respectively, via corresponding Moore-Penrose pseudo-inverses. However, this extension is not required here, and the details are omitted.

Remark 2 As evaluation of the semiconvex transform of a quadratic function via (11) involves computing the maximum of a sum of quadratic functions, the associated operator Υ of (13) naturally features a Schur complement. Similarly, the operator Υ^{-1} of (14) associated with the inverse semiconvex transform (12) also features a Schur complement. Indeed, as the fundamental solution semigroup developed later in Section 4 is founded on such maximizations, Schur complements also naturally arise in its propagation.

3.2 Optimal control problem

In order to construct a max-plus fundamental solution for DRE (1), (2), it is useful to define a corresponding optimal control problem on a finite time horizon $t \in \mathbb{R}_{\geq 0}$ via a value function $W_t : \mathbb{R}^n \rightarrow \mathbb{R}$ given by

$$W_t(x) \doteq (\mathcal{S}_t \Psi)(x) \quad (15)$$

for all $x \in \mathbb{R}^n$. Here, $\Psi : \mathbb{R}^n \rightarrow \mathbb{R}$ is a quadratic terminal payoff defined by $\Psi(x) \doteq \frac{1}{2}x'P_0x$ for all $x \in \mathbb{R}^n$, with $P_0 \in \mathbb{S}_{>M}^{n \times n}$ and $M \in \mathbb{S}^{n \times n}$ as per (2) and (9). This terminal payoff appears in (15) as the argument of a *dynamic programming evolution operator* \mathcal{S}_t defined by

$$\begin{aligned} (\mathcal{S}_t \psi)(x) &\doteq \sup_{w \in \mathcal{L}_2([0, t]; \mathbb{R}^m)} J_\psi(t, x, w), \quad \psi \in \text{dom}(\mathcal{S}_t), \\ \text{dom}(\mathcal{S}_t) &\doteq \left\{ \psi : \mathbb{R}^n \rightarrow \mathbb{R}^- \mid \begin{array}{l} (\mathcal{S}_t \psi)(x) \in \mathbb{R}^- \\ \forall x \in \mathbb{R}^n \end{array} \right\}, \end{aligned} \quad (16)$$

for all $x \in \mathbb{R}^n$. Here, payoff $J_\psi(t, \cdot, \cdot) : \mathbb{R}^n \times \mathcal{L}_2([0, t]; \mathbb{R}^m) \rightarrow \mathbb{R}^-$ is defined by

$$J_\psi(t, x, w) \doteq \int_0^t \frac{1}{2}|y_s|^2 - \frac{1}{2}|w_s|^2 ds + \psi(x_t) \quad (17)$$

for all $x \in \mathbb{R}^n$, $w \in \mathcal{L}_2([0, t]; \mathbb{R}^m)$, in which $x_s \in \mathbb{R}^n$, $w_s \in \mathbb{R}^m$, and $y_s \in \mathbb{R}^p$ denote the state, input, and output (respectively) of the linear system

$$\begin{aligned} \dot{x}_s &= Ax_s + Bw_s, \quad x_0 = x \in \mathbb{R}^n, \\ y_s &= Cx_s, \end{aligned} \quad (18)$$

at time $s \in [0, t]$. It is straightforward to show that the value function W_t of (15) is quadratic, see [1, 7–9], with

$$W_t(x) = (\mathcal{S}_t \Psi)(x) = \frac{1}{2}x'P_t x \quad (19)$$

for all $x \in \mathbb{R}^n$, with $P_t \in \mathbb{S}^{n \times n}$ satisfying DRE (1) subject to the initial condition (2).

3.3 Auxiliary optimal control problem

In addition to the optimal control problem (15), it is useful to introduce an auxiliary optimal control problem defined on the same time horizon but with a different terminal payoff. The solution of this auxiliary problem is **subsequently shown** to be equivalent to the symplectic fundamental solution (in an appropriate sense). To this end, **given** $t \in \mathbb{R}_{>0}$, define the value function $S_t(\cdot, z) : \mathbb{R}^n \rightarrow \mathbb{R}$, $z \in \mathbb{R}^n$, in terms of the dynamic programming evolution operator \mathcal{S}_t of (16) by

$$S_t(x, z) \doteq (\mathcal{S}_t \varphi(\cdot, z))(x) \quad (20)$$

for all $x \in \mathbb{R}^n$. This value function is again quadratic, with

$$S_t(x, z) = \frac{1}{2} \begin{bmatrix} x \\ z \end{bmatrix}' Q_t \begin{bmatrix} x \\ z \end{bmatrix}, \quad (21)$$

for all $x, z \in \mathbb{R}^n$, in which $Q_t \in \mathbb{S}^{2n \times 2n}$ is the unique solution of the DRE

$$\dot{Q}_t = \hat{A}' Q_t + Q_t \hat{A} + Q_t \hat{B} \hat{B}' Q_t + \hat{C}' \hat{C} \quad (22)$$

initialized with

$$Q_0 = \mu(M) \in \mathbb{R}^{2n \times 2n} \quad (23)$$

and M as per (9), (10), for all $t \in [0, t^*(Q_0))$. Here, $t^*(Q_0) \in \mathbb{R}_{>0}^+$ denotes the corresponding maximal horizon of existence (6), with the constant matrices $\hat{A} \in \mathbb{R}^{2n \times 2n}$, $\hat{B} \in \mathbb{R}^{2n \times m}$, and $\hat{C} \in \mathbb{R}^{p \times 2n}$ of (22) defined by

$$\hat{A} \doteq \begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix}, \quad \hat{B} \doteq \begin{bmatrix} B \\ 0 \end{bmatrix}, \quad \hat{C} \doteq [C \ 0]. \quad (24)$$

Using the notation of (3), DRE (22), (23) is equivalent to $Q_t^{11}, Q_t^{22} \in \mathbb{S}^{n \times n}$, $Q_t^{12} \in \mathbb{R}^{n \times n}$ satisfying

$$\dot{Q}_t^{11} = A' Q_t^{11} + Q_t^{11} A + Q_t^{11} B B' Q_t^{11} + C' C, \quad (25)$$

$$\dot{Q}_t^{12} = (A + B B' Q_t^{11})' Q_t^{12}, \quad Q_t^{21} = (Q_t^{12})', \quad (26)$$

$$\dot{Q}_t^{22} = (Q_t^{12})' B B' Q_t^{12}, \quad (27)$$

for all $t \in [0, t^*(M))$, subject to $Q_0^{11} = -Q_0^{12} = Q_0^{22} = M$, with $M \in \mathbb{S}^{n \times n}$ as per (9). As (26), (27) describe (respectively) a linear evolution equation and an integration, any finite escape of Q_t that arises must be due to the dynamics (25), see for example [23, Proposition 3.6(iv)]. That is, the maximal horizon of existence for (22), (25) must be equal, ie. $t^*(Q_0) = t^*(M)$. Assumption 1 further implies that

$$t^*(Q_0) = t^*(M) = +\infty. \quad (28)$$

As DRE (25) is of the same form as (1), the particular solution Q_t of DRE (22), (23) can be characterized explicitly via the symplectic fundamental solution (5), and vice versa.

Theorem 1 *Under Assumption 1, the particular solution Q_t of DRE (22), (23) and the symplectic fundamental solution Σ_t of (5) for DRE (1) are equivalent. That is, there exists an invertible operator $\Xi : \mathbb{S}^{2n \times 2n} \rightarrow \mathbb{S}^{2n \times 2n}$ such that*

$$Q_t = \Xi(\Sigma_t), \quad \Sigma_t = \Xi^{-1}(Q_t) \quad (29)$$

for all $t \in \mathbb{R}_{\geq 0}$.

Proof See Appendix A.1. ■

Theorem 1 demonstrates **that**, under the conditions of Assumption 1, any particular solution of the DRE (1), (2) can be represented equivalently by the symplectic fundamental solution Σ_t of (5), or via the Hessian Q_t of the quadratic value function of the auxiliary optimal control problem (20), (21), see (29). **The** following sufficient condition for Assumption 1 is **thus** useful.

Lemma 1 *Suppose there exists a stabilizing solution $M_0 \in \mathbb{S}_{\geq 0}$ of the algebraic Riccati equation (ARE)*

$$0 = A'M_0 + M_0A + M_0BB'M_0 + C'C. \quad (30)$$

Then, there always exists an invertible $M \in \mathbb{S}^{n \times n}$ satisfying

$$M - M_0 \in \mathbb{S}_{< 0}^{n \times n}, \quad (31)$$

such that Assumption 1 holds.

Proof See Appendix A.2. ■

Remark 3 Lemma 1 provides a constructive approach to validating Assumption 1 directly. It also enables indirect validation via the bounded and strict bounded real lemmas, see for example [3,4]. In particular, stability of A , controllability of (A, B) , observability of (C, A) , and the finite gain property $\|(A, B, C)\|_{\mathcal{H}_\infty} \leq 1$ imply via the bounded real lemma that Assumption 1 holds. Alternatively, stability of A and the strict gain property $\|(A, B, C)\|_{\mathcal{H}_\infty} < 1$ imply via the strict bounded real lemma that Assumption 1 holds.

3.4 Max-plus integral operator representations for (16)

A horizon indexed *max-plus linear max-plus integral operator* defined on a space \mathcal{P} is an operator of the form

$$\begin{aligned} \mathcal{F}_t^\oplus \pi &\doteq \int_{\mathcal{P}}^\oplus F_t(\cdot, \omega) \otimes \pi \circ \chi_t(\cdot, \omega) d\omega, \quad \pi \in \text{dom}(\mathcal{F}_t^\oplus), \\ \text{dom}(\mathcal{F}_t^\oplus) &\doteq \left\{ \pi : \mathcal{P} \rightarrow \mathbb{R}^- \mid \begin{array}{l} (\mathcal{F}_t^\oplus \pi)(x) \in \mathbb{R}^- \\ \forall x \in \mathbb{R}^n \end{array} \right\}, \end{aligned} \quad (32)$$

where $F_t : \mathbb{R}^n \times \mathcal{P} \rightarrow \mathbb{R}^-$ denotes the kernel of the operator, $\chi_t : \mathbb{R}^n \times \mathcal{P} \rightarrow \mathbb{R}^n$ is an auxiliary operator (included here for generality), and $\pi \in \text{dom}(\mathcal{F}_t^\oplus)$ is the **function valued** argument of \mathcal{F}_t^\oplus representing a terminal payoff (or value function) or its semiconvex dual. The dynamic programming evolution operator \mathcal{S}_t of (16) defines a max-plus linear max-plus integral operator of this form, with $\mathcal{P} \doteq \mathcal{L}_2([0, t]; \mathbb{R}^m)$ and

$$F_t(x, w) = I_t(x, w) \doteq \int_0^t \frac{1}{2}|y_s|^2 - \frac{1}{2}|w_s|^2 ds, \quad \chi_t(x, w) \doteq x_t,$$

in which $I_t(x, w)$ is the integrated running payoff associated with initial state $x \in \mathbb{R}^n$ and input $w \in \mathcal{P}$ over the horizon $t \in \mathbb{R}_{\geq 0}$, and $x_t \in \mathbb{R}^n$ is the corresponding terminal state, both defined via (18). That is, for all $x \in \mathbb{R}^n$,

$$(\mathcal{S}_t \psi)(x) = \int_{\mathcal{P}}^{\oplus} I_t(x, w) \otimes \psi(x_t) dw. \quad (33)$$

Similarly, recalling the definition (7) of the max-plus delta function δ^- , the identity max-plus linear max-plus integral operator on $\mathcal{P} \doteq \mathbb{R}^n$ is defined via $F_t(\cdot, y) \doteq \delta^-(\cdot, y)$ and $\chi_t(\cdot, y) \doteq y \in \mathbb{R}^n$ by

$$(\mathcal{I}^{\oplus} \psi)(x) \doteq \int_{\mathbb{R}^n}^{\oplus} \delta^-(x, y) \otimes \psi(y) dy, \quad (34)$$

for all $x \in \mathbb{R}^n$, ie. $\mathcal{I}^{\oplus} \psi = \psi$ for any $\psi \in \text{dom}(\mathcal{I}^{\oplus})$, in which the domain $\text{dom}(\mathcal{I}^{\oplus})$ is defined as per (32).

Theorem 2 *Under Assumption 1, and given the dynamic programming evolution operator \mathcal{S}_t of (16) with $t \in \mathbb{R}_{\geq 0}$ fixed, there exists a max-plus linear max-plus integral operator \mathcal{G}_t^{\oplus} of the form (32) such that*

$$\begin{aligned} \mathcal{S}_t \psi &= \mathcal{G}_t^{\oplus} \psi \doteq \int_{\mathbb{R}^n}^{\oplus} G_t(\cdot, y) \otimes \psi(y) dy, \quad \forall \psi \in \text{dom}(\mathcal{G}_t^{\oplus}), \\ \text{dom}(\mathcal{G}_t^{\oplus}) &\doteq \text{dom}(\mathcal{S}_t), \end{aligned} \quad (35)$$

with kernel $G_t : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^-$ defined for all $x, y \in \mathbb{R}^n$ by

$$G_t(x, y) \doteq (\mathcal{S}_t \delta^-(\cdot, y))(x) = (\mathcal{D}_{\varphi} S_t(x, \cdot))(y), \quad (36)$$

with respect to (7), (11), (16), (20).

Proof Fix arbitrary $t \in \mathbb{R}_{\geq 0}$ and $x, y \in \mathbb{R}^n$. Recalling the definition (20), (21) of S_t ,

$$S_t(x, y) = (\mathcal{S}_t \varphi(\cdot, y))(x) = \frac{1}{2} x' Q_t^{11} x + x' Q_t^{12} y + \frac{1}{2} y' Q_t^{22} y,$$

wherein Assumption 1 and (27) imply that

$$Q_t^{22} \in \mathbb{S}_{\geq M}^{n \times n}. \quad (37)$$

Consequently, $S_t(x, \cdot) \in \text{dom}(\mathcal{D}_{\varphi}) = \mathcal{S}_+^{-M}$ by definition (11) of the semiconvex transform, so that

$$G_t(x, \cdot) \doteq \mathcal{D}_{\varphi} S_t(x, \cdot) \in \mathcal{S}_-^{-M} = \text{dom}(\mathcal{D}_{\varphi}^{-1}) \quad (38)$$

is well-defined. Note in particular that $G_t(x, y) \in \mathbb{R}^-$ by definition (8) of \mathcal{S}_-^{-M} . As $t \in \mathbb{R}_{\geq 0}$ and $x, y \in \mathbb{R}^n$ are arbitrary, a max-plus linear max-plus integral

operator \mathcal{G}_t^\oplus of the form (32) is well-defined via (38) by selecting $F_t \doteq G_t$ and $\chi_t(\cdot, y) \doteq y$. Recalling definitions (12), (20), (34) of \mathcal{D}_φ^{-1} , S_t , \mathcal{I}^\oplus , and (33),

$$\begin{aligned}
S_t(x, y) &= (\mathcal{S}_t \varphi(\cdot, y))(x) = (\mathcal{S}_t \mathcal{I}^\oplus \varphi(\cdot, y))(x) \\
&= \int_{\mathcal{P}}^\oplus I_t(x, w) \otimes \left[\int_{\mathbb{R}^n}^\oplus \delta^-(x_t, \xi) \otimes \varphi(\xi, y) d\xi \right] dw \\
&= \int_{\mathbb{R}^n}^\oplus \left[\int_{\mathcal{P}}^\oplus I_t(x, w) \otimes \delta^-(x_t, \xi) dw \right] \otimes \varphi(\xi, y) d\xi \\
&= \int_{\mathbb{R}^n}^\oplus (\mathcal{S}_t \delta^-(\cdot, \xi))(x) \otimes \varphi(\xi, y) d\xi = \int_{\mathbb{R}^n}^\oplus \varphi(y, \xi) \otimes (\mathcal{S}_t \delta^-(\cdot, \xi))(x) d\xi \\
&= (\mathcal{D}_\varphi^{-1} T_t(x, \cdot))(y) \tag{39}
\end{aligned}$$

where the interchange of max-plus integrals involved corresponds to an interchange of suprema, the second last equality follows by symmetry of φ , ie. $\varphi(\xi, y) = \varphi(y, \xi)$, and $T_t(x, y) \doteq (\mathcal{S}_t \delta^-(\cdot, y))(x)$. Substituting (39) in (38),

$$G_t(x, \cdot) = \mathcal{D}_\varphi S_t(x, \cdot) = \mathcal{D}_\varphi \mathcal{D}_\varphi^{-1} T_t(x, \cdot) = T_t(x, \cdot).$$

That is, (36) holds. For any $\psi \in \text{dom}(S_t)$, a similar argument yields

$$\begin{aligned}
(\mathcal{S}_t \psi)(x) &= (\mathcal{S}_t \mathcal{I}^\oplus \psi)(x) = \int_{\mathcal{P}}^\oplus I_t(x, w) \otimes \left[\int_{\mathbb{R}^n}^\oplus \delta^-(x_t, y) \otimes \psi(y) dy \right] dw \\
&= \int_{\mathbb{R}^n}^\oplus \left[\int_{\mathcal{P}}^\oplus I_t(x, w) \otimes \delta^-(x_t, y) dw \right] \otimes \psi(y) dy \\
&= \int_{\mathbb{R}^n}^\oplus (\mathcal{S}_t \delta^-(\cdot, y))(x) \otimes \psi(y) dy = \int_{\mathbb{R}^n}^\oplus G_t(x, y) \otimes \psi(y) dy \\
&= (\mathcal{G}_t^\oplus \psi)(x).
\end{aligned}$$

That is, (35) holds. ■

Remark 4 The kernel G_t of the max-plus linear max-plus integral operator \mathcal{G}_t^\oplus defined in Theorem 2 can be bounded above by the value function of a third optimal control problem. In particular, applying (36),

$$G_t(x, y) = (\mathcal{S}_t \delta^-(\cdot, y)) \leq (\mathcal{S}_t \psi_0)(x)$$

for all $t \in \mathbb{R}_{\geq 0}$, $x, y \in \mathbb{R}^n$, where $\psi_0 : \mathbb{R}^n \rightarrow \mathbb{R}$ is the zero terminal payoff defined by $\psi_0(x) \doteq 0$ for all $x \in \mathbb{R}^n$. By inspection of (16), $\mathcal{S}_t \psi_0$ is the value function of a standard optimal control problem arising in \mathcal{L}_2 -gain analysis. It is finite valued if there exists a stabilizing solution of ARE (30).

In developing a max-plus fundamental solution for DRE (1), (2) via Theorem 2, it is useful to establish a connection between finiteness of the kernel G_t of (36) and controllability of the underlying dynamics (18).

Lemma 2 *Suppose Assumption 1 holds. Then, (A, B) in (18) is controllable if and only if the kernel G_t of the max-plus linear max-plus integral operator \mathcal{G}_t^\oplus defined by (35) is finite for all $t \in \mathbb{R}_{>0}$. That is,*

$$(A, B) \text{ in (18) is controllable} \iff G_t(x, y) \in \mathbb{R} \forall t \in \mathbb{R}_{>0}, x, y \in \mathbb{R}^n.$$

Proof See Appendix A.3. ■

As a consequence of Lemma 2, the following is assumed for the remainder.

Assumption 2 (A, B) of (18) is controllable.

3.5 Max-plus fundamental solution for DRE (1), (2)

Dynamic programming implies that the set of dynamic programming evolution operators $\{\mathcal{S}_t\}_{t \in \mathbb{R}_{\geq 0}}$ defines the well-known Lax-Oleinik dynamic programming semigroup [12]. Applying Theorem 2, it immediately follows that $\{\mathcal{G}_t^\oplus\}_{t \in \mathbb{R}_{\geq 0}}$ must also define a one-parameter semigroup of **max-plus linear max-plus integral** operators via (35). In particular, $\{\mathcal{G}_t^\oplus\}_{t \in \mathbb{R}_{\geq 0}}$ naturally inherits (from the Lax-Oleinik semigroup) the semigroup and identity properties

$$\mathcal{G}_t^\oplus \mathcal{G}_\tau^\oplus = \mathcal{G}_{t+\tau}^\oplus, \quad \mathcal{G}_0^\oplus = \mathcal{I}^\oplus, \quad (40)$$

for $t, \tau \in \mathbb{R}_{\geq 0}$. This particular semigroup is referred to as the *max-plus primal space fundamental solution semigroup* for the optimal control problem (15), see [9, 22]. The modifier *primal* used here refers to the fact that propagation occurs in the primal space of payoffs. (A corresponding *max-plus dual space fundamental solution semigroup* also exists, where propagation occurs in a dual space defined by the semiconvex transform (11), see Section 3.6 and [7–10, 22].)

In the specific case of the optimal control problem defined by (15), the properties (35), (40) may be used to directly propagate the value function W_t to longer time horizons, with

$$W_{t+\tau} = \mathcal{G}_\tau^\oplus W_t, \quad W_t = \mathcal{G}_t^\oplus \Psi \quad (41)$$

for any $t, \tau, t + \tau \in [0, t^*(P_0))$. In view of (19), (41), a particular solution P_t of DRE (1) satisfying the initial condition (2) can be similarly propagated forward in time. This gives rise to a characterization of P_t in terms of the Hessian of the kernel G_t of the max-plus primal space fundamental solution \mathcal{G}_t^\oplus . This characterization is referred to as a *max-plus primal space fundamental solution* for DRE (1), (2).

Theorem 3 *Under Assumptions 1 and 2, there exists a bijection $\Pi : \mathbb{S}^{2n \times 2n} \rightarrow \mathbb{S}^{2n \times 2n}$ such that the kernel G_t of (36) takes the explicit finite quadratic form*

$$G_t(x, y) = \frac{1}{2} \begin{bmatrix} x \\ y \end{bmatrix}' \Lambda_t \begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}, \quad \Lambda_t \doteq \Pi^{-1}(Q_t), \quad (42)$$

for all $x, y \in \mathbb{R}^n$, $t \in \mathbb{R}_{>0}$, where Q_t is as per (22).

Proof See Appendix A.4. ■

By inspection of Theorems 1 and 3, the controllability Assumption 2 implies that the symplectic fundamental solution Σ_t and the Hessian Λ_t of the max-plus primal space fundamental solution kernel G_t are equivalent. In particular, there exists a bijection $\Pi^{-1} \circ \Xi : \mathbb{R}^{2n \times 2n} \rightarrow \mathbb{S}^{2n \times 2n}$ such that

$$\Lambda_t = \Pi^{-1} \circ \Xi(\Sigma_t), \quad \Sigma_t = \Xi^{-1} \circ \Pi(\Lambda_t) \quad (43)$$

for all $t \in \mathbb{R}_{>0}$. Consequently, it is natural to expect that Λ_t defines an alternative fundamental solution for DRE (1), (2).

Theorem 4 *Suppose that Assumptions 1 and 2 hold. Given any $P_0 \in \mathbb{S}_{>M}^{n \times n}$, the corresponding unique solution P_t of DRE (1), (2) exists and is given explicitly by*

$$P_t = \Lambda_t^{11} - \Lambda_t^{12} (P_0 + \Lambda_t^{22})^{-1} (\Lambda_t^{12})', \quad (44)$$

for all $t \in (0, t^*(P_0))$, where $\Lambda_t \in \mathbb{S}^{2n \times 2n}$ is as per (42), and the maximal horizon of existence is $t^*(P_0) \in \mathbb{R}_{>0}^+$ is

$$t^*(P_0) = \sup \{ t \in \mathbb{R}_{>0} \mid P_0 + \Lambda_t^{22} \in \mathbb{S}_{<0}^{n \times n} \}. \quad (45)$$

Proof See Appendix A.5. ■

By inspection of (4), (5), and (42), (44), it is evident that the symplectic and max-plus fundamental solutions both provide a characterization of all particular solutions of the DRE (1), (2). Furthermore, both provide characterizations of the corresponding finite escape time $t^*(P_0) \in \mathbb{R}_{>0}^+$, see (6), (45). However, by inspection, a crucial difference between these latter characterizations concerns their ease of evaluation, assuming their respective fundamental solutions are known for all time. In particular, the existence or otherwise of a finite escape at time t due to an initial condition $P_0 \in \mathbb{S}_{>M}^{n \times n}$ can be verified using the max-plus characterization (45) by testing the inequality $P_0 + \Lambda_t^{22} \in \mathbb{S}_{<0}^{n \times n}$ once. However, the same verification using the symplectic characterization (6) requires testing invertibility of $\Sigma_s^{11} + \Sigma_s^{12} P_0$ for all $s \in (0, t]$.

3.6 Relationship with the max-plus dual space fundamental solution of [7]

As indicated in Section 3.5, the max-plus fundamental solution defined by Λ_t of (42), (43) and Theorem 4 may be regarded as a max-plus *primal space* fundamental solution [22] for DRE (1), (2), as its evolution is tied directly to the dynamic programming evolution of the value function of the auxiliary optimal control problem (20) in the primal space of **semiconvex payoffs**. In contrast, the max-plus *dual space* fundamental solution of [7] is tied to the dynamic programming evolution of a value function in a dual space defined by the semiconvex transform. In particular, (41) appears analogously in [7] as

$$W_{t+\tau} = \mathcal{D}_\psi^{-1} \mathcal{B}_\tau^\oplus \mathcal{D}_\varphi W_t, \quad W_t = \mathcal{D}_\varphi^{-1} \mathcal{B}_t^\oplus \mathcal{D}_\varphi \Psi, \quad (46)$$

in which \mathcal{D}_φ is the semiconvex transform (11). In (46) and [7], \mathcal{B}_τ^\oplus is a max-plus linear max-plus integral operator of a form corresponding to (35), with

$$\begin{aligned} \mathcal{B}_\tau^\oplus a &\doteq \int_{\mathbb{R}^n}^\oplus B_t(\cdot, z) \otimes a(z) dz \quad \forall a \in \text{dom}(\mathcal{B}_\tau^\oplus), \\ \text{dom}(\mathcal{B}_\tau^\oplus) &\doteq \left\{ a \in \mathcal{S}_-^{-M} \mid \mathcal{B}_\tau^\oplus a \in \mathcal{S}_-^{-M} \right\}, \end{aligned} \quad (47)$$

in which \mathcal{S}_-^{-M} is as per (8), and kernel B_t is defined analogously to (36), by

$$B_t(y, z) \doteq (\mathcal{D}_\varphi S_t(\cdot, z))(y) \quad (48)$$

for all $y, z \in \mathbb{R}^n$, $t \in \mathbb{R}_{>0}$. By inspection of (36) and (48), the kernels G_t and B_t defining the primal and dual space fundamental solutions follow by taking the semiconvex transform of the value function S_t of (20) with respect to either its second or first variable respectively. Furthermore, (15), (41), (46) imply that

$$\mathcal{S}_t = \mathcal{G}_t^\oplus = \mathcal{D}_\varphi^{-1} \mathcal{B}_t^\oplus \mathcal{D}_\varphi, \quad t \in \mathbb{R}_{>0}.$$

This identity, along with the semigroup property of (40) inherited from dynamic programming, naturally imply that \mathcal{B}_t^\oplus satisfies a corresponding semigroup property. The interested reader is referred to [7, 8, 22] for further details.

Remark 5 The primal and dual space fundamental solutions of Theorem 4 and [7] are applicable to DREs of the specific form (1), (2). They correspond to control problems defined via maximization of a payoff, as per (15), (16). For control problems defined via minimization of a cost, as is typically the case for LQR problems, the DRE analogous to (1) takes the form

$$\dot{P}_t = A'P_t + P_tA - P_tBB'P_t + C'C,$$

in which $P_t \in \mathbb{S}^{n \times n}$ analogously describes a particular solution evolved forward from an initial condition P_0 . In this case, it is anticipated that the development of an applicable *min-plus fundamental solution* follows by exploiting the min-plus linearity, semiconcavity, and semigroup properties of the dynamic programming evolution operator corresponding to (16). This development largely parallels that leading to Theorem 4, and the details are omitted.

4 FUNDAMENTAL SOLUTIONS SEMIGROUPS

Both the symplectic fundamental solution $\Sigma_t \in \mathbb{R}^{2n \times 2n}$ and the max-plus primal space fundamental solution $\Lambda_t \in \mathbb{S}^{2n \times 2n}$, specified respectively by (5) and (42), provide a path for establishing existence of a unique particular solution P_t of DRE (1), (2) on the time interval $[0, t] \in \mathbb{R}_{>0}$, $t \in \mathbb{R}_{>0}$, and computing that solution. As the term *fundamental solution* implies, this is possible for any initial data P_0 satisfying (2). Indeed, both fundamental solutions can

be evolved to longer time horizons *independently* of any specific initial data for the DRE, thereby giving rise to [corresponding symplectic](#) and *max-plus fundamental solution semigroups of matrices*. In defining the latter max-plus fundamental solution semigroup, it is useful to define the Schur complement matrix operation \circledast acting on $\Lambda, \hat{\Lambda} \in \mathbb{S}^{2n \times 2n}$ by

$$\begin{aligned} [\Lambda \circledast \hat{\Lambda}]^{11} &\doteq \Lambda^{11} - \Lambda^{12}(\hat{\Lambda}^{11} + \Lambda^{22})^+(\Lambda^{12})', \\ [\Lambda \circledast \hat{\Lambda}]^{12} &\doteq -\Lambda^{12}(\hat{\Lambda}^{11} + \Lambda^{22})^+ \hat{\Lambda}^{12}, \\ [\Lambda \circledast \hat{\Lambda}]^{21} &\doteq ([\Lambda \circledast \hat{\Lambda}]^{12})', \\ [\Lambda \circledast \hat{\Lambda}]^{22} &\doteq \hat{\Lambda}^{22} - (\hat{\Lambda}^{12})'(\hat{\Lambda}^{11} + \Lambda^{22})^+ \hat{\Lambda}^{12}, \end{aligned} \tag{49}$$

using the notation of (3), in which $(\cdot)^+$ denotes the Moore-Penrose inverse.

Theorem 5 *Under Assumptions 1 and 2, the families of matrices $\{\Sigma_t\}_{t \in \mathbb{R}_{>0}}$ and $\{\Lambda_t\}_{t \in \mathbb{R}_{>0}}$ defined by (5) and (42), and related via the bijection $\Pi^{-1} \circ \Xi$ of (43), define a pair of one-parameter semigroups of matrices in $\mathbb{R}^{2n \times 2n}$ satisfying*

$$\Sigma_{t+s} = \Sigma_t \Sigma_s, \quad \Lambda_{t+s} = \Lambda_t \circledast \Lambda_s, \tag{50}$$

for all $t, s \in \mathbb{R}_{>0}$, in which the respective associative binary operations are standard matrix multiplication, and the matrix operation \circledast of (49).

Proof Fix $t, s \in \mathbb{R}_{>0}$. The left-hand semigroup property in (50) is immediate by definition (5) of the symplectic fundamental solution Σ_t . With Assumptions 1 and 2 asserted, [Theorems 1 and 3](#) imply that (43) holds, so that $\Lambda_t, \Lambda_s, \Lambda_{t+s} \in \mathbb{S}^{2n \times 2n}$ are well-defined. Furthermore, [Theorem 2](#) and (40) imply that for any $\psi \in \text{dom}(\mathcal{G}_{t+s}^\oplus) \subset \text{dom}(\mathcal{G}_s^\oplus)$,

$$\begin{aligned} \mathcal{G}_t^\oplus \mathcal{G}_s^\oplus \psi &= \int_{\mathbb{R}^n}^\oplus G_t(\cdot, \eta) \otimes \left[\int_{\mathbb{R}^n}^\oplus G_s(\eta, y) \otimes \psi(y) dy \right] d\eta \\ &= \int_{\mathbb{R}^n}^\oplus \left[\int_{\mathbb{R}^n}^\oplus G_t(\cdot, \eta) \otimes G_s(\eta, y) d\eta \right] \otimes \psi(y) dy \\ &= \mathcal{G}_{t+s}^\oplus \psi = \int_{\mathbb{R}^n}^\oplus G_{t+s}(\cdot, y) \otimes \psi(y) dy. \end{aligned}$$

Fix any $x, y \in \mathbb{R}^n$. Applying a [semiconvex version](#) of [7, Lemma 4.5] to equate the kernels of the left- and right-hand sides above, [Theorem 3](#) also implies that

$$\begin{aligned} \frac{1}{2} \begin{bmatrix} x \\ y \end{bmatrix}' \Lambda_{t+s} \begin{bmatrix} x \\ y \end{bmatrix} &= G_{t+s}(x, y) = \int_{\mathbb{R}^n}^\oplus G_t(x, \eta) \otimes G_s(\eta, y) d\eta \\ &= \int_{\mathbb{R}^n}^\oplus \frac{1}{2} \begin{bmatrix} x \\ \eta \end{bmatrix}' \Lambda_t \begin{bmatrix} x \\ \eta \end{bmatrix} \otimes \frac{1}{2} \begin{bmatrix} \eta \\ y \end{bmatrix}' \Lambda_s \begin{bmatrix} \eta \\ y \end{bmatrix} d\eta = \int_{\mathbb{R}^n}^\oplus \lambda_{x,y}(\eta) d\eta \end{aligned} \tag{51}$$

where $\lambda_{x,y} : \mathbb{R}^n \rightarrow \mathbb{R}$ is defined by

$$\begin{aligned} \lambda_{x,y}(\eta) &\doteq \frac{1}{2} \begin{bmatrix} x \\ y \\ \eta \end{bmatrix}' \left[\begin{array}{cc|c} A_t^{11} & 0 & A_t^{12} \\ 0 & A_s^{22} & (A_s^{12})' \\ \hline (A_t^{12})' & A_s^{12} & A_s^{11} + A_t^{22} \end{array} \right] \begin{bmatrix} x \\ y \\ \eta \end{bmatrix} \\ &= \frac{1}{2} \eta' (A_s^{11} + A_t^{22}) \eta + \eta' \begin{bmatrix} A_t^{12} \\ (A_s^{12})' \end{bmatrix}' \begin{bmatrix} x \\ y \end{bmatrix} + \frac{1}{2} \begin{bmatrix} x \\ y \end{bmatrix}' \begin{bmatrix} A_t^{11} & 0 \\ 0 & A_s^{22} \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} \end{aligned}$$

for all $\eta \in \mathbb{R}^n$. As $A_{t+s} \in \mathbb{R}^{2n \times 2n}$ is well-defined, note that $G_{t+s}(x, y) \in \mathbb{R}$, see also Lemma 2. That is, $\sup_{\eta \in \mathbb{R}^n} \lambda_{x,y}(\eta) \in \mathbb{R}$. Consequently, applying [8, Lemma E.2], the following properties hold:

- 1) $A_s^{11} + A_t^{22} \in \mathbb{S}_{\leq 0}^{n \times n}$;
- 2) the Moore-Penrose pseudo-inverse $(A_s^{11} + A_t^{22})^+ \in \mathbb{S}_{\leq 0}^{n \times n}$ exists; and
- 3) there exists a $\eta^* \in \mathbb{R}^n$ given by

$$\eta^* \doteq -(A_t^{22} + A_s^{11})^+ \begin{bmatrix} A_t^{12} \\ (A_s^{12})' \end{bmatrix}' \begin{bmatrix} x \\ y \end{bmatrix}$$

such that

$$\begin{aligned} \int_{\mathbb{R}^n}^{\oplus} \lambda_{x,y}(\eta) d\eta &= \lambda_{x,y}(\eta^*) = \frac{1}{2} \begin{bmatrix} x \\ y \end{bmatrix}' \begin{bmatrix} A_t^{11} & 0 \\ 0 & A_s^{22} \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} \\ &\quad - \frac{1}{2} \begin{bmatrix} x \\ y \end{bmatrix}' \begin{bmatrix} A_t^{12} \\ (A_s^{12})' \end{bmatrix} (A_s^{11} + A_t^{22})^+ \begin{bmatrix} A_t^{12} \\ (A_s^{12})' \end{bmatrix}' \begin{bmatrix} x \\ y \end{bmatrix}. \end{aligned}$$

Applying this last property in (51) and recalling that $x, y \in \mathbb{R}^n$ are arbitrary,

$$A_{t+s} = \begin{bmatrix} A_t^{11} & 0 \\ 0 & A_s^{22} \end{bmatrix} - \begin{bmatrix} A_t^{12} \\ (A_s^{12})' \end{bmatrix} (A_s^{11} + A_t^{22})^+ \begin{bmatrix} A_t^{12} \\ (A_s^{12})' \end{bmatrix}',$$

which yields the right-hand semigroup property in (50) via (49). \blacksquare

The semigroups properties (50) also naturally define respective notions of exponentiation. In particular,

$$\begin{aligned} \Sigma_t &= (\Sigma_1)^t, & \Sigma_1 &= \exp(\mathcal{H}), \\ A_t &= (A_1)^{\otimes t}, & A_1 &= \Pi^{-1} \circ \Xi \circ \exp(\mathcal{H}), \end{aligned} \tag{52}$$

in which $\mathcal{H}, \Xi, \Pi^{-1}$ are as per (5), (64), (81), and the exponentiations $(\cdot)^t, (\cdot)^{\otimes t}$ denote (respectively) the standard matrix exponentiation, and an exponentiation defined with respect to the \otimes operation of (49), see [7, Section 5] and Remark 6 below. As is the case with standard matrix exponentiation, note that (50), (52) imply that

$$(A_1)^{\otimes(t+s)} = A_{t+s} = A_t \otimes A_s = (A_1)^{\otimes t} \otimes (A_1)^{\otimes s}.$$

for all $t, s \in \mathbb{R}_{>0}$.

Remark 6 [7, Section 5] The [right-hand semigroup property of \(50\)](#) facilitates the definition of \otimes -exponentiation for any positive integer $n \in \mathbb{N}$ by

$$(\Lambda_\tau)^{\otimes n} \doteq \underbrace{\Lambda_\tau \otimes \Lambda_\tau \otimes \cdots \otimes \Lambda_\tau}_{n \text{ times}} = \Lambda_{n\tau} \quad (53)$$

where $\tau \in \mathbb{R}_{>0}$. By inspection, $((\Lambda_\tau)^{\otimes m})^{\otimes n} = (\Lambda_\tau)^{\otimes mn}$ for all $m, n \in \mathbb{N}$. Using this observation, (53) can be extended to positive rational and subsequently positive real exponents. In particular, given $p \in \mathbb{Q}_{>0}$ and coprime $m, n \in \mathbb{N}$ such that $np = m$, (53) implies that $(\Lambda_\tau)^{\otimes m} = \Lambda_{mn(\tau/n)} = ((\Lambda_{\tau/n})^{\otimes m})^{\otimes n} \doteq ((\Lambda_\tau)^{\otimes p})^{\otimes n}$. **That is, the \otimes -exponent $(\Lambda_\tau)^{\otimes p}$ is uniquely defined by**

$$(\Lambda_\tau)^{\otimes p} \doteq (\Lambda_{\tau/n})^{\otimes m} \quad (54)$$

for all $p = m/n \in \mathbb{Q}_{>0}$, $m, n \in \mathbb{N}$ coprime. As \mathbb{Q} is dense in \mathbb{R} , and the map $\tau \mapsto \Lambda_\tau$, $\tau \in \mathbb{R}_{>0}$, is continuous by (5), (42), it immediately follows that $\Lambda_t = \lim_{p \in \mathbb{Q}_{>0}, p \rightarrow t} \Lambda_p$. As Λ_p can be replaced with the \otimes -exponent $(\Lambda_1)^{\otimes p}$ of (54), the \otimes -exponent $(\Lambda_1)^{\otimes t}$ of (52) is uniquely defined by

$$(\Lambda_1)^{\otimes t} \doteq \lim_{p \in \mathbb{Q}_{>0}, p \rightarrow t} (\Lambda_1)^{\otimes p} = \lim_{p \in \mathbb{Q}_{>0}, p \rightarrow t} (\Lambda_{1/n})^{\otimes m} \quad (55)$$

for all $t \in \mathbb{R}_{>0}$, identically to [7]. Note that in the right-hand equality of (55), coprime $m, n \in \mathbb{N}$ are uniquely defined for each $p \in \mathbb{Q}_{>0}$ in the limiting sequence. Where $t \in \mathbb{R}_{>0}$ is irrational, it follows immediately that $m, n \rightarrow \infty$.

5 SOLVING THE DRE (1), (2)

Theorems 4 and 5 together describe [a max-plus primal space fundamental solution semigroup of matrices \$\{\Lambda_t\}_{t \in \mathbb{R}_{>0}}\$](#) for propagating solutions $P_t \in \mathbb{S}^{n \times n}$ of DRE (1) forward in time $t \in \mathbb{R}_{>0}$ from initializations $P_0 \in \mathbb{S}_{>M}^{n \times n}$ as per (2). In particular, (49), (50) describe propagation of [this fundamental solution](#), while (44) specifies how [it](#) may be used to evaluate a particular [solution at a time \$t \in \mathbb{R}_{>0}\$, if it exists there](#). In addition, (45) provides a general characterization of the corresponding maximal horizon of existence $t^*(P_0) \in \mathbb{R}_{>0}^+$. By inspection, this characterization allows easy verification of whether a specific time t falls before a finite escape $t^*(P_0) \in \mathbb{R}_{>0}$, [if it exists](#), by testing if $P_0 + \Lambda_t^{22} \in \mathbb{S}_{<0}^{n \times n}$ at that time. Note that the corresponding verification using the characterization (6) provided by the symplectic fundamental solution (5) requires testing invertibility of a matrix over a range of times.

5.1 Recipe

A recipe that uses the one-parameter max-plus primal space fundamental semigroup $\{\Lambda_t\}_{t \in \mathbb{R}_{>0}}$ to compute the solution P_t of DRE (1) for any initialization $P_0 \in \mathbb{S}_{>M}^{n \times n}$ of the form (2), using a fixed time step $\delta \in \mathbb{R}_{>0}$, [follows from Theorems 4 and 5. This recipe is as follows:](#)

I. Initialize and propagate the semigroup (49), (50)

- ❶ (*Initialize basis*) Select $M \in \mathbb{S}^{n \times n}$ of (9), **guided by** Lemma 1. Check that Assumptions 1 and 2 hold.
- ❷ (*Initialize semigroup*) Fix time step $\delta \in \mathbb{R}_{>0}$ and maximal time horizon $\bar{t} \doteq K\delta \in \mathbb{R}_{>0}$ for some fixed $K \in \mathbb{N}$. Using the matrix operators Ξ , Π^{-1} of (64), (81), initialize an element of the semigroup by

$$A_\delta = \Pi^{-1} \circ \Xi \circ \exp(\mathcal{H}\delta) \in \mathbb{S}^{2n \times 2n}, \quad (56)$$

where $\mathcal{H} \in \mathbb{R}^{2n \times 2n}$ is the Hamiltonian matrix (5).

- ❸ (*Propagate semigroup*) Compute a subset $\{A_{k\delta}\}_{k \in \mathbb{N}_{\leq K}}$ of the semigroup, corresponding to a temporal grid defined by $\delta \in \mathbb{R}_{>0}$, via the evolution

$$A_{(k+1)\delta} = A_\delta \oplus A_{k\delta}, \quad k \in \mathbb{N}_{<K}, \quad (57)$$

as per (50). (See also Remark 7 below.)

II. Solve the DRE (1), (2)

- ❹ (*Initialize a solution*) Select $P_0 \in \mathbb{S}_{>M}^{n \times n}$. Set $k = 1$.
- ❺ (*Test for finite escape*) If $P_0 + A_{k\delta}^{22} \in \mathbb{S}_{<0}^{n \times n}$ as per (45) then evaluate the solution $P_{k\delta}$ at time step k as

$$P_{k\delta} = A_{k\delta}^{11} - A_{k\delta}^{12} (P_0 + A_{k\delta}^{22})^{-1} (A_{k\delta}^{12})' \quad (58)$$

as per (44). Otherwise, record a finite escape time as occurring in the interval $((k-1)\delta, k\delta]$ and exit.

- ❻ (*Iterate*) Increment k . If $k \in \mathbb{N}_{\leq K}$ then go to step ❺. Otherwise, exit.

As indicated, the recipe consists of a total of 6 steps, divided into two parts. Part I concerns the initialization and propagation of a subset of the one parameter semigroup of matrices $\{A_t\}_{t \in \mathbb{R}_{>0}}$ required for computing *any* particular solution of DRE (1), (2) up to a pre-specified time horizon $\bar{t} \in \mathbb{R}_{>0}$. Part II concerns the subsequent evaluation of such a particular solution (and corresponding finite escape, if it exists). Crucially, part I need only be completed once, with the elements of the semigroup computed there used repeatedly in part II in evaluating any particular solutions of interest, without modification. This is demonstrated by example.

Remark 7 For fast propagation of $A_{k\delta}$ to large $k \in \mathbb{N}_{\leq K}$, where $K \doteq 2^N$, $N \in \mathbb{N}$, the *linear* time-index accumulation in (57) can be replaced with time-index *doubling* [10], ie.

$$A_{(2^{\kappa+1})\delta} = A_{(2^\kappa)\delta} \otimes A_{(2^\kappa)\delta}. \quad \kappa \in \{0\} \cup \mathbb{N}_{<N}.$$

5.2 Example – no finite escape

In demonstrating the recipe described above, an example from [7] is considered. In particular, define $A \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{n \times m}$, $C \in \mathbb{R}^{p \times n}$, $n = m = p = 2$, by

$$A \doteq \begin{bmatrix} -2.000 & +1.600 \\ -1.600 & -0.400 \end{bmatrix}, \quad B \doteq \begin{bmatrix} +0.216 & -0.008 \\ -0.008 & +0.216 \end{bmatrix}^{\frac{1}{2}},$$

$$C \doteq \begin{bmatrix} +1.500 & +0.200 \\ +0.200 & +1.600 \end{bmatrix}^{\frac{1}{2}}, \quad M_0 \doteq \begin{bmatrix} +0.651 & -0.310 \\ -0.310 & +1.160 \end{bmatrix},$$

where $M_0 \in \mathbb{S}_{>0}^{n \times n}$ is the corresponding stabilizing solution of ARE (30) as per Lemma 1. In view of (31), select

$$M \doteq \begin{bmatrix} -1.000 & -0.200 \\ -0.200 & -1.000 \end{bmatrix} \in \mathbb{S}_{<M_0}^{n \times n}.$$

Consequently, Assumption 1 holds. Step ❶ is completed via a standard rank calculation to verify that Assumption 2 holds. Theorems 1, 3, and 5 subsequently imply that the one parameter semigroup of matrices $\{A_t\}_{t \in \mathbb{R}_{>0}}$ propagated by (49), (50) is well-defined and may be computed as indicated in steps ❷, ❸. With $\delta \doteq 0.05$, $K \doteq 80$, $\bar{t} \doteq 4$, (56) yields

$$A_\delta = \begin{bmatrix} -83.48 & -3.021 & +92.26 & -4.011 \\ -3.021 & -91.11 & +11.07 & +92.42 \\ +92.26 & +11.07 & -102.6 & -3.420 \\ -4.011 & +92.42 & -3.420 & -94.28 \end{bmatrix}.$$

Subsequently iterating via (57) as per step ❹ yields the required set of matrices $\{A_{k\delta}\}_{k \in \mathbb{N}_{\leq K}}$.




Fig. 1 Max-plus and RK45 solutions of DRE (1), (59).

In order to demonstrate the computation of a particular solution of DRE (1) via steps 4 – 6, select an initialization

$$P_0 \doteq -0.1 I \in \mathbb{S}_{>M}^{n \times n} \quad (59)$$

as per step 4 and (2). Iterating through $k \in \mathbb{N}_{\leq K}$ as per steps 5 and 6, testing for finite escape and applying (58), yields the computed solution $P_{k\delta}$, $k \in \mathbb{N}_{\leq K}$ of DRE (1), (59). This solution, along with corresponding symplectic and RK45 solutions, is illustrated in Figure 1. (Here, the MATLAB™ RK45 solver is used, with absolute and relative tolerances set to 10^{-12} .) All three solutions are in reasonable agreement. No finite escape is observed.

5.3 Example – finite escape

In order to illustrate finite escape phenomenon, the set of matrices $\{A_{k\delta}\}_{k \in \mathbb{N}_{\leq K}}$ computed above is reused to evaluate the particular solution of DRE (1), (2) for the initial condition

$$P_0 \doteq \begin{bmatrix} 2.000 & 0.000 \\ 0.000 & 6.500 \end{bmatrix} \in \mathbb{S}_{>M}^{n \times n}. \quad (60)$$

The problem data is otherwise unchanged. Using the initialization (60) in step 4 and iterating steps 5 and 6 yields the corresponding particular DRE solution. A finite escape is demonstrated to occur within the horizon $\bar{t} = 4$ of computation, with $t^*(P_0) \in (2.8, 2.9)$ established using (45). Figure 2




Fig. 2 Maximum eigenvalue map $t \mapsto \sigma_{\max}(P_0 + A_t^{22})$ and determinant map $t \mapsto \det(\Sigma_t^{11} + \Sigma_t^{12} P_0)$, $t = k\delta$.




Fig. 3 Max-plus and RK45 solutions of DRE (1), (60).




Fig. 4 $\log_{10}(\|\cdot\|_2)$ of errors between solutions of DRE (1), (60).

illustrates the maps $t \mapsto \sigma_{\max}(P_0 + \Lambda_t^{22})$ and $t \mapsto \det(\Sigma_t^{11} + \Sigma_t^{12} P_0)$, $t = k\delta$, $k \in \mathbb{N}_{\leq K}$, in which $\sigma_{\max}, \det : \mathbb{S}^{n \times n} \rightarrow \mathbb{R}$ denote the maximum eigenvalue and determinant maps respectively. Note specifically that zero crossings in both are collocated with the finite escape time, as per (6), (45). Note further that $t \mapsto \sigma_{\max}(P_0 + \Lambda_t^{22})$ defines a monotone non-decreasing function. This monotonicity follows from that used to establish the representation (45) of the finite escape time $t^*(P_0)$, see the proof of Theorem 4. It guarantees that no finite escape occurs prior to this zero crossing.

The computed solution P_t of DRE (1), (60) for $t = k\delta$, $k \in \mathbb{N}_{\leq K}$, is illustrated in Figure 3, along with the corresponding symplectic and RK45

solutions. These solutions are in good agreement, as measured by the absolute errors illustrated in Figure 4. As may be observed in the latter figure, these errors increase immediately prior to the finite escape time as the entries of $P_t \in \mathbb{S}^{n \times n}$ diverge to $\pm\infty$. For brevity, an error analysis is not included.

6 CONCLUSIONS

A **fundamental solution** for a class of differential Riccati equations (DREs) is developed using tools from max-plus and semiconvex analysis. It is shown that this fundamental solution is defined by a corresponding fundamental solution semigroup, which describes the evolution of all particular solutions of the DRE, on all time horizons. A simple characterization of finite escape times for these particular solutions is also provided.

A Proofs

A.1 Proof of Theorem 1

Since $Q_t \in \mathbb{R}^{2n \times 2n}$ satisfies DRE (22), (23) for all $t \in \mathbb{R}_{\geq 0}$, see (28), it may be represented by a corresponding symplectic fundamental solution of the form (5), denoted here by $\hat{\Sigma}_t \in \mathbb{R}^{4n \times 4n}$. In order to apply (5), define $\hat{\mathcal{H}}, \Delta \in \mathbb{R}^{4n \times 4n}$ by

$$\hat{\mathcal{H}} \doteq \left[\begin{array}{c|c} -\hat{A} & -\hat{B}\hat{B}' \\ \hline \hat{C}'\hat{C} & \hat{A}' \end{array} \right], \quad \Delta \doteq \left[\begin{array}{c|c} I & 0 \\ \hline 0 & I \end{array} \right],$$

where $\hat{A} \in \mathbb{R}^{2n \times 2n}$, $\hat{B} \in \mathbb{R}^{2n \times m}$, $\hat{C} \in \mathbb{R}^{p \times 2n}$ are as per (24). Note by inspection that $\Delta = \Delta' = \Delta^{-1}$. By substitution, a straightforward calculation yields that

$$\hat{\mathcal{H}} = \Delta \left[\begin{array}{c|c} \mathcal{H} & 0 \\ \hline 0 & 0 \end{array} \right] \Delta,$$

where $\mathcal{H} \in \mathbb{R}^{2n \times 2n}$ is as per (5). Hence, the symplectic fundamental solution $\hat{\Sigma}_t$ for DRE (22), (23) is, again by (5),

$$\hat{\Sigma}_t = \exp(\hat{\mathcal{H}}t) = \Delta \left[\begin{array}{c|c} \exp(\mathcal{H}t) & 0 \\ \hline 0 & I \end{array} \right] \Delta = \Delta \left[\begin{array}{c|c} \Sigma_t & 0 \\ \hline 0 & I \end{array} \right] \Delta = \left[\begin{array}{c|c} \Sigma_t^{11} & 0 \\ \hline 0 & I \end{array} \right] \left[\begin{array}{c|c} \Sigma_t^{12} & 0 \\ \hline \Sigma_t^{21} & 0 \\ \hline 0 & I \end{array} \right], \quad (61)$$

for all $t \in \mathbb{R}_{\geq 0}$, where the notation of (3) has been applied. Hence, the particular solution Q_t of DRE (22), (23) is given in terms of the symplectic fundamental solution (5), with respect to $\hat{\Sigma}_t$, by

$$Q_t = \hat{Y}_t \hat{X}_t^{-1} \quad (62)$$

for all $t \in [0, t^*(Q_0)) \equiv \mathbb{R}_{\geq 0}$, see (28), in which

$$\left[\begin{array}{c} \hat{X}_t \\ \hat{Y}_t \end{array} \right] \doteq \hat{\Sigma}_t \left[\begin{array}{c} I \\ Q_0 \end{array} \right] = \hat{\Sigma}_t \left[\begin{array}{c} I \\ \mu(M) \end{array} \right] = \left[\begin{array}{c|c} \Sigma_t^{11} & 0 \\ \hline 0 & I \end{array} \right] \left[\begin{array}{c|c} \Sigma_t^{12} & 0 \\ \hline \Sigma_t^{21} & 0 \\ \hline 0 & I \end{array} \right] \left[\begin{array}{c} I & 0 \\ 0 & I \\ +M & -M \\ -M & +M \end{array} \right]$$

$$= \begin{bmatrix} \Sigma_t^{11} + \Sigma_t^{12}M - \Sigma_t^{12}M & \\ 0 & I \\ \Sigma_t^{21} + \Sigma_t^{22}M - \Sigma_t^{22}M & \\ -M & +M \end{bmatrix} \in \mathbb{R}^{4n \times 2n}, \quad (63)$$

and $\mu(M)$ is defined by (10). For any fixed $t \in \mathbb{R}_{\geq 0}$, note in particular that

$$\widehat{X}_t^{-1} = \begin{bmatrix} (\Sigma_t^{11} + \Sigma_t^{12}M)^{-1} & (\Sigma_t^{11} + \Sigma_t^{12}M)^{-1} \Sigma_t^{12}M \\ 0 & I \end{bmatrix},$$

in which $(\Sigma_t^{11} + \Sigma_t^{12}M)^{-1}$ is well-defined as $t^*(M) = +\infty$ by Assumption 1 and (6), (28). That is, \widehat{X}_t^{-1} is well-defined. Its substitution in (62), along with \widehat{Y}_t from (63), yields $Q_t = \widehat{Y}_t \widehat{X}_t^{-1} \doteq \Xi(\Sigma_t)$, where $\Xi : \mathbb{R}^{2n \times 2n} \rightarrow \mathbb{R}^{2n \times 2n}$ is defined by

$$\begin{aligned} \Xi(\Sigma) &\doteq \begin{bmatrix} \Xi^{11}(\Sigma) & \Xi^{12}(\Sigma) \\ \Xi^{21}(\Sigma) & \Xi^{22}(\Sigma) \end{bmatrix}, \\ \text{dom}(\Xi) &\doteq \left\{ \Sigma \in \mathbb{R}^{2n \times 2n} \mid \begin{array}{l} \Sigma^{11} + \Sigma^{12}M \in \mathbb{R}^{n \times n} \\ \text{invertible} \end{array} \right\}, \end{aligned} \quad (64)$$

using the notation of (3), with

$$\begin{aligned} \Xi^{11}(\Sigma) &\doteq (\Sigma^{21} + \Sigma^{22}M)(\Sigma^{11} + \Sigma^{12}M)^{-1}, \\ \Xi^{12}(\Sigma) &\doteq \Xi^{11}(\Sigma) \Sigma^{12}M - \Sigma^{22}M, \\ \Xi^{21}(\Sigma) &\doteq -M(\Sigma^{11} + \Sigma^{12}M)^{-1}, \\ \Xi^{22}(\Sigma) &\doteq \Xi^{21}(\Sigma) \Sigma^{12}M + M. \end{aligned}$$

As M is invertible by Assumption 1, it may be verified directly that Ξ of (64) is invertible, with $\Xi^{-1} : \mathbb{R}^{2n \times 2n} \rightarrow \mathbb{R}^{2n \times 2n}$ given by

$$\begin{aligned} \Xi^{-1}(Q) &\doteq \begin{bmatrix} (\Xi^{-1})^{11}(Q) & (\Xi^{-1})^{12}(Q) \\ (\Xi^{-1})^{21}(Q) & (\Xi^{-1})^{22}(Q) \end{bmatrix}, \\ \text{dom}(\Xi^{-1}) &\doteq \left\{ Q \in \mathbb{R}^{2n \times 2n} \mid Q^{21} \in \mathbb{R}^{n \times n} \text{ invertible} \right\}. \end{aligned} \quad (65)$$

where

$$\begin{aligned} (\Xi^{-1})^{11}(Q) &\doteq -(Q^{21})^{-1}Q^{22}, \\ (\Xi^{-1})^{12}(Q) &\doteq -(Q^{21})^{-1}(M - Q^{22})M^{-1}, \\ (\Xi^{-1})^{21}(Q) &\doteq Q^{11}(\Xi^{-1})^{11}(Q) + Q^{12}, \\ (\Xi^{-1})^{22}(Q) &\doteq Q^{11}(\Xi^{-1})^{12}(Q) - Q^{12}M^{-1}. \end{aligned}$$

That is, (29) holds. ■

A.2 Proof of Lemma 1

Fix $M_0 \in \mathbb{S}_{\geq 0}^{n \times n}$ as the stabilizing solution of ARE (30) indicated in the lemma statement. Let $t^*(M_0) \in \mathbb{R}_{> 0}^+$ denote the maximal horizon of existence (6) of the DRE

$$\dot{R}_t = A'R_t + R_tA + R_tBB'R_t + C'C, \quad R_0 = M_0. \quad (66)$$

As M_0 is the stabilizing solution of ARE (30), note that $R_t \doteq M_0$ is the unique solution of this DRE for all $t \in \mathbb{R}_{\geq 0}$. **Consequently**, $t^*(M_0) = +\infty$. Choose any invertible $M \in \mathbb{S}^{n \times n}$ such that (31) holds, and note that such a choice is always possible. Recalling (25), let $Q_t^{11} \in \mathbb{S}^{n \times n}$, $t \in [0, t^*(M))$ denote the unique solution of DRE (25) initialized with $Q_0^{11} = M$. As

DREs (25) and (66) are identical, [Proposition 1 \(see below\)](#) and (31) imply that solutions Q_t^{11} and R_t satisfy the monotonicity property

$$Q_t^{11} - M_0 = Q_t^{11} - R_t \in \mathbb{S}_{\leq 0}^{n \times n} \quad (67)$$

for all $t \in [0, t^*(M))$. By inspection, this provides an upper bound for Q_t^{11} . In order to determine a lower bound, choose $w_s = 0$ for all $s \in [0, t]$ suboptimal in the definition (20) of $S_t(x, 0)$. Recalling (16), (17), (21),

$$\frac{1}{2}x'Q_t^{11}x = S_t(x, 0) \geq \frac{1}{2}x'O_t x \quad (68)$$

for all $x \in \mathbb{R}^n$, in which $O_t \in \mathbb{S}^{n \times n}$ is well-defined by

$$O_t \doteq \exp(A't) M \exp(At) + \int_0^t \exp(A's) C' C \exp(As) ds$$

for all $t \in \mathbb{R}_{\geq 0}$. Note that $O_t \in \mathbb{S}^{n \times n}$ is finite for all $t \in \mathbb{R}_{\geq 0}$, and provides a lower bound for $Q_t^{11} \in \mathbb{S}^{n \times n}$. Hence, combining (67) and (68),

$$Q_t^{11} \in \mathbb{S}_{\geq O_t}^{n \times n} \cap \mathbb{S}_{\leq M_0}^{n \times n}$$

for all $t \in [0, t^*(M))$. A simple contradiction argument subsequently implies that $Q_t^{11} \in \mathbb{S}^{n \times n}$ is finite for all $t \in \mathbb{R}_{\geq 0}$, so that $t^*(M) = +\infty$. [As \$M^{-1}\$ exists by definition, it follows that Assumption 1 holds, as required.](#) ■

Proposition 1 *Given initializations $P_0, \tilde{P}_0 \in \mathbb{S}^{n \times n}$ satisfying $P_0 - \tilde{P}_0 \in \mathbb{S}_{\leq 0}^{n \times n}$, and $t^* \doteq \min(t^*(P_0), t^*(\tilde{P}_0))$, the respective unique solutions $P_t, \tilde{P}_t \in \mathbb{S}^{2n \times 2n}$ of DRE (1) defined for all $t \in [0, t^*)$, satisfy*

$$P_t - \tilde{P}_t \in \mathbb{S}_{\leq 0}^{n \times n} \quad (69)$$

for all $t \in [0, t^*)$.

Proof Fix $t \in [0, t^*)$. Let $\mathcal{T} : \Delta_{0,t} \rightarrow \mathbb{R}^{n \times n}$ denote the evolution operator associated with the time-dependent ordinary differential equation (ODE)

$$\dot{Y}_s = (\hat{A} + \frac{1}{2}\hat{B}\hat{B}'(P_s + \tilde{P}_s))' Y_s,$$

defined for $s \in [0, t]$, with $\Delta_{0,t} \doteq \{(r, s) \in \mathbb{R}_{\geq 0}^2 \mid 0 \leq r \leq s \leq t\}$. By definition, see for example [23, Proposition 3.6, p.138],

$$\begin{aligned} \mathcal{T}(s, \sigma) &= I, \\ \frac{\partial}{\partial s} \mathcal{T}(s, \sigma) &= (\hat{A} + \frac{1}{2}\hat{B}\hat{B}'(P_s + \tilde{P}_s))' \mathcal{T}(s, \sigma), \\ \frac{\partial}{\partial \sigma} \mathcal{T}(s, \sigma) &= -\mathcal{T}(s, \sigma) (\hat{A} + \frac{1}{2}\hat{B}\hat{B}'(P_\sigma + \tilde{P}_\sigma))', \end{aligned} \quad (70)$$

for all $(s, \sigma) \in \Delta_{0,t}$. Define $\pi : [0, t] \rightarrow \mathbb{S}^{n \times n}$ by

$$\pi_\sigma \doteq \mathcal{T}(t, \sigma) (P_\sigma - \tilde{P}_\sigma) \mathcal{T}(t, \sigma)' \quad (71)$$

for all $\sigma \in [0, t]$. Differentiating with respect to σ ,

$$\begin{aligned} \dot{\pi}_\sigma &= \frac{\partial}{\partial \sigma} \mathcal{T}(t, \sigma) (P_\sigma - \tilde{P}_\sigma) \mathcal{T}(t, \sigma)' + \mathcal{T}(t, \sigma) (\dot{P}_\sigma - \dot{\tilde{P}}_\sigma) \mathcal{T}(t, \sigma)' \\ &\quad + \mathcal{T}(t, \sigma) (P_\sigma - \tilde{P}_\sigma) \frac{\partial}{\partial \sigma} \mathcal{T}(t, \sigma)' \\ &= \mathcal{T}(t, \sigma) \Gamma_\sigma \mathcal{T}(t, \sigma)' \end{aligned} \quad (72)$$

for all $\sigma \in [0, t]$, where

$$\begin{aligned} \Gamma_\sigma &\doteq (\dot{P}_\sigma - \dot{\tilde{P}}_\sigma) - (\hat{A} + \frac{1}{2}\hat{B}\hat{B}'(P_\sigma + \tilde{P}_\sigma))'(P_\sigma - \tilde{P}_\sigma) - (P_\sigma - \tilde{P}_\sigma)(\hat{A} + \frac{1}{2}\hat{B}\hat{B}'(P_\sigma + \tilde{P}_\sigma)) \\ &= 0, \end{aligned}$$

in which the equality with zero follows by virtue of the fact that $P_\sigma, \tilde{P}_\sigma$ both satisfy the DRE (1). Consequently, (72) implies that $\dot{\pi}_\sigma = 0$ for all $\sigma \in [0, t]$, so that integration with respect to $\sigma \in [0, t]$ yields $\pi_t = \pi_0$. Recalling (71), it follows immediately that

$$P_t - \tilde{P}_t = \pi_t = \pi_0 = \mathcal{T}(t, 0)(P_0 - \tilde{P}_0)'\mathcal{T}(t, 0)'$$

Recalling that $P_0 - \tilde{P}_0 \in \mathbb{S}_{\leq 0}^{n \times n}$, and noting that $t \in [0, t^*)$ is arbitrary, yields the required assertion (69). \blacksquare

A.3 Proof of Lemma 2

Suppose that Assumption 1 holds. Fix $x, y \in \mathbb{R}^n$, $t \in \mathbb{R}_{>0}$. Note that $G_t(x, y) \in \mathbb{R}^-$ by Theorem 2.

(*Necessity*) Suppose that $G_t(x, y) \in \mathbb{R}$. Recalling the value function interpretation of $G_t(x, y)$, if the dynamics (18) are not controllable from x to y in time t , it immediately follows by definitions (16), (36) that $G_t(x, y) = -\infty$. Hence, the dynamics (18) must be controllable from x to y in time t . Necessity follows as $x, y \in \mathbb{R}^n$ and $t \in \mathbb{R}_{>0}$ are arbitrary.

(*Sufficiency*) Suppose that dynamics (18) are controllable. Proposition 2 (see below) implies that $Q_t^{22} \in \mathbb{S}_{>M}^{n \times n}$, where Q_t^{22} is as per (27). Hence, $S_t(x, \cdot) \in \mathcal{S}_+^{-M} = \text{dom}(\mathcal{D}_\varphi)$, so that $\mathcal{D}_\varphi S_t(x, \cdot) \in \mathcal{S}_-^{-M}$ is well defined. So, applying the semiconvex transform (11) to $S_t(x, \cdot)$ yields

$$\begin{aligned} (\mathcal{D}_\varphi S_t(x, \cdot))(y) &= -\int_{\mathbb{R}^n}^\oplus \varphi(\xi, y) \otimes (-S_t(x, \xi)) d\xi \\ &= -\int_{\mathbb{R}^n}^\oplus \frac{1}{2} \begin{bmatrix} \xi \\ y \end{bmatrix}' \mu(M) \begin{bmatrix} \xi \\ y \end{bmatrix} - \frac{1}{2} \begin{bmatrix} x \\ \xi \end{bmatrix}' Q_t \begin{bmatrix} x \\ \xi \end{bmatrix} d\xi \\ &= -\int_{\mathbb{R}^n}^\oplus \frac{1}{2} \begin{bmatrix} x \\ y \\ \xi \end{bmatrix}' \begin{bmatrix} -Q_t^{11} & 0 & -Q_t^{12} \\ 0 & +M & -M \\ -(Q_t^{12})' & -M & M - Q_t^{22} \end{bmatrix} \begin{bmatrix} x \\ y \\ \xi \end{bmatrix} d\xi \\ &= -\frac{1}{2} \begin{bmatrix} x \\ y \end{bmatrix}' \begin{bmatrix} -Q_t^{11} & 0 \\ 0 & +M \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \frac{1}{2} \begin{bmatrix} x \\ y \end{bmatrix}' \begin{bmatrix} -Q_t^{12} \\ -M \end{bmatrix} (M - Q_t^{22})^{-1} \begin{bmatrix} -Q_t^{12} \\ -M \end{bmatrix}' \begin{bmatrix} x \\ y \end{bmatrix} \\ &\doteq \frac{1}{2} \begin{bmatrix} x \\ y \end{bmatrix}' \Lambda_t \begin{bmatrix} x \\ y \end{bmatrix}, \end{aligned} \tag{73}$$

in which $(M - Q_t^{22})^{-1}$ is guaranteed to exist, so that $\Lambda_t \in \mathbb{R}^{2n \times 2n}$ exists and is finite by definition. Hence, applying the right-hand equality of (36) in Theorem 2,

$$G_t(x, y) = \frac{1}{2} \begin{bmatrix} x \\ y \end{bmatrix}' \Lambda_t \begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}, \tag{74}$$

which is finite, thereby demonstrating sufficiency. \blacksquare

Proposition 2 Under Assumption 1, controllability of the dynamics (18) implies that $Q_t^{22} \in \mathbb{S}_{>M}^{n \times n}$ for all $t \in \mathbb{R}_{>0}$.

Proof With $M \in \mathbb{S}^{n \times n}$ satisfying Assumption 1, recall that $t^*(M) = +\infty$ as per (28). Fix any $t \in \mathbb{R}_{>0}$. Consequently, the optimal dynamics associated with $S_t(x, y)$ of (20), (21) are well-defined by the time-dependent ODE

$$\dot{x}_s^* = (A + BB'Q_{t-s}^{11})x_s^*, \quad x_0 = x, \quad (75)$$

for all $s \in [0, t]$. Let $\mathcal{V}_t : \Delta_{0,t} \rightarrow \mathbb{R}^{n \times n}$ denote the evolution operator associated with (75), with $\Delta_{0,t} \doteq \{(r, s) \in \mathbb{R}_{\geq 0}^2 \mid 0 \leq r \leq s \leq t\}$. By definition, see for example [23, Proposition 3.6, p.138],

$$\begin{aligned} \mathcal{V}_t(\sigma, \sigma) &= I, \\ \frac{\partial}{\partial s} \mathcal{V}_t(s, \sigma) &= (A + BB'Q_{t-s}^{11})\mathcal{V}_t(s, \sigma), \\ \frac{\partial}{\partial \sigma} \mathcal{V}_t(s, \sigma) &= -\mathcal{V}_t(s, \sigma)(A + BB'Q_{t-\sigma}^{11}), \end{aligned} \quad (76)$$

for all $(s, \sigma) \in \Delta_{0,t}$. Define $\mathcal{U}_t : \Delta_{0,t} \rightarrow \mathbb{R}^{n \times n}$ via (76) by

$$\mathcal{U}_t(r, \tau) \doteq \mathcal{V}_t(t - \tau, t - r)' \quad (77)$$

for all $(r, \tau) \in \Delta_{0,t}$. By inspection of (76), (77),

$$\begin{aligned} \mathcal{U}_t(\tau, \tau) &= I, \\ \frac{\partial}{\partial r} \mathcal{U}_t(r, \tau) &= \left[\frac{\partial}{\partial \sigma} \mathcal{V}_t(s, \sigma) \Big|_{(s, \sigma) = (t - \tau, t - r)} \right]' (-1) \\ &= (A + BB'Q_r^{11})'\mathcal{V}_t(t - \tau, t - r)' = (A + BB'Q_r^{11})'\mathcal{U}_t(r, \tau), \\ \frac{\partial}{\partial \tau} \mathcal{U}_t(r, \tau) &= \left[\frac{\partial}{\partial s} \mathcal{V}_t(s, \sigma) \Big|_{(s, \sigma) = (t - \tau, t - r)} \right]' (-1) \\ &= -\mathcal{V}_t(t - \tau, t - r)'(A + BB'Q_\tau^{11})' = -\mathcal{U}_t(r, \tau)(A + BB'Q_\tau^{11})' \end{aligned} \quad (78)$$

That is, $\mathcal{U}_t : \Delta_{0,t} \rightarrow \mathbb{R}^{n \times n}$ is the evolution operator for the dynamics associated with $(A + BB'Q_s^{11})'$, $s \in [0, t]$. Comparing with (26), it immediately follows that $Q_s^{12} = -\mathcal{U}_t(s, 0)M$ for all $s \in [0, t]$. Hence, (27) implies that

$$\begin{aligned} Q_t^{22} - M &= \int_0^t (Q_s^{12})' BB' Q_s^{12} ds \\ &= \int_0^t M \mathcal{U}_t(s, 0)' BB' \mathcal{U}_t(s, 0) M ds = M C_t M \end{aligned} \quad (79)$$

where $C_t \doteq \int_0^t \mathcal{V}_t(t, t-s) BB' \mathcal{V}_t(t, t-s)' ds \in \mathbb{S}_{\geq 0}^{n \times n}$ is the controllability gramian for the pair $(A + BB'Q_{t-s}^{11}, B)$ on $[0, t]$, by definition of \mathcal{V}_t . However, recall that controllability is preserved under state feedback, see for example [1, p.48]. Hence, (A, B) completely controllable implies that $(A + BB'Q_{t-s}^{11}, B)$ is completely controllable, which in turn implies that C_t is invertible for $t \in \mathbb{R}_{>0}$. That is, $C_t \in \mathbb{S}_{> 0}^{n \times n}$ for all $t \in \mathbb{R}_{>0}$. As M is invertible by Assumption 1, the assertion immediately follows by (79). \blacksquare

A.4 Proof of Theorem 3

Fix any $t \in \mathbb{R}_{>0}$, $x \in \mathbb{R}^n$. Applying Lemma 2, and in particular (73), (74), it follows that $Q_t \in \mathbb{R}^{2n \times 2n}$, $\Lambda_t \in \mathbb{S}^{2n \times 2n}$ of (22), (73) are related via

$$Q_t = \Pi(\Lambda_t), \quad \Lambda_t = \Pi^{-1}(Q_t),$$

with matrix operators $\Pi, \Pi^{-1} : \mathbb{S}^{2n \times 2n} \rightarrow \mathbb{S}^{2n \times 2n}$ defined using the notation of (3) by

$$\Pi(\Lambda) \doteq \begin{bmatrix} \Lambda^{11} - \Lambda^{12}(M + \Lambda^{22})^{-1}(\Lambda^{12})' & \Lambda^{12}(M + \Lambda^{22})^{-1}M \\ M(M + \Lambda^{22})^{-1}(\Lambda^{12})' & M - M(M + \Lambda^{22})^{-1}M \end{bmatrix}$$

$$\text{dom}(\Pi) \doteq \left\{ A \in \mathbb{S}^{2n \times 2n} \mid A^{22} \in \mathbb{S}_{< -M}^{n \times n} \right\}, \quad (80)$$

$$\Pi^{-1}(Q) \doteq \begin{bmatrix} Q^{11} + Q^{12}(M - Q^{22})^{-1}(Q^{12})' & Q^{12}(M - Q^{22})^{-1}M \\ M(M - Q^{22})^{-1}(Q^{12})' & M(M - Q^{22})^{-1}M - M \end{bmatrix}$$

$$\text{dom}(\Pi^{-1}) \doteq \left\{ Q \in \mathbb{S}^{2n \times 2n} \mid Q^{22} \in \mathbb{S}_{> M}^{n \times n} \right\}. \quad (81)$$

In particular, (81) follows by inspection of (73), while (80) may be verified by demonstrating that $\Pi \circ \Pi^{-1}$ is the identity. \blacksquare

A.5 Proof of Theorem 4

Throughout, it is assumed that Assumptions 1 and 2 hold, with $M \in \mathbb{S}^{n \times n}$ specified by the former, as per the theorem statement. Note in particular that $t^*(M) = +\infty$, so that $(\Sigma_t^{11} + \Sigma_t^{12}M)^{-1}$ exists for all $t \in \mathbb{R}_{>0}$, where Σ_t is the symplectic fundamental solution identified in (5). Consequently, $Q_t \in \mathbb{S}^{2n \times 2n}$ is well-defined as the unique solution of DRE (22), (23), for all $t \in \mathbb{R}_{>0}$ by Assumption 1, see Theorem 1 and its proof. Note that $P_0 \in \mathbb{S}_{> M}^{n \times n} = \text{dom}(\Upsilon)$ by hypothesis and (13).

The proof proceeds by demonstrating a sequence of implications concerning the following claims, posed with respect to arbitrary fixed $t \in \mathbb{R}_{>0}$ and $P_0 \in \mathbb{S}_{> M}^{n \times n}$:

- 1) $t \in (0, t^*(P_0))$;
- 2) $\Upsilon(P_0) + Q_s^{22} \in \mathbb{S}_{< 0}^{n \times n}$ for all $s \in (0, t]$;
- 3) $\Upsilon(P_0) + Q_t^{22} \in \mathbb{S}_{< 0}^{n \times n}$;
- 4) $P_0 + A_t^{22} \in \mathbb{S}_{< 0}^{n \times n}$; and
- 5) (44) and (45) hold.

In particular, it is shown that $1) \Leftrightarrow 2) \Leftrightarrow 3) \Leftrightarrow 4) \Rightarrow 5)$.

$2) \Rightarrow 1)$: Suppose that $\Upsilon(P_0) + Q_s^{22} \in \mathbb{S}_{< 0}^{n \times n}$ for all $s \in (0, t]$. Applying (13) and Theorem 1,

$$M^{-1}(\Upsilon(P_0) + Q_s^{22})M^{-1} = (M - P_0)^{-1} - (\Sigma_s^{11} + \Sigma_s^{12}M)^{-1}\Sigma_s^{12} \quad (82)$$

where it may be noted that the inverses on the right-hand side are guaranteed to exist. By hypothesis, the left-hand side is invertible, so that a matrix $K_s \in \mathbb{R}^{n \times n}$ is well-defined for an arbitrary $s \in (0, t]$ by

$$K_s \doteq (\Sigma_s^{11} + \Sigma_s^{12}M)^{-1} + (\Sigma_s^{11} + \Sigma_s^{12}M)^{-1}\Sigma_s^{12} \\ \times [(M - P_0)^{-1} - (\Sigma_s^{11} + \Sigma_s^{12}M)^{-1}\Sigma_s^{12}]^{-1}(\Sigma_s^{11} + \Sigma_s^{12}M)^{-1}.$$

However, the Woodbury Lemma implies that

$$K_s = [(\Sigma_s^{11} + \Sigma_s^{12}M) - \Sigma_s^{12}(M - P_0)]^{-1} = (\Sigma_s^{11} + \Sigma_s^{12}P_0)^{-1}.$$

That is, $\Sigma_s^{11} + \Sigma_s^{12}P_0 \in \mathbb{S}^{n \times n}$ is invertible. Recalling (6), and that $s \in (0, t]$ is arbitrary, immediately implies that 1) holds.

$1) \Rightarrow 2)$: Fix an arbitrary $t \in (0, t^*(P_0))$. Analogously to the proof of Theorem 1, let $\tilde{Q}_s \in \mathbb{S}^{2n \times 2n}$ denote the unique solution of DRE (22) subject to the initialization

$$\tilde{Q}_0 = \mu(P_0) \quad (83)$$

defined, via (10), for all $s \in [0, t^*(\tilde{Q}_0))$, where $t^*(\tilde{Q}_0) \in \mathbb{R}_{>0}$ is the corresponding maximal horizon of existence (6). Analogously to the argument yielding (28), observe that $t^*(\tilde{Q}_0) =$

$t^*(P_0)$, so that $t \in (0, t^*(\tilde{Q}_0))$. An application of the symplectic fundamental solution (4), (5), (61), yields

$$\tilde{Q}_s = \tilde{Y}_s \tilde{X}_s^{-1} \quad (84)$$

for all $s \in [0, t]$, in which

$$\begin{aligned} \begin{bmatrix} \tilde{X}_s \\ \tilde{Y}_s \end{bmatrix} &\doteq \hat{\Sigma}_s \begin{bmatrix} I \\ \mu(P_0) \end{bmatrix} \\ &= \begin{bmatrix} \Sigma_s^{11} & 0 & \Sigma_s^{12} & 0 \\ 0 & I & 0 & 0 \\ \Sigma_s^{21} & 0 & \Sigma_s^{22} & 0 \\ 0 & 0 & 0 & I \end{bmatrix} \begin{bmatrix} I & 0 \\ 0 & I \\ +P_0 & -M \\ -M & +M \end{bmatrix} = \begin{bmatrix} \Sigma_s^{11} + \Sigma_s^{12} P_0 & -\Sigma_s^{12} M \\ 0 & I \\ \Sigma_t^{21} + \Sigma_s^{22} P_0 & -\Sigma_s^{22} M \\ -M & +M \end{bmatrix} \in \mathbb{R}^{4n \times 2n}. \end{aligned}$$

for all $s \in [0, t]$. In particular,

$$\tilde{X}_s^{-1} = \begin{bmatrix} (\Sigma_s^{11} + \Sigma_s^{12} P_0)^{-1} & (\Sigma_s^{11} + \Sigma_s^{12} P_0)^{-1} \Sigma_s^{12} M \\ 0 & I \end{bmatrix},$$

in which $(\Sigma_s^{11} + \Sigma_s^{12} P_0)^{-1}$ is well-defined for all $s \in [0, t]$, as $t \in (0, t^*(P_0))$, see (6). Consequently, recalling (84),

$$\tilde{Q}_s^{22} = M - M(\Sigma_s^{11} + \Sigma_s^{12} P_0)^{-1} \Sigma_s^{12} M \quad (85)$$

is well-defined for all $s \in [0, t]$. Recalling (23), (83), as $\tilde{Q}_0 = \mu(P_0) \geq \mu(M) = Q_0$, Proposition 1 applied to (22) implies that $\tilde{Q}_s - Q_s \in \mathbb{S}_{\geq 0}^{2n \times 2n}$, so that in particular

$$\tilde{Q}_s^{22} - Q_s^{22} \in \mathbb{S}_{\geq 0}^{n \times n} \quad (86)$$

for all $s \in [0, t]$. Fix any $s \in (0, t]$. Rearranging (85) and applying (86) and Proposition 2,

$$\begin{aligned} (\Sigma_s^{11} + \Sigma_s^{12} P_0)^{-1} \Sigma_s^{12} &= M^{-1} (M - \tilde{Q}_s^{22}) M^{-1} \\ &\leq M^{-1} (M - Q_s^{22}) M^{-1} \in \mathbb{S}_{< 0}^{n \times n}. \end{aligned} \quad (87)$$

Theorem 1 and (64) implies via the notation of (3) that

$$Q_s^{22} = [\Xi(\Sigma_s)]^{22} = M - M(\Sigma_s^{11} + \Sigma_s^{12} M)^{-1} \Sigma_s^{12} M. \quad (88)$$

Recall that $\Sigma_s \in \text{dom}(\Xi)$ (ie. the inverse involved is guaranteed to exist) by Assumption 1, as $s \in (0, t^*(M)) \equiv \mathbb{R}_{> 0}$. Furthermore, as $s \in (0, t^*(P_0))$, definition (6) implies that $\Sigma_s^{11} + \Sigma_s^{12} P_0$ is invertible. Hence, a matrix $L_s \in \mathbb{S}^{n \times n}$ is well-defined by

$$\begin{aligned} L_s &\doteq (M - P_0) + (M - P_0)(\Sigma_s^{11} + \Sigma_s^{12} P_0)^{-1} \Sigma_s^{12} (M - P_0) \\ &= (M - P_0) + (M - P_0) [(\Sigma_s^{11} + \Sigma_s^{12} M) - \Sigma_s^{12} (M - P_0)]^{-1} \Sigma_s^{12} (M - P_0). \end{aligned}$$

where the second equality follows by adding and subtracting $\Sigma_s^{12} M$ within the inverse. Applying (87), and the fact that $P_0 \in \mathbb{S}_{> M}^{n \times n}$, note that $L_s \in \mathbb{S}_{< 0}^{n \times n}$ by definition. The Woodbury Lemma subsequently implies that

$$L_s = [(M - P_0)^{-1} - (\Sigma_s^{11} + \Sigma_s^{12} M)^{-1} \Sigma_s^{12}]^{-1} = M(\Upsilon(P_0) + Q_s^{22})^{-1} M$$

where the second equality follows as per (82). Consequently, as $M \in \mathbb{S}^{n \times n}$ is invertible and $L_s \in \mathbb{S}_{< 0}^{n \times n}$,

$$\Upsilon(P_0) + Q_s^{22} = M L_s^{-1} M \in \mathbb{S}_{< 0}^{n \times n}.$$

As $s \in (0, t]$ is arbitrary, claim 2) immediately follows.

2) \Rightarrow 3): By hypothesis, $\Upsilon(P_0) + Q_s^{22} \in \mathbb{S}_{<0}^{n \times n}$ for all $s \in (0, t]$. Selecting $s = t$ yields claim 3) as required.

3) \Rightarrow 2): By hypothesis, $\Upsilon(P_0) + Q_t^{22} \in \mathbb{S}_{<0}^{n \times n}$. Furthermore, $\Upsilon(P_0) \in \mathbb{S}_{<-M}^{n \times n}$ by (13), (14). Hence, $Q_t^{22} \in \mathbb{S}^{n \times n}$, so that $(Q_\sigma^{12})' BB' Q_\sigma^{12}$ must be integrable with respect to $\sigma \in [0, t]$ by definition (27). In particular,

$$Q_t^{22} - M = \int_0^t (Q_\sigma^{12})' BB' Q_\sigma^{12} d\sigma \geq \int_0^s (Q_\sigma^{12})' BB' Q_\sigma^{12} d\sigma = Q_s^{22} - M$$

for any fixed $s \in (0, t]$. Hence, $Q_s^{22} - Q_t^{22} \in \mathbb{S}_{<0}^{n \times n}$, so that

$$\Upsilon(P_0) + Q_s^{22} = (\Upsilon(P_0) + Q_t^{22}) + (Q_s^{22} - Q_t^{22}) \in \mathbb{S}_{<0}^{n \times n}.$$

Recalling that $s \in (0, t]$ is arbitrary yields claim 2) as required.

3) \Rightarrow 4): Recalling (13) and Theorem 3, see (42), (80),

$$\begin{aligned} \Upsilon(P_0) + Q_t^{22} &= (-M - M(P_0 - M)^{-1}M) + (M - M(M + \Lambda_t^{22})^{-1}M) \\ &= M[(M - P_0)^{-1} - (M + \Lambda_t^{22})^{-1}]M. \end{aligned} \quad (89)$$

Recalling that $\Upsilon(P_0) + Q_t^{22} \in \mathbb{S}_{<0}^{n \times n}$ by hypothesis,

$$\begin{aligned} \Upsilon(P_0) + Q_t^{22} \in \mathbb{S}_{<0}^{n \times n} &\Leftrightarrow (M - P_0)^{-1} - (M + \Lambda_t^{22})^{-1} \in \mathbb{S}_{<0}^{n \times n} \\ &\Leftrightarrow (M + \Lambda_t^{22}) - (M - P_0) \in \mathbb{S}_{<0}^{n \times n} \\ &\Leftrightarrow P_0 + \Lambda_t^{22} \in \mathbb{S}_{<0}^{n \times n}. \end{aligned} \quad (90)$$

That is, claim 4) holds.

4) \Rightarrow 3): Note that (89) holds as per the 3) \Rightarrow 4) case above. By hypothesis, $P_0 + \Lambda_t^{22} \in \mathbb{S}_{<0}^{n \times n}$. Hence, the string of equivalences (90) implies that 3) holds.

4) \Rightarrow 5): Recalling (15), (35) and (42), the value function W_t of (15), (19) satisfies

$$\begin{aligned} W_t(x) &= \frac{1}{2}x' P_t x = \int_{\mathbb{R}^n}^{\oplus} G_t(x, y) \otimes \Psi(y) dy \\ &= \int_{\mathbb{R}^n}^{\oplus} \frac{1}{2} \begin{bmatrix} x \\ y \end{bmatrix}' \Lambda_t \begin{bmatrix} x \\ y \end{bmatrix} \otimes \frac{1}{2} y' P_0 y dy = \frac{1}{2} \int_{\mathbb{R}^n}^{\oplus} \begin{bmatrix} x \\ y \end{bmatrix}' \begin{bmatrix} \Lambda_t^{11} & \Lambda_t^{12} \\ (\Lambda_t^{12})' & P_0 + \Lambda_t^{22} \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} dy \end{aligned}$$

for all $x \in \mathbb{R}^n$. By hypothesis, $P_0 + \Lambda_t^{22} \in \mathbb{S}_{<0}^{n \times n}$, so that $(P_0 + \Lambda_t^{22})^{-1}$ exists. Hence, the above max-plus integration explicitly evaluates as

$$\frac{1}{2}x' P_t x = \frac{1}{2}x' [\Lambda_t^{11} - \Lambda_t^{12}(P_0 + \Lambda_t^{22})^{-1}(\Lambda_t^{12})'] x.$$

As $x \in \mathbb{R}^n$ is arbitrary, (44) follows immediately. In addition, as 4) \Leftrightarrow 1), it immediately follows that

$$\sup \left\{ t \in \mathbb{R}_{>0} \mid P_0 + \Lambda_t^{22} \in \mathbb{S}_{<0}^{n \times n} \right\} = \sup \left\{ t \in \mathbb{R}_{>0} \mid t \in (0, t^*(P_0)) \right\} = t^*(P_0).$$

That is, (45) holds. ■

References

1. B. Anderson and J. Moore, *Linear optimal control*. Englewood Cliffs, New Jersey, USA: Prentice-Hall, 1971.
2. J. Doyle, K. Glover, P. Khargonekar, and B. Francis, "State space solutions to standard \mathcal{H}_2 and \mathcal{H}_∞ -control problems," *IEEE Transactions on Automatic Control*, vol. 34, no. 8, pp. 831–847, 1989.

3. I. Petersen, B. Anderson, and E. Jonckheere, "A first-principles solution to the non-singular \mathcal{H}_∞ control problem," *International Journal of Robust & Nonlinear Control*, vol. 1, pp. 171–185, 1991.
4. M. Green and D. Limebeer, *Linear robust control*, ser. Information and systems sciences. Prentice-Hall, 1995.
5. E. Davison and M. Maki, "The numerical solution of the matrix Riccati differential equation," *IEEE Transactions on Automatic Control*, vol. 18, pp. 71–73, 1973.
6. J. Lawson and Y. Lim, "The symplectic semigroup and Riccati differential equations," *J. Dynamical and Control Systems*, vol. 12, no. 1, pp. 49–77, 2006.
7. W. McEneaney, "A new fundamental solution for differential Riccati equations arising in control," *Automatica*, vol. 44, pp. 920–936, 2008.
8. P. Dower and W. McEneaney, "A max-plus dual space fundamental solution for a class of operator differential Riccati equations," *SIAM J. Control & Optimization*, vol. 53, no. 2, pp. 969–1002, 2015.
9. H. Zhang and P. Dower, "Max-plus fundamental solution semigroups for a class of difference Riccati equations," *Automatica*, vol. 52, pp. 103–110, 2015.
10. —, "A max-plus based fundamental solution for a class of discrete time linear regulator problems," *Linear Algebra and its Applications*, vol. 471, pp. 693–729, 2015.
11. W. Fleming and W. McEneaney, "A max-plus-based algorithm for a Hamilton-Jacobi-Bellman equation of nonlinear filtering," *SIAM Journal on Control and Optimization*, vol. 38, no. 3, pp. 683–710, 2000.
12. V. Kolokoltsov and V. Maslov, *Idempotent analysis and applications*. Kluwer Publishing House, 1997.
13. G. Litvinov, V. Maslov, and G. Shpiz, "Idempotent functional analysis: An algebraic approach," *Mathematical Notes*, vol. 69, no. 5, pp. 696–729, 2001.
14. M. Akian, S. Gaubert, and V. N. Kolokoltsov, "Set coverings and invertibility of functional galois connections," in *Idempotent Mathematics and Mathematical Physics*, ser. Contemporary Mathematics, G. L. Litvinov and V. P. Maslov, Eds. American Mathematical Society, 2005, pp. 19–51.
15. G. Cohen, S. Gaubert, and J.-P. Quadrat, "Duality and separation theorems in idempotent semimodules," *Linear Algebra & Applications*, vol. 379, pp. 395–422, 2004.
16. W. McEneaney, *Max-plus methods for nonlinear control and estimation*, ser. Systems & Control: Foundations & Applications. Birkhauser, 2006.
17. T. Sasagawa, "On the finite escape phenomena for matrix Riccati equations," *IEEE Transactions on Automatic Control*, vol. 27, pp. 977–979, 1982.
18. S. Kilicaslan and S. Banks, "Existence of solutions of Riccati differential equations," *Journal of Dynamic Systems, Measurement, and Control*, vol. 134, p. 031001, 2012.
19. J.-J. Moreau, "Inf-convolution, sous-additivité, convexité des fonctions numériques," *J. Math. Pures Appl.*, vol. 9, no. 49, pp. 109–154, 1970.
20. R. Rockafellar, "Conjugate duality and optimization," *SIAM Regional Conf. Series in Applied Math.*, vol. 16, 1974.
21. R. Rockafellar and R. Wets, *Variational Analysis*. Springer-Verlag, 1997.
22. P. Dower, W. McEneaney, and H. Zhang, "Max-plus fundamental solution semigroups for optimal control problems," in *proc. SIAM Conference on Control Theory and Its Applications (Paris, France), 2015*, 2015, pp. 368–375.
23. A. Bensoussan, G. D. Prato, M. Delfour, and S. Mitter, *Representation and control of infinite dimensional systems*, 2nd ed. Birkhäuser, 2007.