

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Cazalet, C;Gomez-Valero, L;Rusniok, C;Lomma, M;Dervins-Ravault, D;Newton, HJ;Sansom, FM;Jarraud, S;Zidane, N;Ma, L;Bouchier, C;Etienne, J;Hartland, EL;Buchrieser, C

Title:

Analysis of the Legionella longbeachae genome and transcriptome uncovers unique strategies to cause Legionnaires' disease

Date:

2010-01-01

Citation:

Cazalet, C., Gomez-Valero, L., Rusniok, C., Lomma, M., Dervins-Ravault, D., Newton, H. J., Sansom, F. M., Jarraud, S., Zidane, N., Ma, L., Bouchier, C., Etienne, J., Hartland, E. L. & Buchrieser, C. (2010). Analysis of the Legionella longbeachae genome and transcriptome uncovers unique strategies to cause Legionnaires' disease. Plos Genetics, 6 (2), <https://doi.org/10.1371/journal.pgen.1000851>.

Persistent Link:

<https://hdl.handle.net/11343/242937>

License:

CC BY

Analysis of the *Legionella longbeachae* Genome and Transcriptome Uncovers Unique Strategies to Cause Legionnaires' Disease

Christel Cazalet^{1,3}, Laura Gomez-Valero^{1,3}, Christophe Rusniok¹, Mariella Lomma¹, Delphine Dervins-Ravault¹, Hayley J. Newton², Fiona M. Sansom², Sophie Jarraud³, Nora Zidane⁴, Laurence Ma⁴, Christiane Bouchier⁴, Jérôme Etienne³, Elizabeth L. Hartland², Carmen Buchrieser^{1*}

1 Institut Pasteur, Biologie des Bactéries Intracellulaires, CNRS URA 2171, Paris, France, **2** Department of Microbiology and Immunology, University of Melbourne, Victoria, Australia, **3** Centre National de Référence des Legionella, Université de Lyon, INSERM U851, Faculté de Médecine, IFR 128, Lyon, France, **4** Institut Pasteur, Plate-forme Génomique, Pasteur Génopole Ile de France, Paris, France

Abstract

Legionella pneumophila and *L. longbeachae* are two species of a large genus of bacteria that are ubiquitous in nature. *L. pneumophila* is mainly found in natural and artificial water circuits while *L. longbeachae* is mainly present in soil. Under the appropriate conditions both species are human pathogens, capable of causing a severe form of pneumonia termed Legionnaires' disease. Here we report the sequencing and analysis of four *L. longbeachae* genomes, one complete genome sequence of *L. longbeachae* strain NSW150 serogroup (Sg) 1, and three draft genome sequences another belonging to Sg1 and two to Sg2. The genome organization and gene content of the four *L. longbeachae* genomes are highly conserved, indicating strong pressure for niche adaptation. Analysis and comparison of *L. longbeachae* strain NSW150 with *L. pneumophila* revealed common but also unexpected features specific to this pathogen. The interaction with host cells shows distinct features from *L. pneumophila*, as *L. longbeachae* possesses a unique repertoire of putative Dot/Icm type IV secretion system substrates, eukaryotic-like and eukaryotic domain proteins, and encodes additional secretion systems. However, analysis of the ability of a *dotA* mutant of *L. longbeachae* NSW150 to replicate in the *Acanthamoeba castellanii* and in a mouse lung infection model showed that the Dot/Icm type IV secretion system is also essential for the virulence of *L. longbeachae*. In contrast to *L. pneumophila*, *L. longbeachae* does not encode flagella, thereby providing a possible explanation for differences in mouse susceptibility to infection between the two pathogens. Furthermore, transcriptome analysis revealed that *L. longbeachae* has a less pronounced biphasic life cycle as compared to *L. pneumophila*, and genome analysis and electron microscopy suggested that *L. longbeachae* is encapsulated. These species-specific differences may account for the different environmental niches and disease epidemiology of these two *Legionella* species.

Citation: Cazalet C, Gomez-Valero L, Rusniok C, Lomma M, Dervins-Ravault D, et al. (2010) Analysis of the *Legionella longbeachae* Genome and Transcriptome Uncovers Unique Strategies to Cause Legionnaires' Disease. PLoS Genet 6(2): e1000851. doi:10.1371/journal.pgen.1000851

Editor: Ivan Matic, Université Paris Descartes, INSERM U571, France

Received: November 18, 2009; **Accepted:** January 20, 2010; **Published:** February 19, 2010

Copyright: © 2010 Cazalet et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work received financial support from the Institut Pasteur, the Centre National de la Recherche (CNRS), the Network of Excellence "Europathogenomics" LSHB-CT-2005-512061, and the Australian National Health and Medical Research Council (NHMRC). ML is holder of a Marie Curie fellowship financed by the European Commission (INTRAPATH project MEST-CT-2005-020715) coordinated by Institut Pasteur and LG-V is holder of a Roux postdoctoral research Fellowship financed by the Institut Pasteur. ELH holds an Australian Research Council Future Fellowship. HJN and FMS hold NHMRC Biomedical Training fellowships. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: carmen.buchrieser@pasteur.fr

These authors contributed equally to this work.

Introduction

Legionella longbeachae is one species of the family *Legionellaceae* that causes legionellosis, an atypical pneumonia that can be fatal if not promptly treated. While *Legionella pneumophila* is the leading cause of legionellosis in the USA and Europe, and is associated with around 91% of the cases worldwide, *L. longbeachae* is responsible for approximately 30% of legionellosis cases in Australia and New Zealand and nearly 50% in South Australia [1] and Thailand [2]. Two serogroups (Sg) are distinguished within *L. longbeachae* but most of the human cases of legionellosis are due to Sg1 strains [3,4]. Interestingly, unlike *L. pneumophila*, which inhabits aquatic environments, *L. longbeachae* is found predominantly in potting soil and is transmitted by inhalation of dust from contaminated soils [4,5].

Little is known about the biology and the genetic basis of virulence of *L. longbeachae* but a few studies suggest considerable differences with respect to *L. pneumophila*. In contrast, the intracellular life cycle of *L. pneumophila* is well characterized (for recent reviews see [6–8]). *L. pneumophila* replicates within alveolar macrophages inside a unique phagosome that excludes both early and late endosomal markers, resists fusion with lysosomes and recruits endoplasmic reticulum and mitochondria. Within this protected vacuole *L. pneumophila* replicates and down-regulates the expression of virulence factors. It has been proposed that nutrient limitation then leads to the transition to transmissive phase bacteria that express many virulence-associated traits allowing the release and infection of new host cells [9]. This biphasic life cycle is observed both *in vitro* and *in vivo* as exponential phase bacteria do

Author Summary

Legionella longbeachae, found in potting soil, and *L. pneumophila*, present in aquatic environments, are opportunistic human pathogens that cause Legionnaires' disease, a severe and often fatal pneumonia. The analysis and comparison of the genome sequences of four *L. longbeachae* genomes together with the study of its gene expression program and virulence pattern in different infection models provides important new insight on the organism's lifestyle and virulence strategies. *L. longbeachae* harbors a unique repertoire of secreted substrates, many of which encode eukaryotic like domains that may help the pathogen to subvert host functions and cause disease. Curiously, *L. longbeachae* may also be able to interact with plants. Several proteins present mainly in plants and phytopathogenic bacteria and several enzymes that might confer the ability to degrade plant material were identified in its genome. Interestingly, *L. longbeachae* encodes a chemotaxis system but no flagella, in contrast *L. pneumophila* encodes flagella but no chemotaxis system. It will be an interesting aspect of future research to understand these peculiarities. Finally, the genome sequence and analysis reported here will aid in understanding how *L. longbeachae* causes disease and will open new possibilities to develop tools for rapid identification and risk prediction of *L. longbeachae* infection.

not express virulence factors and the bacteria fail to evade the destructive lysosomes and are delivered to the endocytic network and destroyed [9,10]. The ability of *L. pneumophila* to replicate intracellularly is triggered at the post-exponential phase together with other virulence traits. Less is known about the intracellular life cycle of *L. longbeachae* and its virulence factors. Unlike *L. pneumophila* the ability of *L. longbeachae* to replicate intracellularly is independent of the bacterial growth phase [11]. Phagosome biogenesis is also different. Like *L. pneumophila*, the *L. longbeachae* phagosome is surrounded by endoplasmic reticulum and evades lysosome fusion but in contrast to *L. pneumophila* containing phagosomes the *L. longbeachae* vacuole acquires early and late endosomal markers [12].

Efficient formation of the *L. pneumophila* replication vacuole requires the Dot/Icm type IV secretion system (T4SS) [13–16]

and probably more than 100 translocated effector proteins that modulate different host cell processes, in particular vesicle trafficking [17–19]. While *L. longbeachae* possesses all genes necessary to code a Dot/Icm T4SS [20], it is not known whether it is also essential for virulence and whether *L. pneumophila* and *L. longbeachae* share common effectors.

Another interesting difference between these two species is their ability to colonize the lungs of mice. While only A/J mice are permissive for replication of *L. pneumophila*, A/J, C57BL/6 and BALB/c mice are all permissive for replication of *L. longbeachae* [12,21]. Resistance of C57BL/6 and BALB/c mice to *L. pneumophila* has been attributed to polymorphisms in Nod-like receptor apoptosis inhibitory protein 5 (*naip5*) allele [22–24]. The current model states that *L. pneumophila* replication is restricted due to flagellin dependent caspase-1 activation through Naip5-Ipaf and early macrophage cell death by pyroptosis. Why *L. longbeachae*, in contrast to *L. pneumophila*, is able to replicate in macrophages of all three different mouse strains is still not understood.

In this study we report the complete genome sequencing and analysis of a clinical *L. longbeachae* Sg1 strain isolated in Australia and compare this genome to three *L. longbeachae* draft genome sequences (one Sg1 and two Sg2 strains) and the published genome sequences of four *L. pneumophila* strains [25–27]. In addition, we performed transcriptome analysis and virulence studies of a T4SS mutant of *L. longbeachae*. This has allowed us to propose answers for the questions raised above and brings exciting new insight into the varying adaptation to different ecological niches and different intracellular life cycles of *Legionella* species.

Results/Discussion

The *L. longbeachae* genomes are highly conserved and are 500 kb larger than those of *L. pneumophila*

The *L. longbeachae* NSW150 genome consists of a 4,077,332-bp chromosome and a 71,826-bp plasmid with an average GC content of 37.11% and 38.19%, respectively (Table 1). A total of 3512 protein-encoding genes are predicted, 2046 (58.3%) of which have been assigned a putative function (Table S1, Figure S1). The *L. longbeachae* chromosome is about 500 kb larger than that of *L. pneumophila* and has a significantly different organization as seen in the synteny plot in Figure 1 and Figure S2. Moreover only 2290 (65.2%) *L. longbeachae* genes are orthologous to *L. pneumophila* genes,

Table 1. General features of the completely sequenced *L. pneumophila* and *L. longbeachae* genomes.

	<i>L. longbeachae</i>	<i>L. pneumophila</i>			
	NSW 150	Paris	Lens	Philadelphia	Corby
Chromosome size (kb) ^a	4077 (71)	3504 (131.8)	3345 (59.8)	3397	3576
G + C content (%)	37.1 (38.2)	38.3 (37.4)	38.4 (38)	38.27	38
G + C content of CDS (%)	37.4	39.1	39.4	38.6	38.6
No. of genes ^a	3660 (75)	3136 (142)	3001 (60)	3002	3259
No. of protein coding genes ^a	3512 (67)	2878 (140)	2878 (60)	2942	3206
Percentage of CDS (%)	84.5	87.9	88	90.2	86.8
Average length of CDS (pb)	1015.2	994.6	935.9	960.7	959.4
No. of 16S/23S/5S	4/4/4	3/3/3	3/3/3	3/3/3	3/3/3
No. transfer RNA	46	44	43	43	43
Plasmids	1	1	1	0	0

^a Updated annotation; CDS = coding sequence; in parenthesis data from plasmids.
doi:10.1371/journal.pgen.1000851.t001




Figure 1. Whole-genome synteny map of *L. longbeachae* strain NSW150 and *L. pneumophila* strain Paris. The linearized chromosomes were aligned and visualized by Lineplot in MAGE. Syntenic relationships comprising at least 8 genes are indicated by green and red lines for genes found on the same strand or on opposite strands, respectively. IS elements (pink), ribosomal operons (blue) and tRNAs (green) are also indicated.
doi:10.1371/journal.pgen.1000851.g001

whereas 1222 (34.8%) are *L. longbeachae* specific with respect to *L. pneumophila* Paris, Lens, Philadelphia and Corby (defined by less than 30% amino acid identity over 80% of the length of the smallest protein, s Table S2). It was previously suggested that plasmid-encoded functions such as a two-component system, are important for *L. longbeachae* virulence [28]. Although no similarity was detected between the *L. longbeachae* plasmid here characterized and the 9kb partial plasmid sequence reported of strain *L. longbeachae* A5H5 [28], similar plasmids seem to circulate among different *Legionella* species, as 30 kb of the plasmid of strains Paris, Lens and NSW150, 18 kb of which encode transfer genes (*traI* – *traA*), encoded ORFs showing high amino acid sequence similarity (Figure S3).

With the aim of gaining further information on genome content and diversity of *L. longbeachae* we selected three additional strains, two isolated in the USA one in Australia for genome sequencing and analysis. *L. longbeachae* strain ATCC39642 (Sg1), strain 98072 (Sg2) and strain C-4E7 (Sg2) were deep sequenced using the Illumina technology and then compared to the genome of strain NSW150. We obtained a coverage of 93–96% for each genome with respect to the NSW150 genome (Table 2). The sequences were assembled into 93, 106 and 89 contigs larger than 0.5kbs that were further analyzed regarding gene content and single

nucleotide polymorphisms (SNP). High quality SNPs were detected by mapping the Illumina reads on the finished NSW150 genome sequence. This revealed a high conservation in genome size, content, organization and a low SNP number among the four *L. longbeachae* genomes (Table 2). Interestingly, in contrast to *L. pneumophila* where strains of the same Sg may have very different gene content [25,29], the two strains of *L. longbeachae* each belonging to Sg1 or Sg2, respectively, showed highly conserved genomes. Comparison of the two Sg1 genomes identified 1611 SNPs of which 1426 are located in only seven chromosomal regions mainly encoding putative mobile elements, whereas the remaining 185 SNPs were evenly distributed around the chromosome (Figure S4). In contrast, the SNP number between two strains of different Sg was higher, with about 16 000 SNPs present between Sg1 and Sg2 strains (Table 1, Figure S4). This represents an overall polymorphism of less than 0.4%, which is significantly lower than the polymorphism of about 2% between *L. pneumophila* Sg1 strains Paris and Philadelphia. The low SNP number and relatively homogeneous distribution of the SNPs around the chromosome (Figure S4) suggest recent expansion for the species *L. longbeachae*.

The *dot/icm* type IVB secretion system is highly conserved, and many other secretion systems are present

L. pneumophila has a rather exceptional number and wide variety of secretion systems for efficient and rapid delivery of effector molecules into the phagocytic host cell underlining the importance of protein secretion for this pathogen. This also holds true for *L. longbeachae*. We identified the genes coding the Lsp type II secretion machinery, however, 45% of the type II secretion system substrates described for *L. pneumophila* [30,31] are absent from *L. longbeachae*. Furthermore, the twin arginine translocation system (TAT) and three putative type I secretion systems (T1SS) are present. However, the Lss T1SS might not be functional in *L. longbeachae* as only LssXYZA are conserved (55 to 82% identity with strain Paris) and the two essential components LssB (ABC transporter-ATP binding) and LssD (HlyD family secretion protein) are missing. In contrast, the two additional putative T1SS, encoded by the genes *llo2283-llo2288* and *llo0441-llo0444* appeared to be functional. Furthermore, two HlyD-like proteins (Llo2901 and Llo0979) localized next to ABC transporters (Llo2900 and Llo0980-Llo0981) were present, but no contiguous

Table 2. General features of the *L. longbeachae* draft genomes obtained by new generation sequencing.

	<i>L. longbeachae</i>			
	NSW 150	ATCC39462	98072	C-4E7
Chromosome size (Kb)	4077 (71)	4096	4018 (133.8)	3979 (133.8)
No. of 16S/23S/5S	4/4/4	4/4/4	4/4/4	4/4/4
G + C content (%)	37.1 (38.2)	37.0	37.0 (37.8)	37 (37.8)
No. of contigs >0.5-300 kb	complete	64	65	63
N50 contig size*	complete	138 kb	129 kb	134 kb
Percentage of coverage**	100%	96.3	93.4	93.1
Number of SNP with NSW150	–	1611	16 853	16 820
Plasmids	1	0	1	1

*N50 contig size, calculated by ordering all contig sizes and then adding the lengths (starting from the longest contig) until the summed length exceeds 50% of the total length of all contigs (half of all bases reside in a contiguous sequence of the given size or more);

**for SNP detection

doi:10.1371/journal.pgen.1000851.t002

outer membrane protein was found. However, these proteins could also be part of T1SS and function together with a genetically unlinked outer membrane component, similar to what is seen for the Hly T1SS of *Escherichia coli* and may thus constitute two additional T1SS. Finally, *L. longbeachae* encodes four type IV secretion systems (T4SS). The Lvh T4ASS of *L. pneumophila* is absent from *L. longbeachae* but we identified three other type-IVA secretion systems. One T4ASS is present on the plasmid and the other two are embedded on putative mobile genomic islands (GI) in the chromosome. *llo1819-llo1929* (GI-1) of around 120 kb is bordered by Ser and Arg tRNAs and carries a gene coding for a phage integrase (*llo1819*). The second cluster (GI-2) of 106 kb spans from the integrase coding gene *llo2859* to *llo2960ab* and is also bordered by a Met tRNA. Most of the proteins encoded on GI-2 are of unknown function. However both islands code for several proteins, which may be dedicated to stress response. On GI-1, Llo1862 and Llo1863/*llo1863* are homologous to DNA polymerase IV subunit C and D respectively, involved in the SOS repair pathway. On GI-2 are the OsmC-like protein Llo2923, the putative universal stress proteins Llo2926, Llo2927, Llo2929 and the predicted transcriptional regulator Llo2913 with S24 peptidase domain. Indeed, the S24 peptidase family includes LexA, a transcriptional repressor of SOS response genes to DNA damage. Several transporters were also identified on GI-2: Llo2918 of the MFS superfamily, the Na/H exchange protein Llo2930 and the putative T1SS proteins Llo2900 and Llo2901 discussed above. It possesses in addition a putative restriction/modification system encoded by *llo2865*, *llo2866* and *llo2867*.

Central to the establishment of the intracellular replicative niche and to *L. pneumophila* virulence is the Dot/Icm type IV secretion system. This T4BSS is also present in *L. longbeachae* and the general organization of the genomic region encoding it is conserved with protein identities of 47 to 92% with respect to that of *L. pneumophila*. This is similar to what has been reported previously for other *Legionella* species [20]. In *L. longbeachae* the *icmR* gene is replaced by the *ligB* gene, however, the encoded proteins have been shown to perform similar functions [32,33]. Here we found that IcmE/DotG of *L. longbeachae* is 477 amino acids larger than that of *L. pneumophila*. DotG is part of the core transmembrane complex of the secretion system and it is composed of three domains: a transmembrane N-terminal domain, a central region composed of 42 repeats of 10 amino acid and a C-terminal region homologous to VirB10. The central region of DotG from *L. longbeachae* comprises approximately 90 repeats. It will be challenging to understand the possible impact of this modification on the function of the type-IV secretion system.

The *dot/icm* type IV secretion system of *L. longbeachae* is essential for virulence in *Acanthamoeba castellanii* and in pulmonary mouse infection

To test whether the Dot/Icm T4SS is essential for virulence of *L. longbeachae* we constructed a deletion mutant in the *L. longbeachae* NSW150 gene *llo0364*, homologous to *dotA* of *L. pneumophila* and tested its ability to replicate compared to the wild type strain in *A. castellanii* and the lungs of A/J mice. We found that *L. longbeachae* NSW150 infects *A. castellanii* in a comparable manner to *L. pneumophila* and that the *dotA* mutant was strongly attenuated for intracellular growth in *A. castellanii*, similar to what is seen for a *L. pneumophila dotA* mutant (Figure 2A). Recently Gobin and colleagues established an experimental model of intratracheal *L. longbeachae* infection in A/J mice [21]. Here we compared the ability of the *L. longbeachae dotA* mutant to compete with wild type *L. longbeachae* in the lungs of A/J mice. In mixed infections, we observed that the *dotA* mutant was outcompeted by the wild type

strain 24 h and 72 h after infection (Figure 2B). The competitive index of the *dotA* mutant was calculated by dividing the ratio of mutant to wild type bacteria after infection with the ratio of mutant to wild type bacteria in the inoculum. A competitive index of less than 0.5 is considered a significant attenuation [34]. The competitive index was less than 0.5 at both time-points indicating rapid loss of the *dotA* mutant following infection. In single infections, the *L. longbeachae dotA* mutant was also dramatically attenuated for replication (Figure 2C). Thus, the Dot/Icm secretion system was essential for the virulence of *L. longbeachae*.

L. longbeachae and *L. pneumophila* encode different sets of secreted Dot/Icm substrates and virulence genes

Despite the high degree of conservation of the Dot/Icm T4SS components between *L. pneumophila* and *L. longbeachae* the Dot/Icm substrates were not highly conserved. Indeed 66% of reported *L. pneumophila* Dot/Icm substrates were absent from *L. longbeachae* (Table 3 and Table S3). Instead, we predicted 51 new putative Dot/Icm substrates specific for *L. longbeachae* that encode eukaryotic-like domains and all but one contained the secretion signal described by Nagai and colleagues [35] and many also the additional criteria defined by Kubori and colleagues [36] (Table 4). Interestingly, the distribution of both, the conserved and the newly identified substrates of *L. longbeachae* among the four sequenced strains was highly conserved (Table 3 and Table 4). Both *L. pneumophila* and *L. longbeachae* replicate within a vacuole that recruits endoplasmic reticulum. Several effector proteins have been shown to contribute to the ability of *L. pneumophila* to manipulate host cell trafficking events resulting in this association. The effector proteins SidJ, RalF, VipA, VipF, SidC, YlfA and LepB which contribute to trafficking or recruitment and retention of vesicles to *L. pneumophila* vacuoles were conserved in *L. longbeachae*, but VipD, SidM/DrrA and LidA which interfere also with these events are absent from the *L. longbeachae* genome; however VipD and SidM/DrrA are also not present in all the *L. pneumophila* genomes sequenced.

Although *L. pneumophila* also communicates with early and late endosomal vesicle trafficking pathways [37–39], a major difference in the phagosome maturation of the two species is that the *L. longbeachae* phagosome acquires early and late endocytic markers. Several proteins identified specifically in the genome of *L. longbeachae* may contribute to these differences. First, *L. longbeachae* encodes a family of Ras-related small GTPases (Llo3288, Llo2329, Llo1716 and Llo2249) (Figure S5), which may also be involved in vesicular trafficking and account for the specificities of the *L. longbeachae* life cycle. Remarkably, Llo3288, Llo2329 and Llo1716 are the first small GTPases of the Rab subfamily described in a prokaryote. *L. pneumophila* is also known to exploit monophosphorylated host phosphoinositides (PI) to anchor the effector proteins SidC, SidM/DrrA, LpnE and LidA to the membrane of the replication vacuole [34,40–44]. *L. longbeachae* may employ an additional strategy to interfere with the host PI as Llo0793 is homologous to a mammalian PI metabolizing enzyme phosphatidylinositol-4-phosphate 5-kinase and it is tempting to speculate that this protein allows direct modulation of the host cell PI levels.

As another strategy to alter host trafficking pathways, *L. pneumophila* is able to target microtubule-dependent vesicular transport. AnkX/AnkN, for example, prevents microtubule-dependent vesicular transport interfering with the fusion of the *L. pneumophila*-containing vacuole with late endosomes [45]. AnkX/AnkN is absent from *L. longbeachae*, however *L. longbeachae* did encode a putative tubulin-tyrosine ligase (TTL) Llo2200, which adds to the 19 bacterial TTL identified to date. TTL catalyzes the ATP-dependent post-translational addition of a




Figure 2. Intracellular growth of the wild-type and the *dotA* mutant strain in mouse and amoeba infection. (A) Intracellular replication of *L. longbeachae* in *Acanthamoeba castellanii*. Blue, wild-type *L. longbeachae* strain NSW150; Red, *dotA*::Km mutant. Results are expressed as log₁₀ CFU. Each time point (in hours, x-axis) represents the mean $\pm$ SD of two independent experiments. Infections were performed at 37°C. (B) CI values from mixed infections of A/J mice. Mice were inoculated with approximately 10⁶ CFU of each strain under investigation and were sacrificed at 24 h or 72 h after infection to examine the bacterial content of their lungs. Competition experiment between *L. longbeachae* and the *dotA*::Km mutant representative of 2 independent experiments. (C) Single infections of A/J mice with *L. longbeachae* wt and the *dotA*::Km mutant strain. Results are expressed as log₁₀ CFU. Note: to maintain numbers in the lung *L. longbeachae* must be replicating. Non-replicating bacteria are cleared in this infection model over 72 h (eg. *dotA* mutant) [21].
doi:10.1371/journal.pgen.1000851.g002

tyrosine to the carboxy terminal end of detyrosinated alpha-tubulin. Although the exact physiological function of alpha-tubulin has so far not been established, it has been linked to altered microtubule structure and function [46]. Besides AnkX/AnkN, a large family of ankyrin repeat constitutes *L. pneumophila* Dot/Icm substrates. Interestingly, 23 of the 29 ankyrin proteins identified in the *L. pneumophila* strains are absent from the *L. longbeachae* genome, however *L. longbeachae* encodes 23 specific ankyrin repeat proteins (Table 4).

L. pneumophila is also able to interfere with the host ubiquitination pathway. The U-box protein LubX, which possesses *in vitro* ubiquitin ligase activity specific for the eukaryotic Cdc2-like kinase Clk1 [36], is absent from *L. longbeachae*. However, *llo0448* encodes a predicted U-box protein. None of the three *L. pneumophila* F-box proteins, which may also exploit this pathway, are conserved in *L. longbeachae*, but we identified two new putative F-box proteins Llo1427 and Llo2109 (Table 4). Thus, although the specific proteins may not be conserved, the eukaryotic-like protein-protein interaction domains found in *L. pneumophila* are also present in *L. longbeachae*.

L. longbeachae also encodes several proteins with eukaryotic domains that are not present in *L. pneumophila*. One is the above-mentioned protein Llo2200 encoding a TTL domain. A second is Llo2327, the first bacterial protein that encodes an Src Homology 2 (SH2) domain. SH2 domains, in eukaryotes, have regulatory functions in various intracellular signaling cascades. Furthermore,

L. longbeachae encodes two proteins (Llo1404 and Llo2643) with pentatricopeptide repeat (PPR) domains. This family seems to be greatly expanded in plants, where they appear to play essential roles in organellar RNA metabolism [47–49] where they appear to play essential roles in RNA/DNA metabolism, where. Only 12 bacterial PPR domain proteins have been identified to date, all encoded by two species, the plant pathogens *Ralstonia solanacearum* and the facultative photosynthetic bacterium *Rhodobacter sphaeroides*.

L. longbeachae encodes putative toxins

Recently, a family of cytotoxic glucosyltransferases produced by *L. pneumophila* (Lgt) and related to the group of clostridial glucosylating cytotoxins has been described [50,51]. The three studied enzymes Lgt1/2/3 target one host molecule, eEF1A, and have been implicated in inhibition of eukaryotic protein synthesis and target-cell death [52]. *L. longbeachae* encodes two putative specific cytotoxic glucosyltransferases Llo1721 and Llo1578. They share only low homology with the *L. pneumophila* Lgt proteins with 23% protein identity over 62% of the protein length and 36% protein identity over 32% of the length, respectively. However, the DXD motif that is critical for enzymatic activity of clostridial enzymes is conserved suggesting that these enzymes might also be active in *L. longbeachae*. We also identified Llo3231 as another putative specific glucosyltransferase with a DXD motif, distantly related to the *L. pneumophila* SetA protein (23% protein identity over 67% of the protein length). SetA is known to cause delay in

Table 3. Distribution of selected Dot/Icm substrates of *L. pneumophila* in the *L. longbeachae* genomes.

<i>L. pneumophila</i>				<i>L. longbeachae</i>				Name	Description
Phila-1	Paris	Lens	Corby	NSW150	A	B	C		
lpg0012	lpp0012	lpl0012	lpc0013	llo0432	+	+	+	cegC1	Ankyrin repeat
lpg0038	lpp0037	lpl0038	lpc0039	–	–	–	–	ankQ/legA10	Ankyrin repeat
lpg0103	lpp0117	lpl0103	lpc0122	llo3312	+	+	+	vipF	GNAT family
lpg0171	lpp0233	lpl0234	–	–	–	–	–	legU1	F-box motif
lpg0234	lpp0304	lpl0288	lpc0309	llo0425	+	+	+	sidE/laiD	Unknown
lpg0257	lpp0327	lpl0310	lpc0334	llo2362	+	+	+	sdeA	Multidrug resistance protein
lpg0276	lpp0350	lpl0328	lpc0353	llo0327	+	+	+	legG2	Ras guanine nucleotide exchange
lpg0376	lpp0443	lpl0419	lpc2967	–	–	–	–	sdhA	GRIP, coiled-coil
lpg0390	lpp0457	lpl0433	lpc2954	llo2824	+	+	+	vipA	Unknown
lpg0402	–	–	–	–	–	–	–	ankY/legA9	Ankyrin, STPK
lpg0403	lpp0469	lpl0445	lpc2941	–	–	–	–	ankG/ankZ/ legA7	Ankyrin
lpg0436	lpp0503	lpl0479	lpc2906	–	–	–	–	ankJ/legA11	Ankyrin
lpg0483	lpp0547	lpl0523	lpc2861	llo2705	+	+	+	ankC/legA12	Ankyrin
lpg0621	lpp0675	lpl0658	lpc2673	–	–	–	–	sidA	Unknown
lpg0642	lpp0696	lpl0679	lpc2651	–	–	–	–	wipB	Unknown
lpg0695	lpp0750	lpl0732	lpc2599	–	–	–	–	ankN/ankX legA8	Ankyrin
lpg0940	lpp1002	lpl0971	lpc2349	–	–	–	–	lidA	Unknown
lpg1227	lpp1235	lpl1235	lpc0696	–	–	–	–	vpdB	Acyl transferase/hydrolase
lpg1328	lpp1283	lpl1282	lpc0743	–	–	–	–	legT	Thaumatococcus domain
lpg1355	lpp1309	–	–	–	–	–	–	sidG	Coiled-coil
lpg1488	lpp1444	lpl1540	lpc0903	–	–	–	–	lgt3/legC5	Coiled-coil
lpg1588	lpp1546	lpl1437	lpc1013	–	–	–	–	legC6	Coiled-coil
lpg1642	lpp1612*	lpl1384	lpc1071	llo1144	+	+	+	sidB	Rtx toxin, lipase
lpg1701	lpp1666	lpl1660	lpc1130	–	–	–	–	ppeA/legC3	Coiled-coil
lpg1718	lpp1683	lpl1682	lpc1152	–	–	–	–	ankI/legA4	Ankyrin
lpg1884	lpp1848	lpl1845	lpc1331	–	–	–	–	ylfB/legC2	Coiled-coil
lpg1950	lpp1932	lpl1919	lpc1423	llo1397	+	+	+	ralF	Sec-7 domain
lpg1953	lpp1935	lpl1922	lpc1426	–	–	–	–	legC4	Coiled-coil
lpg1978	lpp1961	lpl1955	lpc1464	–	–	–	–	setA	Putative Glycosyltransferase
lpg2137	lpp2076	lpl2066	lpc1586	–	–	–	–	legK2	STPK
lpg2144	lpp2082	lpl2072	lpc1593	–	–	–	–	ankB/legAU13ceg27	Ankyrin, F-box
lpg2155	lpp2094	lpl2083	lpc1604	llo3096	+	+	+	sidJ	Unknown
lpg2157	lpp2096	lpl2085	lpc1618	–	–	–	–	sdeC	Unknown
lpg2176	lpp2128	lpl2102	lpc1635	–	–	–	–	legS2	Sphingosine-1-phosphate lyase 1
lpg2222	lpp2174	lpl2147	lpc1689	–	–	–	–	lpnE	Sel-1 repeats
lpg2298	lpp2246	lpl2217	lpc1763	llo1707	+	+	+	ylfA/legC7	Coiled-coil
lpg2300	lpp2248	lpl2219	lpc1765	llo0584	+	+	+	ankH/legA3/ankW	Ankyrin, NFκB inhibitor
lpg2322	lpp2270	lpl2242	lpc1789	llo0570	+	+	+	ankK/legA5	Ankyrin
lpg2452	lpp2517	lpl2370	lpc2026	–	–	–	–	ankF/legA14/ceg31	Ankyrin
lpg2456	lpp2522	lpl2375	lpc2020	llo0365	+	+	+	ankD/legA15	Ankyrin
lpg2464	–	lpl2384	–	–	–	–	–	sidM/drrA	Unknown
lpg2465	–	lpl2385	–	–	–	–	–	sidD	Unknown
lpg2490	lpp2555	lpl2411	lpc1987	llo0796	+	+	+	lepB	Coiled-coil, Rab1 GAP
lpg2508	lpp2576	lpl2430	lpc1963	–	–	–	–	sdjA	Unknown
lpg2511	lpp2579	lpl2433	lpc1959	llo3098	+	+	+	sidC	PI(4)P binding domain
lpg2556	lpp2626	lpl2481	lpc1906	llo2218	+	+	+	legK3	STPK
lpg2584	lpp2637	lpl2507	lpc0561	–	–	–	–	sidF	Unknown
lpg2718	lpp2775	lpl2646	lpc0415	–	–	–	–	wipA	Unknown
lpg2793	lpp2839	lpl2708	lpc3079	–	–	–	–	lepA	Coiled-coil

Table 3. Cont.

<i>L. pneumophila</i>				<i>L. longbeachae</i>				Name	Description
Phila-1	Paris	Lens	Corby	NSW150	A	B	C		
<i>lpg2829</i>	<i>lpp2883</i>	–	–	–	–	–	–	<i>sidH</i>	Unknown
<i>lpg2830</i>	<i>lpp2887</i>	–	–	–	–	–	–	<i>lubX/legU2</i>	U-box motif
<i>lpg2831</i>	<i>lpp2888</i>	<i>lpl4276</i>	–	–	–	–	–	<i>VipD</i>	Patatin-like phospholipase
<i>Lpg2999</i>	<i>lpp3071</i>	<i>lpl2927</i>	<i>lpc3315</i>	–	–	–	–	<i>legP</i>	Astacin protease

*pseudogene, *lpp1612a* et *l612b*; A: *L. longbeachae* strain ATCC39462; B: 98072; C: C-4E7.
doi:10.1371/journal.pgen.1000851.t003

vacuolar trafficking [53], however its glucosylating activity remains to be established. In contrast, *L. longbeachae* does not encode a homologue of the *L. pneumophila* structural toxin protein RtxA, however we identified a homolog of the TcaZ toxin (Llo1558) present in the insect pathogen *Photobacterium luminescens* [54].

Many metabolic features of the genome of *L. longbeachae* reflect its soil habitat

L. longbeachae encodes a variety of proteins probably devoted to the metabolism of compounds present in plant cell walls, going in hand with the fact that that bacterium can be isolated from composted plant material. The main components of the plant cell wall are cellulose, hemicellulose and pectin. Cellulose utilization by microorganisms involves endo-1,4-beta-glucanases, cellobiohydrolases and β -glucosidases, that act synergically to convert cellulose to glucose. Examination of the *L. longbeachae* genome sequence revealed the presence of twelve such cellulolytic enzymes. Five glucanases, four cellobiohydrolases and three β -glucosidases are present. Interestingly, *L. pneumophila* also encodes two putative endo-1,4-beta-glucanases and one putative β -glucosidase but does not encode any cellobiohydrolase.

Within the plant cell wall, the cellulose microfibrils are linked via hemicellulosic tethers to form the cellulose-hemicellulose network, which is embedded in the pectin matrix. To gain access to cellulose in plant material, pectin and hemicellulose hydrolysis is necessary. Interestingly, *L. longbeachae* encodes three pectin lyases (Llo1693, Llo1410, Llo1162). The last two proteins possess a signal peptide and may therefore be secreted. Pectin lyases are virulence factors usually found in phytopathogenic microorganisms that degrade the pectic component of the plant cell wall. In addition to these specific enzymes and similar to *L. pneumophila*, *L. longbeachae* encodes a protein homologous to endo-1,4-beta-xylanase. Endo-1,4-beta-xylanase hydrolyses xylan the most common hemicellulose polymer in the plant kingdom and the second most abundant polysaccharide on earth. So, unlike *L. pneumophila*, which does not possess cellobiohydrolase and pectin lyase, *L. longbeachae* seems to be fully equipped to utilize cellulose as a carbon source (Table 5). Soil bacteria also often hydrolyse chitin by the means of chitinases to use it as a carbon source. Chitin originates mainly from the cell wall of fungi and cuticles of crustaceans or insects. In line with the fact that *L. longbeachae* is isolated from soil, we found two chitinases (Llo0050, Llo1558) that are predicted to be secreted proteins. However, the homologue of ChiA from *L. pneumophila* that was shown to be involved in infection of lungs of A/J mice [31] is absent from *L. longbeachae*.

Interestingly, *L. longbeachae* encodes a putative cyanophycin synthase (Llo2537) and therefore may be able to synthesize cyanophycin. Cyanophycin is an amino acid polymer composed of an aspartic acid backbone and arginine side groups. It serves as a

storage compound for nitrogen, carbon and energy in many cyanobacteria. *Acinetobacter baylyi* strain ADP1 was the first non-cyanobacterial strain shown to synthesize cyanophycin, a metabolic capacity that is still restricted to only few prokaryotes [55–58]. *L. longbeachae* also harbors a putative cyanophycinase (Llo2536) enabling the degradation of cyanophycin to dipeptides and a dipeptidase (Llo2535) necessary to hydrolyze beta-Asp-Arg dipeptides. *L. longbeachae* may thus be able to completely utilize cyanophycin, providing a mechanism for energy supply under substrate-limited conditions.

Genome and electron microscopy analysis indicates that *L. longbeachae* encodes a capsule

In the genome of *L. longbeachae* NSW150 we identified two gene clusters encoding proteins that are predicted to be involved in production of lipopolysaccharide (LPS) and/or capsule (Figure 3). Neither shared homology with the *L. pneumophila* LPS biosynthesis gene cluster. One region of 48 kb spans from *llo3148* to *llo3180* (Figure 3A) and the second of 24 kb from *llo0217* to *llo0236* (Figure 3B). In total they contain 26 genes for synthesis of the nucleotide sugar precursor, 12 genes encoding putative glycosyltransferases, 5 polysaccharide translocation genes including homologs of the *ctrABCD* capsule transport operon of *N. meningitidis*, and 10 genes of unknown function (Table S4). The finding that *L. longbeachae* might be encapsulated was further substantiated by electron microscopy analysis. Figure 4 shows that a capsule-like structure surrounds the bacteria.

Gene clusters encoding the core lipopolysaccharide of *L. pneumophila* and *L. longbeachae* are not conserved; however we identified in the genome of *L. longbeachae* homologs of *L. pneumophila* lipidA biosynthesis genes. Llo2684, Llo1461, Llo2686 and Llo0524 are homologous to LpxA, LpxB, LpxD and WaaM lipidA biosynthesis proteins with respectively 84%, 68%, 60% and 78% of identity. Predictions deduced from the sequence analysis of strain NSW150 did not clarify which region was coding for the LPS and which for the capsule. Further insight into the LPS and capsule encoding regions came from the comparison of this region among the four *L. longbeachae* genomes sequenced. The 24 kb region B is identical between the two Sg1 strains sequenced and identical between the two Sg2 strains analyzed, but the Sg1 and Sg2 strains differed from each other in an approximately 10 kb region carrying glycosyltransferases, methyltransferases, and LPS biosynthesis proteins (Figure S6). In contrast the putative capsule encoding region A was highly conserved among all four strains sequenced except for a region carrying three genes, that differed among all four strains independent of the Sg. However, as it is not known whether the Sg specificity of *L. longbeachae* is defined by its capsule or by LPS, further studies are necessary to clearly define the function of the proteins encoded in these two genomic regions.

Table 4. Putative new type IV secretion substrates specific for *L. longbeachae*.

NSW150	ATCC39462	98072	c-4E7	Motif	A	B	C
llo0037	+	+	+	ankyrin	+	42,86	60,00
llo0087	+	+	+	ankyrin	+	57,14	53,33
llo0115	+	+	+	ankyrin	+	28,57	53,33
llo0246	+	+	+	ankyrin	+	28,57	66,67
llo0990	+	---	---	ankyrin	+	28,57	46,67
llo1043	+	+	+	ankyrin	+	28,57	46,67
llo1142	+	+	+	ankyrin	+	28,57	53,33
llo1168	+	+	+	ankyrin	+	28,57	53,33
llo1371	+	+	+	ankyrin, coiled-coil	+	28,57	66,67
llo1395	+	+	+	ankyrin	+	42,86	53,33
llo1618	+	+	+	ankyrin	+	28,57	66,67
llo1646	+	+	+	ankyrin	+	28,57	40,00
llo1651	+	+	+	ankyrin	+	14,29	60,00
llo1715	+	+	+	ankyrin	+	28,57	40,00
llo1742	+	+	+	ankyrin	+	57,14	46,67
llo1894	+	+	+	ankyrin	+	28,57	66,67
llo2133*	+	+	+	ankyrin	+	0,00	33,33
llo2476	+	+	+	ankyrin	+	14,29	46,67
llo2668	+	+	+	ankyrin	+	14,29	46,67
llo3081	+	+	+	ankyrin, patatin-like phospholipase	+	28,57	60,00
llo3093	+	+	+	ankyrin, STPK	+	0,00	66,67
llo3343	+	+	+	ankyrin	+	14,29	33,33
llo3353	+	+	+	ankyrin, NUDIX hydrolase	+	28,57	53,33
llo0114	+	+	+	LRR	+	14,29	40,00
llo1314	+	+	+	LRR	+	0,00	40,00
llo2165	+	+	+	LRR	+	42,86	66,67
llo2494	+	+	+	LRR	+	28,57	66,67
llo3116	---	---	---	LRR	+	57,14	26,67
llo3118	---	---	---	LRR	+	28,57	66,67
llo1139	+	+	+	STPK	+	14,29	33,33
llo1681	+	+	+	STPK	+	42,86	73,33
llo2132	+	+	+	STPK, coiled-coil	-	14,29	73,33
llo2984	+	+	+	STPK	+	14,29	53,33
llo3049	+	+	+	STPK	+	14,29	66,67
llo1984	+	+	+	STPK	+	14,29	33,33
llo1427	+	+	+	F-Box	+	14,29	66,67
llo2109	+	+	+	F-Box	+	28,57	60,00
llo0448	+	+	+	U-Box	+	28,57	73,33
llo1404	+	+	+	PPR	+	28,57	20,00
llo2643	+	+	+	PPR, coiled-coil	+	28,57	46,67
llo2200	+	+	+	TTL	+	14,29	53,33
llo2327	+	+	+	SH2	+	28,57	73,33
llo2352	+	+	+	PAM2	+	42,86	60,00
llo1196	+	+	+	Snare	+	0,00	73,33
llo2381	+	+	+	Snare	+	42,86	60,00
llo0793	+	+	+	Phosphatidylinositol-4-phosphate 5-kinase	+	28,57	66,67
llo3288	+	+	+	Ras-related small GTPase domain	+	14,29	60,00
llo2329	+	+	+	Ras-related small GTPase, Miro-like domain	+	28,57	60,00
llo2249	+	+	+	Miro-like domains	+	57,14	80,00
llo1716	+	+	+	Ras-related small GTPase, Miro-like domain	+	28,57	73,33
llo1892	+	+	+	Putative Immunoglobulin I-set domain	+	14,29	40,00

(A) Presence of a hydrophobic residue or a proline in positions -3 or -4 according to [35]. (B) Enrichment in amino acids that have small side-chains (alanine, glycine, serine and threonine) at positions -8 to -2 according to [36]. (C) Percentage of Polar aminoacids that are favored at positions -13 to +1 according to [36].
doi:10.1371/journal.pgen.1000851.t004

Table 5. Predicted *L. longbeachae* enzymes that may be involved in cellulose degradation.

Gene	Annotation	SignalP	Predicted localization (PSORTb ⁺)	ATCC39462	98072	c-4E7	Homology with <i>L. pneumophila</i>
<i>llo2355</i>	Putative endo-1,4-beta-glucanase	+	Unknown	100%	99%	99%	–
<i>llo3308</i>	Putative endo-1,4-beta-glucanase	+	Unknown	100%	99%	99%	<i>lpp1893/lpg1918/lpl1882/LPC_1372</i>
<i>llo3305</i>	Putative endo-1,4-beta-glucanase	+	Unknown	100%	99%	99%	<i>lpp0546/lpg0482/lpl0522/LPC_2862</i>
<i>llo1381</i>	Putative endo-1,3(4)-beta-glucanase	+	Unknown	100%	99%	99%	–
<i>llo0032</i>	Putative cellobiohydrolase	+	Unknown	100%	99%	99%	–
<i>llo0965</i>	Putative cellobiohydrolase	+	Extracellular	100%	98%	98%	–
<i>llo1892</i>	Putative cellobiohydrolase	+	Extracellular	100%	99%	99%	–
<i>llo2999</i>	Putative cellobiohydrolase	+	Unknown	100%	99%	99%	–
<i>llo1023</i>	Putative beta-glucosidase	+	Unknown	100%	99%	99%	<i>lpp1193/lpg1191/lpl1199/LPC_0658</i>
<i>llo0330</i>	Putative beta-glucosidase	–	Cytoplasmic	100%	99%	99%	–
<i>llo2462</i>	Putative beta-glucosidase	–	Cytoplasmic	100%	99%	99%	<i>lpp0946/lpg0885/lpl0916/LPC_2408</i>
<i>llo0816</i>	Putative endo-1,4-beta-xylanase	+	Unknown	100%	99%	99%	<i>lpp0767/lpg0712/lpl0749/LPC_2581</i>
<i>llo1693</i>	Putative pectin lyase	–	Unknown	100%	96%*	96%*	–
<i>llo1410</i>	Putative pectin lyase	+	Extracellular	100%	99%	99%	–
<i>llo1162</i>	Putative pectin lyase	+	Extracellular	100%	99%	99%	–

*frameshift at the N-terminus

+ PSORTb bacterial protein localization prediction tool (<http://www.psort.org/psortb/>)
doi:10.1371/journal.pgen.1000851.t005

L. longbeachae does not encode flagella explaining differences in mouse susceptibility as compared to *L. pneumophila*

Cytosolic flagellin of *L. pneumophila* triggers Naip5-dependent caspase-1 activation and subsequent proinflammatory cell death by pyroptosis in C57BL/6 mice rendering these mice resistant to infection with *L. pneumophila* [22–24,59–62]. In contrast, caspase-1 activation does not occur upon infection of C57BL/6 and A/J mice macrophages with *L. longbeachae*, which is then able to replicate. One possible explanation has been that due to a lack of pore-forming activity, *L. longbeachae* flagellin may not have access to the cytoplasm of the macrophage where it is thought to be involved in caspase-1 activation. Alternatively, *L. longbeachae* flagellin may not be recognized by the Naip5 pathway [11]. Genome analysis clarified this issue, as we found that *L. longbeachae* does not carry any flagellar biosynthesis genes except the sigma factor FliA, the regulator FleN, the two-component system FleR/FleS and the flagellar basal body rod modification protein FlgD. Interestingly, as shown in Figure 5, all genes bordering flagellar gene clusters were conserved between *L. longbeachae* and *L. pneumophila*, suggesting deletion of these regions from the *L. longbeachae* genome. Furthermore, not a single homologue of flagellar biosynthesis genes could be identified in other parts of the genome. Analysis of the three additional genome sequences of strains *L. longbeachae* ATCC39642, 98072 and C-4E7 confirmed the results. To further investigate this unexpected result, we designed primers in the conserved flanking genes to analyze these genomic regions in 15 *L. longbeachae* strains. All strains tested, eleven of Sg1 and four of Sg2, displayed the same organization as the sequenced strain (Table S5). According to these results, we propose that *L. longbeachae* fails to activate caspase-1 due to the lack of flagellin, which may also partly explain the differences in mouse susceptibility to *L. pneumophila* and *L. longbeachae* infection. The putative *L. longbeachae* capsule may also contribute to this difference.

Although *L. longbeachae* does not encode flagella, it encodes a putative chemotaxis system. Chemotaxis enables bacteria to find favorable conditions by migrating towards higher concentrations of

attractants. The chemotactic response is mediated by a two-component signal transduction pathway, with the histidine kinase CheA and the response regulator CheY, putatively encoded by the genes *llo3302* and *llo3303* respectively, in the *L. longbeachae* genome. Furthermore, two homologues of the ‘adaptor’ protein CheW (encoded by *llo3298*, *llo3300*) that associate with CheA or cytoplasmic chemosensory receptors are present. Ligand-binding to receptors regulates the autophosphorylation activity of CheA in these complexes. The CheA phosphoryl group is subsequently transferred to CheY, which then diffuses away to the flagellum where it modulates motor rotation. Adaptation to continuous stimulation is mediated by a methyltransferase CheR encoded by *llo3299* in *L. longbeachae*. Together, these proteins represent an evolutionarily conserved core of the chemotaxis pathway, common to many bacteria and archaea [55,63]. A similar chemotaxis system is also present in *L. drancourtii* LLAP12 [64] but it is absent from *L. pneumophila*. The flanking genomic regions are highly conserved among *L. longbeachae* and all *L. pneumophila* strains sequenced, suggesting that *L. pneumophila*, although it encodes flagella has lost the chemotaxis system encoding genes.

We also observed using electron microscopy (Figure 4) that *L. longbeachae* possesses a long pilus-like structure. Indeed, all genes necessary to code for type IV pili are present in the genome of *L. longbeachae* and are, with 63–88% amino acid similarity, highly conserved between *L. longbeachae* and *L. pneumophila*. Taken together genome analysis revealed interesting features of the *Legionella* genomes: both encode pilus-like structures, in contrast *L. longbeachae* encodes a chemotaxis system but no flagella, and *L. pneumophila* encodes flagella but no chemotaxis system. It will be an interesting aspect of future research to understand these particular features of the two *Legionella* species.

The regulatory repertoire of *L. longbeachae* suggests different adaptation mechanisms as compared to *L. pneumophila*

Similar to the *L. pneumophila* genomes and consistent with its intracellular lifestyle, the regulatory repertoire of *L. longbeachae* is




Figure 3. Putative capsule and LPS encoding loci in the genome of *L. longbeachae*. (A) 48 kb chromosomal region highly conserved in the four *L. longbeachae* genomes sequenced putatively encoding the capsular biosynthesis genes. (B) 24 kb chromosomal region differing between Sg1 and Sg2 isolates putatively encoding the lipopolysaccharide biosynthesis genes of *L. longbeachae*. Colors indicate different classes of genes: magenta, synthesis pathway of nucleoside sugar precursors; blue, glycosyltransferase; yellow transportation; grey, genes of unknown. doi:10.1371/journal.pgen.1000851.g003




Figure 4. Electron microscopy showing the presence of capsule like structures. Transmission electron micrographs of *L. longbeachae* cells cultured in BYE broth to post exponential growth phase (OD600 3.8). Black arrows, putative capsule structures, red Arrow, putative pili. doi:10.1371/journal.pgen.1000851.g004

rather small. Genome analysis identified 121 transcriptional regulators (113–116 in the four sequenced *L. pneumophila* genomes), which represent only 3.0% of the predicted genes (Table S6). Similar to *L. pneumophila*, *L. longbeachae* encodes six putative sigma factors, RpoD, RpoH, RpoS, RpoN, FliA and the ECF-type sigma factor RpoE.

The most abundant class of regulators of *L. pneumophila* is the GGDEF/EAL family (24 or 23 in all *L. pneumophila* genomes sequenced). This is significantly different in *L. longbeachae*, as we identified only 14 GGDEF/EAL domain-containing regulators, despite the larger size of the *L. longbeachae* genome. Furthermore, this group of regulators may fulfill specific functions in *L. longbeachae*, since most of the regulators possess no orthologs in the *L. pneumophila* genomes (Table S6). The function of these regulators in *L. pneumophila* and *L. longbeachae* is unknown, but in other bacteria these regulators play a role in aggregation, biofilm formation, twitching motility or flagella regulation. In *L. pneumophila* it was suggested, as deduced from gene expression analysis, that some of the GGDEF/EAL regulators may play a role in modulating flagella expression [65,66], thus the lower number of GGDEF/EAL domain-containing proteins of *L. longbeachae* may in part be related to the missing flagellum.

Another difference in the regulatory repertoire of the two *Legionella* species was observed for two component systems. There




Figure 5. Alignment of the chromosomal regions of *L. pneumophila* and *L. longbeachae* coding the flagella biosynthesis genes. The comparison shows that all except the regulatory genes are missing in *L. longbeachae*. Red, conserved regulator encoding genes, grey arrows orthologous genes among the genomes, white arrows, non orthologues genes.
doi:10.1371/journal.pgen.1000851.g005

are 14 response regulators and 13 histidine kinases in *L. pneumophila*, and 17 response regulators and 16 histidine kinases in the *L. longbeachae* genome, but only half of the *L. longbeachae* response regulators possess an ortholog in *L. pneumophila*. For example the recently described two-component system LqsS/LqsR that is part of a quorum sensing system in *L. pneumophila* is missing in *L. longbeachae* [67–69]. Two-component systems are involved in signal transduction pathways that enable bacteria to sense, respond, and adapt to a wide range of environments, stressors, and growth conditions [70]. Different two-component systems may be linked to the different environments to which *L. longbeachae* has to adapt compared to *L. pneumophila*.

In *L. longbeachae*, cyclic AMP may also transduce cellular signals as the genome encodes eight class III adenylate cyclases (Llo0181, Llo1751, Llo2196, Llo1669, Llo0753, Llo1197, Llo1216, Llo3304) of which only one (Llo0181) is also conserved in *L. pneumophila*. LadC, an adenylate cyclase of *L. pneumophila* that was shown to have a significant role in the initiation of infection *in vitro* and *in vivo* [71], is absent from *L. longbeachae*. As shown for *Pseudomonas aeruginosa*, these class III adenylate cyclases may sense environmental signals ranging from nutritional content of the surrounding media to the presence of host cells and control virulence gene expression accordingly [72]. Furthermore, 13 proteins containing cAMP binding motifs were identified, only one of which is shared with *L. pneumophila*, again indicating specific regulatory circuits for *L. longbeachae*. This high number of proteins that may sense cAMP indicates the potential importance of this signaling molecule in *L. longbeachae*.

In contrast, the regulators shown to be important for growth phase and life cycle dependent gene expression, such as the two component system LetA/LetS (Llo2653/llo1235), the RNA-binding protein CsrA (Llo2071), the two small RNAs RsmY and RsmZ regulating CsrA [66,73], SpoT (Llo0908) and RelA (Llo1756) are conserved in *L. longbeachae*. Likewise, the two-

component systems PmrAB (Llo1159/Llo1158) and CpxRA (Llo1781/Llo1782) that regulate the Dot/Icm T4SS system and some of its substrates are both conserved in *L. longbeachae* [74–76].

Global gene expression analysis reveals differences in the *L. longbeachae* and *L. pneumophila* life cycles

It has been shown in several studies that *L. pneumophila* exhibits at least two developmental stages, a replicative/avirulent and a transmissive/virulent phase that are each characterized by the expression of specific traits [9]. These stages are also reflected in a major shift in the gene expression program of *L. pneumophila* between the two phases of its life cycle [65]. In order to investigate, whether *L. longbeachae* had a similar biphasic life cycle we studied its gene expression program in exponential and post exponential growth phase *in vitro*. A multiple-genome microarray was constructed containing 10 692 gene-specific oligonucleotides representing 3567 genes predicted in the genome and on the plasmid and 3010 oligonucleotides specific for intergenic regions. RNA of *in vitro* grown bacteria was sampled at OD 2.5 (exponential growth) and at OD 3.7 (post exponential growth) and the global gene expression program was compared.

Overall, 187 genes in *L. longbeachae* were upregulated in the exponential (E) phase (likewise, downregulated in the postexponential phase, Table S7), and 313 genes were upregulated in the postexponential (PE) phase (downregulated in the E phase, Table S8). Real-time PCR analysis of selected genes validated the microarray results (data not shown). If we compare these results to those obtained for *L. pneumophila* grown *in vitro* [65], we observed several differences. In *L. pneumophila* strain Paris 543 genes are upregulated in E phase. Of the genes present in both genomes 270 are only upregulated in *L. pneumophila* but not in *L. longbeachae*. The 117 genes that are upregulated in both species in exponential phase include many ribosomal proteins, the genes belonging to the ATP synthase machinery (*atp* genes), the NADH deshydrogenase

(*nuo* genes), most of the genes involved in translocation systems (*sec* genes) and several enzymatic activities (Table S7). However, several metabolic pathways clearly induced in E phase in *L. pneumophila* are not induced in *L. longbeachae*. These include the formyl THF biosynthesis, the purine and pyrimidine and the tetrahydrofolate biosynthesis pathways. Furthermore, genes coding for several chaperones (DnaJ, DnaK or GroES), the regulatory protein RecX and several proteins related to starvation and stress are not upregulated in E phase *L. longbeachae*. There are only 11 genes specific for *L. longbeachae* and induced in E phase, all of which code proteins for which no function could be predicted.

In PE phase 313 genes are upregulated in *L. longbeachae*, of which only 53 are also among the 441 PE phase genes of *L. pneumophila*. Interestingly, 208 of the genes upregulated in PE in *L. longbeachae* have no orthologs in *L. pneumophila*, and for 70% of these no function could be predicted. Thus the response of *L. longbeachae* to PE phase growth is distinct from that of *L. pneumophila*. In particular we observed differences in the expression profiles of many factors known to be involved in *L. pneumophila* virulence. For example, of the genes coding putative substrates of the Dot/Icm secretion system only few, *sidC* (llo3098), *sdhB* (llo2439), *sidE* homologue (llo2210), *sdeC/laiC* (llo3092) and *sdeB/laiB* (llo3095) are upregulated in post-exponential phase. However, several of the newly identified putative substrates are induced in *L. longbeachae* in PE phase. These comprise seven proteins homologous to Sid proteins of *L. pneumophila* (Llo0424, Llo0426, Llo2210, Llo2439, Llo3092, Llo3095 and Llo3098), three genes coding homologues of EnhA (llo0852, llo1475 and llo2343), three ankyrin proteins (Llo0115, Llo1646 and Llo1715) and a putative serine threonine kinase (Llo1139). However, clear differences in gene expression between *L. pneumophila* and *L. longbeachae* exist and the switch from replicative to transmissive phase seems to be less pronounced in *L. longbeachae* than in *L. pneumophila*. Interestingly, the genes coding the stationary phase sigma factor RpoS and the sigma factor 28 (FliA) and CsrA, all involved in the regulation of the biphasic life cycle of *L. pneumophila* are not differentially regulated in *L. longbeachae*. In contrast, seven GGDEF/EAL domain-containing regulators (llo0090, llo1253, llo1377, llo2005, llo3125, llo3392 and llo3414) and four cAMP binding proteins (llo3395, llo2387, llo2141 and llo1336) are induced in PE phase. Thus cyclic di-GMP and cAMP may be important signaling molecules for regulating PE phase traits of *L. longbeachae*. According to our transcriptome analysis, the switch in the lifecycle of *L. longbeachae* appears less pronounced as compared to *L. pneumophila*, and regulation may be achieved mainly by secondary messenger molecules.

Concluding remarks

L. longbeachae is the second leading cause of Legionnaires' disease in the world and a major cause of pneumonia in Australia and New Zealand. Yet, still very little is known about its virulence strategies and the genetic basis of virulence and niche adaptation. Analysis of the genome sequences of four *L. longbeachae* strains and its comparison with the published *L. pneumophila* genomes has uncovered important differences in the genetic repertoire of the two species and suggests different strategies for intracellular replication and niche adaptation.

Similar to *L. pneumophila*, *L. longbeachae* encodes a type IVB secretion system homologous to the Dot/Icm system. Inactivation of the type IV secretion system, through deletion of the *dotA* gene, showed that it is essential for virulence, as the *dotA* mutant had a severe growth defect in *A. castellanii* infection and could not establish an infection in the lungs of A/J mice. Despite this resemblance to *L. pneumophila*, the secreted effectors are very different as only 44% of the known *L. pneumophila* substrates were conserved in *L. longbeachae*.

However, like *L. pneumophila*, many of them have eukaryotic domains or resemble eukaryotic proteins. Thus a large cohort of eukaryotic-like proteins was also a specific feature of the *L. longbeachae* genomes. An emerging theme in bacterial virulence is the evolution of virulence factors that can mimic the activities of Ras small GTPases (for a review see [77]). Small GTPases regulate unique biological functions of the cell as diverse as cell division/differentiation, actin cytoskeleton rearrangements, intracellular membrane trafficking. *L. pneumophila* produces the effector proteins RalF [78] and SidM/DrrA [40,41] that activate small G-protein signaling cascades and interfere with host membrane trafficking. Here we identified *L. longbeachae* specific proteins belonging to the Rab subfamily of Ras small GTPases. These are the first prokaryotic Rab GTPases described and they may account for some of the differences in phagosome maturation between *L. longbeachae* and *L. pneumophila*. Overall, more than 3% of the *L. pneumophila* genome is thought to encode T4SS substrates that fulfill various functions, such as interfering with small GTPases of the early secretory pathway, disrupting phosphoinositide signaling or targeting microtubule-dependent vesicular transport. They may represent new strategies to interfere with host cell processes and may partly explain variations in the replication cycle of the two species.

An intriguing and unresolved question has been the susceptibility of C57BL/6 mice to *L. longbeachae* infection but their resistance to *L. pneumophila* infection. Only A/J mice that carry a particular Naip-5 allele are susceptible to *L. pneumophila* infection. Genome analysis has provided some insight into this question through the observation that *L. longbeachae* does not encode flagella, and thus does not trigger Naip5-dependent caspase-1 activation and subsequent proinflammatory cell death by pyroptosis [22–24,59–62]. In contrast, *L. longbeachae* encodes a capsule that might be implicated in the recognition by the host immune system and which may provide some protection against killing by phagocytes. In *L. pneumophila*, expression of flagella is a hallmark of transmissive, virulent bacteria and a marker of its biphasic life cycle. In line with the absence of flagella, *L. longbeachae* also seems to have a less pronounced life cycle switch, as transcriptome analysis revealed a less dramatic change in gene expression compared to *L. pneumophila*. This result might explain the fact that intracellular proliferation of *L. longbeachae* is independent of the growth phase [11].

Previously we and others hypothesized, that *L. pneumophila* had acquired DNA by horizontal transfer or by convergent evolution during its co-evolution with free-living amoebae [25,79] and that *L. pneumophila* uses molecular mimicry to subvert host functions [8,80]. Presumably, *L. longbeachae* is not only able to interact with protozoa but also with plants, as several proteins present in plants and several enzymes which might confer the ability to degrade plant material were identified in the *L. longbeachae* genome.

Interestingly, the comparison of the genome sequence of four strains of *L. longbeachae* identified high gene content conservation unlike *L. pneumophila*. Furthermore, between strains of the same serogroup very few SNPs are present, most of them located in few plasticity zones, indicating recent expansion of this species. Based on these genome sequences, future comparative and functional studies will allow definition of the common and distinct survival tactics of pathogenic *Legionella* spp. and may open new ways to combat *L. pneumophila* and *L. longbeachae* infections.

Materials and Methods

Ethics statement

All animal experiments were conducted with approval from the University of Melbourne Animal Ethics committee application ID 0704867.3.

DNA preparation and sequencing techniques

L. longbeachae strain NSW150 was grown on BCYE agar at 37°C for 3 days and chromosomal DNA was isolated by standard protocols. Cloning, sequencing and assembly were done as described [81]. One library (inserts of 1–3 kb) was generated by random mechanical shearing of genomic DNA, followed by cloning of the fragments into pcDNA-2.1 (Invitrogen). A scaffold was obtained by end-sequencing clones from a BAC library constructed as described [82] using pIndigoBac (Epicentre) as a vector. Plasmid DNA purification was done with a TempliPhi DNA sequencing template amplification kit (Amersham Biosciences). Sequencing reactions were done with an ABI PRISM BigDye Terminator cycle sequencing ready reactions kit and a 3730 XL Genetic Analyzer (Applied Biosystems). We obtained and assembled 40299 sequences and performed finishing by adding 1125 additional sequences, as described earlier [81]. For draft genome sequencing of strains ATCC39642, 98072 and C-4E7 Illumina, shotgun libraries were generated from 5 µg of genomic DNA each using the standard Illumina protocols. Sequencing was carried out on an Illumina Genome Analyzer II as paired-end 36bp reads, following the manufacturer's protocols and with the Illumina PhiX sample used as control. Image analysis and base calling was performed by the Genome Analyser pipeline version 1.3 with default parameters.

Annotation and sequence analysis

Definition of coding sequences and annotation were done as described [81] by using CAAT-box software [83] and MAGE (Magnifying Genomes) [84]. All predicted coding sequences were examined visually. Function predictions were based on BLASTp similarity searches and on the analysis of motifs using the PFAM, Prosite and SMART databases. We identified orthologous genes by reciprocal best-match BLAST and FASTA comparisons. Pseudogenes had one or more mutations that would prevent complete translation. Analysis of the three drafts genome sequences obtained by the Illumina technique was done as follows. First, to precisely determine the average insert size of mate-paired reads, we mapped the reads of each strain to the NSW150 sequence. Then, this value was used to give good mate-pair information to the de novo assembler. Short-reads were assembled *de novo* into contigs (without reference to any other sequence) using Velvet (version 0.7.55) [85]. To increase specificity and length of the generated contigs, we used the hash length (k-mer) of 27. Subsequently Mauve (version 2.3.0) [86] was used to build super contigs by aligning the de novo obtained contigs on the finished NSW150 sequence. Finally, for SNP discovery the program Maq (version 0.7.1) [87] was used for mapping the Solexa reads to the NSW150 reference. To detect high confidence SNPs, we only kept those SNPs that had a coverage of 10x to 300x. SNPs with a frequency lower than 80% were removed.

Construction of a *dotA* mutant in strain *L. longbeachae* NSW150

To construct the *dotA* mutant strain, the chromosomal region containing the *dotA* gene was PCR-amplified with the primers *dotA*-for CTCGCGCATTTGGAACCTTTAT and *dotA*-rev TTCGCTCATAAACCGCTCTT. The product was cloned into the pGEM-T Easy vector (Promega) yielding pGEM-*dotA*. We performed inverse PCR on pGEM-*dotA* with primers *dotA*inv-for CGCGGATCCCCGCATTTAATACGCCAAAC and *dotA*inv-rev CGCGGATCCAAGGTTTTGCGTTGGATAGG containing *Bam*HI overhangs allowing internal deletion of 2582 bp in *dotA*. PCR product was digested with *Bam*HI and ligated to the

kanamycin resistance cassette, which was amplified via PCR from the plasmid pGEM-Kan^R, using primers containing *Bam*HI restriction sites at the ends (Kan-*Bam*HI-for TGCAGGTCGACT-CAGAGGAT Kan-*Bam*HI-rev CGCGGATCCGAGCTCGG-TACC). Linearized vector was electroporated in *L. longbeachae* to obtain *dotA*::Km mutant.

Acanthamoeba castellanii infection assay

For *in vivo* growth of *L. longbeachae* and its *dotA* deletion mutant in *A. castellanii* we followed a protocol previously described [65]. In brief, three days old cultures of *A. castellanii* were washed in infection buffer (PYG 712 medium without proteose peptone, glucose, and yeast extract) and adjusted to 10⁵–10⁶ cells per ml. Stationary phase *Legionella* grown on BCYE agar, diluted in water were mixed with *A. castellanii* at a MOI of 0.1. After allowing invasion for 1 hour at 37°C the *A. castellanii* layer was washed twice with infection buffer (start point of time-course experiment). Intracellular multiplication was monitored using a 300 µl sample, which was centrifuged (14 000 rpm) and vortexed to break up amoeba. The number of colony forming units (CFU) of *Legionellae* was determined by plating on BCYE agar. Each infection was carried out in duplicates.

Pulmonary infection of A/J mice with *L. longbeachae*

The comparative virulence of *L. longbeachae* NSW150 and the *dotA*::Km derivative within A/J mice was examined via competition assays and in single infections, as described previously [21,34]. Briefly, 6- to 8-week-old female A/J mice (Jackson Laboratory, ME) were anesthetized and inoculated intratracheally with approximately 10⁵ CFU of each *L. longbeachae* strain under investigation. At 24 and 72 h following inoculation, mice were sacrificed and their lung tissue isolated. Tissue was homogenized, and complete host cell lysis was achieved by incubation in 0.1% saponin for 15 min at 37°C. Serial dilutions of the homogenate were plated onto both plain and antibiotic-selective BCYE agar to determine the number of viable bacteria and the ratio of wild-type to mutant bacteria colonizing the lung in mixed infections.

Electron microscopy

Bacteria were transferred to Formvar-carbon-coated copper grids after glow discharged for 3', stained with 1% uranyl acetate for 35sec, air dried and observed under a Jeol 1200EXII transmission electron microscope (Jeol, Tokyo, Japan) operated at 80kV. Digital acquisition was performed with a Mega View camera and the Analysis pro software version 3,1 (ELOISE, Roissy, France).

PCR analysis

PCR for the regions containing the flagella biosynthesis coding genes in strain *L. pneumophila* Paris and *L. longbeachae* NSW150 were amplified with genomic DNA of strain Paris and NSW150 respectively. Primers were designed using the Primer 3 software to amplify a specific fragment of about 1–3kb respectively for each region (melting temperatures 58°C). Amplification reactions were performed in a 50-µl reaction volume containing 6 ng of chromosomal DNA. The size of each PCR product was verified on agarose gels. Primers used are listed in Table S9.

Transcriptome analysis

L. longbeachae strain NSW150 was cultured in N-(2-acetamido)-2-aminoethanesulphonic acid (ACES)-buffered yeast extract broth or on ACES-buffered charcoal–yeast (BCYE) extract agar at 37°C. Total RNA was extracted as previously described [88]. *L.*

longbeachae was harvested for RNA isolation at the exponential (OD 2.5) and post-exponential phase (OD 3.7). RNA was prepared from three independent cultures and each RNA sample was hybridized twice to the microarrays (dye swap). RNA was reverse-transcribed with Superscript indirect cDNA kit (Invitrogen) and labeled with Cy5 or Cy3 (Amersham Biosciences) according to the supplier's instructions. The microarray containing 13 710 60mer oligonucleotides specific for 3567 predicted genes of the genome, the plasmid and all intergenic regions longer than 200nts has been designed using the program OligoArray (<http://berry.engin.umich.edu/oligoarray/>). Based on these sequences a custom oligonucleotide array was manufactured (Agilent Technologies) with a final density of 15K. For hybridization, Cy3 and Cy5 target quantities were normalized at 150 pmol. Arrays were scanned using an Axon 4000B scanner with fixed PMT (PMT = 550 for Cy3 and 650 for Cy5). Data were acquired and analyzed by Genepix Pro 5.0 (Axon Instrument). Spots were excluded from analysis in case of high local background fluorescence slide abnormalities or weak intensity. Data normalization and differential analysis were conducted using the R software (<http://www.r-project.org>). For each gene 3 probes were present on the microarray. Data for which at least 2 of the 3 probes gave a significant and non-contradictory result were taken into account. A loess normalization [89] was performed on a slide-by-slide basis (BioConductor package marray; <http://www.bioconductor.org/packages/bioc/html/marray.html>). Differential analysis was carried out separately for each comparison between two time points, using the VM method (VarMixt package [90], together with the Benjamini and Yekutieli [91] p-value adjustment method. The cut off for the expression ratio was set to either superior/equal to 2 or inferior/equal to 0.5 and the general ratio of expression of each gene was calculated as the average expression ratio from the different significant probes.

URLs

The sequence and the annotation of the *L. longbeachae* NSW150 genome is accessible at the LegioList Web Server (<http://genolist.pasteur.fr/LegioList> and <http://genolist.pasteur.fr/>) and under the EMBL/Genbank Accession number: FN650140 the *L. longbeachae* NSW150 plasmid under the EMBL/Genbank Accession number: FN650141. Due to new regulations for genome sequence submissions to EMBL/Genbank the gene names (locus_tag), which are e.g. llo0001 in the article and in the Institut Pasteur databases had to be changed to LLO_0001 in the sequence submission. According to the standards for genome sequences published by Chain and colleagues [92] the *L. pneumophila* NSW150 genome sequence can be defined as "Finished" and the three Solexa genome sequence drafts can be defined as "High-Quality Draft". The complete dataset for the transcriptome analysis is available at <http://genoscript.pasteur.fr> in a MIAME compliance public database maintained at the Institut Pasteur and was submitted to the ArrayExpress database maintained at <http://www.ebi.ac.uk/microarray-as/ae/> under the Accession number: A-MEXP-1779.

Supporting Information

Figure S1 Classification of the *L. longbeachae* CDS in the different COG groups. 2,506 CDS are classified in at least one COG group. Since several genes are assigned to multiple categories, the total number of assignments is greater than the number of ORFs in the genome.
Found at: doi:10.1371/journal.pgen.1000851.s001 (11.39 MB TIF)

Figure S2 Synteny plot of the chromosomes of *L. pneumophila* strain Paris and *L. longbeachae* NSW150. The plot was created using the mummer software package (<http://mummer.sourceforge.net/>).
Found at: doi:10.1371/journal.pgen.1000851.s002 (6.88 MB TIF)

Figure S3 Comparison of the plasmids identified in *L. longbeachae* and *L. pneumophila*. (A) Synteny LinePlot between the *L. longbeachae* plasmid and the plasmids of *L. pneumophila* strain Lens and Paris, respectively. Orthologous genes are defined by bi-directional blastP best hits (BDBH) or a blastP alignment threshold of 35% sequence identity over 80% of the length of the smaller protein. The gap parameter, representing the maximum number of consecutive genes that are not involved in a synteny group was 3. (B) Percentage of aminoacid identity among Tra proteins of the *L. longbeachae* and the *L. pneumophila* strain Lens as compared to the Tra region of strain Paris. (C) Venn diagram showing the common and specific gene content of the plasmids of *L. pneumophila* strains Paris, Lens and *L. longbeachae* NSW150.
Found at: doi:10.1371/journal.pgen.1000851.s003 (17.22 MB TIF)

Figure S4 Distribution of SNPs along the chromosome of *L. longbeachae* ATCC39462 (Sg1) and C-4E7 (Sg2) with respect to the completely sequence genome of *L. longbeachae* NSW 150 (Sg1). Outer circle, Mapping of SNPs between *L. longbeachae* Sg1 (NSW150) and Sg2 (C-4E7), central circle in green, sequence coverage of mapped reads of strain ATCC39462 on the NSW150 genome, inner circle; SNP distribution among the two Sg1 strains sequenced. 1426 SNPs are located in 7 genomic regions; region 1: *llo0557-llo0587* containing 112 SNPs; region 2: *llo0643-llo0653*, carries an integrase gene and contains 152 SNPs; region 3: *llo0814-llo0841* containing 38 SNPs; region 4: *llo0943-llo0952*, carries an integrase gene and contains 152 SNPs; region 5: *llo1813-llo1886*, carries many tra- like genes and contains 651 SNPs; region 6: *llo2119-llo2142*, contains 89 SNPs, region 7: *llo3148-llo3180*, carries genes encoding the putative capsule and contains 166 SNPs.
Found at: doi:10.1371/journal.pgen.1000851.s004 (10.54 MB TIF)

Figure S5 Aminoacid alignment of the RAS-domains of different *L. longbeachae* proteins identified in the genome of strain NSW150. PFAM was used to align the different sequences (<http://pfam.sanger.ac.uk/>).
Found at: doi:10.1371/journal.pgen.1000851.s005 (14.63 MB TIF)

Figure S6 Alignment of the putative LPS-encoding region of *L. longbeachae* Sg1 and Sg2 using the ARTEMIS comparison tool. Note the nearly perfect alignment of the four segments with only two regions differing between Sg1 and Sg2. Furthermore, the putative LPS-coding region of the two strains of the same Sg line perfectly up with a over 90% nucleotide identity. Specific regions and the predicted proteins encoded are depicted below.
Found at: doi:10.1371/journal.pgen.1000851.s006 (8.74 MB TIF)

Table S1 *L. longbeachae* NSW150 protein coding genes and their distribution within functional categories.
Found at: doi:10.1371/journal.pgen.1000851.s007 (0.03 MB DOC)

Table S2 Specific genes of *L. longbeachae* without orthologues in any of the four sequenced *L. pneumophila* genomes.
Found at: doi:10.1371/journal.pgen.1000851.s008 (0.50 MB DOC)

Table S3 Distribution of known and predicted Dot/Icm substrates of *L. pneumophila* in *L. longbeachae*.

Found at: doi:10.1371/journal.pgen.1000851.s009 (0.31 MB DOC)

Table S4 Putative capsule and LPS encoding genes in *L. longbeachae* and its comparison to *L. pneumophila* Paris.

Found at: doi:10.1371/journal.pgen.1000851.s010 (0.14 MB DOC)

Table S5 Analysis of the FlgD, FlcR/S, and FlhA/FlhN encoding regions in *L. longbeachae*.

Found at: doi:10.1371/journal.pgen.1000851.s011 (0.05 MB DOC)

Table S6 Transcriptional regulators identified in *L. longbeachae* and their orthologs in *L. pneumophila*.

Found at: doi:10.1371/journal.pgen.1000851.s012 (0.30 MB DOC)

Table S7 Genes upregulated in *L. longbeachae* in exponential growth phase.

Found at: doi:10.1371/journal.pgen.1000851.s013 (0.23 MB DOC)

Table S8 Genes upregulated in *L. longbeachae* in post-exponential growth phase.

References

- Yu VL, Plouffe JF, Pastoris MC, Stout JE, Schousboe M, et al. (2002) Distribution of *Legionella* species and serogroups isolated by culture in patients with sporadic community-acquired legionellosis: an international collaborative survey. *J Infect Dis* 186: 127–128.
- Phares CR, Wangroongsarb P, Chantra S, Paveenkitiporn W, Tondella ML, et al. (2007) Epidemiology of severe pneumonia caused by *Legionella longbeachae*, *Mycoplasma pneumoniae*, and *Chlamydia pneumoniae*: 1-year, population-based surveillance for severe pneumonia in Thailand. *Clin Infect Dis* 45: e147–155.
- Bibb WF, Sorg RJ, Thomason BM, Hicklin MD, Steigerwalt AG, et al. (1981) Recognition of a second serogroup of *Legionella longbeachae*. *J Clin Microbiol* 14: 674–677.
- Cameron S, Roder D, Walker C, Feldheim J (1991) Epidemiological characteristics of *Legionella* infection in South Australia: implications for disease control. *Aust N Z J Med* 21: 65–70.
- Steele TW, Moore CV, Sangster N (1990) Distribution of *Legionella longbeachae* serogroup 1 and other *Legionellae* in potting soils in Australia. *Appl Environ Microbiol* 56: 2984–2988.
- Isberg RR, O'Connor TJ, Heidtman M (2009) The *Legionella pneumophila* replication vacuole: making a cosy niche inside host cells. *Nat Rev Microbiol* 7: 13–24.
- Shin S, Roy CR (2008) Host cell processes that influence the intracellular survival of *Legionella pneumophila*. *Cell Microbiol* 10: 1209–1220.
- Nora T, Lomma M, Gomez-Valero L, Buchrieser C (2009) Molecular mimicry: an important virulence strategy employed by *Legionella pneumophila* to subvert host functions. *Future Microbiol* 4.
- Molofsky AB, Swanson MS (2004) Differentiate to thrive: lessons from the *Legionella pneumophila* life cycle. *Mol Microbiol* 53: 29–40.
- Byrne B, Swanson MS (1998) Expression of *Legionella pneumophila* virulence traits in response to growth conditions. *Infect Immun* 66: 3029–3034.
- Asare R, Abu Kwaik Y (2007) Early trafficking and intracellular replication of *Legionella longbeachae* within an ER-derived late endosome-like phagosome. *Cell Microbiol* 9: 1571–1587.
- Asare R, Santic M, Gobin I, Doric M, Suttles J, et al. (2007) Genetic susceptibility and caspase activation in mouse and human macrophages are distinct for *Legionella longbeachae* and *L. pneumophila*. *Infect Immun* 75: 1933–1945.
- Berger KH, Isberg RR (1993) Two distinct defects in intracellular growth complemented by a single genetic locus in *Legionella pneumophila*. *Mol Microbiol* 7: 7–19.
- Roy CR, Berger KH, Isberg RR (1998) *Legionella pneumophila* DotA protein is required for early phagosome trafficking decisions that occur within minutes of bacterial uptake. *Mol Microbiol* 28: 663–674.
- Segal G, Purcell M, Shuman HA (1998) Host cell killing and bacterial conjugation require overlapping sets of genes within a 22-kb region of the *Legionella pneumophila* genome. *Proc Natl Acad Sci U S A* 95: 1669–1674.
- Vogel JP, Andrews HL, Wong SK, Isberg RR (1998) Conjugative transfer by the virulence system of *Legionella pneumophila*. *Science* 279: 873–876.
- Burstein D, Zusman T, Degtyar E, Viner R, Segal G, et al. (2009) Genome-scale identification of *Legionella pneumophila* effectors using a machine learning approach. *PLoS Pathog* 5: e1000508. doi:10.1371/journal.ppat.1000508.
- Ensminger AW, Isberg RR (2009) *Legionella pneumophila* Dot/Icm translocated substrates: a sum of parts. *Curr Opin Microbiol* 12: 67–73.
- Ninio S, Roy CR (2007) Effector proteins translocated by *Legionella pneumophila*: strength in numbers. *Trends Microbiol* 15: 372–380.
- Morozova I, Qu X, Shi S, Asamani G, Greenberg JE, et al. (2004) Comparative sequence analysis of the *icm/dot* genes in *Legionella*. *Plasmid* 51: 127–147.
- Gobin I, Susa M, Begic G, Hartland EL, Doric M (2009) Experimental *Legionella longbeachae* infection in intratracheally inoculated mice. *J Med Microbiol* 58: 723–730.
- Molofsky AB, Byrne BG, Whitfield NN, Madigan CA, Fuse ET, et al. (2006) Cytosolic recognition of flagellin by mouse macrophages restricts *Legionella pneumophila* infection. *J Exp Med* 17: 1093–1104.
- Ren T, Zamboni DS, Roy CR, Dietrich WF, Vance RE (2006) Flagellin-deficient *Legionella* mutants evade caspase-1- and Naip5-mediated macrophage immunity. *PLoS Pathog* 2: e18. doi:10.1371/journal.ppat.0020018.
- Wright EK, Goodart SA, Gowney JD, Hadinoto V, Endrizzi MG, et al. (2003) Naip5 affects host susceptibility to the intracellular pathogen *Legionella pneumophila*. *Curr Biol* 13: 27–36.
- Cazalet C, Rusniok C, Bruggemann H, Zidane N, Magnier A, et al. (2004) Evidence in the *Legionella pneumophila* genome for exploitation of host cell functions and high genome plasticity. *Nat Genet* 36: 1165–1173.
- Chien M, Morozova I, Shi S, Sheng H, Chen J, et al. (2004) The genomic sequence of the accidental pathogen *Legionella pneumophila*. *Science* 305: 1966–1968.
- Steinert M, Heuner K, Buchrieser C, Albert-Weissenberger C, Glöckner G (2007) *Legionella* pathogenicity: genome structure, regulatory networks and the host cell response. *Int J Med Microbiol* 297: 577–587.
- Doyle RM, Heuzenroeder MW (2002) A mutation in an *ompR*-like gene on a *Legionella longbeachae* serogroup 1 plasmid attenuates virulence. *Int J Med Microbiol* 292: 227–239.
- Cazalet C, Jarraud S, Ghavi-Helm Y, Kunst F, Glaser P, et al. (2008) Multigenome analysis identifies a worldwide distributed epidemic *Legionella pneumophila* clone that emerged within a highly diverse species. *Genome Res* 18: 431–441.
- Cianciotto NP (2009) Many substrates and functions of type II secretion: lessons learned from *Legionella pneumophila*. *Future Microbiol* 4: 797–805.
- DeRoy S, Dao J, Soderberg M, Rossier O, Cianciotto NP (2006) *Legionella pneumophila* type II secretome reveals unique exoproteins and a chitinase that promotes bacterial persistence in the lung. *Proc Natl Acad Sci U S A* 103: 19146–19151.
- Feldman M, Segal G (2004) A specific genomic location within the *icm/dot* pathogenesis region of different *Legionella* species encodes functionally similar but nonhomologous virulence proteins. *Infect Immun* 72: 4503–4511.
- Feldman M, Zusman T, Hagag S, Segal G (2005) Coevolution between nonhomologous but functionally similar proteins and their conserved partners in the *Legionella* pathogenesis system. *Proc Natl Acad Sci U S A* 102: 12206–12211.
- Newton HJ, Sansom FM, Dao J, McAlister AD, Sloan J, et al. (2007) Sel1 repeat protein LpnE is a *Legionella pneumophila* virulence determinant that influences vacuolar trafficking. *Infect Immun* 75: 5575–5585.
- Nagai H, Cambronne ED, Kagan JC, Amor JC, Kahn RA, et al. (2005) A C-terminal translocation signal required for Dot/Icm-dependent delivery of the *Legionella* RalF protein to host cells. *Proc Natl Acad Sci U S A* 102: 826–831.

36. Kubori T, Hyakutake A, Nagai H (2008) *Legionella* translocates an E3 ubiquitin ligase that has multiple U-boxes with distinct functions. *Mol Microbiol* 67: 1307–1319.
37. Shevchuk O, Batzilla C, Hagele S, Kusch H, Engelmann S, et al. (2009) Proteomic analysis of Legionella-containing phagosomes isolated from Dictyostelium. *Int J Med Microbiol* 299: 489–508.
38. Urwyler S, Brombacher E, Hilbi H (2009) Endosomal and secretory markers of the *Legionella*-containing vacuole. *Commun Integr Biol* 2: 107–109.
39. Urwyler S, Nyfeler Y, Ragaz C, Lee H, Mueller LN, et al. (2009) Proteome analysis of *Legionella* vacuoles purified by magnetic immunoseparation reveals secretory and endosomal GTPases. *Traffic* 10: 76–87.
40. Machner MP, Isberg RR (2006) Targeting of host Rab GTPase function by the intravacuolar pathogen *Legionella pneumophila*. *Dev Cell* 11: 47–56.
41. Murata T, Delprato A, Ingmundson A, Toomre DK, Lambright DG, et al. (2006) The *Legionella pneumophila* effector protein DrrA is a Rab1 guanine nucleotide-exchange factor. *Nat Cell Biol* 8: 971–977.
42. Weber SS, Ragaz C, Hilbi H (2009) The inositol polyphosphate 5-phosphatase OCLR1 restricts intracellular growth of *Legionella*, localizes to the replicative vacuole and binds to the bacterial effector LpnE. *Cell Microbiol* 11: 442–460.
43. Weber SS, Ragaz C, Reus K, Nyfeler Y, Hilbi H (2006) *Legionella pneumophila* exploits PI(4)P to anchor secreted effector proteins to the replicative vacuole. *PLoS Pathog* 2: e46. doi:10.1371/journal.ppat.0020046.
44. Brombacher E, Urwyler S, Ragaz C, Weber SS, Kami K, et al. (2008) The Rab1 guanine nucleotide exchange factor SidM is a major PtdIns(4)P-binding effector protein of *Legionella pneumophila*. *J Biol Chem* 284: 4846–4856.
45. Pan X, Lührmann A, Satoh A, Laskowski-Arce MA, Roy CR (2008) Ankyrin repeat proteins comprise a diverse family of bacterial type IV effectors. *Science* 320: 1651–1654.
46. Eiserich JP, Estévez AG, Bamberg TV, Ye YZ, Chumley PH, et al. (1999) Microtubule dysfunction by posttranslational nitrotyrosination of alpha-tubulin: a nitric oxide-dependent mechanism of cellular injury. *Proc Natl Acad Sci U S A* 96: 6365–6370.
47. Lurin C, Andrés C, Aubourg S, Bellaoui M, Bitton F, et al. (2004) Genome-wide analysis of *Arabidopsis* pentatricopeptide repeat proteins reveals their essential role in organelle biogenesis. *Plant Cell* 16: 2089–2103.
48. Nakamura T, Schuster G, Sugiura M, Sugita M (2004) Chloroplast RNA-binding and pentatricopeptide repeat proteins. *Biochem Soc Trans* 32: 571–574.
49. Schmitz-Linneweber C, Small I (2008) Pentatricopeptide repeat proteins: a socket set for organelle gene expression. *Trends Plant Sci* 13: 663–670.
50. Belyi Y, Stahl M, Sovkova I, Kaden P, Luy B, et al. (2009) Region of elongation factor 1A1 involved in substrate recognition by *Legionella pneumophila* glucosyltransferase Lgt1: identification of Lgt1 as a retaining glucosyltransferase. *J Biol Chem* 284: 20167–20174.
51. Belyi Y, Tabakova I, Stahl M, Aktories K (2008) Lgt: a family of cytotoxic glucosyltransferases produced by *Legionella pneumophila*. *J Bacteriol* 190: 3026–3035.
52. Belyi Y, Niggeweg R, Opitz B, Vogelsang M, Hippenstiel S, et al. (2006) *Legionella pneumophila* glucosyltransferase inhibits host elongation factor 1A. *Proc Natl Acad Sci U S A* 103: 16953–16958.
53. Heidtman M, Chen EJ, Moy MY, Isberg RR (2009) Large-scale identification of *Legionella pneumophila* Dot/Icm substrates that modulate host cell vesicle trafficking pathways. *Cell Microbiol* 11: 230–248.
54. Duchaud E, Rusniok C, Frangeul L, Buchrieser C, Givaudan A, et al. (2003) The genome sequence of the entomopathogenic bacterium *Photorhabdus luminescens*. *Nat Biotechnol* 21: 1307–1313.
55. Hazelbauer GL, Falke JJ, Parkinson JS (2008) Bacterial chemoreceptors: high-performance signaling in networked arrays. *Trends Biochem Sci* 33: 9–19.
56. Krehenbrink M, Oppermann-Sanio FB, Steinbuechel A (2002) Evaluation of non-cyanobacterial genome sequences for occurrence of genes encoding proteins homologous to cyanophycin synthetase and cloning of an active cyanophycin synthetase from *Acinetobacter* sp. strain DSM 587. *Arch Microbiol* 177: 371–380.
57. Fuser G, Steinbuechel A (2007) Analysis of genome sequences for genes of cyanophycin metabolism: identifying putative cyanophycin metabolizing prokaryotes. *Macromol Biosci* 7: 278–296.
58. de Berardinis V, Durot M, Weissenbach J, Salanoubat M (2009) *Acinetobacter bayly* ADP1 as a model for metabolic system biology. *Curr Opin Microbiol* 12: 568–576.
59. Diez E, Lee SH, Gauthier S, Yaraghi Z, Tremblay M, et al. (2003) Birc1c is the gene within the Lgn1 locus associated with resistance to *Legionella pneumophila*. *Nat Genet* 33: 55–60.
60. Lamkanfi M, Amer A, Kanneganti TD, Munoz-Planillo R, Chen G, et al. (2007) The Nod-like receptor family member Naip5/Birc1c restricts *Legionella pneumophila* growth independently of caspase-1 activation. *J Immunol* 178: 8022–8027.
61. Lightfield KL, Persson J, Brubaker SW, Witte CE, von Moltke J, et al. (2008) Critical function for Naip5 in inflammasome activation by a conserved carboxy-terminal domain of flagellin. *Nat Immunol* 9: 1171–1178.
62. Zamboni DS, Kobayashi KS, Kohlsdorf T, Ogura Y, Long EM, et al. (2006) The Birc1c cytosolic pattern-recognition receptor contributes to the detection and control of *Legionella pneumophila* infection. *Nat Immunol* 7: 318–325.
63. Kentner D, Sourjik V (2006) Spatial organization of the bacterial chemotaxis system. *Curr Opin Microbiol* 9: 619–624.
64. La Scola B, Birtles RJ, Greub G, Harrison TJ, Ratcliff RM, et al. (2004) *Legionella drancourtii* sp. nov., a strictly intracellular amoebal pathogen. *Int J Syst Evol Microbiol* 54: 699–703.
65. Brüggemann H, Hagman A, Jules M, Sismeiro O, Dillies M, et al. (2006) Virulence strategies for infecting phagocytes deduced from the *in vivo* transcriptional program of *Legionella pneumophila*. *Cell Microbiol* 8: 1228–1240.
66. Sahr T, Brüggemann H, Jules M, Lomma M, Albert-Weissenberger C, et al. (2009) Two small ncRNAs jointly govern virulence and transmission in *Legionella pneumophila*. *Mol Microbiol* 72: 741–762.
67. Tiaden A, Spirig T, Carranza P, Brüggemann H, Riedel K, et al. (2008) Synergistic contribution of the *Legionella pneumophila* lqs genes to pathogen-host interactions. *J Bacteriol* 190: 7532–7547.
68. Spirig T, Tiaden A, Kiefer P, Buchrieser C, Vorholt JA, et al. (2008) The *Legionella* autoinducer synthase LqsA produces an alpha-hydroxyketone signaling molecule. *J Biol Chem* 283: 18113–18123.
69. Tiaden A, Spirig T, Weber SS, Brüggemann H, Bosshard R, et al. (2007) The *Legionella pneumophila* response regulator LqsR promotes virulence as an element of the regulatory network controlled by RpoS and LetA. *Cell Microbiol* 9: 2903–2920.
70. Skerker JM, Prasol MS, Perchuk BS, Biondi EG, Laub MT (2005) Two-component signal transduction pathways regulating growth and cell cycle progression in a bacterium: a system-level analysis. *PLoS Biol* 3: e334. doi:10.1371/journal.pbio.0030334.
71. Newton HJ, Sansom FM, Dao J, Cazalet C, Brüggemann H, et al. (2008) Significant Role for ladC in Initiation of *Legionella pneumophila* Infection. *Infect Immun* 76: 3075–3085.
72. Lory S, Wolfgang M, Lee V, Smith R (2004) The multi-talented bacterial adenylate cyclases. *Int J Med Microbiol* 293: 479–482.
73. Rasis M, Segal G (2009) The Lta-RsmYZ-CsrA regulatory cascade, together with RpoS and PmrA, post-transcriptionally regulates stationary phase activation of *Legionella pneumophila* Icm/Dot effectors. *Mol Microbiol* 72: 995–1010.
74. Gal-Mor O, Segal G (2003) Identification of CpxR as a positive regulator of icm and dot virulence genes of *Legionella pneumophila*. *J Bacteriol* 185: 4908–4919.
75. Zusman T, Aloni G, Halperin E, Kotzer H, Degtyar E, et al. (2007) The response regulator PmrA is a major regulator of the icm/dot type IV secretion system in *Legionella pneumophila* and *Coxiella burnetii*. *Mol Microbiol* 63: 1508–1523.
76. Altman E, Segal G (2008) The response regulator CpxR directly regulates expression of several *Legionella pneumophila* icm/dot components as well as new translocated substrates. *J Bacteriol* 190: 1985–1996.
77. Alto NM (2008) Mimicking small G-proteins: an emerging theme from the bacterial virulence arsenal. *Cell Microbiol* 10: 566–575.
78. Nagai H, Kagan JC, Zhu X, Kahn RA, Roy CR (2002) A bacterial guanine nucleotide exchange factor activates ARF on *Legionella* phagosomes. *Science* 295: 679–682.
79. de Felipe KS, Pampou S, Jovanovic OS, Pericone CD, Ye SF, et al. (2005) Evidence for acquisition of *Legionella* type IV secretion substrates via interdomain horizontal gene transfer. *J Bacteriol* 187: 7716–7726.
80. Brüggemann H, Cazalet C, Buchrieser C (2006) Adaptation of *Legionella pneumophila* to the host environment: role of protein secretion, effectors and eukaryotic-like proteins. *Curr Opin Microbiol* 9: 86–94. Epub 2006 Jan 2006.
81. Glaser P, Frangeul L, Buchrieser C, Rusniok C, Amend A, et al. (2001) Comparative genomics of *Listeria* species. *Science* 294: 849–852.
82. Buchrieser C, Rusniok C, Frangeul L, Couve E, Billault A, et al. (1999) The 102-kilobase pgm locus of *Yersinia pestis*: sequence analysis and comparison of selected regions among different *Yersinia pestis* and *Yersinia pseudotuberculosis* strains. *Infect Immun* 67: 4851–4861.
83. Frangeul L, Glaser P, Rusniok C, Buchrieser C, Duchaud E, et al. (2004) CAAT-Box, Contigs-Assembly and Annotation tool-box for genome sequencing projects. *Bioinformatics* 20: 790–797.
84. Vallenet D, Labarre L, Rouy Z, Barbe V, Bocs S, et al. (2006) MaGe: a microbial genome annotation system supported by synteny results. *Nucleic Acids Res* 34: 53–65.
85. Zerbino DR, Birney E (2008) Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 18: 821–829.
86. Darling AC, Mau B, Blattner FR, Perna NT (2004) Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res* 14: 1394–1403.
87. Li H, Ruan J, Durbin R (2008) Mapping short DNA sequencing reads and calling variants using mapping quality scores. *Genome Res* 18: 1851–1858.
88. Milohanic E, Glaser P, Coppee JY, Frangeul L, Vega Y, et al. (2003) Transcriptome analysis of *Listeria monocytogenes* identifies three groups of genes differently regulated by PrfA. *Mol Microbiol* 47: 1613–1625.
89. Yang YH, Dudoit S, Luu P, Lin DM, Peng V, et al. (2002) Normalization for cDNA microarray data: a robust composite method addressing single and multiple slide systematic variation. *Nucleic Acids Res* 30: e15.
90. Delmar P, Robin S, Daudin JJ (2005) VarMixt: efficient variance modelling for the differential analysis of replicated gene expression data. *Bioinformatics* 21: 502–508.
91. Reiner A, Yekutieli D, Benjamini Y (2003) Identifying differentially expressed genes using false discovery rate controlling procedures. *Bioinformatics* 19: 368–375.
92. Chain PS, Grahnam DV, Fulton RS, Fitzgerald MG, Hostetler J, et al. (2009) Genomics. Genome project standards in a new era of sequencing. *Science* 326: 236–237.