

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Doolan, BJ;Vu, M;Varigos, GA;Hogan, C;Ross, G;Sood, S;Winship, I

Title:

Warfarin ineffective as symptomatic therapy for erythropoietic protoporphyria

Date:

2019-11-01

Citation:

Doolan, B. J., Vu, M., Varigos, G. A., Hogan, C., Ross, G., Sood, S. & Winship, I. (2019). Warfarin ineffective as symptomatic therapy for erythropoietic protoporphyria. *Australasian Journal of Dermatology*, 60 (4), pp.e360-e361. <https://doi.org/10.1111/ajd.13033>.

Persistent Link:

<https://hdl.handle.net/11343/285647>

Manuscript Title Page:

Article Title:

Warfarin ineffective as symptomatic therapy for erythropoietic protoporphyria

Brent J Doolan¹, Mi Vu¹, George A Varigos^{1,2}, Chris Hogan³, Gayle Ross¹, Siddharth Sood^{2,4}, Ingrid Winship^{2,5}

1. Department of Dermatology, The Royal Melbourne Hospital, Melbourne, Victoria, Australia

2. Department of Medicine, The University of Melbourne, Melbourne, Victoria, Australia

3. Department of Haematology, The Royal Melbourne Hospital, Melbourne, Victoria, Australia

4. Department of Gastroenterology and Hepatology, The Royal Melbourne Hospital, Melbourne, Victoria, Australia

5. Department of Genetic Medicine, The Royal Melbourne Hospital, Melbourne, Victoria, Australia

Author Information: (Correspondence via Brent Doolan)

Brent J Doolan, BSc MBBS MPH&TM

Department of Dermatology

The Royal Melbourne Hospital, Melbourne, Victoria, Australia 3050

Ph: +61 3 9342 4531

E: brentdoolanis@gmail.com

Mi Vu, BPharm MBBS

Department of Dermatology

The Royal Melbourne Hospital, Melbourne, Victoria, Australia 3050

Ph: +61 3 9342 4531

E: hoa-mi.vu@mh.org.au

George A Varigos, MBBS FACD

Department of Dermatology

The Royal Melbourne Hospital, Melbourne, Victoria, Australia 3050

Department of Medicine

The University of Melbourne, Melbourne, Victoria, Australia 3010

Ph: +61 3 9342 4531

E: George.varigos@mh.org.au

Chris Hogan, MBBS FRACP

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as [doi: 10.1111/AJD.13033](https://doi.org/10.1111/AJD.13033)

This article is protected by copyright. All rights reserved

Department of Haematology

The Royal Melbourne Hospital, Melbourne, Victoria, Australia 3050

Ph: +61 3 9342 7695

E: chris.hogan@mh.org.au

Gayle Ross, MBBS FACD

Department of Dermatology

The Royal Melbourne Hospital, Melbourne, Victoria, Australia 3050

Ph: +61 3 9342 4531

E: skindepthderm@gmail.com

Siddarth Sood, MBBS FRACP

Department of Gastroenterology and Hepatology

The Royal Melbourne Hospital, Melbourne, Victoria, Australia 3050

Department of Medicine

The University of Melbourne, Melbourne, Victoria, Australia 3010

Ph: +61 3 9342 7470

E: Siddharth.Sood@mh.org.au

Ingrid Winship, MBBS PhD FACD FRACP

Department of Genetic Medicine

The Royal Melbourne Hospital, Melbourne, Victoria, Australia 3050

Department of Medicine

The University of Melbourne, Melbourne, Victoria, Australia 3010

Ph: +61 3 9342 7151

E: ingrid.winship@mh.org.au

Word count: 600 (Excluding references and table)

Tables: 1

Figures: 0

Running Title: Warfarin ineffective for erythropoietic protoporphyria

Financial and Material Support: None identified

Acknowledgements Section:

Professor Sanjoy Paul (Melbourne EpiCentre) has kindly provided statistical analysis of data. Dr William Cranwell, Dr Virginia Pilcher and Annette Phemister were involved in the recruitment and assessment of patients.

Conflicts of Interest/Disclosures:

All authors declare that they have no financial, personal or other conflicts of interest in relation or affiliations to this research and its publication that could influence the author's decisions, work or manuscript. There was no funding supplied from any source.

Previous Presentations:

This study was presented at the 2nd Annual Australasian Porphyria Conference at The Royal Children's Hospital, Melbourne, Australia; November 23, 2018.

DR. BRENT J DOOLAN (Orcid ID : 0000-0002-9497-0504)

DR. MI VU (Orcid ID : 0000-0002-4093-4567)

Article type : Research Letter

Warfarin ineffective as symptomatic therapy for erythropoietic protoporphyria

Autosomal recessive erythropoietic protoporphyria (EPP) is an inherited cutaneous porphyria, with an incidence of approximately 1:200,000 people.¹ It arises most commonly due to a mutation in the *FECH* gene, causing reduced activity of the ferrochelatase enzyme that catalyses the insertion of iron into protoporphyrin IX (PPIX) to form haem.¹ As a result, PPIX accumulates within the erythrocytes, liver and skin. Retention within the skin causes acute, profoundly debilitating photosensitivity when exposed to sunlight.¹ Conventional treatments have been largely ineffective, with sun avoidance the main management, resulting in restriction of daily activities and poor quality of life.¹ Alfamelanotide (human α -melanocyte-stimulating hormone) implants have been approved in some European countries as a possible treatment option for EPP with minor side effects, but currently this is not approved for use in Australia.² Management also seeks to prevent the progression of possible liver disease to liver failure.

We originally reported a case study in a patient who had complete resolution of EPP symptoms, with administration of therapeutic warfarin following aortic valve replacement.³ His symptoms recurred after ceasing warfarin prior to an elective surgical procedure, and resolved upon recommencement. Given this outcome, we sought to evaluate the efficacy and safety of warfarin therapy in providing symptomatic relief to individuals with EPP.

Australians with EPP (*FECH* gene mutation) were recruited from the multi-disciplinary porphyria service at The Royal Melbourne Hospital. Participants were assessed for warfarin sensitivity by screening for; CYP2C9 gene polymorphisms (known to affect plasma levels of warfarin and its metabolites) and VKORC1 gene polymorphisms (known to influence the anticoagulant effect). A single-blinded, crossover study was applied. Participants were administered daily doses of warfarin, with change to the fixed dose every 3-months for a total of 12 months. Participants were initially commenced on 1mg of warfarin, then 0mg, then 0.5mg, with a 1-week washout period in between change of dose. A further open-label period, aiming for therapeutic international normalised ratio (INR 1.5-2.5) was also undertaken. Participants were seen monthly and assessed for sun tolerability, pain and completed a daily sun diary, dermatology quality of life questionnaire (DLQI) and a validated EPP symptomatic scoring questionnaire, adapted from Biolcati *et al.*⁴ Monthly laboratory measures included coagulation analysis and red blood cell (RBC) porphyrin and total plasma porphyrin levels.

Five participants were recruited (Table 1); with three intermediate metabolisers of warfarin. Primary endpoints were analysed based on week 12 results (1mg warfarin dosing) and results of open-label therapeutic warfarin dosing. Two participants (normal warfarin metabolisers) entered into the open-label period with a maximum INR of 1.7 and 2.2. A comparison was undertaken between outcome measures at baseline (0mg of warfarin) and maximum dose of warfarin administered for each participant (Table 2). Mean compliance of daily medication administration for all participants was $94.6 \pm 3.2\%$ (SD). There was no associated improvement in EPP symptoms from analysis of quality of life questionnaires. Biochemical RBC and total plasma porphyrin levels showed no improvement in either low dose or therapeutic doses of warfarin.

We were limited in the capacity of sunlight exposure throughout phases of this study, where seasonal changes (winter) resulted in decreased visible light wavelengths and sunlight contact with skin, which is important in the pathogenesis of the skin related EPP symptoms. We attempted to minimise this effect, by instigating the open-label extension period during the intense sunlight of the Australian summer months. Limitations also were noted in

relation to poor compliance of the daily skin diaries, with mean daily compliance of $21 \pm 4.8\%$ (SD). The intermediate warfarin metaboliser participants elected not to enrol into the extension study, due to a perceived lack of efficacy during the blinded period. It remains unknown whether these participants may have benefited from therapeutic doses, as the patient we described in the initial case report was also an intermediate metaboliser of warfarin. In the extension study, we were able to obtain safe INR values of 1.9, which may have been lower than the therapeutic dose that allowed resolution of symptoms in the original case report, which aimed for a target INR of 2.5.³

Whilst our original case study suggested an antioxidant and anti-inflammatory effect of warfarin in mitigating the symptoms of EPP, this pilot study was not able to replicate this. Given the paucity of effective treatments, further investigation into warfarin and other antioxidant agents should still be considered, especially given the outcome of the initial case study.

Word Count: 726

The study was approved by The Melbourne Health Human Research Ethics Committee (HREC/15/MH/231) and conforms to the recognised standards according to the Declaration of Helsinki.

References:

1. Bissell DM, Anderson KE, Bonkovsky HL. Porphyrria. *N. Engl. J. Med.* 2017; 377(9): 862-72. doi: 10.1056/NEJMra1608634.
2. Langendonk JG, Balwani M, Anderson KE et al. Afamelanotide for erythropoietic protoporphyria. *N Eng J Med.* 2015 Jul 2;373(1):48-59. doi: 10.1056/NEJMoa1411481
3. Winship I, Ross G, Nicoll A, et al. Antioxidant effect of warfarin therapy: a possible symptomatic treatment for erythropoietic protoporphyria. *Arch. Dermatol.* 2009; 145(8): 960-61. doi: 10.1001/archdermatol.2009.165

4. Biolcati G, Marchesini E, Sorge F, et al. Long-term observational study of afamelanotide in 115 patients with erythropoietic protoporphyria. *Br. J. Dermatol.* 2015; 172(6): 1601-12. doi: 10.1111/bjd.13598

Tables:

Table 1: Erythropoietic protoporphyria patient characteristics					
Characteristic	P01	P02	P03	P04	P06*
Sex	Male	Female	Female	Female	Male
Age (years)	41.8	51.3	39.3	24.2	47.6
Body Mass Index (kg/m ²)	27.1	23.9	33.8	25.1	30.2
Years since diagnosis	38	47	26	14	42
Photosensitivity?	Y	Y	Y	Y	Y
Intermediate metaboliser to Warfarin?	Y°	N	N	N	Y°
Period on study (weeks)	10	72	51	65	25
*Participant P05 withdrew before commencement of study; Y = Yes; N = No;					
° Intermediate metaboliser to both CYP2C9 and VKORC1 gene polymorphisms					

Table 2: Individual results of erythropoietic protoporphyria patients comparing outcomes measures of placebo to maximum dosing periods

Characteristic	P01	P02	P03	P04	P06*
Baseline period (0mg warfarin)					
Season	Summer	Summer	Summer	Summer	Autumn
INR	1.0	1.0	0.9	0.9	1.0
DLQI Score	9	4	3	9	17
EPP QoL Score	26	25	12	27	23
RBC Porphyrin ($\mu\text{mol/L}$)	69.9	43	41.6	12.8	79
Total Plasma Porphyrin (mg/L)	891	634	698	324	1062
Maximum warfarin dose					
Season	Winter	Summer	Autumn	Summer	Autumn
Warfarin dose (mg)	1.0	8.0	1.0	6.0	1.0
INR	1.3	1.7	1.1	2.2	1.1
DLQI Score	6	1	5	2	8
EPP QoL Score	29	29	14	18	22
RBC Porphyrin ($\mu\text{mol/L}$)	75.6	37.5	64.0	12.9	82
Total Plasma Porphyrin (mg/L)	702	262	680	230	522
*P05 withdrew before study commencement; INR = International Normalised Ratio; DLQI = Dermatology Life Quality Index; EPP QoL = Erythropoietic Protoporphyrin Quality of Life; RBC = Red Blood Cell					