



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Liu, ZSJ;Sattabongkot, J;White, M;Chotirat, S;Kumpitak, C;Takashima, E;Harbers, M;Tham, WH;Healer, J;Chitnis, CE;Tsuboi, T;Mueller, I;Longley, RJ

Title:

Naturally acquired antibody kinetics against Plasmodium vivax antigens in people from a low malaria transmission region in western Thailand

Date:

2022-12-01

Citation:

Liu, Z. S. J., Sattabongkot, J., White, M., Chotirat, S., Kumpitak, C., Takashima, E., Harbers, M., Tham, W. H., Healer, J., Chitnis, C. E., Tsuboi, T., Mueller, I. & Longley, R. J. (2022). Naturally acquired antibody kinetics against Plasmodium vivax antigens in people from a low malaria transmission region in western Thailand. BMC Medicine, 20 (1), <https://doi.org/10.1186/s12916-022-02281-9>.

Persistent Link:

<https://hdl.handle.net/11343/306313>

License:

[CC BY](#)

RESEARCH ARTICLE

Open Access



Naturally acquired antibody kinetics against *Plasmodium vivax* antigens in people from a low malaria transmission region in western Thailand

Zoe Shih-Jung Liu^{1,2,3}, Jetsumon Sattabongkot⁴, Michael White⁵, Sadudee Chotirat⁴, Chalermpon Kumpitak⁴, Eizo Takashima⁶, Matthias Harbers⁷, Wai-Hong Tham^{1,2}, Julie Healer^{1,2}, Chetan E. Chitnis⁸, Takafumi Tsuboi⁶, Ivo Mueller^{1,2†} and Rhea J. Longley^{1,2*†}

Abstract

Background: *Plasmodium vivax* (*P. vivax*) is the dominant *Plasmodium* spp. causing the disease malaria in low-transmission regions outside of Africa. These regions often feature high proportions of asymptomatic patients with sub-microscopic parasitaemia and relapses. Naturally acquired antibody responses are induced after *Plasmodium* infection, providing partial protection against high parasitaemia and clinical episodes. However, previous work has failed to address the presence and maintenance of such antibody responses to *P. vivax* particularly in low-transmission regions.

Methods: We followed 34 patients in western Thailand after symptomatic *P. vivax* infections to monitor antibody kinetics over 9 months, during which no recurrent infections occurred. We assessed total IgG, IgG subclass and IgM levels to up to 52 *P. vivax* proteins every 2–4 weeks using a multiplexed Luminex[®] assay and identified protein-specific variation in antibody longevity. Mathematical modelling was used to generate the estimated half-life of antibodies, long-, and short-lived antibody-secreting cells.

Results: Generally, an increase in antibody level was observed within 1-week post symptomatic infection, followed by an exponential decay of different rates. We observed mostly IgG1 dominance and IgG3 sub-dominance in this population. IgM responses followed similar kinetic patterns to IgG, with some proteins unexpectedly inducing long-lived IgM responses. We also monitored antibody responses against 27 IgG-immunogenic antigens in 30 asymptomatic individuals from a similar region. Our results demonstrate that most antigens induced robust and long-lived total IgG responses following asymptomatic infections in the absence of (detected) boosting infections.

Conclusions: Our work provides new insights into the development and maintenance of naturally acquired immunity to *P. vivax* and will guide the potential use of serology to indicate immune status and/or identify populations at risk.

Background

Malaria is an infectious disease caused by parasites under the genus of *Plasmodium* and is transmitted via female *Anopheles* mosquitoes. *Plasmodium falciparum* (*P. falciparum*) and *Plasmodium vivax* (*P. vivax*) are the two most common *Plasmodium* parasites that infect humans,

*Correspondence: Longley.r@wehi.edu.au

†Ivo Mueller and Rhea J Longley are joint senior authors.

¹ The Walter and Eliza Hall Institute of Medical Research, Parkville, Victoria 3052, Australia

Full list of author information is available at the end of the article



causing significant morbidity and mortality worldwide. In particular, *P. vivax* is the most widely distributed *Plasmodium* species outside sub-Saharan Africa, contributing to more than 4.5 million cases in 2020. In southeast Asia, *P. vivax* accounts for 39% of the region's malaria burden [1]. The capacity of *P. vivax* to cause negative impacts on human health and, in some cases, death, is significant, especially for children under 5 years of age [2, 3].

P. vivax has several distinct features in its life cycle which make this species of *Plasmodium* a major challenge for elimination. The parasites are injected into the blood stream by a female *Anopheles* mosquito and quickly travel to the liver for maturation [4]. An unknown proportion of the liver-stage parasites remain in the hepatocytes and enter dormancy for months up to years until they are reactivated (by currently unknown signals), re-initiating the blood-stage symptoms [5]. These arrested liver-stage parasites are known as hypnozoites, and the delayed blood-stage infections they cause are manifesting relapses of the disease [6, 7]. Relapses can contribute up to 80% of all blood-stage infections [4, 5], but no current diagnostic tools can detect individuals who have hypnozoites in their livers but no current blood-stage parasites [8]. Primaquine and tafenoquine are available to clear hypnozoites [9], but their use is contraindicated in individuals with G6PD deficiency [10–12]. Additionally, a high proportion of low-density asymptomatic *P. vivax* infections have been reported in multiple southeast Asian regions [13–15]. There is increasing evidence for other hidden reservoirs of *P. vivax* parasites, including in the spleen [16, 17] and bone marrow [18, 19], which may indeed contribute to the low-density peripheral blood-stage infections of *P. vivax*. These low-density asymptomatic individuals are often missed during screening and do not present to medical facilities, further contributing to ongoing transmission. New tools specifically targeting *P. vivax* are needed, particularly to identify hypnozoite carriers, or ideally, prevent hypnozoites from forming in the first place. An increased understanding of naturally acquired immune responses to the parasite could result in innovative approaches to both these challenges, including sero-surveillance [20] and the identification of novel vaccine candidates, along with the design of vaccines that induce long-lived immunity [21].

Naturally acquired immunity is elicited upon natural exposure to pathogens and can provide significant insights into the immune status of populations [22]. In the case of infectious diseases, naturally acquired immunity is often correlated with partial protection against clinical symptoms or associated with past levels of exposure to the pathogen [23, 24]. In the context of *P. vivax* malaria, studies of naturally acquired immunity have provided novel insights such as the identification

of targets of immunity for development of vaccines and sero-diagnostics [25]. Whilst the acquisition of naturally acquired total IgG responses against *P. vivax* has been relatively well studied, there have been only limited studies assessing longevity of these responses. Understanding longevity is important for both applying sero-surveillance tools that have an actionable outcome (i.e. infer on recent not past infections) and developing vaccines that induce durable immunity. It has generally been assumed that *Plasmodium*-specific antibody responses are short-lived in the absence of new boosting infections, as has been described for *P. falciparum* [23]. However, multiple studies have now reported the presence of long-lived *P. vivax*-specific antibody responses, even in the absence of ongoing exposure [26–29]. Additionally, there is a lack of findings on the responses of IgG subclasses in low-transmission regions compared to those with higher transmission [30, 31]. Most of these studies have focused on only a limited number of *P. vivax* antigens and thus have not assessed factors contributing to antibody longevity, with the exception of our work demonstrating that *P. vivax*-specific IgG responses were protein-specific and longer-lived in individuals from higher transmission regions [26].

To further our understandings of naturally acquired immunity and to explore its potential translational implications, we studied the acquisition and maintenance of antibody responses against a panel of 52 *P. vivax* antigens following both symptomatic and asymptomatic *P. vivax* infections in low-transmission regions in Thailand [27, 32]. Factors potentially influencing the persistence of antibody responses were also investigated. These included the presence of signal peptides (SP) [33, 34], *Plasmodium* exported element (PEXEL) motifs [35, 36], transmembrane domains (TMDs) [37], glycosylphosphatidylinositol (GPI) anchors [38, 39], predicted domains [40–42] and low complexity regions (LCRs) [43–45]. These insights will be critical for applying recently developed sero-surveillance markers [46], and testing the inference that they can identify individuals with recent *P. vivax* infections, as well as improving our understanding of what drives a long-lived antibody response and how this could be exploited for selection of candidate antigens or improving vaccine design.

Methods

Study populations

Thirty-four symptomatic patients of all ages (see Table 1) with confirmed *P. vivax* infection were enrolled in a 9-month longitudinal study in Tha Song Yang, Tak Province, Thailand, as previously described [46]. These individuals were a sub-set from a larger study [27] and selected due to the absence of recurrent *Plasmodium*

Table 1 Demographic characteristics of the two study populations

| Cohort | Thai symptomatics (n = 34) | Thai asymptomatics (n = 30) |
|--|---|---|
| Symptoms | Symptomatic | Asymptomatic |
| Age (years) ^a | 29 (7–71) | 26.5 (4–72) ^b |
| Proportion male | 58.8% | 63.3% ^b |
| Self-reported past malaria exposure (yes/no) | 52.9% had past exposure | 56.7% had past exposure |
| Study duration | 9 months | 14 months |
| Time points with antibody measurements | 17 | 10–14 |
| <i>P. vivax</i> infection ^c | Enrolment (w0) | Enrolment—6 months |
| Recurrent infection | No | No |
| Treatment | Chloroquine (3 days) & Primaquine (14 days) | N/A |
| Parasite density at enrolment (copies/mL) ^{a,c} | 2.5×10^5 (2.85×10^4 – 6.84×10^6) | 1.03×10^2 (2.44×10^{-1} – 9.58×10^5) |
| <i>P. vivax</i> infections remained detectable (months) ^{c,d} | N/A | 1 (1–4) |

^a Data are shown as median and range

^b No age or gender data available for four asymptomatic individuals

^c Indicated by positive *P. vivax*-specific qPCR or PCR results

^d Data are shown as median and range

infections during follow-up. Treatment was administered according to Thai national guidelines upon confirmation of diagnosis. This included both chloroquine (25 mg base/kg body weight over 3 days) and primaquine (15 mg daily over 14 days) under directly observed therapy [27]. Blood samples were collected via finger prick at 17 time points post-infection and treatment: time of infection (week 0), week 1, week 2, then every fortnight for 6 months, then every month until the 9th month. Light microscopy and quantitative *P. vivax*-specific PCR were performed on all samples to confirm the absence of recurrent *Plasmodium* infections during the follow-up.

To compare antibody kinetic profiles with the symptomatic individuals, 30 individuals with asymptomatic *P. vivax* infections were selected from a larger longitudinal cohort study conducted in the Kanchanaburi and Ratchaburi provinces of western Thailand in 2013–2014 [32]. As these individuals had asymptomatic infections, they were not treated, as this was not an indication for treatment at the time of the study (and given PCRs were performed at a later time in the laboratory in Bangkok) [32]. Plasma samples were collected via finger prick at enrolment and approximately every month for a year (a total of 14 visits). The 30 individuals (see Table 1) were detected as *P. vivax*-positive by PCR at early time points (between enrolment and 6 months) of this yearlong study, and each volunteer had at least 10 samples available post-infection. The infections were classified as asymptomatic as the individuals had no fever (<37.5°C) and no self-reported malaria symptoms at the time of infection. There was no difference in age between the individuals in the symptomatic and asymptomatic cohorts (*p* value > 0.99 by Mann-Whitney *U* test).

Plasma samples of 274 malaria-free individuals from the Australian Red Cross (ARC, *n* = 100) in Melbourne, Australia, the Thai Red Cross (TRC, *n* = 72) in Bangkok, Thailand, and the Volunteer Blood Donor Registry (VBDR, *n* = 102) in Melbourne, Australia, were collected as previously described [46].

Antigen selection and expression

Fifty-two *P. vivax* antigens known to be immunogenic in low-transmission regions such as Thailand and Brazil were selected for this study, based on our previous results [26, 46]. Thirty-nine antigens were expressed using a wheat germ cell-free (WGCF) system (CellFree Sciences, Matsuyama, Japan), whilst 13 were expressed using *E. coli*. This panel of antigens included both those that can act as markers of recent exposure [46] and those currently under consideration as vaccine candidates [4, 47–51]. All antigens were expressed and purified as previously described in detail [46]. See Additional File 2: Table S2 for the full list of 52 *P. vivax* proteins.

Antigen conjugation

The conjugation of purified *P. vivax* antigens to non-magnetic COOH microspheres (Bio-Rad Laboratories, Inc., California, USA) was performed as described [46]. The concentration of each antigen used for conjugation was optimised to achieve a standard curve with log-linearity and is recorded in Additional File 2: Table S2. In brief, 2.5×10^6 microspheres with unique internal colours were activated before optimal concentration of an antigen was added and incubated either overnight at 4°C or for 2 h at room temperature in the dark before excess antigen were

removed. The standard curve was generated using a positive control plasma pool, as detailed below.

Total IgG antibody measurements

Total IgG antibody measurements were performed using a multiplexed bead-based assay as previously described [46]. Briefly, 500 antigen-conjugated microspheres per well were added into a MultiScreen Solvinert 96 Well Filter Plate (Merk Millipore) and incubated at room temperature with plasma samples diluted in a 1:100 ratio. Antigen-coupled microspheres were then incubated with a PE-conjugated anti-human secondary antibody at 1:100 dilution (donkey F (ab')₂ anti-human IgG Fc, 1 mg/ml, Jackson ImmunoResearch Laboratories, Inc.). For the clinical Tak cohort, IgG levels at all 17 time points were measured. For the asymptomatic Kanchanaburi cohort, IgG levels were measured at time of infection and at a minimum of 10 later time points. Plasma samples from hyper-immune individuals in Papua New Guinea (PNG) were pooled and used as the positive control from which 2-fold serially diluted standard curves were established, as described [46].

The assays were performed on either a Bio-Plex 200[®] (Bio-Rad Laboratories, Inc.) or Luminex[®] 200[™] (Luminex[®] Corporation, Texas, USA) machine as previously described [52]. Median fluorescence intensity (MFI) values were converted into arbitrary relative antibody unit (RAU) based on the antigen-specific standard curves on each plate, to adjust for any plate-plate variations. The conversion was performed using a 5-parameter logistic regression model [46].

IgG subclass and IgM antibody measurements


IgM antibody measurements were performed at all 17 time points in the symptomatic cohort, and positivity was determined at the peak of response (1-week post-infection). The same multiplexed assay was used as described above for IgG, with the following modifications: samples were diluted at 1:200 and donkey F (ab')₂ anti-human IgM Fc_{5m} at 1:400 dilution (1 mg/ml, Jackson ImmunoResearch Laboratories, Inc.) was used as the secondary antibody for detection.

To investigate the IgG subclass profile in the symptomatic cohort, measurements were performed at 7 time points: w0, w2, w4, and then every 8 weeks until 9 months. Week 2, rather than peak responses at week 1, was selected due to plasma availability. Again, the multiplexed assay was used, with the following modifications. To maximise signals and minimise non-specific background, samples were diluted at 1:50 and antigen-coupled microspheres were incubated with secondary antibody diluted at 1:100 each: mouse anti-human IgG1 hinge (0.1 mg/ mL, clone 4E3, SouthernBiotech); IgG2 Fc (0.1 mg/ mL, clone HP6002, SouthernBiotech); IgG3 hinge (0.1 mg/ mL clone HP6050, SouthernBiotech); IgG4 Fc (0.1 mg/ mL, clone HP6025, SouthernBiotech).

The study design and antibody analyses of the symptomatic and asymptomatic longitudinal cohorts is shown and described in Fig. 1.

Antigen feature selection

To explore the potential factors influencing estimated antibody half-lives, 6 antigen features were selected for



analysis: signal peptides, transmembrane domains, GPI anchors, PEXEL motifs, predicted domains and low-complexity regions. The presence of these features in a sequence was as deposited in PlasmoDB [53] (<https://plasmodb.org/plasmo/app>) or as predicted using the online servers: Transmembrane Helices Hidden Markov Model (TMHMM) Server version 2.0 [54] (<http://www.cbs.dtu.dk/services/TMHMM/>), PredGPI [55] (<http://gpcr.biocomp.unibo.it/predgpi/pred.htm>), InterPro [56] (<https://www.ebi.ac.uk/interpro/>), and SEG [57].

Mathematical modelling

Following infection, we assume that the proliferation and differentiation of B cells leads to a boost in antibody secreting cells (ASCs) of size β , commencing at a time δ before the first sample. A proportion ρ of these ASCs are assumed to be short-lived with half-life d_s , with a proportion $1 - \rho$ being long-lived with half-life d_l . It is assumed that all ASCs secrete IgG molecules which decay with a half-life d_a . The antibody level of an individual at time t after first sample is given by:

$$Ab_j(t) = Ab_0 e^{-r_l t} + \beta_j \left(\rho_j \frac{e^{-r_s(t-\delta)} - e^{-r_a(t-\delta)}}{r_a - r_s} + (1 - \rho_j) \frac{e^{-r_l(t-\delta)} - e^{-r_a(t-\delta)}}{r_a - r_l} \right)$$

where $r_a = \log(2)/d_a$ is the rate of decay of IgG molecules, $r_s = \log(2)/d_s$ is the rate of decay of short-lived plasma B cells, and $r_l = \log(2)/d_l$ is the rate of decay of long-lived plasma B cells. It is assumed that the measured antibody level was Ab_0 which decays exponentially at rate r_l .

The model was fitted to longitudinal antibody level measurements from all participants. Mixed effects methods were used to capture the natural variation in antibody kinetics between individual participants, whilst estimating the average value and variance of the immune parameters across the entire population of individuals. The models were fitted in a Bayesian framework using Markov chain Monte Carlo (MCMC) methods. Mixed effects methods allow individual-level parameters to be estimated for each participant separately, with these individual-level (or mixed effects) parameters being drawn from global distributions. For example, for each participant n the half-life of the short-lived ASCs may be estimated as d_s^n (an individual-level parameter). These N estimates of the local parameters d_s^n will be drawn from a probability distribution. A log-Normal distribution is suitable as it has positive support on $[0, \infty)$. Thus, we have $\log(d_s^n) \sim N(\mu_s, \sigma_s^2)$. The mean d_s and the variance Σ_s^2 of the estimates of d_s^n are given by $d_s = e^{\mu_s + \frac{\sigma_s^2}{2}}$ and $\Sigma_s^2 = (e^{\sigma_s^2} - 1)e^{2\mu_s + \sigma_s^2}$. The relationship between the parameters describing the population-level distribution

and the parameters for each individual in the population are depicted in the schematic diagram in Additional File 1: Fig. S1.

The model likelihood was calculated via the following process. For individual n we have data on observed antibody levels $A^n = \{a_1, \dots, a_K\}$ at times $T^n = \{t_1, \dots, t_K\}$. We denote $D^n = (A^n, T^n)$ to be the vector of data for individual n . For individual n , the parameters $Ab_0^n, \beta^n, \delta^n, d_s^n, d_l^n, d_a^n$ and ρ^n are estimated. These parameters are denoted $\theta^n = \{Ab_0^n, \beta^n, \delta^n, d_s^n, d_l^n, d_a^n, \rho^n\}$. The model predicted antibody levels will be $\{A(t_1), A(t_2), \dots, A(t_K)\}$. We assume log-Normally distributed measurement error such that the difference between $\log(a_i)$ and $\log(Ab(t_i))$ is Normally distributed with variance σ_{obs}^2 . For model predicted antibody levels $Ab(t_i)$ the data likelihood for individual n is given by

$$L_{mod}^n(\theta^n | D^n) = \prod_{k \in K} \frac{e^{-\frac{(\log(a_i) - \log(Ab(t_i)))^2}{2\sigma_{obs}^2}}}{a_k \sigma_{obs} \sqrt{2\pi}}$$

As described above, for each individual there are 7 parameters to be estimated: $\theta^n = \{Ab_0^n, \beta^n, \delta^n, d_s^n, d_l^n, d_a^n, \rho^n\}$. The mixed effects likelihood can be written as follows:

$$L_{mix}^n(\theta^n | D^n) = \left(\frac{e^{-\frac{(\log(Ab_0^n) - \mu_{Ab,0})^2}{2\sigma_{Ab,0}^2}}}{\sqrt{2\pi} Ab_0^n \sigma_{Ab,0}} \right) \left(\frac{e^{-\frac{(\log(\beta^n) - \mu_\beta)^2}{2\sigma_\beta^2}}}{\sqrt{2\pi} \beta^n \sigma_\beta} \right)$$

$$\left(\frac{e^{-\frac{(\log(d_s^n) - \mu_{d,s})^2}{2\sigma_{d,s}^2}}}{\sqrt{2\pi} d_s^n \sigma_{d,s}} \right) \left(\frac{e^{-\frac{(\log(d_l^n) - \mu_{d,l})^2}{2\sigma_{d,l}^2}}}{\sqrt{2\pi} d_l^n \sigma_{d,l}} \right) \left(\frac{e^{-\frac{(\log(d_a^n) - \mu_{d,a})^2}{2\sigma_{d,a}^2}}}{\sqrt{2\pi} d_a^n \sigma_{d,a}} \right) \left(\frac{e^{-\frac{(\log(\frac{\rho^n}{1-\rho^n}) - \mu_\rho)^2}{2\sigma_\rho^2}}}{\sqrt{2\pi} \rho^n (1 - \rho^n) \sigma_\rho} \right)$$

As the proportion of the ASCs that are long-lived must be bounded by 0 and 1, the individual-level parameters ρ^n are assumed to be drawn from logit-Normal distributions. Note that the pre-existing antibody level Ab_0^n will be variable, depending on a large number of covariates such as age and past exposure. We therefore do not attempt to constrain pre-existing antibody levels using mixed effects.

Denote $D = \{D^1, \dots, D^N\}$ to be the vector of data for all N participants. We denote θ to be the combined vector of population-level parameters and individual-level parameters to be estimated. The total likelihood is obtained by multiplying the likelihood for each participant

$$L_{total}(\theta|D) = \prod_{n \in N} L_{mix}^n(\theta^n|D^n) L_{mod}^n(\theta^n|D^n)$$

The model was fitted to the data using MCMC methods using software for Bayesian statistical inference of non-linear mixed-effects models. This utilises a Metropolis-within-Gibbs sampler, whereby population-level parameters are updated using a Gibbs sampler, and individual-level and observational parameters are updated using a Metropolis-Hastings sampler.

Statistical analyses

Statistical tests, including Fisher's exact test, Mann-Whitney test and Spearman's correlation test, were performed using R version 4.0.4 and GraphPad Prism version 7.0. Linear and logistic regression models to establish associations between antigen features and antibody responses were performed using Stata version 12.1.

Ethics statement

Collection of the Thai symptomatic longitudinal study in Tha Song Yang, Tak Province, was approved by The Ethics Committee of the Faculty of Tropical Medicine, Mahidol University, Thailand (MUTM 2014-025-01 and -02). For the Thai asymptomatic cohort study in the Kanchanaburi and Ratchaburi provinces, collection of samples was approved by The Ethics Committee of the Faculty of Tropical Medicine, Mahidol University, Thailand (MUTM 2013-027-01). The Human Research Ethics Committee (HREC) at the Walter and Eliza Hall Institute of Medical Research (WEHI) approved the use of these samples in Melbourne (#14/02). The collection and/or use of the 3 malaria-free control panels in Melbourne were approved by the HREC at WEHI (#14/02).

Results

Naturally acquired total IgG kinetics are antigen-specific and dominated by subclass IgG1 following symptomatic *P. vivax* infections


We first determined the levels of antigen-specific total IgG antibody in Thai symptomatic patients ($n = 34$) at the time of *P. vivax* infection and throughout 9 months of follow-up in comparison to antibody measurements in three large malaria-naïve control panels ($n = 274$). Antibody responses to the 52 *P. vivax* antigens were measured at baseline (week 0), weeks 1 and 2, then every 2 weeks for 6 months, followed by every month until 9 months after infection (total of 17 measurements). These individuals had no recurrent *Plasmodium* infections during the 9-month follow-up (as determined by *P. vivax*-specific PCR). An overview of the study design and the antibody measurements performed is shown in Fig. 1 and Table 1. The 52 *P. vivax* antigens included both pre-erythrocytic

and blood-stage expressed proteins, and were selected as they are known to be immunogenic in *P. vivax*-endemic regions, can act as markers of exposure to infection, or are of interest for vaccine development (see Methods and Additional File 2: Table S2 for further details).

Total IgG antibody responses against all 52 antigens were adjusted by subtracting the negative control baseline consisting of malaria-free individuals from three separate sources (Fig. 2) (the total IgG response for each individual antigen prior to adjustment is shown in Additional File 1: Fig. S2). We observed that IgG antibody responses to most *P. vivax* antigens followed a general pattern consisting of an initial peak in total IgG within 2 weeks post-infection and a subsequent decay of varying rates towards the baseline (Fig. 2). Subsequently, we classified 27 antigens with median total IgG of more than 1 SD above the median negative control baseline at 1-week post-infection as "IgG-immunogenic" for extended analyses (Additional File 1: Fig. S2). All 27 IgG-immunogenic proteins had responses detectable above the baseline at the 9-month final visit, indicating the seropositivity against these *P. vivax* proteins is long-lived. Total IgG responses were also assessed at an individual person level, with variation evident between study participants (Additional File 1: Fig. S3). These observations could be due to differing effects of age, past exposure and history of infections with other pathogens. However, the sample size was not designed for detecting individual-level differences in IgG acquisition and maintenance.

As the magnitude of the IgG response cannot be directly compared between different *P. vivax* antigens using our assay (due to differing concentrations of antigen coupled to beads), we defined immunogenicity in comparison to the hyperimmune PNG pool that was included as a positive control on each plate, as per our previous work [48]. We calculated the proportion of Thai participants, at week 1 post-infection, reaching 1, 5, 10, 25 and 50% of the antibody response in the PNG pool (Additional File 2: Table S1). The top 5 most immunogenic proteins were PVX_000930 (sexual stage antigen s16), PVX_096995 (tryptophan-rich antigen, Pv-fam-a), PVX_097715 (hypothetical protein), PVX_112670 (unspecified protein) and PVX_094255 (reticulocyte binding protein 2b, RBP2b) (two constructs tested), with >20 of the 34 individuals reaching 50% of the antibody response in the PNG pool.


To determine the predominance and longevity of IgG subclass responses, IgG1, 2, 3 and 4 responses to 26 IgG-immunogenic antigens were measured at the time of *P. vivax* infection, week 2 post-infection and then every 2-8 weeks throughout the study follow-up (PVX_123685 was excluded from the panel by mistake) (Fig. 3). Similar to total IgG, as antigen-specific baseline signals



varied significantly between antigens, data were adjusted with the negative control baseline by subtraction. We observed that after adjustment IgG1 and IgG3 responses were the predominant IgG subclasses detected following symptomatic *P. vivax* infections in this western Thai population (Fig. 3), with no or minimal detection of IgG2 and IgG4. IgG1 and IgG3 responses closely followed the same pattern as total IgG: an early peak followed by a decay of different rates. Of the two detectable subclasses, IgG1 was the dominant response and the IgG1 magnitude most strongly mirrored the total IgG response over time. In contrast, we found that hyper-immune adults from PNG (used as the positive control pool) had similar levels of IgG1 and IgG3 to multiple *P. vivax* proteins and that the level of IgG3 was higher compared to that of the Thai symptomatic patients to most proteins (Additional File 1: Fig. S4).

Naturally acquired antigen-specific IgM responses are longer-lived than total IgG following symptomatic *P. vivax* infections

Whilst IgM responses are traditionally thought to be transient following infections, more recent evidence has suggested they can be long-lived [58] or follow a similar kinetic to IgG [58, 59], at least in the context of *P. falciparum* malaria, though this is likely dependent on the antigen assessed and transmission setting [60]. We first assessed IgM immunogenicity at 1-week post-infection in the symptomatic cohort and classified 15 antigens as IgM-positive (more than 2 SD above baseline), including both IgG-immunogenic ($n = 8$) and non-IgG-immunogenic ($n = 7$) antigens (Additional File 1: Fig. S5). For the 15 IgM-positive antigens we assessed the kinetics over 9 months and compared the longevity to that of IgG (Fig. 4), both adjusted against corresponding baseline



signals of the negative control panels. Unadjusted kinetic profiles of IgM responses for each antigen are shown in Additional File 1: Fig. S6.

All antigens exhibited IgM kinetic patterns similar to total IgG, with variability depending on the particular antigen, consisting of an initial peak at week 1 post-infection and gradual decay over the 9 months of study follow-up (Fig. 4). However, the peak at week 1 post infection was generally of lesser magnitude than IgG, and thus, the decay for IgG was steeper, with IgM reaching its minimal level more quickly. IgM signals against all 15 antigens were maintained above the median of negative control panels for more than 6 months, and the magnitude of the response was relatively similar for all proteins. In comparison with total IgG, variations in responses between volunteers were also less prominent in IgM (Fig. 4). The relationship between IgG and IgM responses did differ between proteins, with 8 antigens exhibiting substantially higher IgM than total IgG responses throughout the study (PVX_088820, PVX_091710, PVX_092990, PVX_097680, PVX_098585, PVX_112670, PVX_117880 and PVX_125728), 4 antigens exhibiting stronger total IgG than IgM responses (PVX_000930, PVX_087885, PVX_095055 and PVX_096995) and 3 antigens where total IgG and IgM were comparable (PVX_082670, PVX_082700 and PVX_082735). The level of background IgM antibodies did vary within the malaria-naïve negative control samples tested (Additional File 1: Fig. S5), so whilst the data in Fig. 4 has been adjusted against the

median of the negative controls, this may not adequately account for the background cross-reactivity common for IgM and should be considered when interpreting these results. We thus focused our subsequent analyses on total IgG and IgG subclass results.

Antigens with PEXEL motifs and no transmembrane domains are associated with longer-lived total IgG and IgG1 responses, respectively

To explore the potential sources that could account for the differences in maintenance of IgG and IgG subclass responses against *P. vivax* antigens, we examined the impact of several antigenic features that could be associated with long-lived antibody responses. These included the presence or absence of predicted signal peptides, transmembrane domains, GPI anchors, PEXEL motifs, predicted domains and the number of low complexity regions (Additional File 2: Table S2). The majority of the 52 *P. vivax* antigens were predicted to contain signal peptides (79%), no transmembrane domains (58%), no GPI anchor (92%) and no PEXEL motif (92%). Half of the antigens were predicted to have domains. On average, these antigens had one predicted domain which was most commonly an epidermal growth factor (EGF) or Tryptophan-Threonine-rich *Plasmodium* antigen (Pfam) domain at the C-terminus. All antigens were predicted to contain LCRs (range = 1–17, mean = 5.98; median = 4), except for PVX_098915, which had none.




Fig. 4 Comparison of adjusted total IgG and IgM against 15 *P. vivax* antigens following symptomatic infections. Fifteen antigens with IgM positivity at 1-week post-infection (> 2 SD above baseline median) were selected, and their IgM responses over a 9-month period were measured and compared to total IgG responses plotted on the same scale. Each line and colour represent an antibody class (either IgG or IgM) and its specific antibody kinetics following exposure. Data were adjusted by subtracting the median of pooled negative control panels ($n = 274$). Data are expressed as the median \pm 95% CI

In order to have a quantitative measure of antibody longevity for this analysis, we used mathematical modelling to estimate the decay rate (half-life) of antigen-specific antibodies per antigen beyond the 9-months study in the Thai symptomatic cohort ($n = 34$). The model of antibody kinetics was based on previous work [61, 62], with improvements as detailed in the methods. The model assumes that the antigenic stimulus from the *Plasmodium* infection results in the proliferation and differentiation of B cells into both short- and long-lived ASCs (or plasma cells) that secrete Ig molecules. The short- and long-lived ASCs decay at different rates resulting in the bi-phasic decay in antibody levels over time, such as observed in this study with total IgG (Fig. 2). The model estimates the half-life of secreted antibody molecules and both short- and long-lived ASCs and takes into consideration the antigen-specific background signals measured in malaria-naïve negative control panels ($n = 274$) (Additional File 1: Fig. S7).

To assess if the presence of these antigenic features was linked to the estimated decay rate of antibodies in this cohort, we performed non-parametric Wilcoxon tests between antigens with and without the antigenic features. As shown in Fig. 5A, *P. vivax* antigens with PEXEL

motifs were associated with longer estimated half-life of total IgG molecules (medians = 25.72 days versus 18.20 days, $p = 0.038$). Additionally, antigens with no predicted TMD were associated with an estimated half-life of IgG1-producing long-lived ASCs more than three times longer than proteins with TMD domains (medians = 1305.71 days versus 407.12 days, $p = 0.037$) (Fig. 5B). No significant associations were observed for other antigenic features and estimated decay rates. Similarly, no significant associations were found after performing linear regression between the number of predicted domains or LCRs and estimated half-lives generated from modelling (data not shown).

Antigen-specific total IgG responses are longer-lived following asymptomatic compared to symptomatic *P. vivax* infections

Total IgG, IgG subclass and IgM measurements were made in Thai individuals following symptomatic *P. vivax* infections. As these individuals were symptomatic, they have not yet gained sufficient immunity to clinical disease. We were therefore interested to determine whether total IgG kinetics differed in individuals from a similar region of western Thailand who had already acquired





Fig. 5 The association between presence of PEXEL motifs and estimated half-life of total IgG antibodies, and the presence of TMD and the estimated half-life of IgG1-producing long-lived ASCs in the Thai symptomatic population. The kinetics of total IgG antibody against 52 IgG-immunogenic *P. vivax* antigens were determined following symptomatic *P. vivax* infections in Thai individuals in the absence of recurrent infections for 9 months. Mathematical modelling was then used to generate the estimated half-life of total IgG and IgG1-producing long-lived ASCs. The presence of PEXEL and TMDs was predicted for each *P. vivax* antigen using the online servers: PlasmoDB (<https://plasmodb.org/plasmo/app>) and Transmembrane Helices Hidden Markov Model (TMHMM) Server version 2.0 (<http://www.cbs.dtu.dk/services/TMHMM/>). Wilcoxon tests (non-parametric) were performed to compare between antigens with (Yes) and without (No) PEXEL or TMD. *p* value less than 0.05 was considered significant

immunity to symptomatic disease. Thirty individuals were selected from an existing cohort [32] who had an asymptomatic *P. vivax* infection with data available from 14 visits in total (each visit one-month apart). Each individual had data from at least 10 visits post infection and were PCR-negative for *Plasmodium* at all of these subsequent visits. Total IgG responses were measured against the 27 *P. vivax* antigens found to be immunogenic (total IgG) in the symptomatic cohort. Kinetic profiles against these 27 antigens are shown in Fig. 6 and the antigen-specific profiles in Additional File 1: Fig. S8 (with direct comparison between the symptomatic and asymptomatic cohorts).

Twenty-one of 27 *P. vivax* antigens were found to be IgG-immunogenic (more than 1 SD above baseline at the time of infection, w0) following an asymptomatic *P. vivax* infection, all of which showed a long-lived total IgG profile (detectable IgG above baseline at 10-months post-infection). In comparison, 25 proteins in the symptomatic cohort had detectable IgG at the final visit (9 months post-infection) above the median negative control baseline, with a greater drop in magnitude than observed in the asymptomatic cohort. Individual variation in responses to antigens was larger than observed in the symptomatic patients, indicated by the longer error bars in Additional File 1: Fig. S8. This is likely related to greater differences in levels of past exposure in these

individuals, compared to most symptomatic patients having 0 or 1 past infections (self-reported history). For *P. vivax* antigens that were non-immunogenic in the asymptomatic *P. vivax* infected individuals ($n = 6$), 3 antigens (PVX_097715, PVX_090970, PVX_090265) showed high seropositivity in the symptomatic Thai cohort in which at least half ($n \geq 17$) of the participants reached 50% of the antibody response in the PNG pool as shown in Additional File 2: Table S1. The lack of response in the asymptomatic individuals could be due to the reduced parasite load; the median parasite density in the asymptomatic individuals (median = 1.0×10^2) was 2.5×10^3 times lower than the symptomatic (median = 2.5×10^5) (Table 1). There was no clear peak in response at the first time point post-infection (Additional File 1: Fig. S8), as there was for the symptomatic cohort, which could relate to the timing (monthly sampling rather than 1–2 weekly) or because the onset of infection is less well-defined in the asymptomatic cohort.

We also calculated the proportion of asymptomatic participants reaching 1, 5, 10, 25 and 50% of the antibody response in the PNG pool at week 0 since infection (Additional File 2: Table S3). The top 5 most immunogenic proteins in this asymptomatic cohort were PVX_094255A (reticulocyte binding protein 2b, RBP2b) (two constructs tested), PVX_087885 (rhoptry associated membrane antigen, putative), PVX_090265



(tryptophan-rich antigen, Pv-fam-a), PVX_099980 (MSP1-19), and PVX_095055 (Rh5 interacting protein, putative), with > 15 of the 30 participants reaching 50% of the antibody response in the PNG pool.

Discussion

Antibody responses against a large number of *P. vivax* proteins are acquired following *P. vivax* infections; however, little is known about how long these antibody responses last in different settings and what factors might influence the longevity. In the present study, we used human samples from a low transmission region in Thailand to develop a model of the dynamics of naturally acquired antibody responses against 52 *P. vivax* antigens, including well-studied vaccine candidates and novel targets, following both symptomatic and asymptomatic *P. vivax* infections. We show that *P. vivax* antigens are able to induce total IgG responses of varying longevities, dependent on the *P. vivax* antigen target, with a general pattern of bi-phasic decay. Of the 52 *P. vivax* antigens, 27 were highly immunogenic at the peak of the antibody response 1-week post clinical *P. vivax* infection, with all of these having detectable IgG levels above baseline at 9-months post-infection despite a decline in the

magnitude. Identification of the peak antibody response 1–2 weeks post clinical symptoms is in line with prior studies of antibody kinetics following *P. falciparum* [59, 61] and *P. knowlesi* [63] infections. We observed that for most antigens IgG1 was the dominant subclass and that the decay of IgG1 mirrored that of total IgG. Of interest, we also detected long-lived antigen-specific IgM responses to 15 *P. vivax* proteins, with a similar kinetic pattern after symptomatic infections. When comparing antibody kinetics following symptomatic or asymptomatic *P. vivax* infections, we show that responses following asymptomatic infections are maintained at a higher magnitude for longer, in line with our previous observations that antibody longevity increases in individuals with higher levels of past exposure [26].

To investigate the source of variation in IgG antibody longevity against different *P. vivax* antigens, we first improved upon an existing mathematical model of antibody kinetics to quantify an estimate of the half-life of secreted antibody molecules and both short- and long-lived ASCs against the 27 immunogenic proteins for total IgG, IgG1 and IgG3. We then used these quantified antibody longevity variables to assess the impact of the presence of several major protein characteristics that could

influence the immune response, such as SPs, TMDs, GPI anchors and PEXEL motifs. Only two significant associations were found: (1) *P. vivax* antigens with PEXEL motifs were associated with longer estimated half-lives of total IgG molecules and (2) *P. vivax* antigens with no predicted TMD were associated with longer estimated half-lives of IgG1-producing long-lived ASCs. PEXEL motifs are responsible for trafficking and secreting of several *Plasmodium* surface antigens [64]; thus, proteins that contain PEXEL motifs could be capable of inducing robust immune responses, accounting for the association with longer estimated half-lives of total IgG molecules. However, there was no association between the presence of PEXEL motifs and longevity of short- or long-lived ASCs, and the difference in longevity of the IgG molecules was small (i.e. 25.72 days versus 18.20 days). We had hypothesised that the presence of TMDs would suggest the *P. vivax* antigens were membrane-bound and thus highly accessible to the immune system. However, we found that antigens with no TMD had longer half-lives (>3-fold) for IgG1-secreting long-lived ASCs. There was no significant relationship between the presence of TMD and total IgG or IgG3-secreting long-lived ASCs. The impact of these protein features warrants further examination, for example through the assessment of other factors that could influence antibody longevity such as regions of structural disorder [65], the presence of predicted B cell epitopes, the sequence polymorphism of the antigens, and how these interact or could confound the current findings of associations with PEXEL and TMDs. Interestingly, the only other study we could identify that has aimed to assess potential protein features and their association with antibody longevity in the context of *Plasmodium* infections, hypothesised that the tandem repeated epitope sequences in the *P. falciparum* protein MSP2 could account for the short antibody longevity observed to this protein [66] compared to the other antigens they assessed. Tandem repeats are common in regions of protein disorder, and regions of protein disorder are enriched in exported proteins [65], making this a key feature for future assessment.

Interestingly, whilst we expected the 52 *P. vivax* antigens to be mostly immunogenic based on our prior studies [26, 46], we found that only 27 induced IgG and 15 induced IgM seropositive responses at week 1 following infection. Potential reasons are that we have used purified antigens compared to our large-scale analysis of more than 300 crude *P. vivax* antigens [26] and that our sample size is much smaller than our analysis of 829 individuals from malaria-endemic regions using identical purified antigens [46]. We also set a relatively stringent cut-off of the median plus two times standard deviation for IgM, to account for the variability observed in the malaria-naïve

negative controls. There is some evidence of non-specific IgM binding specific *P. falciparum* proteins, and this could be further assessed for *P. vivax* [67, 68]; accounting for such a phenomenon may reveal more antigens with *P. vivax*-specific IgM responses above background. Other studies have also demonstrated immunogenicity against a number of our *P. vivax* antigens identified as non-immunogenic in the current study, supporting the notion that the lack of immune response identified is related to the small sample size, genetic variations between individuals and lack of past exposure in the symptomatic Thai patients. For example, whilst the *P. vivax* antigen GAMA was categorised as non-immunogenic in our study, Changrob et al. reported high seroprevalence of more than 55% even a year after recovery from acute infection in southern Thailand [69]. We classed 6 tryptophan-rich antigens as non-immunogenic, among which 2 (PVX_112675 and PVX_092990) were found to be highly immunogenic and capable of inducing long-lived antibody responses that lasted for at least 5 years post-infection in individuals from Korea, Myanmar and China [70]. Various reasons could account for such differences, in addition to those previously mentioned, including differences in study design (possibly boosting infections), protein construct sequence, detection methodology, past history of infection and region-specific *P. vivax* transmission levels.

Our data indicates that cytophilic antibodies (IgG1 and IgG3) are the dominant IgG subclass response induced by our large panel of *P. vivax* antigens, at least following symptomatic *P. vivax* infections in this low-transmission region. This is consistent with previous research highlighting IgG1 and IgG3 as the key subclasses following malaria, regardless of the *Plasmodium* species or geographical location [48, 59, 71–73]. It has previously been proposed that *P. falciparum* infection initially induces non-cytophilic followed by cytophilic antibodies [74], but this was not what we observed following *P. vivax* infections in individuals with symptomatic infections in western Thailand (who presumably have had limited past exposure, supported by their self-reported malaria history). Whether or not this observation is species-specific remains unclear. In comparison to previously published studies mostly utilising cross-sectional cohorts, our longitudinal data and relatively frequent sampling allowed us to demonstrate that the dominance of IgG subclasses remains constant over time following *P. vivax* infection, at least in the absence of any boosting infections. We did not observe any evidence of subclass switch, which is proposed to occur with an increase in exposure and age [72], in this low-transmission region. However, in our positive control pool consisting of hyperimmune individuals from PNG, we did observe the presence of both IgG1

and IgG3 against most *P. vivax* antigens, compared to IgG1 alone for our Thai volunteers, suggesting that with increasing exposure and in higher transmission levels this switch occurs. Interestingly, the subclass switch was hypothesised to contribute to the longevity of total IgG upon exposure to malaria antigens [75, 76] but was not observed in our study where the study site is of relatively low transmission.

IgM antibodies have been identified as a short-lived response due to the general notion that IgM is rapidly replaced by IgG after a short period of time exposed to a pathogen [77]. In the context of *P. vivax* infections, Park and colleagues reported *P. vivax* MSP1-specific IgM responses lasting less than 3 months [78]. However, our findings indicate that IgM responses can be long-lived (for at least 9 months), consistent with more recently published reports [79–81]. Whilst the underlying mechanism for induction and maintenance of long-lived IgM is unclear, IgM-producing long-lived plasma cells have been shown to reside in the spleen in a mouse model [80]. These cells undergo somatic hypermutation upon antigenic stimulation, independent of germinal centre and T cell help, suggesting an affinity maturation process that is unique and distinct from that of IgG [80, 82]. Our study is the first to address *P. vivax*-specific IgM kinetics in detail and to demonstrate that IgM can be maintained for at least 9 months in the absence of boosting infections. The direct comparison between total IgG and IgM kinetics also provides unique insights into the antigen-specific difference in the temporal dynamics between the two antibody responses. However, caution is warranted when interpreting these results as evidence in other disease models has shown that IgM can potentially be cross-reactive to highly IgG-immunogenic epitopes [83, 84]. Confounding factors that could potentially influence the longevity of *P. vivax*-specific IgM responses should also be taken into consideration, such as the presence of other pathogens displaying structurally similar antigens between participants in this cohort.

We also measured IgG antibodies following a PCR-positive but asymptomatic *P. vivax* infection in a similar region of western Thailand as a comparison to the symptomatic cohort in the nearby region. Defining antibody responses in asymptomatic individuals is challenging given that case detection and recruitment become more dependent on molecular surveillance tools that are less accessible than microscopy in field-settings. Yet, it is important to assess longevity of naturally acquired immune responses following asymptomatic infections given the high proportion of *P. vivax* infections that are asymptomatic (i.e. [85, 86]). We were able to utilise samples previously collected as part of

an observational longitudinal cohort study, in which most infections were asymptomatic [32], to address this question. We have previously shown that total IgG responses against a panel of 11 *P. vivax* antigens are induced following asymptomatic *P. vivax* infections [87]. Our current results confirm these findings, against a larger panel of *P. vivax* antigens, and we were able to further extend these results to show that these responses can be maintained for an extended period of time following clearance of the infection (> 10 months). Whilst there have been previous studies assessing antibody longevity following *P. vivax* infections [29, 69, 88, 89], ours is the first to do so to a large panel of *P. vivax* antigens with much more frequent sampling after an infection. This enables us to establish an antibody kinetic profile of higher resolution and with high certainty of no recurrent boosting infections during follow-up. In addition, the malaria-naïve control panels in our study are exceptionally larger than most previously published work, allowing more accurate prediction of seroprevalence in the target population.

Our study has some limitations. First, antibody levels measured at enrolment (week 0) are already elevated above the baseline measured in malaria-naïve control panels. This may be due to a delay in care seeking, or because some individuals in our study have had past exposure to malaria. It is also currently impossible to determine whether the symptomatic infections detected at week 0 in our individuals are new infections from mosquito bites or due to relapses from previous infections. The source of infection may play a role in antibody boosting and maintenance. Second, due to the longer interval between sampling (4 weeks), there is the possibility of missed infections present in the asymptomatic cohort, which could contribute to the greater variation observed between individuals, especially towards the end of the study after adjusted for time of infection as the sample size reduced. Third, whilst of interest, our study was not designed to look at differences in antibody acquisition and longevity at the individual level (i.e. with age and past exposure). Additionally, the IgG subclass and IgM profiles were not obtained in the asymptomatic cohort and thus further investigations on IgM responses following *P. vivax* asymptomatic infections will be required. Finally, we only measured total IgG responses in the asymptomatic cohort against the 27 *P. vivax* proteins identified as immunogenic in the symptomatic cohort; the profile against those non-immunogenic proteins may have been different in the individuals with greater past exposure, particularly given the breadth of the response is expected to expand with increasing exposure [90].

Conclusions

In the present study, we have provided new insights that contribute to our understanding of *P. vivax*-specific antibody acquisition and maintenance in low-transmission settings such as western Thailand. Our findings have highlighted the antigen- and time-specific nature of *P. vivax* humoral responses, and these responses could be further assessed in the context of using antibodies as markers of past exposure or immune status [23]. Examination of serological status in populations could facilitate identification of transmission hotspots for timely management and appropriate adjustment of pre-existing intervention programs or identify populations at risk of resurgence due to waning immunity. We have recently identified 8 *P. vivax* antigens that can be used as serological markers of recent exposure to *P. vivax* infection, reaching 80% sensitivity and 80% specificity for classification of infection in the prior 9-months [46]. Our current findings, with detailed understanding of antibody kinetics, could be incorporated into the existing algorithm to potentially further increase the accuracy of these serological exposure markers. Our findings also provide evidence that IgG subclasses would be unlikely to improve these markers, which are based on total IgG, given the dominant IgG1 and IgG3 response follows the same pattern of longevity. In addition, we also show that multiple *P. vivax* antigens in our panel, that are not current vaccine candidates, are highly immunogenic and induce long-lived antibody responses following asymptomatic *P. vivax* infections. This finding will broaden our current scope of ‘vaccine candidates’ and facilitate exploration of untraditional targets. The finding of long-lived IgM is in support of recent work in *P. falciparum* [58] and suggests IgM should also be assessed in studies looking at markers of immunity. We have also identified two possible antigenic features associated with long-lived IgG antibody responses, including PEXEL motifs which facilitate transport of antigens to the cell surface. IgG responses were also longer lived in the asymptomatic cohort, people with presumably higher levels of past exposure. Moving forward, investigations on antigenic sequence features and host cellular responses and their associations with antibody maintenance are of high priority. Such information will help understand what drives a long-lived immune response and whether this can be exploited to design vaccines with longer durability.

Abbreviations

P. vivax: *Plasmodium vivax*; *P. falciparum*: *Plasmodium falciparum*; G6PD: Glucose-6-phosphate dehydrogenase; Ig: Immunoglobulin; SP: Signal peptide; PEXEL: *Plasmodium* exported element; TMD: Transmembrane domain; GPI: Glycosylphosphatidylinositol; LCR: Low complexity region; PCR: Polymerase chain reaction; ARC: Australian Red Cross; TRC: Thai Red Cross; VBDR: Volunteer Blood Donor Registry; WGCF: Wheat germ cell-free; PNG: Papua New Guinea;

MFI: Median fluorescence intensity; RAU: Relative antibody unit; ASC: Antibody secreting cell; MCMC: Markov chain Monte Carlo; SD: Standard deviation; RBP: Reticulocyte binding protein; EGF: Epidermal growth factor.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12916-022-02281-9>.

Additional File 1: Fig S1. Schematic representation of the relationship between the population-level and individual-level parameters for the half-life of the short-lived ASCs. **Fig S2.** Total IgG kinetic profiles against 52 *P. vivax* antigens in 34 Thai patients with symptomatic *P. vivax* infections. **Fig S3.** Adjusted total IgG kinetic profiles at an individual level following symptomatic infections. **Fig S4.** Comparison of the adjusted peak IgG subclass magnitude between symptomatic Thai patients and the PNG hyperimmune pool. **Fig S5.** Comparison of the peak IgM magnitude between symptomatic Thai patients and malaria-free individuals. **Fig S6.** Unadjusted IgM kinetics against 15 *P. vivax* antigens in 34 Thai individuals following symptomatic *P. vivax* infections. **Fig S7.** Distribution of data for estimated half-lives of antibodies and ASCs against IgG-immunogenic antigens following symptomatic *P. vivax* infections in the Thai symptomatic cohort. **Fig S8.** Comparison of adjusted total IgG against 27 *P. vivax* antigens following symptomatic or asymptomatic *P. vivax* infections.

Additional file 2: Table S1. Immunogenicity of total IgG against 52 *P. vivax* antigens at 1-week post-infection in Thai symptomatic volunteers. **Table S2.** *P. vivax* proteins used in this study. **Table S3.** Immunogenicity of total IgG against 27 IgG-immunogenic *P. vivax* antigens at 0-week since infection in Thai asymptomatic volunteers.

Acknowledgements

We gratefully acknowledge the extensive field teams that contributed to sample collection and qPCR assays in the original cohort studies. We thank all of the individuals who participated in each of the studies and the Australian and Thai Red Cross for donation of blood samples. We thank the Volunteer Blood Donor Registry at WEHI for donation of plasma samples and Lina Laskos for their collection and advice. We thank Christopher King (Case Western Reserve University) for provision of the PNG control plasma pool. We thank Fumie Matsuura (CellFree Sciences), Christele Huon (Institut Pasteur) and Jakub Gruszczyk (WEHI) for contributing to expression of proteins. We thank Connie Li-Wai-Suen for writing the R script used for converting MFI values to relative antibody units.

Authors' contributions

Study planning and project conception—JS; IM; R/L. Antigenic protein production—ET; MH; W-HT; JH; CEC; TT. Sample management—ZS-JL; JS; SC; CK; IM; R/L. Mathematical modelling—MW. Data analysis—ZS-JL; R/L. Figure/table preparation—ZS-JL; R/L. Manuscript preparation—ZS-JL; R/L. Manuscript edits and proofing—ZS-JL; JS; MW; SC; CK; ET; MH; W-HT; JH; CEC; TT; IM; R/L. All authors read and approved the final manuscript.

Funding

Funding was provided by the National Health and Medical Research Council (#1092789, #1134989, and #1043345 to IM, #1143187 to W-HT, and #1173210 to RL), the Global Health Innovative Technology Fund (<https://www.ghitfund.org/>) (T2015-142 to IM) and the National Institute of Allergy and Infectious Diseases (NIH grant 5R01 AI 104822 to JS). We acknowledge support from the National Research Council of Thailand and the Victorian State Government Operational Infrastructure Support and Australian Government NHMRC IRIISS. W-HT was a Howard Hughes Medical Institute-Wellcome Trust International Research Scholar (<https://www.hhmi.org/programs/biomedical-research/international-programs/~208693/Z/17/Z>). ZS-JL was supported by the Melbourne Research Scholarship, University of Melbourne.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

Collection of the Thai symptomatic longitudinal study in Tha Song Yang, Tak Province, was approved by The Ethics Committee of the Faculty of Tropical Medicine, Mahidol University, Thailand (MUTM 2014-025-01 and -02). For the Thai asymptomatic cohort study in the Kanchanaburi and Ratchaburi provinces, collection of samples was approved by The Ethics Committee of the Faculty of Tropical Medicine, Mahidol University, Thailand (MUTM 2013-027-01). The Human Research Ethics Committee (HREC) at the Walter and Eliza Hall Institute of Medical Research (WEHI) approved the use of these samples in Melbourne (#14/02). The collection and/or use of the 3 malaria-free control panels in Melbourne were approved by the HREC at WEHI (#14/02). All individuals provided informed consent and/or assent.

Consent for publication

Not applicable.

Competing interests

R.J.L., M.W., T.T., and I.M. are inventors on patent PCT/US17/67926 on a system, method, apparatus and diagnostic test for *Plasmodium vivax*. MH was employed by company CellFree Sciences Co., Ltd. The remaining authors declare no conflict of interest.

Author details

¹The Walter and Eliza Hall Institute of Medical Research, Parkville, Victoria 3052, Australia. ²Department of Medical Biology, University of Melbourne, Parkville, Victoria 3010, Australia. ³Current affiliation: Deakin University, School of Medicine, IMPACT, Institute for Innovation in Physical and Mental Health and Clinical Translation, Geelong, Victoria 3220, Australia. ⁴Mahidol Vivax Research Unit, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand. ⁵Infectious Disease Epidemiology and Analytics G5 Unit, Department of Global Health, Institut Pasteur, Paris, France. ⁶Proteo-Science Center, Ehime University, Matsuyama, Japan. ⁷CellFree Sciences Co., Ltd., Yokohama, Japan and RIKEN Centre for Integrative Medical Sciences, Yokohama, Japan. ⁸Malaria Parasite Biology and Vaccines, Department of Parasites & Insect Vectors, Institut Pasteur, Paris, France.

Received: 2 November 2021 Accepted: 2 February 2022

Published online: 09 March 2022

References

- WHO. World malaria report; 2021. p. 2021.
- Mendis K, Sina BJ, Marchesini P, Carter R. The neglected burden of *Plasmodium vivax* malaria. *Am J Trop Med Hyg.* 2001;64(1-2 Suppl):97–106.
- Baird JK. Evidence and implications of mortality associated with acute *Plasmodium vivax* malaria. *Clin Microbiol Rev.* 2013;26(1):36–57.
- Robinson LJ, Wampfler R, Betuela I, Karl S, White MT, Li Wai Suen CSN, et al. Strategies for understanding and reducing the *Plasmodium vivax* and *Plasmodium ovale* hypnozoite reservoir in Papua New Guinean children: a randomised placebo-controlled trial and mathematical model. *Plos Med.* 2015;12(10):e1001891.
- Commons RJ, Simpson JA, Watson J, White NJ, Price RN. Estimating the proportion of *Plasmodium vivax* recurrences caused by relapse: a systematic review and meta-analysis. *Am J Trop Med Hyg.* 2020;103(3):1094–9.
- Hulden L, Hulden L. Activation of the hypnozoite: a part of *Plasmodium vivax* life cycle and survival. *Malaria J.* 2011;10(1):90.
- White N. Determinants of relapse periodicity in *Plasmodium vivax* malaria. *Malaria J.* 2011;10(1):297.
- Olliaro PL, Barnwell JW, Barry A, Mendis K, Mueller I, Reeder JC, et al. Implications of *Plasmodium vivax* biology for control, elimination, and research. *Am J Trop Med Hyg.* 2016;95(6_Suppl):4–14.
- Llanos-Cuentas A, Lacerda MVG, Hien TT, Vélez ID, Namaik-Larp C, Chu CS, et al. Tafenoquine versus Primaquine to prevent relapse of *Plasmodium vivax* malaria. *N Engl J Med.* 2019;380(3):229–41.
- Baird JK, Rieckmann KHJ. Can primaquine therapy for vivax malaria be improved? *Trends Parasitol.* 2003;19(3):115–20.
- Howes RE, Battle KE, Satyagraha AW, Baird JK, Hay SI. G6PD deficiency: global distribution, genetic variants and primaquine therapy. *Adv Parasitol.* 2013;81:133–201 Elsevier.
- Kondrashin A, Baranova AM, Ashley EA, Recht J, White NJ, Sergiev VP. Mass primaquine treatment to eliminate vivax malaria: lessons from the past. *Malaria J.* 2014;13(1):51.
- Nguitragool W, Mueller I, Kumpitak C, Saeseu T, Bantuchai S, Yorsaeng R, et al. Very high carriage of gametocytes in asymptomatic low-density *Plasmodium falciparum* and *P. vivax* infections in western Thailand. *Parasit Vectors.* 2017;10(1):512.
- Hutagalung JJTLGH. Prevalence of asymptomatic submicroscopic malaria in eastern Indonesia: a cross sectional survey and spatial analysis. *Lancet Glob Health.* 2017;5:S13.
- Nguyen T-N, von Seidlein L, Nguyen T-V, Truong P-N, Do Hung S, Pham H-T, et al. The persistence and oscillations of submicroscopic *Plasmodium falciparum* and *Plasmodium vivax* infections over time in Vietnam: an open cohort study. *Lancet Infect Dis.* 2018;18(5):565–72.
- Kho S, Qotrunnada L, Leonardo L, Andries B, Wardani PA, Fricot A, et al. Hidden biomass of intact malaria parasites in the human spleen. *N Engl J Med.* 2021;384(21):2067–9.
- Kho S, Qotrunnada L, Leonardo L, Andries B, Wardani PA, Fricot A, et al. Evaluation of splenic accumulation and colocalization of immature reticulocytes and *Plasmodium vivax* in asymptomatic malaria: a prospective human splenectomy study. *Plos Med.* 2021;18(5):e1003632.
- Brito MAM, Baro B, Raiol TC, Ayllon-Hermida A, Safe IP, Deroost K, et al. Morphological and transcriptional changes in human bone marrow during natural *Plasmodium vivax* malaria infections. *J Infect Dis.* 2020.
- Obaldia N 3rd, Meibalan E, Sa JM, Ma S, Clark MA, Mejia P, et al. Bone marrow is a major parasite reservoir in *Plasmodium vivax* infection. *mBio.* 2018;9(3):e00625–18.
- Tayipto Y, Liu Z, Mueller I, Longley RJ. Serology for *Plasmodium vivax* surveillance: a novel approach to accelerate towards elimination. *Parasitol Int.* 2021;87:102492.
- Reyes-Sandoval A. *Plasmodium vivax* pre-erythrocytic vaccines. *Parasitol Int.* 2021;84:102411.
- Doolan DL, Dobaño C, Baird JK. Acquired immunity to malaria. *Clin Microbiol Rev.* 2009;22(1):13–36.
- Fowkes FJ, Boeuf P, Beeson JG. Immunity to malaria in an era of declining malaria transmission. *Parasitology.* 2016;143(2):139–53.
- Pinkevych M, Petravic J, Chelimo K, Kazura JW, Moormann AM, Davenport MP. The dynamics of naturally acquired immunity to *Plasmodium falciparum* infection. *Plos Comput Biol.* 2012;8(10):e1002729.
- Longley RJ, Sattabongkot J, Mueller I. Insights into the naturally acquired immune response to *Plasmodium vivax* malaria. *Parasitology.* 2016;143(2):154–70.
- Longley RJ, White MT, Takashima E, Morita M, Kanoi BN, Li Wai Suen CSN, et al. Naturally acquired antibody responses to more than 300 *Plasmodium vivax* proteins in three geographic regions. *Plos Negl Trop Dis.* 2017;11(9):e0005888.
- Longley RJ, Sriporote P, Chobson P, Saeseu T, Sukasem C, Phuanukoonnon S, et al. High efficacy of primaquine treatment for *Plasmodium vivax* in Western Thailand. *Am J Trop Med Hyg.* 2016;95(5):1086–9.
- Morais CG, Soares IS, Carvalho LH, Fontes CJ, Krettli AU, Braga EMJTA, et al. Antibodies to *Plasmodium vivax* apical membrane antigen 1: persistence and correlation with malaria transmission intensity. *Am J Trop Med Hyg.* 2006;75(4):582–7.
- Wipasa J, Suphavilai C, Okeell LC, Cook J, Corran PH, Thaiika K, et al. Long-lived antibody and B Cell memory responses to the human malaria parasites, *Plasmodium falciparum* and *Plasmodium vivax*. *PLoS Pathog.* 2010;6(2):e1000770.
- Franca CT, He WQ, Gruszczyk J, Lim NT, Lin E, Kiniboro B, et al. *Plasmodium vivax* reticulocyte binding proteins are key targets of naturally acquired immunity in young Papua New Guinean children. *Plos Negl Trop Dis.* 2016;10(9):e0005014.
- He WQ, Shakri AR, Bhardwaj R, França CT, Stanisic DI, Healer J, et al. Antibody responses to *Plasmodium vivax* Duffy binding and Erythrocyte binding proteins predict risk of infection and are associated with protection from clinical Malaria. *Plos Negl Trop Dis.* 2019;13(2):e0006987.
- Nguitragool W, Karl S, White M, Koepfli C, Felger I, Singhasivanon P, et al. Highly heterogeneous residual malaria risk in western Thailand. *Int J Parasitol.* 2019;49(6):455–62.

33. Vatakis D, McMillan M. The signal peptide sequence impacts the immune response elicited by a DNA epitope vaccine. *Clin Vaccine Immunol.* 2011;18(10):1776–80.
34. Kovjazin R, Carmon L. The use of signal peptide domains as vaccine candidates. *Hum Vaccin Immunother.* 2014;10(9):2733–40.
35. Haase S, de Koning-Ward TF. New insights into protein export in malaria parasites. *Cell Microbiol.* 2010;12(5):580–7.
36. Hiller NL, Bhattacharjee S, van Ooij C, Liolios K, Harrison T, Lopez-Estraño C, et al. A host-targeting signal in virulence proteins reveals a secretome in malarial infection. *Science.* 2004;306(5703):1934–7.
37. Kovjazin R, Volovitz I, Daon Y, Vider-Shalit T, Azran R, Tsaban L, et al. Signal peptides and trans-membrane regions are broadly immunogenic and have high CD8+ T cell epitope densities: Implications for vaccine development. *Mol Immunol.* 2011;48(8):1009–18.
38. Naik RS, Branch OH, Woods AS, Vijaykumar M, Perkins DJ, Nahlen BL, et al. Glycosylphosphatidylinositol anchors of *Plasmodium falciparum*: molecular characterization and naturally elicited antibody response that may provide immunity to malaria pathogenesis. *J Exp Med.* 2000;192(11):1563–76.
39. Cheng Y, Wang B, Lu F, Han JH, Ahmed MA, Han ET. Immunological characterization of *Plasmodium vivax* Pv32, a novel predicted GPI-anchored merozoite surface protein. *Malar J.* 2018;17(1):273.
40. Shi Q, Lynch MM, Romero M, Burns JM Jr. Enhanced protection against malaria by a chimeric merozoite surface protein vaccine. *Infect Immun.* 2007;75(3):1349–58.
41. Balam S, Olugbile S, Servis C, Diakitè M, D'Alessandro A, Frank G, et al. *Plasmodium falciparum* merozoite surface protein 2: epitope mapping and fine specificity of human antibody response against non-polymorphic domains. *Malar J.* 2014;13:510.
42. de Silva HD, Saleh S, Kovacevic S, Wang L, Black CG, Plebanski M, et al. The antibody response to *Plasmodium falciparum* Merozoite Surface Protein 4: comparative assessment of specificity and growth inhibitory antibody activity to infection-acquired and immunization-induced epitopes. *Malar J.* 2011;10:266.
43. Toll-Riera M, Radó-Trilla N, Martys F, Albà MM. Role of low-complexity sequences in the formation of novel protein coding sequences. *Mol Biol Evol.* 2012;29(3):883–6.
44. Mendes TA, Lobo FP, Rodrigues TS, Rodrigues-Luiz GF, daRocha WD, Fujiwara RT, et al. Repeat-enriched proteins are related to host cell invasion and immune evasion in parasitic protozoa. *Mol Biol Evol.* 2013;30(4):951–63.
45. Chaudhry SR, Lwin N, Phelan D, Escalante AA, Battistuzzi FU. Comparative analysis of low complexity regions in *Plasmodia*. *Sci Rep.* 2018;8(1):335.
46. Longley RJ, White MT, Takashima E, Brewster J, Morita M, Harbers M, et al. Development and validation of serological markers for detecting recent *Plasmodium vivax* infection. *Nat Med.* 2020;26(5):741–9.
47. Hietanen J, Chim-Ong A, Chiramanewong T, Gruszczak J, Roobsoong W, Tham W-H, et al. Gene models, expression repertoire, and immune response of *Plasmodium vivax* reticulocyte binding proteins. *Infect Immun.* 2016;84(3):677–85.
48. França CT, White MT, He W-Q, Hostetler JB, Brewster J, Frato G, et al. Identification of highly-protective combinations of *Plasmodium vivax* recombinant proteins for vaccine development. *eLife.* 2017;6:e28673.
49. Cole-Tobian JL, Michon P, Biasor M, Richards JS, Beeson JG, Mueller I, et al. Strain-specific duffy binding protein antibodies correlate with protection against infection with homologous compared to heterologous *Plasmodium vivax* strains in Papua New Guinean children. *Infect Immun.* 2009;77(9):4009–17.
50. Healer J, Thompson JK, Riglar DT, Wilson DW, Chiu Y-H, Miura K, et al. Vaccination with conserved regions of erythrocyte-binding antigens induces neutralizing antibodies against multiple strains of *Plasmodium falciparum*. *Plos One.* 2013;8(9):e72504.
51. Mueller I, Galinski MR, Baird JK, Carlton JM, Kochar DK, Alonso PL, et al. Key gaps in the knowledge of *Plasmodium vivax*, a neglected human malaria parasite. *Lancet Infect Dis.* 2009;9(9):555–66.
52. Mazhari R, Brewster J, Fong R, Bourke C, Liu ZSJ, Takashima E, et al. A comparison of non-magnetic and magnetic beads for measuring IgG antibodies against *Plasmodium vivax* antigens in a multiplexed bead-based assay using Luminex technology (Bio-Plex 200 or MAGPIX). *Plos One.* 2020;15(12):e0238010.
53. Bahl A, Brunk B, Crabtree J, Fraunholz MJ, Gajria B, Grant GR, et al. PlasmoDB: the *Plasmodium* genome resource. A database integrating experimental and computational data. *Nucleic Acids Res.* 2003;31(11):212–5.
54. Krogh A, Larsson B, Von Heijne G, Sonnhammer ELJJ. Predicting trans-membrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol.* 2001;305(3):567–80.
55. Pierleoni A, Martelli PL, Casadio R. PredGPI: a GPI-anchor predictor. *BMC Bioinform.* 2008;9(1):392.
56. Mitchell AL, Attwood TK, Babbitt PC, Blum M, Bork P, Bridge A, et al. InterPro in 2019: improving coverage, classification and access to protein sequence annotations. *Nucleic Acids Res.* 2018;47(D1):D351–D60.
57. Wootton JCJC. Non-globular domains in protein sequences: automated segmentation using complexity measures. *Comput Chem.* 1994;18(3):269–85.
58. Boyle MJ, Chan JA, Handayani I, Reiling L, Feng G, Hilton A, et al. IgM in human immunity to *Plasmodium falciparum* malaria. *Sci Adv.* 2019;5(9):eaax4489.
59. Rogier E, Nace D, Dimbu PR, Wakeman B, Pohl J, Beeson JG, et al. Framework for characterizing longitudinal antibody response in children after *Plasmodium falciparum* infection. *Front Immunol.* 2021;12:617951.
60. Walker MR, Knudsen AS, Partey FD, Bassi MR, Frank AM, Castberg FC, et al. Acquisition and decay of IgM and IgG responses to merozoite antigens after *Plasmodium falciparum* malaria in Ghanaian children. *Plos One.* 2020;15(12):e0243943.
61. Yman V, White MT, Asghar M, Sundling C, Sondén K, Draper SJ, et al. Antibody responses to merozoite antigens after natural *Plasmodium falciparum* infection: kinetics and longevity in absence of re-exposure. *BMC Med.* 2019;17(1):22.
62. White MT, Griffin JT, Akpogheneta O, Conway DJ, Koram KA, Riley EM, et al. Dynamics of the antibody response to *Plasmodium falciparum* infection in African children. *J Infect Dis.* 2014;210(7):1115–22.
63. Herman LS, Fornace K, Phelan J, Grigg MJ, Anstey NM, William T, et al. Identification and validation of a novel panel of *Plasmodium knowlesi* biomarkers of serological exposure. *Plos Negl Trop Dis.* 2018;12(6):e0006457.
64. Boddey JA, Moritz RL, Simpson RJ, Cowman AF. Role of the *Plasmodium* export element in trafficking parasite proteins to the infected erythrocyte. *Traffic.* 2009;10(3):285–99.
65. Guy AJ, Irani V, MacRaild CA, Anders RF, Norton RS, Beeson JG, et al. Insights into the immunological properties of intrinsically disordered malaria proteins using proteome scale predictions. *Plos One.* 2015;10(10):e0141729.
66. Akpogheneta OJ, Duah NO, Tetteh KK, Dunyo S, Lanar DE, Pinder M, et al. Duration of naturally acquired antibody responses to blood-stage *Plasmodium falciparum* is age dependent and antigen specific. *Infect Immun.* 2008;76(4):1748–55.
67. Crosnier C, Iqbal Z, Knuepfer E, Maciucă S, Perrin AJ, Kamuyu G, et al. Binding of *Plasmodium falciparum* merozoite surface proteins DBLMS1 and DBLMS2 to human immunoglobulin M is conserved among broadly diverged sequence variants. *J Biol Chem.* 2016;291(27):14285–99.
68. Quintana MP, Ecklu-Mensah G, Tcherniuk SO, Ditlev SB, Oleinikov AV, Hviid L, et al. Comprehensive analysis of Fc-mediated IgM binding to the *Plasmodium falciparum* erythrocyte membrane protein 1 family in three parasite clones. *Sci Rep.* 2019;9(1):6050.
69. Changrob S, Han J-H, Ha K-S, Park WS, Hong S-H, Chootong P, et al. Immunogenicity of glycosylphosphatidylinositol-anchored micronemal antigen in natural *Plasmodium vivax* exposure. *Malaria J.* 2017;16(1):348.
70. Wang B, Lu F, Cheng Y, Chen J-H, Jeon H-Y, Ha K-S, et al. Immunoprofiling of the tryptophan-rich antigen family in *Plasmodium vivax*. *Infect Immun.* 2015;83(8):3083–95.
71. Morais CG, Soares IS, Carvalho LH, Fontes CJF, Krettli AU, Braga EM. IgG isotype to C-terminal 19 kDa of *Plasmodium vivax* merozoite surface protein 1 among subjects with different levels of exposure to malaria in Brazil. *Parasitol Res.* 2005;95(6):420–6.
72. Weaver R, Reiling L, Feng G, Drew DR, Mueller I, Siba PM, et al. The association between naturally acquired IgG subclass specific antibodies to the PFRH5 invasion complex and protection from *Plasmodium falciparum* malaria. *Sci Rep.* 2016;6:33094.
73. Saavedra-Langer R, Marapara J, Valle-Campos A, Durand S, Vásquez-Chasnamote ME, Silva H, et al. IgG subclass responses to excreted-secreted

- antigens of *Plasmodium falciparum* in a low-transmission malaria area of the Peruvian Amazon. *Malaria J.* 2018;17(1):328.
74. Ferreira M, Kimura E, Katzin A, Santos-Neto L, Ferrari J, Villalobos J, et al. The IgG-subclass distribution of naturally acquired antibodies to *Plasmodium falciparum*, in relation to malaria exposure and severity. *Ann Trop Med Parasitol.* 1998;92(3):245–56.
 75. Tongren JE, Drakeley CJ, McDonald SL, Reyburn HG, Manjurano A, Nkya WM, et al. Target antigen, age, and duration of antigen exposure independently regulate immunoglobulin G subclass switching in malaria. *Infect Immun.* 2006;74(1):257–64.
 76. Stanisic DI, Richards JS, McCallum FJ, Michon P, King CL, Schoepflin S, et al. Immunoglobulin G subclass-specific responses against *Plasmodium falciparum* merozoite antigens are associated with control of parasitemia and protection from symptomatic illness. *Infect Immun.* 2009;77(3):1165–74.
 77. Ho F, Lortan JE, MacLennan IC, Khan M. Distinct short-lived and long-lived antibody-producing cell populations. *Eur J Immunol.* 1986;16(10):1297–301.
 78. Park JW, Moon SH, Yeom JS, Lim KJ, Sohn MJ, Jung WC, et al. Naturally acquired antibody responses to the C-terminal region of merozoite surface protein 1 of *Plasmodium vivax* in Korea. *Clin Diagn Lab Immunol.* 2001;8(1):14–20.
 79. Skountzou I, Satyabhama L, Stavropoulou A, Ashraf Z, Esser ES, Vassilieva E, et al. Influenza virus-specific neutralizing IgM antibodies persist for a lifetime. *Clin Vacc Immunol.* 2014;21(11):1481–9.
 80. Bohannon C, Powers R, Satyabhama L, Cui A, Tipton C, Michaeli M, et al. Long-lived antigen-induced IgM plasma cells demonstrate somatic mutations and contribute to long-term protection. *Nat Commun.* 2016;7:11826.
 81. Zenklusen I, Jongo S, Abdulla S, Ramadhani K, Lee Sim BK, Cardamone H, et al. Immunization of malaria-preexposed volunteers with PfSPZ vaccine elicits long-lived IgM invasion-inhibitory and complement-fixing antibodies. *J Infect Dis.* 2018;217(10):1569–78.
 82. Krishnamurthy AT, Thouvenel CD, Portugal S, Keitany GJ, Kim KS, Holder A, et al. Somatic hypermutated *Plasmodium*-specific IgM(+) memory B Cells are rapid, plastic, early responders upon malaria rechallenge. *Immunity.* 2016;45(2):402–14.
 83. MacNeil A, Reed Z, Rollin PE. Serologic cross-reactivity of human IgM and IgG antibodies to five species of Ebola virus. *Plos Negl Trop Dis.* 2011;5(6):e1175.
 84. Papa A, Karabaxoglou D, Kansouzidou AJV. Acute West Nile virus neuroinvasive infections: Cross-reactivity with dengue virus and tick-borne encephalitis virus. *J Med Virol.* 2011;83(10):1861–5.
 85. Baum E, Sattabongkot J, Sirichaisinthop J, Kiattibutr K, Davies DH, Jain A, et al. Submicroscopic and asymptomatic *Plasmodium falciparum* and *Plasmodium vivax* infections are common in western Thailand-molecular and serological evidence. *Malaria J.* 2015;14(1):95.
 86. Baum E, Sattabongkot J, Sirichaisinthop J, Kiattibutr K, Jain A, Taghavian O, et al. Common asymptomatic and submicroscopic malaria infections in Western Thailand revealed in longitudinal molecular and serological studies: a challenge to malaria elimination. *Malar J.* 2016;15(1):333.
 87. Longley RJ, França CT, White MT, Kumpitak C, Sa-angchai P, Gruszczyk J, et al. Asymptomatic *Plasmodium vivax* infections induce robust IgG responses to multiple blood-stage proteins in a low-transmission region of western Thailand. *Malaria J.* 2017;16(1):178.
 88. McLean AR, Boel ME, McGready R, Ataide R, Drew D, Tsuboi T, et al. Antibody responses to *Plasmodium falciparum* and *Plasmodium vivax* blood-stage and sporozoite antigens in the postpartum period. *Sci Rep.* 2016;6:32159.
 89. Min HMK, Changrob S, Soe PT, Han JH, Muh F, Lee S-K, et al. Immunogenicity of the *Plasmodium vivax* merozoite surface protein 1 paralog in the induction of naturally acquired antibody and memory B cell responses. *Malaria J.* 2017;16(1):354.
 90. Arévalo-Herrera M, Lopez-Perez M, Dotsey E, Jain A, Rubiano K, Felgner PL, et al. Antibody profiling in naïve and semi-immune individuals experimentally challenged with *Plasmodium vivax* sporozoites. *Plos Negl Trop Dis.* 2016;10(3):e0004563.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

