



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Lam, SK;Hao, X;Lin, E;Han, X;Norton, R;Mosier, AR;Seneweera, S;Chen, D

Title:

Effect of elevated carbon dioxide on growth and nitrogen fixation of two soybean cultivars in northern China

Date:

2012-07-01

Citation:

Lam, S. K., Hao, X., Lin, E., Han, X., Norton, R., Mosier, A. R., Seneweera, S. & Chen, D. (2012). Effect of elevated carbon dioxide on growth and nitrogen fixation of two soybean cultivars in northern China. *Biology and Fertility of Soils*, 48 (5), pp.603-606. <https://doi.org/10.1007/s00374-011-0648-z>.

Persistent Link:

<https://hdl.handle.net/11343/283117>

Cover page

Title: Effect of elevated carbon dioxide on growth and nitrogen fixation of two soybean cultivars in northern China

List of authors: Shu Kee Lam^a, Xingyu Hao^{b,c}, Erda Lin^b, Xue Han^b, Rob Norton^{a, d}, Arvin R. Mosier^a, Saman Seneweera^a and Deli Chen^{a*}

Institute or laboratory of origin:

^a Melbourne School of Land and Environment, The University of Melbourne, Victoria 3010, Australia

^b Key Laboratory of Ministry of Agriculture on Agro-environment and Climate Change; Agro-Environment and Sustainable Development Institute, Chinese Academy of Agricultural Sciences, Beijing 100081, China

^c College of Agronomy, Shanxi Agricultural University, Taigu 030801, Shanxi, China

^d International Plant Nutrition Institute, 54 Florence Street, Horsham, Victoria 3400, Australia

***Corresponding author:** Deli Chen (telephone: +61 3 83448148; fax: +61 3 83445037 and e-mail: delichen@unimelb.edu.au)

Abstract

The effect of elevated carbon dioxide (CO_2) concentration on symbiotic nitrogen (N_2) fixation in soybean under open-air conditions has not been reported. Two soybean cultivars (*Glycine max* (L.) Merr. cv. Zhonghuang 13 and cv. Zhonghuang 35) were grown to maturity under ambient ($415 \pm 16 \mu\text{mol mol}^{-1}$) and elevated ($550 \pm 17 \mu\text{mol mol}^{-1}$) [CO_2] at the Free-Air Carbon dioxide Enrichment (FACE) experimental facility in northern China. Elevated [CO_2] increased above- and below-ground biomass by 16-18% and 11-20%, respectively, but had no significant effect on the tissue C/N ratio at maturity. Elevated [CO_2] increased the percentage of N derived from the atmosphere (%Ndfa, estimated by natural abundance) from 59 to 79% for Zhonghuang 13, and the amount of N fixed from 166 to 275 kg N ha^{-1} , but had no significant effect on either parameter for Zhonghuang 35. These results suggest that variation in N_2 fixation ability in response to elevated [CO_2] should be used as key trait for selecting cultivars for future climate with respect to meeting the higher N demand driven by a carbon rich atmosphere.

Keywords

Free-air carbon dioxide enrichment (FACE); symbiotic nitrogen fixation; natural abundance; soybean cultivar

Introduction

The atmospheric carbon dioxide (CO₂) concentration has increased from 280 μmol mol⁻¹ in 1800 to 390 μmol mol⁻¹ now, and is expected to reach about 550 μmol mol⁻¹ in 2050 (Houghton et al. 2001). The response of symbiotic N₂ fixation in legumes to elevated [CO₂] is crucial to C sequestration in future CO₂ environments because continued C sequestration is contingent on additional N input (Hungate et al. 2003). A recent review suggests that symbiotic N₂ fixation at elevated [CO₂] could be enhanced as a result of [CO₂]-induced increase in nodule size, nodule number per plant, or specific nitrogenase activity (Rogers et al. 2009). Increased heterotrophic N₂ fixation in paddy soils under elevated [CO₂] has been attributed to enhanced nitrogenase activity (Das et al. 2011). Soybean (*Glycine max* (L.) Merr.) is an important oil seed crop legume and a major source of protein for human consumption and stockfeed. Soybean production and area harvested have increased by 7.6 and 3.8 times, respectively, during the past half century (Masuda and Goldsmith 2009). In closed environment studies, elevated [CO₂] increased soybean nodule dry mass and nitrogenase activity (Finn and Brun 1982; Matsunami et al. 2009; Miyagi et al. 2007; Serraj et al. 1998), but had no effect on the nodule number and mass under well-watered condition (Serraj et al. 1998). There are no reports on the effect of elevated [CO₂] on symbiotic N₂ fixation in soybean under fully open-air conditions. The present study was therefore conducted on a soybean field in a soybean-wheat rotation in Changping (40°10'N, 116°14'E),

Beijing, China using Free-Air Carbon Dioxide (FACE) facility to investigate the effect of elevated [CO₂] on the growth and N₂ fixation of two soybean cultivars.

Materials and Methods

The FACE system consists of 12 octagonal experimental areas, six elevated ($550 \pm 17 \mu\text{mol mol}^{-1}$) and six ambient ($415 \pm 16 \mu\text{mol mol}^{-1}$), each with a diameter of 4 m. The experimental areas were separated by at least 14 m to minimize cross-contamination of CO₂ between the areas. The soil (0-0.20 m) is a clay loam with a pH (soil:water ratio of 1:5) of 8.4 and contained 1.06% organic C, 0.11% total N, 25.5 mg kg⁻¹ mineral N (6.8 mg kg⁻¹ ammonium-N and 18.7 mg kg⁻¹ nitrate-N; 2M KCl extraction), 50 mg kg⁻¹ available phosphorus (Bray 1) and 140 mg kg⁻¹ ammonium acetate extractable potassium. The long term average rainfall and temperature during the soybean growing season are 475 mm and 21.3°C, respectively. The present experiment was conducted from mid June to October in 2009, under a randomized complete block design, with [CO₂] and cultivar as main effects with three replicates. The seeds of two cultivars of soybean (*Glycine max* (L.) Merr. cv. Zhonghuang 13 and cv. Zhonghuang 35) were sown on 17 June 2009 at row spacing of 0.5 m and with basal fertilizer application of diammonium phosphate and potassium chloride at 4.8 kg N ha⁻¹, 7.2 kg P ha⁻¹ and 37.3 kg K ha⁻¹. Zhonghuang 13 is a high-protein cultivar while Zhonghuang 35 a high-oil cultivar. Soybean seeds were not artificially inoculated according

to local farming practice. The plots were thinned after crop emergence to ensure uniform plant density. Irrigation of 37.5 mm equivalent rainfall was applied on 9 July to ease the drought stress. At full maturity (reproductive stage R8) on 6 October, 3 m² of soybean plants were harvested, and seven plants were randomly selected and the soil surrounding (10 cm radius) each plant was excavated to a depth of 25 cm to ensure all roots were recovered. Each of these seven plants was separated into above-ground and below-ground (including roots and nodules) parts, which were oven-dried at 70°C for 48 h, weighed, ground into powder and analyzed for total C (%), total N (%) and $\delta^{15}\text{N}$ values by isotope ratio mass spectrometry (IRMS) (Hydra 20-20, SerCon). All the above-ground parts harvested from the 3 m² were oven-dried at 70°C for 48 h to obtain yield and biomass in kg ha⁻¹. Below-ground biomass (in kg ha⁻¹) was calculated using the above/below-ground ratio of the seven plants. ¹⁵N natural abundance technique (Unkovich et al. 1997) was adopted for the calculation of the percentage of soybean N derived from the atmosphere (%Ndfa). Wheat (*Triticum aestivum* L.) plants harvested from each treatment area were used as reference plant materials, and their $\delta^{15}\text{N}$ values determined by IRMS for the calculation of %Ndfa. The amount of N fixed in above-ground parts was calculated as the product of %Ndfa and total N content of the above-ground parts. Data were analysed with MINITAB 14 statistical package using a General Linear Model analysis of variance.

Results and Discussion

Plant biomass and tissue quality

Elevated [CO₂] significantly increased above-ground (16-18%) and below-ground biomass (11-20%) for both cultivars (Table 1), which is comparable to the corresponding increases (17 and 30%) reported from the SoyFACE experiment in the US (Morgan et al. 2005; Rodriguez 2004). Under elevated [CO₂], there was a significant ($p < 0.05$) increase in the yield of Zhonghuang 35 from 2172 to 2744 kg ha⁻¹, and a trend of increase for Zhonghuang 13 from 2006 and 2089 kg ha⁻¹ ($p > 0.05$) (Hao et al. unpublished). The [CO₂]-induced increase in biomass has been attributed to a higher canopy C assimilation under elevated [CO₂] (Ainsworth et al. 2002). Elevated [CO₂] had no significant effect on the concentrations of total C and N of the above-ground and below-ground parts of both soybean cultivars at maturity and their C/N ratio (Table 1). Others also reported that the C/N ratio of soybean straw at maturity was unaffected by elevated [CO₂] (Prior et al. 2004; Torbert et al. 2004), although higher C/N ratio in fully expanded leaf was observed under elevated [CO₂] at vegetative and reproductive stages of soybean (Ainsworth et al. 2007). The increases in C accumulation (17%; $p < 0.05$) and N uptake (21%; $p < 0.05$) of the above-ground parts of soybean under CO₂ enrichment (Table 1) are consistent with Prévost et al. (2010). In the present study, the higher N acquisition of soybean due to the higher biomass accumulation

under elevated [CO₂] was probably met by symbiotically fixed N and N from the soil.

(Insert Table 1)

Nitrogen fixation

The %Ndfa value calculated for Zhonghuang 13 (69%) was higher ($p < 0.001$) than Zhonghuang 35 (27%) when averaged across [CO₂] treatments, corresponding to 220 and 71 kg N ha⁻¹ fixed from the atmosphere. The %Ndfa of the two cultivars is comparable to the upper (69%) and lower (36%) quartiles of a meta-analysis of N₂ fixation in soybean at ambient [CO₂] (Salvagiotti et al. 2008). The difference in %Ndfa between cultivars, which implies the variation in N₂ fixation ability among genotypes, was also reported in other soybean N₂ fixation studies at ambient [CO₂] (Danso et al. 1987; Houngnandan et al. 2008; Matsunami et al. 2009). Elevated [CO₂] increased %Ndfa for Zhonghuang 13 from 59 to 79% ($p = 0.07$) (Fig. 1a), and the amount of N fixed from 166 to 275 kg N ha⁻¹ ($p < 0.05$) (Fig. 1b), but had no significant effect on either parameter for Zhonghuang 35 (Fig. 1). In the present study, the soybean plants were not artificially inoculated but exposed to a mixture of rhizobacterial strains in the field. Difference in the efficiency and the response to elevated [CO₂] of these strains associated with the two cultivars could be attributed to differences in the establishment and functioning of an effective symbiosis, which depends on the genetic background of determinants in both host plant and rhizobacteria species (Keyser and Li 1992).

The increase in %Ndfa for Zhonghuang 13 under elevated [CO₂] was unlikely attributed to the improved water-use efficiency at elevated [CO₂] (Leakey et al. 2009) nor the tolerance of N₂ fixation to drought stress under elevated [CO₂] for soybean (Serraj 2003; Serraj et al. 1998) because water supply was substantiated by irrigation to ease drought stress for all treatments. The lack of response to elevated [CO₂] of N₂ fixation in Zhonghuang 35 suggests that this cultivar relied more on soil mineral N to meet the higher N demand under elevated [CO₂] than Zhonghuang 13. This implies the rhizobacterium associated with Zhonghuang 35 was more sensitive to the soil nitrate level (25.5 mg N kg⁻¹) at the site, as high mineral N level was reported to inhibit the nitrogenase activity of soybean nodules (Arrese-Igor et al. 1997; Kanayama et al. 1990; Streeter and Wong 1988).

(Insert Fig. 1)

In summary, elevated [CO₂] increased the above- and below-ground biomass of the two soybean cultivars tested, but there is genetic variability in symbiotic N₂ fixation between these cultivars in response to elevated [CO₂]. The factors manipulating the variation in N₂ fixation ability under elevated [CO₂], such as nodule development, root exudation, nitrogenase activity and related gene expression, need to be assessed to explore the underlying mechanisms for the variation. This could be exploited to improve N supply in cropping systems in future high carbon dioxide atmospheres.

Acknowledgements

This work was supported by the Australian Greenhouse Office, the Australian Centre for Agricultural Research (ACIAR) and a Riady Scholarship. The authors wish to thank Xiaogang Dong, Ji Gao, Zongpeng Yang and Qiubing Ye for field assistance, and Jianlei Sun for assistance with chemical analyses.

References

- Ainsworth EA, Davey PA, Bernacchi CJ, Dermody OC, Heaton EA, Moore DJ, Morgan PB, Naidu SL, Ra H-sY, Zhu X-g, Curtis PS, Long SP (2002) A meta-analysis of elevated [CO₂] effects on soybean (*Glycine max*) physiology, growth and yield. *Glob Change Biol* 8:695-709
- Ainsworth EA, Rogers A, Leakey ADB, Heady LE, Gibon Y, Stitt M, Schurr U (2007) Does elevated atmospheric [CO₂] alter diurnal C uptake and the balance of C and N metabolites in growing and fully expanded soybean leaves? *J Exp Bot* 58:579-591
- Arrese-Igor C, Minchin FR, Gordon AJ, Nath AK (1997) Possible causes of the physiological decline in soybean nitrogen fixation in the presence of nitrate. *J Exp Bot* 48:905-913
- Danso SKA, Hera C, Douka C (1987) Nitrogen fixation in soybean as influenced by cultivar and Rhizobium strain. *Plant Soil* 99:163-174
- Das S, Bhattacharyya P, Adhya TK (2011) Impact of elevated CO₂, flooding, and temperature interaction on heterotrophic nitrogen fixation in tropical rice soils. *Biol Fertil Soils* 47:25-30
- Finn GA, Brun WA (1982) Effect of atmospheric CO₂ enrichment on growth, nonstructural carbohydrate content, and root nodule activity in soybean. *Plant Physiol* 69:327-331
- Houghton JT, Ding Y, Griggs DJ, Nogueir M, van der Winden PJ, Dai X, Maskell K, Johnson CA (2001) *Climate Change 2001: the scientific basis. Contribution of Working Group*

I to the Third Assessment Report of the Intergovernmental Panel on Climate Change.

Cambridge University Press, Cambridge

Houngnandan P, Yemadje RGH, Oikeh SO, Djidohokpin CF, Boeckx P, Cleemput OV (2008)

Improved estimation of biological nitrogen fixation of soybean cultivars (*Glycine max*

L. Merrill) using ¹⁵N natural abundance technique. *Biol Fertil Soils* 45:175-183

Hungate BA, Dukes JS, Shaw MR, Luo Y, Field CB (2003) Nitrogen and climate change.

Science 302:1512-1513

Kanayama Y, Watanabe I, Yamamoto Y (1990) Inhibition of nitrogen fixation in soybean

plants supplied with nitrate I. nitrite accumulation and formation of

nitrosylhemoglobin in nodules. *Plant Cell Physiol* 31:341-346

Keyser HH, Li F (1992) Potential for increasing biological nitrogen fixation in soybean. *Plant*

Soil 141:119-135

Leakey ADB, Ainsworth EA, Bernacchi CJ, Rogers A, Long SP, Ort DR (2009) Elevated

CO₂ effects on plant carbon, nitrogen, and water relations: six important lessons from

FACE. *J Exp Bot* 60:2859-2876

Masuda T, Goldsmith PD (2009) World soybean production: area harvested, yield, and

long-term projections. *Int Food Agribus Manage Rev* 12:143-162

Matsunami T, Otera M, Amemiya S, Kokubun M, Okada M (2009) Effect of CO₂

concentration, temperature and N fertilization on biomass production of soybean

- genotypes differing in N fixation capacity. *Plant Prod Sci* 12:156-167
- Miyagi KM, Kinugasa T, Hikosaka K, Hirose T (2007) Elevated CO₂ concentration, nitrogen use, and seed production in annual plants. *Glob Change Biol* 13:2161-2170
- Morgan PB, Bollero GA, Nelson RL, Dohleman FG, Long SP (2005) Smaller than predicted increase in aboveground net primary production and yield of field-grown soybean under fully open-air [CO₂] elevation. *Glob Change Biol* 11:1856-1865
- Prévost D, Bertrand A, Juge C, Chalifour FP (2010) Elevated CO₂ induces differences in nodulation of soybean depending on bradyrhizobial strain and method of inoculation. *Plant Soil* 331:115-127
- Prior SA, Torbert HA, Runion GB, Rogers HH (2004) Elevated atmospheric CO₂ in agroecosystems: residue decomposition in the field. *Environ Manage* 33:S344-S354
- Rodriguez VV (2004) Soybean root biomass under elevated CO₂ and O₃ concentration subject to FACE conditions. Dissertation, University of Illinois at Urbana-Champaign
- Rogers A, Ainsworth EA, Leakey ADB (2009) Will elevated carbon dioxide concentration amplify the benefits of nitrogen fixation in legumes? *Plant Physiol* 151:1009-1016
- Salvagiotti F, Cassman KG, Specht JE, Walters DT, Weiss A, Dobermann A (2008) Nitrogen uptake, fixation and response to fertilizer N in soybeans: A review. *Field Crops Res* 108:1-13
- Serraj R (2003) Atmospheric CO₂ increase benefits symbiotic N₂ fixation by legumes under

drought. *Curr Sci* 85:1341-1343

Serraj R, Sinclair TR, Allen LH (1998) Soybean nodulation and N₂ fixation response to drought under carbon dioxide enrichment. *Plant Cell Environ* 21:491-500

Streeter J, Wong PP (1988) Inhibition of legume nodule formation and N₂ fixation by nitrate. *Crit Rev Plant Sci* 7:1-23

Torbert HA, Prior SA, Rogers HH, Runion GB (2004) Elevated atmospheric CO₂ effects on N fertilization in grain sorghum and soybean. *Field Crops Res* 88:57-67

Unkovich MJ, Pate JS, Sanford P (1997) Nitrogen fixation by annual legumes in Australian Mediterranean agriculture. *Aust J Agric Res* 48:267-293

Figure caption

Fig. 1 Effect of elevated [CO₂] on (a) %Ndfa and (b) amount of N fixed in the above-ground parts of two soybean cultivars. Each data point represents the mean of three replicates. Vertical bar indicates least significant difference (LSD) ($p = 0.05$)




Fig. 1

Table

Table 1 Effect of elevated [CO₂] on the biomass, total C and N, C/N ratio, biomass C and N uptake of two soybean cultivars at full maturity.

Treatment	Biomass (kg ha ⁻¹)	Total C (%)	Total N (%)	C/N	Biomass C (kg ha ⁻¹)	N uptake (kg ha ⁻¹)
Above-ground parts						
Zhonghuang 13						
Ambient [CO ₂]	7684 (146)	46.2 (0.3)	3.69 (0.36)	12.7 (1.1)	3549 (51)	283 (22)
Elevated [CO ₂]	9092 (255)	47.0 (1.0)	3.84 (0.13)	12.2 (0.2)	4269 (129)	349 (16)
Zhonghuang 35						
Ambient [CO ₂]	7645 (834)	47.9 (0.2)	3.25 (0.11)	14.8 (0.5)	3659 (382)	249 (30)
Elevated [CO ₂]	8840 (223)	47.5 (0.6)	3.34 (0.10)	14.2 (0.3)	4197 (147)	295 (16)
[CO ₂]	*	NS	NS	NS	*	*
Cultivar	NS	NS	*	**	NS	NS
[CO ₂] × Cultivar	NS	NS	NS	NS	NS	NS
Below-ground parts						
Zhonghuang 13						
Ambient [CO ₂]	792 (28)	44.8 (0.3)	0.64 (0.10)	74.5 (13.5)	355 (14)	5.08 (0.91)
Elevated [CO ₂]	880 (55)	43.9 (0.0)	0.70 (0.05)	63.0 (4.4)	386 (24)	6.24 (0.76)
Zhonghuang 35						
Ambient [CO ₂]	764 (32)	45.6 (0.2)	0.54 (0.03)	85.4 (4.6)	348 (16)	4.09 (0.26)
Elevated [CO ₂]	917 (22)	44.8 (0.9)	0.53 (0.07)	86.7 (11.2)	410 (9)	4.90 (0.67)
[CO ₂]	*	NS	NS	NS	*	NS
Cultivar	NS	NS	NS	NS	NS	NS
[CO ₂] × Cultivar	NS	NS	NS	NS	NS	NS

Values are means of the three replicates (SE) for each treatment. Significant effects are

indicated as $*p < 0.05$ and $**p < 0.01$. NS, not significant.

The values of biomass C and N uptake reported in this table were calculated before rounding of decimal place. They are not necessarily the perfect product of the biomass and %C and %N listed in this table.