



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Mariani, M;Fletcher, MS;Holz, A;Nyman, P

Title:

ENSO controls interannual fire activity in southeast Australia

Date:

2016-10-28

Citation:

Mariani, M., Fletcher, M. S., Holz, A. & Nyman, P. (2016). ENSO controls interannual fire activity in southeast Australia. *Geophysical Research Letters*, 43 (20), pp.10-900. <https://doi.org/10.1002/2016GL070572>.

Persistent Link:

<https://hdl.handle.net/11343/291858>

ENSO controls inter-annual fire activity in southeast Australia

Mariani M.¹, Fletcher M.-S.¹, Holz A.², Nyman P.³

¹ School of Geography, University of Melbourne, Australia

² Department of Geography, School of the Environment, Portland State University, Portland, OR, USA

³ School of Ecosystem and Forest Science, University of Melbourne, Australia

ABSTRACT

El Niño Southern Oscillation (ENSO) is the main mode controlling the variability in the ocean-atmosphere system in the South Pacific. While the ENSO influence on rainfall regimes in the South Pacific is well documented, its role in driving spatiotemporal trends in fire activity in this region has not been rigorously investigated. This is particularly the case for the highly flammable and densely populated southeast Australian sector, where ENSO is a major control over climatic variability. Here we conduct the first region-wide analysis of how ENSO controls fire activity in southeast Australia. We identify a significant relationship between ENSO and both fire frequency and area burnt. Critically, wavelet analyses reveal that, despite substantial temporal variability in the ENSO system, ENSO exerts a persistent and significant influence on southeast Australian fire activity. Our analysis has direct application for developing robust predictive capacity for the increasingly important efforts at fire management.

INTRODUCTION

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1002/2016GL070572](https://doi.org/10.1002/2016GL070572)

Fire is a ubiquitous Earth system process that began soon after the appearance of terrestrial plants 420 million years ago [Scott & Glasspool, 2006]. At the global scale, fire affects ecosystem patterns and processes, determines vegetation distribution [Bond & Keeley, 2005; Bond *et al.*, 2005], impacts the climate system [Bowman *et al.*, 2009] and contributes to the carbon cycle [Liu *et al.*, 2015; Santín *et al.*, 2016]. Moreover, climate-driven changes in fire activity are a significant threat for human populations living in flammable biomes [e.g. Parisien *et al.*, 2016] and for ecosystem services and function [Calder *et al.*, 2015]. The predicted increase in fire activity in many regions on Earth in response to climate change [Parisien and Moritz, 2009; Moritz *et al.*, 2012; Westerling *et al.*, 2006], therefore, represents a significant challenge for attempts of a sustainable management of the Earth system. Despite the clear importance of fire and recognition of climate as a key component controlling this process, the relationship between climate and fire is still poorly understood in many regions on Earth. In southeast Australia, one of the most fire-prone areas on Earth [Hennessey *et al.*, 2005], the main climate mode controlling key determinants of fire weather (moisture and temperature variability) is the El Niño Southern Oscillation (ENSO) [Risbey *et al.*, 2009]. In general, El Niño events, the warm phase of ENSO, starve southeast Australia of rainfall and promote drought and wildfire [Kiem and Franks, 2004; Verdon *et al.*, 2004; Nicholls and Lucas, 2007; Murphy and Timbal, 2008; Hill *et al.*, 2009]. The current amplification of El Niño activity due to anthropogenic climate change [Power *et al.*, 2013], then, heralds a serious threat to southeast Australia's water security, remnant fire-sensitive ecosystems and the ever-expanding flammable bush-urban interface [Guilyardi 2006; Lenton *et al.* 2008]. Here, we present a novel analysis of climate-fire dynamics in southeast Australia in an attempt to elucidate the dominant climatic drivers of fire in both space and time in this heavily populated region.

Author Manuscript

An enhanced understanding of the climatic precursors of increased fire activity can lead to improved predictive power [e.g. Verdon et al. 2004; Nicholls and Lucas 2007; Harris et al., 2014]. This is particularly important for areas such as southeast Australia where the quasi-periodic frequency of oscillations in key climate phenomena, such as ENSO, make fire prediction a difficult task. ENSO is a particularly important control over fire regimes across the entire Pacific region, and the spatiotemporal variability of ENSO is evident from the range of analyses that attempt to elucidate the role of ENSO in governing fire activity. Indeed, significant correlation between ENSO and fire activity has been reported for the Florida everglades between 1948–99 [Beckage et al., 2003], California during the last 150 years [Herweijer et al., 2007], tropical Mexico between 1984–99 [Román-Cuesta et al., 2003], southwest United States between 1700–1905 [Swetnam and Betancourt, 1990], Indonesia over the last 250 years [van der Kaars et al., 2010] and southern South America over the most recent centuries [Veblen et al., 1999; Holz and Veblen, 2012; Holz et al., 2012a]. While there is evidence that El Niño events increase the chance of fire weather in southeast Australia [e.g. Fox-Hughes et al., 2014; Grose et al., 2014; Williams et al., 2001], few analyses from this region assess the influence of ENSO over actual fire occurrence (area burnt and/or number of fires). Of these, anecdotal evidence of an ENSO control over fire activity has been reported from the island of Tasmania, where fires over the historical period are significantly related to sea surface temperatures (SSTs) in the Coral Sea, a location influenced by ENSO [Nicholls & Lucas, 2007]. More concrete evidence comes from the south coast of the region (state of Victoria), where a significant correlation between ENSO and both area burnt and number of fires has been reported between 1972–2012 [Harris et al., 2014]. While ENSO is most often considered the dominant control over southeast Australian climate and fire activity, the Indian Ocean Dipole (IOD) and the Southern Annular Mode (SAM), both important controls over climatic variability in this region, have been shown to influence

fire activity within the southeast Australian sector [e.g. Cai et al., 2009; Mariani and Fletcher, 2016], suggesting that multiple modes of climatic variability must be considered when attempting to determine the drivers of fire activity in this region.

A critical limitation of previous analyses of ENSO-fire relationships in southeast Australia is the lack of a regional synthesis of actual fire activity conducted through a rigorous climatological framework. A case-in-point is the work on the role of ENSO in governing fire activity in Tasmania [Nicholls & Lucas, 2007]. Tasmania is a topographically complex mid-latitude island in which the climatic determinants of fire weather and activity are polarised between ENSO in the north and east and the SAM in the south and west [Mariani and Fletcher, 2016]. SAM relates to changes in the position and intensity of the circumpolar southern westerlies, a system that, while periodically synergistic with ENSO through time [Fogt et al., 2011], primarily acts independently to ENSO. Indeed, recent evidence reveals that fire activity in the south and west of Tasmania is determined by SAM, with ENSO having little explanatory power over fire activity through the recent and deeper past [Mariani & Fletcher 2016]. Further, analyses of the climatic determinants of fire activity in southeast Australia have seldom employed actual fire data [Harris et al., 2014; Nicholls & Lucas 2007; Mariani & Fletcher 2016], rather most employ indices of fire weather to ascertain the link between climate and the potential for fire occurrence [Fox-Hughes et al., 2014; Grose et al., 2014; Williams et al., 2001]. Moreover, of those that do employ actual fire occurrence data, only those focused on the island of Tasmania exclude planned burns from their analysis [Nicholls & Lucas, 2007; Mariani & Fletcher 2016], an important source of noise that reduces the efficacy of the fire occurrence dataset for correlation with climate variability.

Here, we present a multi-scale analysis of the role of ENSO in governing actual fire activity in southeast Australia. Our specific aims are to: 1) explore the relationship between ENSO and fire occurrence in the densely populated and highly flammable southeast of Australia; 2) synthesize the seasonal importance of ENSO in determining inter-annual fire activity in this region and 3) test whether or not ENSO and fire occurrences have coherent time-frequency domains. We conducted our climate-fire analysis using the Southern Oscillation Index (SOI), an index of ENSO variability that strongly affects precipitation in southeast Australia [Risbey et al., 2009]. We employed observed fire history data filtered to include un-planned fires only and conducted correlation function and wavelet analyses to determine long-term seasonal-scale associations and time frequencies coherence between SOI and fire activity through time.

METHODS

To visualize the spatial pattern of the influence of ENSO on southeast Australian precipitation we created a correlation map between annual and seasonal precipitation totals and the Southern Oscillation Index (SOI). SOI is calculated as the normalised difference between mean sea level pressure at Tahiti and Darwin and it is negative during El Niño events and positive during La Niña events [Trenberth, 1984]. Although there are several indices for ENSO (e.g. Niño-4, Niño-3.4, ENSO-Modoki Index –EMI), we used the SOI because has shown to have the highest correlation values with rainfall in Australia compared to the other indices [Risbey et al., 2009]. This is probably because the SOI is more closely related to the rainfall process in southeast Australia through its relationship with large-scale surface pressure, compared to ocean-based indices which rely on sea surface temperature (SST) [Risbey et al., 2009]. We calculated Pearson correlation coefficients (r) between

seasonal/annual rainfall during the period 1961–1990 for 1208 meteorological stations across the southeast Australia and the seasonal/annual SOI. Precipitation data was obtained from the Australian Bureau of Meteorology (BOM). The r values from the stations were spatially interpolated using the Universal Kriging method in ArcMap 10.2 [ESRI-Environmental Systems Resource Institute, 2011, Redlands, California].

To account for fuel limitations, an essential parameter in determining fire activity [Krawchuck et al. 2009; Cochrane, 2003; Pausas and Ribeiro, 2013; Bradstock, 2010; McWethy et al., 2013; Boer et al., 2016], we constrain our analysis to the cool temperate forest biome of southeast Australia (latitude between 28°S and 44°S; longitude between 140°E and 155°E), where biomass is considered to be always abundant and in which climate is the principal determinant of temporal changes in fire activity [Bradstock et al., 2014]. This screening removed fires from the Australian Alps, as much of this landscape approaches biomass limitation with respect to fire [Bradstock et al., 2014]. Observed fire occurrence data were obtained through local administrative databases and span the period between the fire seasons of 1951/1952 to 2013/2014 (hereon, we define a fire-season year by the year in which the fire season ends – e.g. fire season 1951/1952 = 1952). We determine the “fire season” as the period between December and March, during which the majority of fire events occur [Williamson et al., 2016]. For Tasmania, only the eastern and north side of the state was considered, as fires activity and precipitation in the west of this island have been linked to variability in the SAM, with no identified relationship between fire activity and ENSO [Mariani and Fletcher, 2016].

Fire activity is represented in this study as two metrics: number of occurrences and area burnt, representing different components of a fire regime. Area burned represents the outcome of a fire season and includes ignitions as well as fire weather and fuels that determine fire spread. Fire

occurrence is mainly linked to ignitions and is more sensitive to artefacts linked to detection. The fire activity data (number of fires and area burnt) were first filtered to include all fires except deliberate management fires and then normalized by the standard deviations (z-scores). Z-scores values higher (lower) than 0.5 (-0.5) were used to identify significant “fire years” (“non-fire years”) [Mariani and Fletcher, 2016]. The list of years used in the analysis is shown in the Supplementary Table S1. A persistent increase in the number of fires toward the present likely reflects improving detection and recording through time, thus, we applied linear detrending to the time series to extract the inter-annual variability. Figure 1 presents the location of all fires recorded in southeast Australia from 1951 to 2014.

To identify a relationship between the annual and seasonal SOI and fire activity in the southeast Australia, we performed Superposed Epoch Analysis (SEA) analysis in R [R core team, 2012 - dplR package]. This analysis allows for assessing the significance of the departure from the mean for a given set of key event years (e.g. fire years) and lagged years using bootstrapped confidence intervals [Lough and Fritts, 1987]. To analyse the relationship between fire activity and the SOI, we used significant “fire years” and “non-fire years” during the 1951-2014 period and annual and seasonal SOI values. The seasonal indices of SOI were defined as: Winter (June-July-August of the fire season end year), Spring (September-October-November of the fire season end year), Summer (December of the previous year and January and February of the fire season end year), Autumn (March-April-May of the fire season end year). Given the importance of the SAM and IOD in the climate/fire dynamics of the Southern Hemisphere [e.g. Cai et al., 2009; Risbey et al., 2009; Hill et al., 2009; Garreaud et al., 2009; Mariani and Fletcher, 2016], the same analysis using SEA has been also

conducted with these climate modes to test whether they have a significant relationship with fire activity in the study region.

To test whether the fire frequency and area burnt and the SOI show a similar periodicity, wavelet coherence analysis was undertaken using the `wt()` and `wtc()` functions respectively in the R package “Biwavelet” [Gouhier, 2013]. For this analysis, the spring-summer period was chosen as it represents the season with the highest significant correlation (p value <0.05) between the SOI and the two fire activity metrics (Supplementary Table S2). Wavelet analyses have often been applied to climate data [e.g., Meyers et al. 1993; Lau and Weng 1995; Wang and Wang 1996] and provides a useful tool to reveal frequency localization in climate signals in time. The Morlet continuous wavelet transform was applied and the data were padded with zeros at each end to reduce wraparound effects [Torrence and Webster, 1999].

RESULTS

A total of 25,390 un-planned fires with a total area of 37,291,374 ha were recorded in the temperate biome region of southeast Australia during the period 1951–2014 (Figure 1a). A total of 8 significant fire years were determined for fire occurrence, whereas 22 significant fire years were found using the area burnt metric (Figure 2c,d). The total of significant non-fire years was 10 and 19 for the number of fires and area burnt, respectively (Figure 2c,d). The list of years used in the SEA is presented in Supplementary Table S1. The spatial climate correlation analysis shows a complex pattern of correlation between SOI and rainfall anomalies across southeast Australia using annual and seasonal values (Figure 1b-f). Strongest positive correlations between SOI and rainfall across most of the study region occur in winter and spring, while weak negative correlations occur in

summer and autumn. An exception to this is the northeast of the study region, in which the strongest positive correlations occur during summer and a neutral/negative correlation is present during winter and spring (Figure 1c,d). Together, this variable seasonal pattern renders the annual SOI-rainfall correlations signature near neutral in most of the region, except north-eastern Tasmania (Figure 1a). Importantly, negative correlations shown in the maps are always weak (i.e. r value < -0.3 ; p -value > 0.05), indicating a probable non-significance of these correlations during the reference period (1961-1990).

Testing both number of fires and area burnt metrics, the SEA revealed a statistically significant (p value < 0.05) negative SOI departure (El Niño conditions) occurring in winter and spring of the preceding year and during summer and autumn of the same year (Figure 3a,b), i.e. fire activity is significantly related to SOI in the full seasonal cycle leading up to and including a fire season. To support this result, we show that non-fire years (fire seasons with an anomalously low fire occurrence) correspond to a significant (p value < 0.05) positive departure in SOI (La Niña conditions) during the same seasons (Figure 3a,b). At an inter-annual scale, significant negative annual SOI departures (El Niño) were found during the fire years using the number of fires, whereas area burnt shows negative but not significant departures for annual SOI (Figure 3b). During the non-fire years for area burnt and number of fires, a significant positive SOI departure (La Niña) was found at lag zero. The analyses of both SAM and IOD seasonal and annual indices with the two fire activity metrics did not produce significant results (see Supplementary Figure S1 and S2), while analyses using both Niño 3.4 and the EMI are consistent with the SOI results, thus reinforcing the robustness of our results (Supplementary Figure S3 and S4).

Wavelet coherence indicates that negative SOI (El Niño) occurred in concert with positive anomalies in fire occurrence and area burnt at the respective time frequency during the time periods (Figure 4). The number of fires metric shows a strong anti-correlated coherence pattern (i.e. a leftward arrow indicated x and y are anti-correlated) with the spring-summer SOI with a short period frequency (1-4 years) between 1963 and 1978, while area burnt shows a strong anti-phase frequency between 2-5 years over the same 1963-1978 period. During the 1998-2010 period a strong anti-phase coherence pattern with the spring-summer SOI with a periodicity between 1-5 years is observed in the area burnt coherence plot. A persistent strong anti-phase coherence pattern with periodicity between 10-18 years was found both in the area burnt and fire occurrence metrics throughout the entire analysis period. Areas outside the 'cone of influence', where edge effects are present, have not been considered for the interpretation of the results.

DISCUSSION

Our analysis constitutes the first synthesis of the fire-climate teleconnections over the last 60 years across temperate southeast Australia, a highly flammable and densely populated region that is subject to frequent catastrophic wildfires that have major societal, cultural and economic impacts. The SEA clearly indicates a multi-seasonal control of ENSO on fire activity in this region, most likely due to ENSO related precipitation anomalies. The significant negative departure of SOI during winter and spring of the year preceding the fire season (Figure 3), and the strong relationship between precipitation and ENSO (Figure 1 and 2), indicates that a reduction in water availability in this period is crucial for successful ignition during the following fire season. Likewise, another important factor emerging from this analysis is the ENSO state occurring during the fire season: an El Niño condition present during summer and autumn is more likely to result in an increase in both number of fires

and area burnt. Moreover, significant non-fire years are correlated with La Niña (high moisture) conditions during the preceding winter and spring and current summer and autumn (Figure 3a,b). Interestingly, the results for non-fire years using the number of fires show a significant SOI positive departure at lag -1 and 0. This may relate to the general persistence of a La Niña-conditions during consecutive years (Figure 2b). Additionally, we find no evidence for a dominant role of the IOD in governing fire activity across the temperate forest biome of southeast Australia through our analysis period, in direct contrast to the study of Cai et al. (2009). The lack of consistency between our results and that of Cai et al (2009) likely results from the low number of fire seasons ($n=21$), narrow analysis period (1950-2008) and use of spring-summer Niño 3.4 ENSO index, an index shown to have lower correlation values with Australian rainfall relative to the SOI [Risbey et al., 2009], in the latter analysis.

The strong coherency we have identified in our wavelet analysis at a 12 year periodicity between spring-summer El Niño events and fire activity (area burnt and fire number) throughout most of the analysis period (1963-2000) (Figure 4) indicates, for the first time, a clear and persistent decadal-scale modulation of fire activity in southeast Australia by ENSO. Given the irregular frequency found in the ENSO time-series [Soon-II and Wang, 1999; Wang et al., 2001], it is remarkable that the observed decadal-scale modulation of fire activity by ENSO is fixed throughout the analysis period, highlighting the persistently dominant influence of ENSO over the climate of this highly flammable region, also given the predicted increase in extreme El Niños by the end of this century [Cai et al., 2014]. Further, the wavelet analysis also reveals a clear correlation between temporal shifts in ENSO variability and actual fire activity in southeast Australia. The higher frequency coherence (1-5 years) between 1963-1978 also corresponds to a period of maximum ENSO frequency [Soon-II and Wang,

1999]. Thus, our results reveal that southeast Australia is highly sensitive to shifts in the frequency of variability in the ENSO system and underscores the potential threat of increased fire activity in this region in response to the recent intensification of the ENSO system [Power et al., 2013]. Interestingly, there is an apparent decoupling between the coherence of area burnt and number of fires and the SOI (Figure 4) – i.e. ENSO seems having a control on the spread of fires between 1995-2003 but another factor is controlling the ignition during this period. It is possible that cultural factors, such as increased awareness, predictive capacity and subsequent fire-suppression efforts (e.g. construction of fire breaks, increased hazard reduction burning and increased public awareness) actively suppressed ignition events in the latter part of the record. Importantly, the impact of ENSO on preconditioning the landscape for fire spread once a fire starts appears to have been less affected by these cultural adaptations.

Models of future fire activity predict an increase in fire activity under projections of climate change scenarios in temperate forest biomes, such as southeast Australia [Moritz et al., 2012]. Moreover, global temperature increase has the potential to alter the thermal balance in the equatorial Pacific Ocean, driving an increase in the strength and frequency of El Niño events [Timmermann et al., 1999; Guilyardi 2006], a process that is, indeed, already underway (Power et al 2013). Our identification of both a persistent and non-stationary influence of ENSO over fire activity in southeast Australia has direct and significant implications for managing ecosystem processes and human populations within the ENSO climate domain. Moreover, extreme variability in the Australian climate complicates attempts to understand the impacts of anthropogenic climate change [Jolly et al., 2015], and our analysis provides information on the magnitude and temporal variability of climate-driven fire activity that will significantly enhance the detection of persistent changes in a

highly variable system. Our identification of a multi-seasonal link between supra-annual ENSO dynamics and fire activity in southeast Australia reveals that the relationship between ENSO and fire must be considered at multiple scales and for all seasons when planning for future fire activity in this densely populated region.

Our results provide additional evidence for the long-term influence of ENSO over the Earth system. Over the past millennia, ENSO has been implicated in a variety of processes that includes terrestrial ecosystem dynamics in eastern Australia [Donders et al., 2007] and southwest Tasmania [Fletcher et al., 2014; Fletcher et al., 2015]; storm patterns in New Zealand [Gomez et al., 2011], coral reef systems in the eastern Pacific [Toth et al., 2012]; human cultural change [Sandweiss, 2001; Magilligan and Goldstein, 2001; Turney et al., 2006] and fire activity in temperate southern South America [Whitlock et al., 2007; Holz et al., 2012b] and Tasmania [Fletcher et al., 2015; Rees et al., 2015]. Importantly, modern ENSO variability is muted, relative to the last few millennia [e.g. Moy et al., 2002; Conroy et al., 2008; Yan et al., 2011], and it is critical that we gain an understanding of how ENSO variability drives changes locally, regionally and globally if we are to sustainably manage the Earth systems.

CONCLUSION

Under a warming world scenario and a predicted future increase in fire activity across temperate biomes [Moritz et al., 2012] and ENSO activity is projected to intensify [Timmermann et al., 1999; Guilyardi, 2006; Lenton et al., 2008; Power et al., 2013; Kim et al., 2014]. This research constitutes the first attempt in disentangling the role of SOI in driving fire activity across the entire south-

eastern Australian region during the past 60 years using observed wildfire activity datasets. We reveal that El Niño phases (negative SOI) are significantly linked with fire occurrence (number of fires) and area burnt in this region at the inter-annual scale throughout all seasons (i.e. winter and spring of the year preceding the fire season and during summer and autumn of the fire season year), and a coherent pattern of decadal scale frequencies over the spring-summer period. The results obtained in this study reveal the seasonal ENSO states as important parameters to consider in fire forecasting and in ecological and palaeoecological interpretations.

ACKNOWLEDGEMENTS

Research was supported by ARC grants DI110100019 and IN140100050. Andrés Holz was in part supported by the Australian Research Council (Grant # DP110101950) and the US National Science Foundation Awards (Grant # 0966472). We thank the Victorian Department of Environment, Land, Water and Planning, the New South Wales Office of Environment and Heritage, the ACT Parks and Wildfire and TasList (Government of Tasmania) for providing the fire occurrence data. We also thank Karl Braganza and Alex Evans from the Australian Bureau of Meteorology for providing the rainfall data from all the stations across southeast Australia. Data elaborated with these analyses are available upon request to MM (mmariani@student.unimelb.edu.au).

REFERENCES

- An, S.-I., and B. Wang (2000), Interdecadal change of the structure of the ENSO Mode and its impact on the ENSO frequency*, *Journal of Climate*, 13(12), 2044-2055.
- Ashok, K., S. K. Behera, S. A. Rao, H. Weng, and T. Yamagata (2007), El Niño Modoki and its possible teleconnection, *Journal of Geophysical Research: Oceans*, 112(C11).

Beckage, B., W. J. Platt, M. G. Slocum, and B. Panko (2003), Influence of the El Niño Southern Oscillation on fire regimes in the Florida Everglades, *Ecology*, 84(12), 3124-3130.

Boer, M. M., D. M. Bowman, B. P. Murphy, G. J. Cary, M. A. Cochrane, R. J. Fensham, M. A. Krawchuk, O. F. Price, V. R. De Dios, and R. J. Williams (2016), Future changes in climatic water balance determine potential for transformational shifts in Australian fire regimes, *Environmental Research Letters*, 11(6), 065002.

Bond, W. J., and J. E. Keeley (2005), Fire as a global 'herbivore': the ecology and evolution of flammable ecosystems, *Trends in ecology & evolution*, 20(7), 387-394.

Bond, W. J., F. I. Woodward, and G. F. Midgley (2005), The global distribution of ecosystems in a world without fire, *New Phytologist*, 165(2), 525-537.

Bowman, D. M. J. S., et al. (2009), Fire in the Earth System, *Science*, 324(5926), 481-484.

Bradstock, R., T. Penman, M. Boer, O. Price, and H. Clarke (2014), Divergent responses of fire to recent warming and drying across south-eastern Australia, *Global change biology*, 20(5), 1412-1428.

Bradstock, R. A. (2010), A biogeographic model of fire regimes in Australia: current and future implications, *Global Ecology and Biogeography*, 19, 145-158.

Cai, W., T. Cowan, and M. Raupach (2009), Positive Indian Ocean dipole events precondition southeast Australia bushfires, *Geophysical Research Letters*, 36(19).

Cai, W., S. Borlace, M. Lengaigne, P. Van Rensch, M. Collins, G. Vecchi, A. Timmermann, A. Santoso, M. J. McPhaden, and L. Wu (2014), Increasing frequency of extreme El Niño events due to greenhouse warming, *Nature climate change*, 4(2), 111-116.

Calder, W. S., D. Parker, C. J. Stopka, G. Jiménez-Moreno, and B. N. Shuman (2015), Medieval warming initiated exceptionally large wildfire outbreaks in the Rocky Mountains, *Proceedings of the National Academy of Sciences*, 112(43), 13261-13266.

Cochrane, M. A. (2003), Fire science for rainforests, *Nature*, 421(6926), 913-919.

Conroy, J. L., J. T. Overpeck, J. E. Cole, T. M. Shanahan, and M. Steinitz-Kannan (2008), Holocene changes in eastern tropical Pacific climate inferred from a Galápagos lake sediment record, *Quaternary Science Reviews*, 27(11-12), 1166-1180.

Donders, T. H., S. Haberle, G. Hope, F. Wagner, and H. Visscher (2007), Pollen evidence for the transition of the Eastern Australian climate system from the post-glacial to the present-day ENSO mode, *Quaternary Science Reviews*, 26, 1621-1637.

ESRI 2011. ArcGIS Desktop: Release 10. Redlands, CA: Environmental Systems Research Institute.

Fletcher, M.-S., A. Benson, H. Heijnis, P. S. Gadd, L. C. Cwynar, and A. B. H. Rees (2015), Changes in biomass burning mark the onset an ENSO-influenced climate regime at 42°S in southwest Tasmania, Australia, *Quaternary Science Reviews*, 122(0), 222-232.

Fletcher, M.-S., B. B. Wolfe, C. Whitlock, D. Pompeani, H. Heijnis, S. Haberle, P. Gadd, and D. M. J. S. Bowman (2014), The legacy of mid Holocene fire on a Tasmanian montane landscape, *Journal of Biogeography*, 41.3 (2014): 476-488.

Fogt, R. L., D. H. Bromwich, and K. M. Hines (2011), Understanding the SAM influence on the South Pacific ENSO teleconnection, *Climate Dynamics*, 36(7-8), 1555-1576.

Fox-Hughes, P., R. Harris, G. Lee, M. Grose, and N. Bindoff (2014), Future fire danger climatology for Tasmania, Australia, using a dynamically downscaled regional climate model, *International journal of wildland fire*, 23(3), 309-321.

Garreaud, R. D., M. Vuille, R. Compagnucci, and J. Marengo (2009), Present-day South American climate, *Palaeogeography Palaeoclimatology Palaeoecology*, 281(3-4), 180-195.

Gomez, B., J. Carter, A. R. Orpin, K. M. Cobb, M. J. Page, N. A. Trustrum, and A. S. Palmer (2011), ENSO-SAM interactions during the middle and late Holocene, *The Holocene*, 0959683611405241.

Gouhier, T. (2013), Biwavelet: conduct univariate and bivariate wavelet analyses (ver. 0.14), edited.

Guilyardi, E. (2006), El Niño–mean state–seasonal cycle interactions in a multi-model ensemble, *Climate Dynamics*, 26(4), 329-348.

Grose, M. R., P. Fox-Hughes, R. M. Harris, and N. L. Bindoff (2014), Changes to the drivers of fire weather with a warming climate—a case study of southeast Tasmania, *Climatic Change*, 124(1-2), 255-260.

Harris, S., N. Nicholls, and N. Tapper (2014), Corrigendum to: Forecasting fire activity in Victoria, Australia, using antecedent climate variables and ENSO indices, *International journal of wildland fire*, 23(2), 290-293.

Hennessey, K., C. Lucas, N. Nicholls, J. Bathols, R. Suppiah, and J. Ricketts (2005), Climate change impacts on fire-weather in south-east Australia, pp.88

Hill, K. J., A. Santoso, and M. H. England (2009), Interannual Tasmanian Rainfall Variability Associated with Large-Scale Climate Modes, *Journal of Climate*, 22, 4383-4397.

Holz, A., and T. T. Veblen (2012), Wildfire activity in rainforests in western Patagonia linked to the Southern Annular Mode, *International Journal of Wildland Fire*, 21(2), 114-126.

Holz, A., T. Kitzberger, J. Paritsis, and T. T. Veblen (2012), Ecological and climatic controls of modern wildfire activity patterns across southwestern South America, *Ecosphere*, 3(11), 1-25.

Holz, A., S. Haberle, T. Veblen, R. D. Pol-Holz, and J. Southon (2012), Fire history in western Patagonia from paired tree-ring fire-scar and charcoal records, *Climate of the Past*, 8(2), 451-466.

Jolly, W. M., M. A. Cochrane, P. H. Freeborn, Z. A. Holden, T. J. Brown, G. J. Williamson, and D. M. Bowman (2015), Climate-induced variations in global wildfire danger from 1979 to 2013, *Nature communications*, 6.

Kiem, A. S., and S. W. Franks (2004), Multi-decadal variability of drought risk, eastern Australia, *Hydrological Processes*, 18(11), 2039-2050.

Kim, S.-J., J. Lü, and B.-M. Kim (2014), The Southern Annular Mode (SAM) in PMIP2 simulations of the last glacial maximum, *Advances in Atmospheric Sciences*, 31(4), 863-878.

Krawchuk, M. A., M. A. Moritz, M.-A. Parisien, J. Van Dorn, and K. Hayhoe (2009), Global pyrogeography: the current and future distribution of wildfire, *PloS one*, 4(4), e5102.

Lau, K., and H. Weng (1995), Climate signal detection using wavelet transform: How to make a time series sing, *Bulletin of the American Meteorological Society*, 76(12), 2391-2402.

Lenton, T. M., H. Held, E. Kriegler, J. W. Hall, W. Lucht, S. Rahmstorf, and H. J. Schellnhuber (2008), Tipping elements in the Earth's climate system, *Proceedings of the National Academy of Sciences*, 105(6), 1786-1793.

Liu, W., J. Lu, L. R. Leung, S.-P. Xie, Z. Liu, and J. Zhu (2015), The de-correlation of westerly winds and westerly-wind stress over the Southern Ocean during the Last Glacial Maximum, *Climate Dynamics*, 1-12.

Lough, J., and H. Fritts (1987), An assessment of the possible effects of volcanic eruptions on North American climate using tree-ring data, 1602 to 1900 AD, *Climatic Change*, 10(3), 219-239.

Magilligan, F. J., and P. S. Goldstein (2001), El Niño floods and culture change: A late Holocene flood history for the Rio Moquegua, southern Peru, *Geology*, 29(5), 431-434.

Mariani, M., and M. S. Fletcher (2016), The Southern Annular Mode determines inter-annual and centennial-scale fire activity in temperate southwest Tasmania, Australia, *Geophysical Research Letters*, 43(4), 1702-1709

Marshall, G. J. (2003), Trends in the Southern Annular Mode from observations and reanalyses, *Journal of Climate*, 16, 24.

McWethy, D., P. Higuera, C. Whitlock, T. Veblen, D. Bowman, G. Cary, S. Haberle, R. Keane, B. Maxwell, and M. McGlone (2013), A conceptual framework for predicting temperate ecosystem sensitivity to human impacts on fire regimes, *Global Ecology and Biogeography*, 22(8), 900-912.

Meyers, S. D., B. G. Kelly, and J. J. O'Brien (1993), An introduction to wavelet analysis in oceanography and meteorology: With application to the dispersion of Yanai waves, *Monthly weather review*, 121(10), 2858-2866.

Moritz, M. A., M.-A. Parisien, E. Batllori, M. A. Krawchuk, J. Van Dorn, D. J. Ganz, and K. Hayhoe (2012), Climate change and disruptions to global fire activity, *Ecosphere*, 3(6), 1-22.

Moy, C. M., S. O. Seltzer, D. T. Rodbell, and D. M. Anderson (2002), Variability of El Niño/Southern Oscillation activity at millennial timescales during the Holocene, *Nature*, 420, 162-165.

Murphy, B. F., and B. Timbal (2008), A review of recent climate variability and climate change in southeastern Australia, *International Journal of Climatology*, 28(7), 859-879.

Nicholls, N., and C. Lucas (2007), Interannual variations of area burnt in Tasmanian bushfires: relationships with climate and predictability, *International Journal of Wildland Fire*, 16(5), 540-546.

Parisien, M.-A. (2016), Science can map a solution to a fast-burning problem, *Nature*, 534(7607), 297.

Parisien, M.-A., and M. A. Moritz (2009), Environmental controls on the distribution of wildfire at multiple spatial scales, *Ecological Monographs*, 79(1), 127-154.

Pausas, J. G., and E. Ribeiro (2013), The global fire–productivity relationship, *Global Ecology and Biogeography*, 22(6), 728-736.

Power, S., F. Delage, C. Chung, G. Kociuba, and K. Keay (2013), Robust twenty-first-century projections of El Niño and related precipitation variability, *Nature*, 502(7472), 541-545.

Rees, A. B., L. C. Cwynar, and M.-S. Fletcher (2015), Southern Westerly Winds submit to the ENSO regime: A multiproxy paleohydrology record from Lake Dobson, Tasmania, *Quaternary Science Reviews*, 126, 254-263.

Risbey, J. S., M. J. Pook, P. C. McIntosh, M. C. Wheeler, and H. H. Hendon (2009), On the remote drivers of rainfall variability in Australia, *Monthly Weather Review*, 137(10), 3233-3253.

Román-Cuesca, R. M., M. Gracia, and J. Retana (2003), Environmental and human factors influencing fire trends in ENSO and non-ENSO years in tropical Mexico, *Ecological Applications*, 13(4), 1177-1192.

Sandweiss, D. H., K. A. Maasch, R. L. Burger, J. B. Richardson, H. B. Rollins, and A. Clement (2001), Variation in Holocene El Niño frequencies: Climate records and cultural consequences in ancient Peru, *Geology*, 29(7), 603.

Santín, C., S. H. Doerr, E. S. Kane, C. A. Masiello, M. Ohlson, J. M. Rosa, C. M. Preston, and T. Dittmar (2016), Towards a global assessment of pyrogenic carbon from vegetation fires, *Global change biology*, 22(1), 76-91.

Saji, N., B. Goswami, P. Vinayachandran, and T. Yamagata (1999), A dipole mode in the tropical Indian Ocean, *Nature*, 401(6751), 360-363.

Scott, A. C., and I. J. Glasspool (2006), The diversification of Paleozoic fire systems and fluctuations in atmospheric oxygen concentration, *Proceedings of the National Academy of Sciences*, 103(29), 10861-10865.

Swetnam, T. W., and J. L. Betancourt (1990), Fire-southern oscillation relations in the southwestern United States, *Science (Washington)*, 249(4972), 1017-1020.

Timmermann, A., M. Latif, A. Grötzner, and R. Voss (1999), Modes of climate variability as simulated by a coupled general circulation model. Part I: ENSO-like climate variability and its low-frequency modulation, *Climate Dynamics*, 15(8), 605-618.

Torrence, C., and P. J. Webster (1999), Interdecadal changes in the ENSO-monsoon system, *Journal of Climate*, 12(8), 2679-2690.

Toth, L. T., R. B. Aronson, S. V. Vollmer, J. W. Hobbs, D. H. Urrego, H. Cheng, I. C. Enochs, D. J. Combosch, R. van Woesik, and I. G. Macintyre (2012), ENSO drove 2500-year collapse of eastern Pacific coral reefs, *Science*, 337(6090), 81-84.

Trenberth, K. E. (1984), Signal versus noise in the Southern Oscillation, *Monthly Weather Review*, 112(2), 326-332.

Turney, C. S. M., S. Haberle, D. Fink, A. P. Kershaw, M. Barbetti, T. T. Barrows, M. Black, T. J. Cohen, T. Correge, and P. P. Hesse (2006), Integration of ice-core, marine and terrestrial records for the Australian Last Glacial Maximum and Termination: a contribution from the OZ INTIMATE group, *Journal of Quaternary Science*, 21(7), 751-761.

van der Knaap, S., N. Tapper, and E. J. Cook (2010), Observed relationships between El Niño-Southern Oscillation rainfall variability and vegetation and fire history on Halmahera, Maluku, Indonesia, *Global Change Biology*, 16(6), 1705-1714.

Veblen, T. T., T. Kitzberger, R. Villalba, and J. Donnegan (1999), Fire history in northern Patagonia: the roles of humans and climatic variation, *Ecological Monographs*, 69(1), 47-67.

Verdon, D. C., A. S. Kiem, and S. W. Franks (2004), Multi-decadal variability of forest fire risk-eastern Australia, *International Journal of Wildland Fire*, 13(2), 165-171.

Wang, B., and Y. Wang (1996), Temporal structure of the Southern Oscillation as revealed by waveform and wavelet analysis, *Journal of Climate*, 9(7), 1586-1598.

Wang, C. (2001), On the ENSO mechanisms, *Advances in Atmospheric Sciences*, 18(5), 674-691.

Westerling, A. L., H. G. Hidalgo, D. R. Cayan, and T. W. Swetnam (2006), Warming and earlier spring increase western US forest wildfire activity, *science*, 313(5789), 940-943.

Whitlock, C., P. I. Moreno, and P. Bartlein (2007), Climatic controls of Holocene fire patterns in southern South America, *Quat. Res.*, 68(1), 28-36.

Williams, A. A., D. J. Karoly, and N. Tapper (2001), The sensitivity of Australian fire danger to climate change, *Climatic change*, 49(1-2), 171-191.

Williamson, G. J., L. D. Prior, W. M. Jolly, M. A. Cochrane, B. P. Murphy, and D. M. Bowman (2016), Measurement of inter-and intra-annual variability of landscape fire activity at a continental scale: the Australian case, *Environmental Research Letters*, 11(3), 035003.

Yan, H., L. Sun, Y. Wang, W. Huang, S. Qiu, and C. Yang (2011), A record of the Southern Oscillation Index for the past 2,000 years from precipitation proxies, *Nature Geoscience*, 4(9), 611-614.

Author Manuscript

FIGURES

Figure 1. Location of the 25,390 fire occurrences recorded in southeast Australia (a) and correlation map between annual (b) and seasonal (c,d,e,f) rainfall and the SOI during the reference period 1961-1990. Grid resolution is $0.05^\circ \times 0.05^\circ$. Significance has been tested across the 1208 stations used for the interpolation across southeast Australia and only areas with a $p\text{-value} < 0.05$ are shown.

Figure 2. Summary figure for the period 1951-2014 showing a) Rainfall anomaly for the Southeastern Australian region (data from BOM); b) Annual SOI, red (blue) bars represent El Niño (La Niña) anomalies; c) Z-scores of the area burnt values and d) Detrended z-scores of the number of fires occurrences.

Figure 3. Results from the superposed epoch analysis (SEA) between annual and seasonal SOI and a) Number of fires b) Area burnt during fire years and non-fire years. Dark bars indicate lags with a significant SOI departure ($p\text{ value} < 0.05$).

Figure 4. Results from the wavelet coherence analysis for the period 1951-2013 showing a) coherence between spring-summer SOI and number of fires; b) coherence between spring-summer SOI and area burnt. Black solid lines indicate areas exceeding the 90% confidence intervals. Leftward arrows indicate that x and y are anti-correlated: i.e. negative SOI (El Niño) occur in concert with high number of events and with positive anomalies in area burnt at the respective time frequency during the time periods shown in the figures. Dark grey solid line represents the 'cone of influence', where edge effects become important.




Figure 1. Location of the 25,390 fire occurrences recorded in southeast Australia (a) and correlation map between annual (b) and seasonal (c,d,e,f) rainfall and the SOI during the reference period 1961-1990. Grid resolution is $0.05^\circ \times 0.05^\circ$. Significance has been tested across the 1208 stations used for the interpolation across southeast Australia and only areas with a p -value < 0.05 are shown.




Figure 2. Summary figure for the period 1951-2014 showing a) Rainfall anomaly for the southeastern Australian region (data from BOM); b) Annual SOI, red (blue) bars represent El Niño (La Niña)

AU

anomalies; c) Z-scores of the area burnt values and d) Detrended z-scores of the number of fires occurrences.

Author Manuscript




Figure 3. Results from the superposed epoch analysis (SEA) between annual and seasonal SOI and a) Number of fires b) Area burnt during fire years and non-fire years. Dark bars indicate lags with a significant SOI departure (p value < 0.05). Annual (Jan-Dec); Winter(JJA); Spring(MAM); Summer(DJF); Autumn (SON).

Author Manuscript





Figure 4 Results from the wavelet coherence analysis for the period 1951-2013 showing a) coherence between spring-summer SOI and number of fires; b) coherence between spring-summer SOI and area burnt. Black solid lines indicate areas exceeding the 90% confidence intervals. Leftward arrows indicate that x and y are anti-correlated: i.e. negative SOI (El Niño) occur in concert with high number of events and with positive anomalies in area burnt at the respective time frequency during


the time periods shown in the figures. Dark grey solid line represents the 'cone of influence', where edge effects become important. Age is expressed as calendar years CE (x-axis).

Author Manuscript


anuscript




2016GL070572-f01-z-.tif



2016GL070572-f02-z-.tif



2016GL070572-f03-z.tif



2016GL070572-f04-z.tif