



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Tan, JZA;Gleeson, PA

Title:

Cargo sorting at the trans-golgi network for shunting into specific transport routes: Role of arf small g proteins and adaptor complexes

Date:

2019-06-01

Citation:

Tan, J. Z. A. & Gleeson, P. A. (2019). Cargo sorting at the trans-golgi network for shunting into specific transport routes: Role of arf small g proteins and adaptor complexes. *Cells*, 8 (6), <https://doi.org/10.3390/cells8060531>.

Persistent Link:

<https://hdl.handle.net/11343/251244>

License:

[CC BY](#)

Review

# Cargo Sorting at the *trans*-Golgi Network for Shunting into Specific Transport Routes: Role of Arf Small G Proteins and Adaptor Complexes

Jing Zhi Anson Tan  and Paul Anthony Gleeson \* 

Department of Biochemistry and Molecular Biology and Bio21 Molecular Science and Biotechnology Institute, The University of Melbourne, Melbourne, VIC 3010, Australia; j.tan49@student.unimelb.edu.au

\* Correspondence: pgleeson@unimelb.edu.au; Tel.: +61-3-8344-2354; Fax: +61-3-9348-1428

Academic Editor: Hesso Farhan

Received: 30 April 2019; Accepted: 30 May 2019; Published: 3 June 2019




**Abstract:** The *trans*-Golgi network (TGN) is responsible for selectively recruiting newly synthesized cargo into transport carriers for delivery to their appropriate destination. In addition, the TGN is responsible for receiving and recycling cargo from endosomes. The membrane organization of the TGN facilitates the sorting of cargoes into distinct populations of transport vesicles. There have been significant advances in defining the molecular mechanism involved in the recognition of membrane cargoes for recruitment into different populations of transport carriers. This machinery includes cargo adaptors of the adaptor protein (AP) complex family, and monomeric Golgi-localized  $\gamma$  ear-containing Arf-binding protein (GGA) family, small G proteins, coat proteins, as well as accessory factors to promote budding and fission of transport vesicles. Here, we review this literature with a particular focus on the transport pathway(s) mediated by the individual cargo adaptors and the cargo motifs recognized by these adaptors. Defects in these cargo adaptors lead to a wide variety of diseases.

**Keywords:** *trans*-Golgi network; adaptor proteins; AP-1; AP-3; AP-4; GGAs; protein sorting; post-Golgi transport


## 1. Introduction

In mammalian cells, the *trans*-Golgi network (TGN) has a distinct morphology, compared to the earlier Golgi cisternae, in that the TGN comprises networks of tubular, branching, and reticulating membranes [1,2]. The TGN has been typically viewed as a major sorting hub where cargoes are sorted into distinct transport carriers for trafficking to post-Golgi compartments and the plasma membrane (PM) [3,4] (Figure 1A). Moreover, the TGN also receives transport carriers from the endocytic trafficking pathway [5–7] (Figure 1A). Therefore, the TGN is a central organelle that intersects with both the biosynthetic and endocytic pathways. Given the multiple trafficking routes, transport machinery is employed to maintain the integrity of cargo sorting into the transport carriers. In addition, there is evidence that the cargoes may be segregated into specific TGN membrane micro-domains, and these microdomains are governed by small G proteins and TGN golgins [8–10]. Transport machinery, including small G proteins of the ADP ribosylation factor (Arf) family, cytosolic cargo adaptor proteins, coat proteins, and accessory proteins are key players in regulating cargo sorting at the TGN (reviewed in [3,4,6]) (Figures 1 and 2). Central to the sorting process is the interaction of cytosolic adaptor proteins with specific motifs on the cytoplasmic domains of membrane cargo proteins. These adaptor proteins are recruited to the TGN membranes by active membrane-associated Arf small G proteins, which then allow them to bind cargoes (Figure 2). In some cases, soluble cargoes indirectly associate with adaptor proteins through specific transmembrane cargo receptors [3,4,6]. Sorting signals, such as the classical tyrosine-based YXX $\Phi$  ( $\Phi$  denotes a bulky hydrophobic amino acid

and X denotes any amino acid) and dileucine-based (DE)XXX(LL) signals, located at the cytoplasmic tail of cargoes or cargo receptors are important for the recognition and binding by adaptor proteins [11,12] (Figure 2). This interaction then allows for selective inclusion of cargoes into their respective transport carriers [11–13] (Figure 2B). Some adaptor proteins recruit coat proteins, such as clathrin, which then polymerize to form electron-dense coated membrane vesicles [14]. Adaptor proteins also recruit accessory proteins to drive membrane curvature and scission, which leads to vesicle formation [15–20]. Upon the release of transport vesicles from the membranes, coat proteins are dissociated and recycled for additional rounds of vesicle formation.



**Figure 1.** Membrane trafficking pathways at the *trans*-Golgi network (TGN): (A) a schematic showing the post-Golgi trafficking pathways (red arrows) and the retrograde recycling pathways (black arrows) from the plasma membrane and endosomal/lysosomal compartments back to the TGN; (B) the distinct trafficking pathways at the TGN are regulated by specific cargo adaptor proteins including adaptor protein (AP) complexes (AP-1 in red, AP-3 in blue, AP-4 in orange); and (C) monomeric Golgi-localized,  $\gamma$  ear-containing, Arf-binding proteins (GGA1-3 in brown).




**Figure 2.** Cargo sorting at the TGN: (A) schematic showing the process of membrane recruitment of the adaptor protein complex, AP-1, and the binding of sorting motifs on the cytoplasmic tail of cargoes. GTP-bound membrane-associated small G protein Arf1 recruits cytosolic AP-1 to the TGN membranes. Arf1 also induces conformational change in AP-1, from closed to an open conformation, to allow the binding of cargo sorting motif(s). The  $\mu 1$  subunit (green) binds tyrosine-based sorting motif (YXX $\Phi$ , green cargo) and the  $\sigma 1$ - $\gamma$  subunits bind dileucine sorting motif (DE)XXXL(LI), brown cargo). (B) Schematic representation of the membrane recruitment of AP complexes by active small G proteins, followed by binding of AP complex to the cytoplasmic tail of cargo proteins. Subsequently, the hinge-ear extensions of each AP complex mediate the recruitment of coat and accessory proteins to drive membrane curvature and vesicle formation.

## 2. Small G Proteins of the ADP Ribosylation Factor (Arf) Family

The ADP ribosylation factor (Arf) family of small G proteins plays a vital role in post-Golgi sorting and trafficking, as well as maintaining the structure of the Golgi apparatus as reviewed in [8,9]. At the TGN, Arf proteins regulate the membrane recruitment of various cytosolic cargo adaptors, such as the adaptor protein (AP) complexes, and lipid-modifying enzymes, such as phosphatidylinositol (PtdIns) kinases, to initiate membrane trafficking [8,9]. The activity of Arf proteins is regulated between their membrane-associated, GTP-bound active state, and the cytosolic, GDP-bound inactive state [8,9,21] (Figure 3). The GTP and GDP states of Arfs are mediated by the Arf family guanine nucleotide-exchange factors (GEFs) and guanine nucleotide-activating proteins (GAP), respectively [8,9,22] (Figure 2A).

All Arfs are recruited to membranes through an N-terminal amphipathic helix from a hydrophobic pocket [8,9]. Upon the exchange of GDP for GTP, Arfs undergo conformational rearrangement in switch 1 and 2 and the interswitch regions [21] (Figure 3B). This conformational change leads to the displacement of the myristoylated N-terminal amphipathic helix from the hydrophobic pocket [21] (Figure 3B). The exposed myristoylated amphipathic N-terminal helix then inserts into the membrane [21,23] (Figure 3B). These features distinguish Arfs from other Ras superfamily small G proteins (Ras, Rho, Rab, and Ran), which harbor a long hypervariable C-terminal domain for membrane targeting and anchoring [9,23,24].



**Figure 3.** Regulation of Arf family small G proteins: (A) The state of the Arf small G proteins GTP binding and hydrolysis is regulated by Arf family guanine nucleotide exchange factors (GEFs) and GTPase-activating proteins (GAPs), respectively. The membrane-associated GTP-Arf then recruits effectors such as cargo adaptor proteins, membrane lipid-modifying enzymes, and additional GEFs. (B) GDP-Arf associates with the surface of the membrane in a reversible manner. Exchange of GDP for GTP in Arf causes a conformational change of the switch (SW1 and SW2) and interswitch regions to enter the hydrophobic pocket occupied by the myristoylated N-terminal helix (red). GTP-Arf then associates tightly with the membrane via the exposed N-terminal helix.

In humans, there are five Arf small G proteins and more than 20 Arf-like (Arl) small G proteins within the Arf family [9,25], and several of them are localized at the TGN [8,9]. All five Arfs are ubiquitously expressed and can be further divided into Class I (Arf1 and Arf3, Arf2 is not found in humans), Class II (Arf4 and Arf5), and Class III (Arf6) based on sequence homology [8,9]. The Class I Arfs, which share 97% sequence identity, are the most abundantly expressed Arf small G proteins [26–28]. The Class II Arfs, which are 90% identical, share 81% sequence identity with Class I Arfs [29]. Unlike the Class I and II Arfs, Arf6 is the only Class III Arf and it shares the least sequence identity (66%–70%) with other Arf family members [8,9]. Arf6 does not associate with the Golgi and is the only Arf that is localized at the PM, where it regulates events such as endocytosis [8,9,28].

On the other hand, the Class I and II Arfs are localized to the Golgi apparatus, including the TGN [26–29]. The individual knockdowns of Class I and Class II Arfs showed no obvious impact on Golgi morphology, suggesting functional redundancy in some aspects of membrane trafficking [29]. However, the double knockdowns of Class I and II Arfs, in every combination, revealed insights into specific defects in secretory and endocytic trafficking [29]. Along the secretory trafficking pathway, the depletion of Arf1 + Arf3 dramatically altered the morphology of ER-Golgi intermediate compartment (ERGIC) and accumulated vesicular stomatitis virus-G (VSVG) in large ERGIC-positive puncta, indicating a block in trafficking to the cis-Golgi [29]. On the other hand, Arf1 and Arf4 depletion caused tubulation and vesiculation of the Golgi, and blocked VSVG trafficking from the ER [29]. However, not all export pathways were blocked upon the depletion of Arf1 and Arf4. For example, the trafficking of the amyloid precursor protein (APP) from the Golgi was not affected by the silencing of Arf1 and Arf4 [30]. These studies reveal that different combinations of Arfs are likely to regulate distinct trafficking pathways.

The Class I and II Arfs are also likely to regulate post-Golgi trafficking as they are also localized to the TGN [26–28,31]. The cargo adaptor protein complex 1 (AP-1) is dissociated from the TGN only by the depletion of both Arf1 and Arf4, however, not by the single depletion of either Arf1 or Arf4 [32]. Given that AP-1 is required for post-Golgi trafficking of several cargoes (discussed below), Arf1 and Arf4 may either function collaboratively or distinctly to regulate post-Golgi trafficking through the recruitment of AP-1. Interestingly, the specific recruitment of GDP-bound, but not GTP-bound, Arf4 and Arf5, by calcium-dependent activator protein for secretion 1 (CAPS1) to the TGN has been implicated in the trafficking of dense-core vesicles (DCV) from the Golgi network in rat PC12 cells [31]. Arf4 and Arf5 are likely to also regulate post-Golgi trafficking in neurons through CAPS, which has been shown to regulate DCV secretion of a subset of neurotransmitters in the Purkinje neurons [31,33]. The depletion of either CAPS1 or Arf4/Arf5 leads to the accumulation of a DCV marker, chromogranin, in the TGN and reduced DCV secretion [31]. In addition to the combined functions of Class I and II Arfs at the Golgi, the GTP-bound Arfs may also regulate post-Golgi trafficking of specific cargoes via the recruitment of adaptor proteins. The GTP-bound Arf1 and Arf4 have been shown to regulate the transport of mannose 6-phosphate receptors (MPRs) [34,35] and ciliary cargoes [36] from the TGN, respectively.

In addition to the recruitment of adaptor proteins to the membranes, Arfs also induce conformational changes in adaptor proteins to promote their binding to sorting signals on cargoes [37]. Structural analysis reveals that cytosolic AP-1 exists as a closed conformation and this closed conformation is unable to bind to cargo sorting signals [37] (Figure 2A). Upon the recruitment of AP-1 to the membrane, Arf1 drives the conformational opening of the core of AP-1 such that the cargo binding sites are now exposed for cargo binding [37] (Figure 2A). There are three Arf1 binding sites on AP-1 that are involved in membrane recruitment and activation of cargo binding [37] (Figure 2A). Two of the Arf1 binding sites, which involve the switch region and are required for AP-1 membrane association, are located on the N-terminus of each large subunit ( $\beta$ 1 and  $\gamma$ ) of AP-1 [37] (Figure 2A). Simultaneously, a third proposed binding site, which is between the back side of Arf1 and the central part of AP-1 $\gamma$  subunit trunk domain, leads to the assembly of 2:2 dimer of AP-1 and Arf1, and the full activation of AP-1 for cargo binding [37].

### 3. Roles of Phospholipids in Regulating Protein Sorting

Phosphoinositides (PIs) play a central role in maintaining the identity of the Golgi membranes, which is necessary for proper membrane trafficking and sorting. Each specific PI, phosphatidylinositol mono-, bis-, and trisphosphate, is generated via reversible phosphorylation at positions three, four, and five hydroxyl groups on the inositol ring of the PI [38,39]. This process is driven by distinct pool of PI kinases (PIKs) and PI phosphatases that are localized on membrane organelles [38,39]. Phosphatidylinositol 4-phosphate (PI4P) is predominantly enriched in TGN membranes, and together with Arf small G proteins, are required for mediating the recruitment of the cargo adaptors, including AP-1 and GGAs [40,41].


The distribution and levels of PI4P are regulated by PI4 kinases (PI4Ks) and phosphatases that are present at the TGN [3,40]. There are four PI4Ks: PI4KII $\alpha$ , PI4KII $\beta$ , PI4KIII $\alpha$ , and PI4KIII $\beta$  in mammalian cells, with PI4KII $\alpha$  and PI4KIII $\beta$  representing the bulk of the PI4P production at the Golgi [38,39]. The knockdown of PI4KII $\alpha$  reduces PI4P levels at the TGN and causes the dissociation of AP-1 and GGAs into the cytosol, thus blocking membrane trafficking [40,41]. On the other hand, PI4KIII $\beta$  is required for maintaining the structural integrity of the Golgi since an expression of a dominant negative PI4KIII $\beta$  dramatically altered the Golgi organization [42]. The recruitment and membrane association of PI4KIII $\beta$  to the Golgi is regulated by GTP-bound Arf1 [42] and another Arf1 binding partner, neuronal calcium sensor 1 (NCS1) [43,44]. Moreover, the interaction of Arf1 and NCS1 controls PI4KIII $\beta$  activity and regulates both constitutive and regulated secretion [43,44]. Since the stimulation of PI4P production leads to the membrane recruitment of cargo adaptors, this PI pathway facilitates the coordinated sorting and trafficking of cargoes at the TGN.

The PI4P phosphatase, Sac1, has also been shown to regulate PI4P at the Golgi and is required for growth-dependent secretion in primary human fibroblasts [45]. In quiescent cells, Sac1 accumulates at the Golgi and eliminates PI4P, thereby downregulating anterograde trafficking [45]. In contrast, in human fibroblasts grown in the presence of high serum, Sac1 relocates to the ER which results in increased levels of PI4P at the Golgi, hence promoting anterograde trafficking [45].

#### 4. Cargo Adaptors at the TGN

Cargo adaptors include the complexes of the adaptor protein (AP) family, as well as the highly-conserved monomeric adaptors, Golgi-localized  $\gamma$  ear-containing Arf-binding proteins (GGAs), which collectively play a central role in the sorting of proteins at the TGN [3,46] (Figure 1B,C). AP complexes are the most well-characterized cargo adaptors and several of these are located at the TGN. Currently, there is a family of five homologous AP complexes identified in higher eukaryotes (Figure 4) which are: AP-1, AP-3, and AP-4 that are associated with the TGN; AP-2 that regulates endocytosis at the PM; and AP-5, the most recently identified member of the AP complex family, that facilitates protein retrieval, at the late endosomes/lysosomes, back to the Golgi [18,19,47]. The AP complexes are made up of two large subunits (one of  $\beta$ 1-5 and one of either  $\alpha$ ,  $\gamma$ ,  $\delta$ ,  $\epsilon$ ,  $\zeta$ ; ~100 kDa), one medium subunit ( $\mu$ 1-5, ~50 kDa), and one small subunit ( $\sigma$ 1-5, ~20 kDa) [18,19,47] (Figure 4). The N-terminal trunk domains of the two large subunits, the medium subunit, and the small subunit form the large globular core of the AP complex [13,20,37,48,49]. The C-terminus of each large subunit forms a long-flexible hinge region and a globular ear (appendage) domain that extend as a long projection from the core [13,20,48,49] (Figure 4). The core of the AP complex is important for membrane recruitment via binding to Arf small G proteins and phospholipids, as well as sorting signal recognition in the cargo proteins [12,13,18,19] (Figure 2A,B). On the other hand, the hinge region recruits coat proteins (i.e., clathrin), and the ear domain recruits accessory proteins which are important for membrane curvature and vesicle formation [12,13,18,19] (Figure 2A,B).

The highly-conserved GGA proteins function as ARF-dependent, monomeric clathrin adaptors to facilitate cargo sorting at the TGN. GGAs have homologous domains to the large subunits of AP complex [35,50–53] (Figure 4). In mammalian cells, there are three ubiquitously expressed GGAs (GGA1–3) that are localized at the TGN [50–52]. All GGAs have four domains which are comprised of the VHS (Vps27, Hrs, and STAM) domain, GAT (GGA and TOM1) domain, a flexible hinge-like region, and a C-terminal ear domain that is homologous to the AP-1  $\gamma$ 1-ear domain [50–52]. The VHS domain of GGAs has been shown to bind acidic-cluster dileucine sorting signals of cargoes, such as MPRs and sortilin [34,54]. Moreover, the VHS domain of each GGA has different binding preferences [34]. GGAs are recruited to the TGN membranes through binding of its GAT domain to membrane-associated GTP-Arf1 [55]. Similar to the AP complexes, the hinge-like domain of GGAs is required for clathrin recruitment [35,53], and the ear domain has been proposed to recruit accessory proteins [46,52].



**Figure 4.** Cargo adaptor proteins: Five adaptor protein (AP) complexes, AP-1 (red), AP-2 (green), AP-3 (blue), AP-4 (orange), and AP-5 (pink), have been identified to date in higher eukaryotes. Each heterotetrameric AP complex comprises two ~100 kDa large subunits ( $\beta 1$ – $5$ , and either  $\alpha$ ,  $\gamma$ ,  $\delta$ ,  $\epsilon$ , or  $\zeta$ ), one ~50 kDa medium subunit ( $\mu 1$ – $5$ ), and one ~20 kDa small subunit ( $\sigma 1$ – $5$ ). Together, they form the AP core, which is important for membrane recruitment and cargo sorting motif recognition. The C-termini of both large subunits of each AP complex give rise to the hinge and ear domains for further recruitment of accessory proteins. AP-1 is localized at the TGN/recycling endosomes and regulates bidirectional transport. AP-1 also regulates basolateral sorting in polarized cells. AP-2 is responsible for endocytosis of cargoes from the cell surface. AP-3 is localized at the TGN/early endosomes and regulates transport to the late endosomes/lysosomes. AP-4 is localized at the TGN and regulates cargo trafficking from the TGN to the early endosomes. AP-5 is localized at the late endosomes/lysosomes, and the trafficking pathway it regulates is still unclear. The three Golgi-localized  $\gamma$ -ear containing Arf binding isoforms, GGA1, GGA2, and GGA3 (brown), are monomeric and have similar structural protein folding to the ear domain of AP-1 $\gamma$  subunit. GGAs are localized at the TGN and endosomes.

#### 4.1. Adaptor Protein (AP) Complexes at the TGN

AP-1, AP-3, and AP-4 are localized at the TGN and are predicted to generate distinct transport vesicles [18–20]. AP-1 is associated with clathrin-coated vesicles (CCVs) [53] and binds clathrin via the clathrin-binding motif which is located at the flexible hinge of  $\beta 1$  subunit [56,57]. AP-3  $\beta 3$  subunit also contains a clathrin-binding motif, but AP-3 subunits are not enriched in CCVs prepared from brain or the liver [53,58]. Therefore, it remains unclear whether AP-3 is functionally capable of recruiting clathrin. On the other hand, AP-4 lacks the clathrin-binding motif and is not associated with CCVs [59,60]. Mass spectrometric protein quantification of HeLa cell lysates revealed that the number of AP-1 (~4500) and AP-3 (~4500) vesicles per cell were 30 times more abundant than AP-4 vesicles (~150) [18]. The differences in abundance suggest that AP-1 and AP-3 likely represent the major cargo trafficking pathways and the AP-4 pathway is relatively minor. Nonetheless, given that the loss of AP-4 produced drastic clinical phenotypes in humans, AP-4 may represent a specialized TGN trafficking route for a specific set of proteins that collectively represent a minor fraction of all the membrane cargoes which exit the TGN [18].

##### AP-1:

The best characterized TGN-localized AP complex is AP-1. There are two isoforms of AP-1 that are  $\mu 1$  subunit,  $\mu 1A$ , and  $\mu 1B$ , which give rise to AP-1A and AP-1B, respectively [61,62]. Whilst AP-1A is expressed ubiquitously, AP-1B is only expressed in a subset of polarized epithelial cells in mammals [61, 62]. A classical AP-1 mediated trafficking pathway is the delivery of newly synthesized lysosomal hydrolases bound to the MPRs from the TGN to the endosomes [63,64]. The loss of AP-1 leads to the secretion of lysosomal hydrolases instead of delivery to the lysosomes [64]. Interestingly, the loss of AP-1 also results in the accumulation of MPRs at the endosomes [64]. In order to determine the role of AP-1 in the endosomal trafficking of MPRs, a knocksideways technique to reroute the TGN-localized AP-1 to the mitochondria was developed [65]. CCVs isolated from the AP-1 knocksideways cells revealed that the MPRs (CD-MPR and CI-MPR) were significantly more depleted than the lysosomal hydrolases [65]. This finding suggests that AP-1 is also required for the trafficking of ligand-free MPRs, most likely in the retrograde pathway, back to the TGN for another round of binding and delivery

of newly synthesized lysosomal hydrolases [65]. Collectively, these data support a role for AP-1 in regulating bidirectional trafficking between the TGN and endosomes.

AP-1 is also implicated with the sorting of cargoes into polarized epithelial cells and neurons [12,62,66,67]. The roles of AP-1A and AP-1B in epithelial basolateral sorting have been investigated extensively [62,67,68]. Early studies indicated that AP-1A and AP-1B differed in their localization. AP-1A is localized at the TGN and AP-1B is localized at the recycling endosomes. It was proposed that AP-1A regulates biosynthetic sorting at the TGN and AP-1B is involved in recycling of basolateral proteins from the recycling endosomes [69,70]. However, a recent study, using high-resolution microscopy, has challenged these earlier findings and demonstrated that AP-1A and AP-1B colocalize to a similar extent at the TGN and recycling endosomes in polarized Madin-Darby Canine Kidney (MDCK) epithelial cells [62]. In addition, double knockdown of AP-1A and AP-1B, but not AP-1A or AP-1B single knockdown, dramatically reduced the delivery of the newly synthesized transferrin receptor (TfR) and low-density lipoprotein receptor (LDLR) from the TGN to the basolateral PM [67]. These results suggest that both AP-1A and AP-1B compensate for each other in regulating the exit of cargoes from the TGN. However, AP-1A is likely to regulate biosynthetic trafficking to the PM independent of the recycling endosomes, since the knockdown of AP-1A, but not AP-1B, increased the trafficking of LDLR and TfR from the TGN to the recycling endosomes [67]. Interestingly, knockdown of AP-1B, but not AP-1A, significantly decreased the steady-state basolateral polarity of LDLR and TfR [67], suggesting a specific role of AP-1B in the maintenance of basolateral polarity in epithelial cells. In a follow-up study, yeast 2-hybrid and pull-down experiments demonstrated that AP-1B binds noncanonical tyrosine-based and acidic clusters sorting motifs of LDLR more strongly than AP-1A [62]. Therefore, the differential expression of AP-1B in epithelial cells may promote the sorting and maintenance of basolateral polarity of proteins that are not efficiently regulated by AP-1A. All in all, both AP-1A and AP-1B have complementary and non-overlapping roles in sorting at the TGN and epithelial basolateral polarity.

AP-1A regulates the polarized sorting of cargoes, such as the TfR, the coxsackievirus and adenovirus receptor (CAR), and the neuron-specific receptors (metabotropic glutamate receptor 1 (mGluR1), and the N-methyl-D-aspartate (NMDA)-type ionotropic glutamate receptors NR2A and NR2B), to the somatodendritic domain in rat hippocampal neurons [12,66]. AP-1 is localized to the TGN/recycling endosomes and dendrites [66], and live-cell imaging showed that the bidirectional tubular carriers moving between soma and dendrites are decorated with AP-1 [66]. AP-1 functions by excluding the cargo proteins from the transport carriers that are destined for axons [66]. The disruption of AP-1 and cargo interaction by overexpressing a dominant negative  $\mu$ 1A-W408S mutant resulted in misincorporation of TfR into the axonal transport carriers at the TGN/recycling endosomes in the soma/neuronal body [66]. Moreover, the overexpression of  $\mu$ 1A-W408S also led to mis-sorting of mGluR1, NR2A, and NR2B from the somatodendritic domain to the axons [66]. These findings highlight the importance of AP-1 in regulating somatodendritic sorting and polarity in neurons. The role of AP-1 in embryonic development is also indispensable since the knockout of the genes encoding  $\gamma$  or  $\mu$ 1 subunit in mice causes early embryonic lethality [64,71]. In addition, mutation of  $\sigma$ 1 subunit in humans is associated with X-linked mental retardation and several neurological disorders, further highlighting the importance of AP-1 in neurological development [72–74].

### AP-3:

AP-3, similar to AP-1, also exists in two isoforms, i.e., AP-3A ( $\beta$ 3A/ $\mu$ 3A) and AP-3B ( $\beta$ 3B/ $\mu$ 3B), where, AP-3A is expressed ubiquitously and AP-3B is neuronal-specific [58,75–77]. Immunofluorescent data revealed that the majority of endogenous AP-3 is associated with punctate structures that lack TfR and it was proposed that these structures were the late endosome/lysosomes [58]. In addition, some AP-3 is associated in the perinuclear region, which likely represents the Golgi [58]. The role of AP-3 in cargo sorting to lysosomes, subsequently, came from studies identifying mutations in  $\beta$ 3A in patients with Hermansky-Pudlak syndrome (HPS), which is a genetic disorder with defective lysosome-related organelles [78]. The AP-3-deficient fibroblasts from these patients showed an increase

in cell surface expression of lysosomal-associated membrane proteins CD63, LAMP-1, and LAMP-2, as compared with fibroblasts from a healthy individual, which showed localization at the late endosome/lysosome [78]. A null allele of another AP-3 subunit,  $\delta$ , was identified in an HPS mouse model, mocha [79]. The lack of AP-3 in mocha was established as the cause of pleiotropic hematologic, hypopigmentation, and neurological phenotypes in this mouse line [79]. The human and mouse phenotypes from AP-3 deficiency have one thing in common, a defective endosomal-lysosomal sorting pathway [79]. In a later study, mocha fibroblasts were also shown to display increased cell surface expression of CD63 and LAMP1 [80], similar to fibroblasts from human patients with HPS described by [78]. Using immunoelectron microscopy, LAMP-1 and LAMP-2 were often found to coincide in endosomal-associated tubules that were decorated with AP-3 in a human liver carcinoma cell line, HepG2 cells [80]. In addition, AP-3 was found in budding vesicles that were distinct to AP-1 budding vesicles at the endosomal tubular membranes [78]. These endosomal AP-3 budding vesicles were also smaller than the AP-1 budding vesicles at the TGN [80]. These studies suggest that AP-1 and AP-3 regulate distinct trafficking routes from the endosomes. Therefore, AP-3 is likely to be responsible for cargo sorting from the endosomes to lysosomes, and lysosomal-related organelles such as melanosomes.

#### AP-4:

Unlike AP-1 and AP-3, AP-4 is not as well characterized. AP-4 is the most recently discovered AP complex associated with the TGN [59,60]. It has been reported that AP-4 shows a distinct immunofluorescence staining pattern as compared with AP-1 at the TGN [59,60], suggesting that both AP complexes are likely to regulate distinct post-Golgi sorting and trafficking. Despite its relatively low abundance compared to AP-1 and AP-3 in HeLa cells [18], AP-4 is expressed in all human tissues [59,60], thus suggesting important functions in membrane trafficking. Indeed, several independent studies have reported mutations in all four subunits of AP-4 genes encoding  $\beta 4$  [81–85],  $\epsilon$  [81,86–88],  $\mu 4$  [84,87,89], and  $\sigma 4$  [81,90] in human patients. These patients displayed severe intellectual disability and progressive spastic paraplegia, with early onset of spasticity [81–84,86–90]. These studies also highlight the importance of each AP-4 subunit in forming a functional AP-4 complex.

In a post-mortem analysis of an AP-4 $\mu$  deficient patient, histology of the brain tissue revealed neuronal malformations, neuroaxonal degeneration, and white matter loss [89]. Given the severity of the neurological phenotype associated with the loss of AP-4, it is likely that AP-4 plays an important role in brain development and neuronal trafficking. Surprisingly, in contrast to the observed phenotypes in humans, AP-4  $\beta 4^{-/-}$  mice were healthy and fertile with no obvious phenotypes detected, including in the brain [91]. A poor rotor performance was the only reported fitness phenotype, hence, suggesting defects in motor neurons [91]. Nonetheless, AP-4  $\beta 4^{-/-}$  mice revealed a functional role of AP-4 in polarized sorting in neurons [91]. The sorting of  $\alpha$ -amino-3-hydroxy-5-methyl-5-isoxazolepropionic acid (AMPA) receptor to the somatodendritic domain in neurons by AP-4 requires indirect association of AP-4 with the AMPA receptor via transmembrane AMPA receptor regulatory proteins (TARPs) [91]. In AP-4  $\beta 4^{-/-}$  hippocampal neurons and Purkinje cells, the AMPA receptor and TARPs are missorted into LC3-II (a lipidated form of protein light chain 3) positive autophagosomes that accumulated in the swollen axons [91]. In addition, the  $\delta 2$  subtype of glutamate receptor, which had been shown to bind AP-4  $\mu 4$  in a previous study [92], also accumulated in the axons of AP-4  $\beta 4^{-/-}$  Purkinje cells [91]. Conversely, the distribution of several other somatodendritic receptors, including TfR, mGluR1, and NMDA-type glutamate receptor NR1, were not affected [91]. Given that AP-1A regulates the somatodendritic sorting of TfR and mGluR1 in rat hippocampal neurons [66], these results demonstrate the selectivity of AP-4 in the post-Golgi sorting and trafficking of cargoes that are distinct for AP-1 [66,91].

In vitro and Y2H analyses have shown that AP-4  $\mu 4$  binds to canonical tyrosine-based motifs of cargoes including CD63 [60], LAMP-1, and LAMP-2 [93]. However, the binding affinities of AP-4  $\mu 4$  to these cargoes are weak, and the depletion of AP-4 did not alter the distribution of these proteins to

lysosomes [94]. Therefore, in contrast to AP-3, AP-4 is unlikely to regulate TGN to lysosomal trafficking. AP-4  $\mu$ 4 has also been shown to strongly bind synthetic peptides encoding the tyrosine-based sorting motif in LDLR, and the basolateral sorting motif in CD-MPR by surface plasmon resonance binding assays [95]. However, the depletion of AP-4  $\mu$ 4 in MDCK cells only partially affected basolateral sorting of LDLR and CD-MPR [95]. Given that AP-1 has been shown to also regulate the trafficking of LDLR and MPRs as described above, AP-4 may function in parallel with AP-1. Consistent with this observation, AP-4 vesicle formation has been shown to be enhanced when CCVs components were depleted [18,96]. Moreover, the level of AP-4 vesicles also increased when CCVs formation was altered in AP-1 knocksideways studies from Robinson's laboratory [18,65]. Therefore, AP-4 is likely to be a specialized cargo adaptor that regulates the TGN export of a selective set of cargoes and, in addition, may sort a subset of cargoes that are normally transported in CCVs.

An important role of AP-4 in regulating trafficking from the TGN to endosomes was uncovered in a study from the Bonifacino's laboratory [97]. AP-4  $\mu$ 4 was shown to bind to amyloid precursor protein (APP) directly, which represents a novel AP-cargo interaction [97]. AP-4 is required for the post-Golgi trafficking of APP to the endosomes in HeLa cells and neurons, a transport step which regulates APP processing [30,97,98].

More recently, AP-4 has been shown to mediate export of a multi-pass membrane core autophagy protein, ATG9A (autophagy-related protein 9A), from the TGN to endosomes and/or preautophagosomes, and is thought to promote early stages of autophagosome maturation [99–101]. Moreover, a recent study from the Robinson's laboratory identified two additional proteins, SERINC1 and SERINC3 (serine incorporator 1 and 3), which co-traffic with ATG9A in AP-4 derived vesicles [101]. These vesicles were also found to associate with two newly identified cytosolic AP-4 accessory proteins, RUSC1 and RUSC3 (run and SH3-domain containing protein), which are thought to be required for microtubule plus end-directed trafficking [100].

#### 4.2. Golgi-Localized, $\gamma$ Ear-Containing, Arf-Binding Proteins (GGAs)

The function, and potential redundancy, of the highly conserved GGAs (GGA1-3) remain only partially defined. Evidence for a non-overlapping function amongst the GGAs comes from analyses of GGA null mice, which showed that the loss of GGA2, but not GGA1 or GGA3, is embryonic lethal [102,103]. GGAs are reported to interact and regulate the trafficking of cargoes such as MPRs and sortilin from the TGN to the endosomes [34,54]. Interestingly, GGAs and AP-1 have been shown to function cooperatively in the sorting of MPRs and their bound lysosomal enzymes into CCVs at the TGN [65,104]. However, only AP-1 is required for the trafficking of ligand-free MPRs from the endosomes back to the TGN [65]. These studies suggest that GGAs and AP-1 have cooperative roles in regulating anterograde trafficking from the TGN but distinct roles in endosomal sorting.

In addition to membrane trafficking from the TGN, GGAs have also been shown to regulate trafficking from other endosomal compartments. Firstly, GGA1 has recently been shown to mediate the recycling of cargoes such as BACE1 from the early endosomes to the recycling endosomes [105]. Secondly, GGA2 was reported to interact with epidermal growth factor receptor (EGFR) in endosomal structures [106]. The depletion of GGA2 leads to missorting of EGFR to the lysosomes via post-Golgi compartments, indicating that GGA2 is required for sustaining the level of EGFR during cell growth [106]. An increase in turnover of EGFR may explain the developmental arrest of GGA2 null mice. Lastly, the hinge region in GGA3 has two binding sites for ubiquitin and has been shown to play a role in targeting ubiquitinated proteins for lysosomal degradation [107]. For example, GGA3 regulates the lysosomal targeting and turnover of BACE1 as the depletion of GGA3 increased intracellular levels of BACE1 and elevated A $\beta$  levels [108,109].

### 5. Cargo Sorting Signals

The cytosolic domain of cargoes consists of short, linear sequences of amino acids that function as sorting signals. The recognition of sorting signals by specific cargo adaptor proteins allows for


selective incorporation of cargoes into transport carriers along the biosynthetic and endocytic trafficking pathways. The tyrosine-based (YXX $\Phi$ ) and dileucine-based (DE)XXXL(LI) motifs (X denotes any amino acid and  $\Phi$  denotes a bulky hydrophobic amino acid, i.e., isoleucine, leucine, methionine, phenylalanine or valine) are the best-characterized sorting signals (Table 1). The YXX $\Phi$  signals are recognized by the  $\mu$  subunit of AP-1 through 4 while the  $\mu$  subunit of AP-5 has yet to be identified to bind tyrosine-based sorting signals [110,111]. On the other hand, (DE)XXXL(LI) signals are recognized by hemicomplexes of AP-1 ( $\gamma$ - $\sigma$ 1), A-P2 ( $\alpha$ - $\sigma$ 2), and AP-3 ( $\delta$ - $\sigma$ 3) [110,111] (Table 1). There is no evidence suggesting that AP-4 or AP-5 binds dileucine-based signals. The AP complexes exhibit different binding affinities for individual (DE)XXXL(LI) signals. For example, AP-1 and AP-2, but not AP-3, bind the DDQRDLI signal of human major histocompatibility complex class II invariant chain (Ii) [112]. In contrast, AP-3, but not AP-2, binds the DERAPLI and EEKQPLL signals of LIMP-II and Tyrosinase, respectively [113,114] (Table 1). Therefore, these findings highlight the selectivity of sorting signals by AP complexes.

**Table 1.** Cargo sorting signals.

Motifs	Cargoes	Functions	Cargo Adaptor Proteins	Reference
YXX $\Phi$	TGN38, TfR, Furin, MPRs, EGFR, CD63, LAMP1, LAMP2	Endocytosis, Endosomal and Lysosomal targeting	$\mu$ subunits of AP complexes	[48,79,93,110,114–118]
	TfR, CAR	TGN $\rightarrow$ somatodendritic domains in neurons	$\mu$ 1A (AP-1A)	[66]
(DE)XXXL(LI)	Invariant chain	TGN $\rightarrow$ endosomes	Hemicomplexes of AP-1 $\gamma$ - $\sigma$ 1	[112,119]
	LIMP-II, Tyrosinase	Lysosomal targeting	Hemicomplexes of AP-3 $\delta$ - $\sigma$ 3	[113,114]
	BACE1	Endocytosis	Hemicomplexes of AP-2 $\alpha$ - $\sigma$ 2	[120]
YXX $\Phi$ E	APP family members, ATG9A	TGN $\rightarrow$ endosomes	$\mu$ 4 (AP-4)	[97,99]
Noncanonical Tyrosine-based motif	LDLR	TGN $\rightarrow$ basolateral PM	$\mu$ 1B (AP-1B)	[62]
	TARP (AMPA receptor)	TGN $\rightarrow$ somatodendritic domains in neurons	$\mu$ 4 (AP-4)	[91]
Phenylalanine-based motif	$\delta$ 2 glutamate receptor	TGN $\rightarrow$ somatodendritic domains in neurons	$\mu$ 4 (AP-4)	[92]
Acidic clusters-DXXLL	MPRs	TGN $\leftrightarrow$ endosomes	GGAs	[34]
	BACE1	Endosomal recycling	GGAs	[105,121,122]
Acidic clusters	Nef protein of HIV-1	Endocytosis: Downregulation of MHC-I	$\mu$ 1 (AP-1)	[123]
	Furin, CD-MPR	TGN sorting		[124,125]

Using X-ray crystallographic analyses, the binding mechanisms of  $\mu$ 1A [126],  $\mu$ 2 [48,115],  $\mu$ 3A [116], and  $\mu$ 4 [60,93] to YXX $\Phi$  sorting signals have been determined (Figure 5A–D). The N-terminal domain of all the  $\mu$  subunits is required for an AP complex assembly while the C-terminal domain binds the YXX $\Phi$  signals [127] (Figure 5A–D). The 16  $\beta$ -strands within the C-terminal domain of the  $\mu$ 1–4 subunits are organized into subdomains A and B, giving rise to the immunoglobulin-like  $\beta$ -sandwich fold (Figure 5A–D). In  $\mu$ 2, the  $\beta$ 1 and  $\beta$ 16 strands in subdomain A binds YXX $\Phi$  where the Y and  $\Phi$  residues fit into two hydrophobic pockets (Figure 5B). The crystal structure of  $\mu$ 1A was solved in a complex with the class 1 major histocompatibility complex (MHC-I) cytosolic tail, which lacks a  $\Phi$  residue (Figure 5A). Although  $\mu$ 1A binds the Y residue similarly to  $\mu$ 2, the other regions of  $\mu$ 1A also establish additional interactions with the MHC-I tail. The YXX $\Phi$  signal of TGN38 binds  $\mu$ 3A at a site that is equivalent on  $\mu$ 2 (Figure 5C), however, with fewer contacts for stabilization [116]. In contrast to the  $\mu$ 1–3 subunits,  $\mu$ 4 exhibits the most distinct binding mechanism characterized to date for the YXX $\Phi$  signals (Figure 5D). Although the binding of  $\mu$ 4 to canonical YXX $\Phi$  signals is weak [93], it exhibits strong binding affinity for a unique subset of YXX $\Phi$  signals fitting the YXX $\Phi$ E motif [97,99,128]. The YXX $\Phi$ E motif is found in the cytosolic tails of the APP family members: APP (YKFFE), APLP1 (YKYLE), and APLP2 (YRFLE), as well as the autophagy-related protein 9A (ATG9A, YQRLE) [97,99,128]. Strikingly, the crystal

structure of the  $\mu$ 4 C-terminal domain, in a complex with APP's YKFFE cytosolic signal, revealed a unique binding site located on  $\beta$ 4,  $\beta$ 5, and  $\beta$ 6 strands in the subdomain A of  $\mu$ 4 [97,128] (Figure 5D). This binding site, which also has hydrophobic pockets for Y and  $\Phi$  residues, is located opposite to that of the YXX $\Phi$  binding sites on  $\mu$ 1–3 [97,128] (Figure 5A–D). Moreover, the crystal structure of  $\mu$ 4 also predicts the presence of a similar  $\mu$ 2 binding site and mutations in this site abolished the binding of the LAMP-2 canonical YXX $\Phi$  signal [93,97,128]. Thus, the subdomain A of  $\mu$ 4 is likely to have two binding sites, one for YXX $\Phi$  and one for YXX $\Phi$ E. Collectively, these studies provided valuable insights that illustrate that the binding affinities and mechanism of cargo signals by  $\mu$ 1–4 subunits are distinct, therefore, highlighting the presence of cargo selectivity by AP complexes for regulating cargo transport to specific intracellular locations.



**Figure 5.** Tyrosine sorting motifs binding sites on AP complex. Crystal structures of the C-terminal domains of  $\mu$  subunits of AP-1 through 4 in association with tyrosine-based sorting motifs (underlined). The C-terminal domains of  $\mu$ 1–4 are further subdivided into subdomain A (blue) and subdomain B (red). (A) Mouse AP-1  $\mu$ 1 C-terminal domain in complex with class 1 major histocompatibility complex (MHC-1) peptide SYSQAA (PDB: 4EN2), (B) rat AP-2  $\mu$ 2 C-terminal domain in complex with epidermal growth factor receptor (EGFR) peptide FYRAL (PDB: 1BW8), (C) rat AP-3  $\mu$ 3A C-terminal domain in complex with TGN38 peptide SDYQRL (PDB: 4IKN), (D) human AP-4  $\mu$ 4 C-terminal domain in complex with amyloid precursor protein (APP) peptide TYKFFEQ (PDB: 3L81).

The binding of noncanonical tyrosine-based signals by the  $\mu$ 1 and  $\mu$ 4 subunits (Table 1) are essential for the polarized sorting of cargoes to the basolateral PM of epithelial cells and somatodendritic PM of neuronal cells, respectively [62,91]. In addition,  $\mu$ 4 is shown to bind a phenylalanine-based motif (Table 1), which is required for cargo sorting to the somatodendritic PM of neurons [92]. Another class of sorting signals, the acidic cluster-dileucine DXXLL motif, is recognized only by the GGAs via its VHS domain, described in [34,54,110] (Table 1). The acidic cluster-DXXLL motif is required for endosomal recycling of BACE1 (see later sections) [105,121,122] and trafficking of cargoes such as CD- and CI-MPRs between the TGN and endosomes [34,65]. Lastly, less information is known about acidic cluster sorting signals, but they are proposed to regulate sorting in the TGN [124] (Table 1).

This motif is also often found in transmembrane proteins (i.e., furin) that are localized at the TGN in steady-state [110]. A recent study from the Robinson laboratory identified  $\mu$ 1 as an acidic cluster sorting machine [123]. In vitro studies from the same laboratory showed that the basic patch (Lys274, Lys298, Lys302, Arg303, and Arg304) of  $\mu$ 1 is required for the binding of acidic cluster signals [123]. Thus, these studies show the capability of cargo adaptor proteins to recognize a wide array of cargo sorting signals in a selective manner.

## 6. Accessory Proteins of AP Complexes and the Formation of Transport Vesicles

Membrane curvature followed by membrane fission from the cytosolic face of a continuous membrane (i.e., TGN) is required for the formation and release of transport vesicles, and thus allows vesicular trafficking to proceed [129,130]. The cargo adaptors by themselves do not drive membrane curvature or bud formation. This process is controlled by various accessory proteins [129,130]. The highly-conserved epsin N-terminal homology (ENTH) domain (~130–150 residues) is found in epsin family proteins (epsin1–3) [131,132], epsin-related protein (epsinR/enthoprotin) [129,132,133], and tepsin (tetra-epsin/epsin for AP-4) [134]. The formation of an amphipathic  $\alpha$ -helix in the ENTH domain binds phosphoinositide (PI) and partially inserts into the membranes [17,135]. The ENTH domain of epsin1 is capable of promoting the formation of tubules from liposomes derived from total brain lipids [17]. Furthermore, the addition of the ENTH domain to liposomes containing 10% PI<sub>4,5</sub>P<sub>2</sub> (a PI that is enriched at the PM) was shown to induce tubulation as well as fragmentation into small vesicles [17]. These observations are supported by a biophysical computational analysis conducted to analyze the impact of the ENTH domain membrane insertion on a continuous membrane bilayer [16]. This analysis, which takes into account the molecular features of lipids and proteins, predicted that the partial insertion of proteins into hydrophobic bilayers is sufficient to drive membrane fission and results in the formation of separate vesicles from continuous membranes [16]. These predictions were further validated using an in vitro quantitative liposome vesiculation assay, which showed that the ENTH domain causes membrane vesiculation and tubulation, and the degree of vesiculation also correlated strongly with the number of amphipathic helices [16]. The transient overexpression of epsin1 in BSC1 cells depleted of dynamin, i.e., a membrane-remodelling GTPase required for endocytic membrane fission events [136], rescued CCV formation in a transferrin uptake assay [16]. Together, these results suggest that the ENTH domain is capable of driving membrane curvature and fission.

Epsin1, the best-characterized epsin family member, associates with the PM by binding to PI<sub>4,5</sub>P<sub>2</sub> [17]. Epsin1 also binds AP-2 via the appendage domains of  $\alpha$  [137] and  $\beta$ 2 [138] large subunits to stimulate a clathrin assembly and drive clathrin-mediated endocytosis at the PM [17]. In contrast, EpsinR is recruited to the TGN membranes in an Arf-dependent manner and preferentially binds PI4P [133,139]. EpsinR is associated with AP-1 by binding to the  $\gamma$  subunit appendage domain and is involved in CCV budding at the TGN [139]. However, unlike epsin1, epsinR does not stimulate a clathrin assembly [139]. Tepsin is the only AP-4 accessory protein identified to date [96,140]. Interestingly, while the membrane recruitment of epsin1 and epsinR are independent of AP-2 and AP-1, respectively, tepsin requires its AP complex, AP-4, for localization to the membranes [96]. Tepsin contains two C-terminal motifs that bind to the appendage domains of AP-4 $\epsilon$  [140] and  $\beta$ 4 [141]. The loss of AP-4 abolished the membrane recruitment of tepsin to the TGN in fibroblasts derived from human patients with loss-of-function mutations in AP-4  $\sigma$ 4 [90]. In line with these observations, a recent study using X-ray crystallography revealed that the ENTH domain of tepsin (tENTH) lacks the amphipathic  $\alpha$ -helix and a PI binding pocket, which are found in the ENTH domains of both epsin1 and epsinR [134]. Thus, explaining why tepsin requires AP-4 for membrane recruitment. Phylogenetics and comparative genomic analyses suggest that tepsin diverged from epsins ~1.5 million years ago, and probably supports different biological functions [134]. Interestingly, tepsin contains a unique tepsin VHS-like domain (tVHS), which is not found in epsin1 and epsinR [134]. Structural data revealed that tVHS lacks an  $\alpha$ -helix-8 that is found in VHS domains, suggesting that it lacks common VHS

functions such as the binding of dileucine-based motifs [134]. Nonetheless, it is predicted that both tENTH and tVHS domains are likely to associate with a protein partner [134]. Further studies are required to fully understand the role of tepsin and its tENTH and tVHS domains in the generation of transport carriers.

## 7. Challenges and Future Directions

Many studies to date have analyzed cargo sorting in transfected cells with high levels of expression of individual cargoes. The impact of overexpression may influence the selectivity of recruitment into defined transport pathways, particularly at the TGN which has multiple transport routes. An important advancement is the development of systems to analyze the trafficking of endogenous cargo in both immortalized and primary cells. A recent innovative approach to analyze endogenous cargoes involves the development of a proteomic technique known as dynamic organellar maps, which provides a powerful complement to imaging-based analyses [142,143]. This technique combines subcellular fractionation with quantitative mass spectrometry to determine the abundance and distribution of individual proteins in the different organelle compartments in an unbiased manner [142,143]. The dynamic organellar maps technique also enables the spatial distribution of cargo proteins in each intracellular organelle to be quantified upon the loss of key transport machinery, either through genetic editing or in an inherited disease. This approach allows rapid assessment of the impact of depletion on a transport machinery component on the distribution of the entire cohort of cargo proteins that share a particular pathway. This technique should be particularly amenable to the study of minor transport pathways, such as the AP-4 mediated pathway, since AP-4 is present in low levels as compared with the other adaptor complexes.

A recent study by the Robinson laboratory utilized this approach to uncover the role of AP-5 in regulating late endosomes to Golgi recycling of various proteins including the CI-MPR, Golgi phosphoprotein 1 (GOLM1), and the Golgi integral membrane protein 4 (GOLIM4) [144]. Moreover, using the same approach, an important role of AP-4 in autophagy has been uncovered [100]. Therefore, the dynamic organellar maps technique holds considerable promise for the analyses, in an unbiased manner, of currently poorly characterized transport pathways and the impact of perturbation on the trafficking machinery.

Much of the work to date has analyzed trafficking pathways in cultured immortalized cells. Another important challenge is to define transport pathways in primary cells. For example, in melanocytes, AP-1 can partially rescue the requirement of AP-3 for the trafficking of tyrosinase from distinct tubular vesicular domains in endosome to melanosomes [145]. This study would not be possible in nonspecialized cells, which lack melanosomes. The dynamic organellar maps technique has been utilized to generate a quantitative comparison of the organellar organization in neurons [143]. Another powerful approach of this tool is to compare the trafficking studies conducted previously in cultured cells with primary cells.

In addition, the dynamics of membrane trafficking needs to be better defined, in particular, the TGN where multiple transport pathways are generated from the same compartment. The development of a technique to synchronize the trafficking of a newly synthesized protein from the Golgi, known as the retention using selective hooks (RUSH) system [146], will enable direct analysis of the spatial partitioning of cargo destined for different transport pathways. Indeed, a recent study using the RUSH system has demonstrated a role for the transmembrane and luminal domains in the segregation of cargo proteins in the Golgi prior to recruitment by specific transport carriers [147].

An interesting issue is the potential roles of cargo adaptors in functions other than the selection of cargo into transport carriers. For example, AP-1 interacts with kinesins to promote retrograde transport along the axon in neurons [148,149]. Identifying additional non-traditional roles of adaptor proteins could be very important. A number of pathogens can hijack trafficking pathways by selective interactions with adaptor protein complexes, for example HIV Vpu hijacks AP-1 dependent

trafficking [150]. It will be of considerable interest to assess whether the other adaptor protein complexes can also be selectively targeted by intracellular pathogens.

Finally, technical strategies to overcome the limitations of knockouts and knockdowns, where compensatory pathways and indirect effects may dominate, will be very important. The rapid inactivation of proteins, such as the knocksideways approach [65], will continue to be invaluable to reduce the time period between the loss of protein and the functional analyses. Moreover, the application of these technologies to in vivo whole organisms, will provide the capacity to directly examine the impact of inactivating adaptor complexes on physiological responses.

**Author Contributions:** J.Z.A.T. and P.A.G. contributed to the planning, writing, and design of the graphics.

**Funding:** This work was supported by funding from the National Health and Medical Research Council (APP1082600). JZAT also acknowledges support by the University of Melbourne International Graduate Scholarships.

**Conflicts of Interest:** The authors have no conflict to declare.

## Abbreviations

TGN	<i>trans</i> -Golgi network
AP	adaptor protein
GGA	Golgi-localized $\gamma$ -ear containing Arf binding proteins

## References

1. Ladinsky, M.S.; Mastrorade, D.N.; McIntosh, J.R.; Howell, K.E.; Staehelin, L.A. Golgi structure in three dimensions: Functional insights from the normal rat kidney cell. *J. Cell Biol.* **1999**, *144*, 1135–1149. [[CrossRef](#)] [[PubMed](#)]
2. Klumperman, J. Architecture of the mammalian golgi. *Cold Spring Harb. Perspect. Biol.* **2011**, *3*. [[CrossRef](#)] [[PubMed](#)]
3. Guo, Y.; Sirkis, D.W.; Schekman, R. Protein sorting at the trans-Golgi network. *Annu. Rev. Cell Dev. Biol.* **2014**, *30*, 169–206. [[CrossRef](#)] [[PubMed](#)]
4. De Matteis, M.A.; Luini, A. Exiting the Golgi complex. *Nat. Rev. Mol. Cell Biol.* **2008**, *9*, 273–284. [[CrossRef](#)] [[PubMed](#)]
5. Herrmann, J.M.; Spang, A. Intracellular parcel service: Current issues in intracellular membrane trafficking. *Methods Mol. Biol.* **2015**, *1270*, 1–12. [[PubMed](#)]
6. De Matteis, M.A.; Luini, A. Mendelian disorders of membrane trafficking. *N. Engl. J. Med.* **2011**, *365*, 927–938. [[CrossRef](#)] [[PubMed](#)]
7. Derby, M.C.; Gleeson, P.A. New insights into membrane trafficking and protein sorting. *Int. Rev. Cytol.* **2007**, *261*, 47–116.
8. Donaldson, J.G.; Jackson, C.L. Arf family G proteins and their regulators: Roles in membrane transport, development and disease. *Nat. Rev. Mol. Cell Biol.* **2011**, *12*, 362–375. [[CrossRef](#)]
9. Gillingham, A.K.; Munro, S. The small G proteins of the arf family and their regulators. *Annu. Rev. Cell Dev. Biol.* **2007**, *23*, 579–611. [[CrossRef](#)]
10. Gleeson, P.A.; Lock, J.G.; Luke, M.R.; Stow, J.L. Domains of the TGN: Coats, tethers and G proteins. *Traffic* **2004**, *5*, 315–326. [[CrossRef](#)]
11. Kirchhausen, T.; Bonifacino, J.S.; Riezman, H. Linking cargo to vesicle formation: Receptor tail interactions with coat proteins. *Curr. Opin. Cell Biol.* **1997**, *9*, 488–495. [[CrossRef](#)]
12. Bonifacino, J.S. Adaptor proteins involved in polarized sorting. *J. Cell Biol.* **2014**, *204*, 7–17. [[CrossRef](#)] [[PubMed](#)]
13. Traub, L.M.; Bonifacino, J.S. Cargo recognition in clathrin-mediated endocytosis. *Cold Spring Harb. Perspect. Biol.* **2013**, *5*, a016790. [[CrossRef](#)] [[PubMed](#)]
14. Faini, M.; Beck, R.; Wieland, F.T.; Briggs, J.A. Vesicle coats: Structure, function, and general principles of assembly. *Trends Cell Biol.* **2013**, *23*, 279–288. [[CrossRef](#)] [[PubMed](#)]

15. Hirst, J.; Edgar, J.R.; Borner, G.H.; Li, S.; Sahlender, D.A.; Antrobus, R.; Robinson, M.S. Contributions of epsinR and gadinin to clathrin-mediated intracellular trafficking. *Mol. Biol. Cell* **2015**, *26*, 3085–3103. [[CrossRef](#)]
16. Boucrot, E.; Pick, A.; Camdere, G.; Liska, N.; Evergren, E.; McMahon, H.T.; Kozlov, M.M. Membrane fission is promoted by insertion of amphipathic helices and is restricted by crescent bar domains. *Cell* **2012**, *149*, 124–136. [[CrossRef](#)]
17. Ford, M.G.; Mills, I.G.; Peter, B.J.; Vallis, Y.; Praefcke, G.J.; Evans, P.R.; McMahon, H.T. Curvature of clathrin-coated pits driven by epsin. *Nature* **2002**, *419*, 361–366. [[CrossRef](#)]
18. Hirst, J.; Irving, C.; Borner, G.H. Adaptor protein complexes ap-4 and ap-5: New players in endosomal trafficking and progressive spastic paraplegia. *Traffic* **2013**, *14*, 153–164. [[CrossRef](#)]
19. Robinson, M.S. Adaptable adaptors for coated vesicles. *Trends Cell Biol.* **2004**, *14*, 167–174. [[CrossRef](#)]
20. Robinson, M.S.; Bonifacino, J.S. Adaptor-related proteins. *Curr. Opin. Cell Biol.* **2001**, *13*, 444–453. [[CrossRef](#)]
21. Goldberg, J. Structural basis for activation of Arf GTPase: Mechanisms of guanine nucleotide exchange and GTP-myristoyl switching. *Cell* **1998**, *95*, 237–248. [[CrossRef](#)]
22. Bos, J.L.; Rehmann, H.; Wittinghofer, A. GEFs and GAPs: Critical elements in the control of small G proteins. *Cell* **2007**, *129*, 865–877. [[CrossRef](#)] [[PubMed](#)]
23. Antonny, B.; Beraud-Dufour, S.; Chardin, P.; Chabre, M. N-terminal hydrophobic residues of the G-protein ADP-ribosylation factor-1 insert into membrane phospholipids upon GDP to GTP exchange. *Biochemistry* **1997**, *36*, 4675–4684. [[CrossRef](#)] [[PubMed](#)]
24. Chavrier, P.; Gorvel, J.P.; Stelzer, E.; Simons, K.; Gruenberg, J.; Zerial, M. Hypervariable c-terminal domain of rab proteins acts as a targeting signal. *Nature* **1991**, *353*, 769–772. [[CrossRef](#)] [[PubMed](#)]
25. Kahn, R.A.; Cherfils, J.; Elias, M.; Lovering, R.C.; Munro, S.; Schurmann, A. Nomenclature for the human Arf family of GTP-binding proteins: ARF, ARL, and SAR proteins. *J. Cell Biol.* **2006**, *172*, 645–650. [[CrossRef](#)] [[PubMed](#)]
26. Manolea, E.; Chun, J.; Chen, D.W.; Clarke, I.; Summerfeldt, N.; Dacks, J.B.; Melancon, P. Arf3 is activated uniquely at the trans-golgi network by brefeldin A-inhibited guanine nucleotide exchange factors. *Mol. Biol. Cell* **2010**, *21*, 1836–1849. [[CrossRef](#)]
27. Hosaka, M.; Toda, K.; Takatsu, H.; Torii, S.; Murakami, K.; Nakayama, K. Structure and intracellular localization of mouse ADP-ribosylation factors type 1 to type 6 (Arf1-Arf6). *J. Biochem.* **1996**, *120*, 813–819. [[CrossRef](#)]
28. Cavenagh, M.M.; Whitney, J.A.; Carroll, K.; Zhang, C.; Boman, A.L.; Rosenwald, A.G.; Mellman, I.; Kahn, R.A. Intracellular distribution of Arf proteins in mammalian cells. Arf6 is uniquely localized to the plasma membrane. *J. Biol. Chem.* **1996**, *271*, 21767–21774. [[CrossRef](#)]
29. Volpicelli-Daley, L.A.; Li, Y.; Zhang, C.J.; Kahn, R.A. Isoform-selective effects of the depletion of ADP-ribosylation factors 1-5 on membrane traffic. *Mol. Biol. Cell* **2005**, *16*, 4495–4508. [[CrossRef](#)]
30. Toh, W.H.; Tan, J.Z.; Zulkefli, K.L.; Houghton, F.J.; Gleeson, P.A. Amyloid precursor protein traffics from the Golgi directly to early endosomes in an arl5b- and ap4-dependent pathway. *Traffic* **2017**, *18*, 159–175. [[CrossRef](#)]
31. Sadakata, T.; Shinoda, Y.; Sekine, Y.; Saruta, C.; Itakura, M.; Takahashi, M.; Furuichi, T. Interaction of calcium-dependent activator protein for secretion 1 (caps1) with the class ii ADP-ribosylation factor small GTPases is required for dense-core vesicle trafficking in the trans-Golgi network. *J. Biol. Chem.* **2010**, *285*, 38710–38719. [[CrossRef](#)] [[PubMed](#)]
32. Nakai, W.; Kondo, Y.; Saitoh, A.; Naito, T.; Nakayama, K.; Shin, H.W. Arf1 and Arf4 regulate recycling endosomal morphology and retrograde transport from endosomes to the Golgi apparatus. *Mol. Biol. Cell* **2013**, *24*, 2570–2581. [[CrossRef](#)] [[PubMed](#)]
33. Sadakata, T.; Itakura, M.; Kozaki, S.; Sekine, Y.; Takahashi, M.; Furuichi, T. Differential distributions of the Ca<sup>2+</sup>-dependent activator protein for secretion family proteins (caps2 and caps1) in the mouse brain. *J. Comp. Neurol.* **2006**, *495*, 735–753. [[CrossRef](#)] [[PubMed](#)]
34. Puertollano, R.; Aguilar, R.C.; Gorshkova, I.; Crouch, R.J.; Bonifacino, J.S. Sorting of mannose 6-phosphate receptors mediated by the GGAs. *Science* **2001**, *292*, 1712–1716. [[CrossRef](#)] [[PubMed](#)]
35. Puertollano, R.; Randazzo, P.A.; Presley, J.F.; Hartnell, L.M.; Bonifacino, J.S. The GGAs promote Arf-dependent recruitment of clathrin to TGN. *Cell* **2001**, *105*, 93–102. [[CrossRef](#)]

36. Mazelova, J.; Astuto-Gribble, L.; Inoue, H.; Tam, B.M.; Schonteich, E.; Prekeris, R.; Moritz, O.L.; Randazzo, P.A.; Deretic, D. Ciliary targeting motif VxPx directs assembly of a trafficking module through Arf4. *EMBO J.* **2009**, *28*, 183–192. [[CrossRef](#)] [[PubMed](#)]
37. Ren, X.; Farias, G.G.; Canagarajah, B.J.; Bonifacino, J.S.; Hurley, J.H. Structural basis for recruitment and activation of the AP-1 clathrin adaptor complex by Arf1. *Cell* **2013**, *152*, 755–767. [[CrossRef](#)] [[PubMed](#)]
38. D'Angelo, G.; Vicinanza, M.; Wilson, C.; De Matteis, M.A. Phosphoinositides in Golgi complex function. *Subcell. Biochem.* **2012**, *59*, 255–270. [[PubMed](#)]
39. Vicinanza, M.; D'Angelo, G.; Di Campli, A.; De Matteis, M.A. Function and dysfunction of the PI system in membrane trafficking. *EMBO J.* **2008**, *27*, 2457–2470. [[CrossRef](#)] [[PubMed](#)]
40. Wang, Y.J.; Wang, J.; Sun, H.Q.; Martinez, M.; Sun, Y.X.; Macia, E.; Kirchhausen, T.; Albanesi, J.P.; Roth, M.G.; Yin, H.L. Phosphatidylinositol 4 phosphate regulates targeting of clathrin adaptor AP-1 complexes to the Golgi. *Cell* **2003**, *114*, 299–310. [[CrossRef](#)]
41. Wang, J.; Sun, H.Q.; Macia, E.; Kirchhausen, T.; Watson, H.; Bonifacino, J.S.; Yin, H.L. PI4P promotes the recruitment of the GGA adaptor proteins to the *trans*-Golgi network and regulates their recognition of the ubiquitin sorting signal. *Mol. Biol. Cell* **2007**, *18*, 2646–2655. [[CrossRef](#)] [[PubMed](#)]
42. Godi, A.; Pertile, P.; Meyers, R.; Marra, P.; Di Tullio, G.; Iurisci, C.; Luini, A.; Corda, D.; De Matteis, M.A. ARF mediates recruitment of Ptdins-4-OH kinase-beta and stimulates synthesis of Ptdins(4,5)P2 on the Golgi complex. *Nat. Cell Biol.* **1999**, *1*, 280–287. [[CrossRef](#)] [[PubMed](#)]
43. Zhao, X.; Varnai, P.; Tuymetova, G.; Balla, A.; Toth, Z.E.; Oker-Blom, C.; Roder, J.; Jeromin, A.; Balla, T. Interaction of neuronal calcium sensor-1 (ncs-1) with phosphatidylinositol 4-kinase beta stimulates lipid kinase activity and affects membrane trafficking in Cos-7 cells. *J. Biol. Chem.* **2001**, *276*, 40183–40189. [[CrossRef](#)] [[PubMed](#)]
44. Haynes, L.P.; Thomas, G.M.; Burgoyne, R.D. Interaction of neuronal calcium sensor-1 and ADP-ribosylation factor 1 allows bidirectional control of phosphatidylinositol 4-kinase beta and *trans*-Golgi network-plasma membrane traffic. *J. Biol. Chem.* **2005**, *280*, 6047–6054. [[CrossRef](#)] [[PubMed](#)]
45. Blagoveshchenskaya, A.; Cheong, F.Y.; Rohde, H.M.; Glover, G.; Knodler, A.; Nicolson, T.; Boehmelt, G.; Mayinger, P. Integration of Golgi trafficking and growth factor signaling by the lipid phosphatase sac1. *J. Cell Biol.* **2008**, *180*, 803–812. [[CrossRef](#)]
46. Bonifacino, J.S. The gga proteins: Adaptors on the move. *Nat. Rev. Mol. Cell Biol.* **2004**, *5*, 23–32. [[CrossRef](#)]
47. Guardia, C.M.; De Pace, R.; Mattera, R.; Bonifacino, J.S. Neuronal functions of adaptor complexes involved in protein sorting. *Curr. Opin. Neurobiol.* **2018**, *51*, 103–110. [[CrossRef](#)]
48. Collins, B.M.; McCoy, A.J.; Kent, H.M.; Evans, P.R.; Owen, D.J. Molecular architecture and functional model of the endocytic AP2 complex. *Cell* **2002**, *109*, 523–535. [[CrossRef](#)]
49. Heldwein, E.E.; Macia, E.; Wang, J.; Yin, H.L.; Kirchhausen, T.; Harrison, S.C. Crystal structure of the clathrin adaptor protein 1 core. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 14108–14113. [[CrossRef](#)]
50. Dell'Angelica, E.C.; Puertollano, R.; Mullins, C.; Aguilar, R.C.; Vargas, J.D.; Hartnell, L.M.; Bonifacino, J.S. GGAs: A family of ADP ribosylation factor-binding proteins related to adaptors and associated with the Golgi complex. *J. Cell Biol.* **2000**, *149*, 81–93. [[CrossRef](#)]
51. Hirst, J.; Lui, W.W.Y.; Bright, N.A.; Totty, N.; Seaman, M.N.J.; Robinson, M.S. A family of proteins with gamma-adaptin and VHS domains that facilitate trafficking between the *trans*-Golgi network and the vacuole/lysosome. *J. Cell Biol.* **2000**, *149*, 67–79. [[CrossRef](#)] [[PubMed](#)]
52. Takatsu, H.; Yoshino, K.; Nakayama, K. Adaptor gamma ear homology domain conserved in gamma-adaptin and GGA proteins that interact with gamma-synergins. *Biochem. Biophys. Res. Commun.* **2000**, *271*, 719–725. [[CrossRef](#)] [[PubMed](#)]
53. Robinson, M.S. Forty years of clathrin-coated vesicles. *Traffic* **2015**, *16*, 1210–1238. [[CrossRef](#)] [[PubMed](#)]
54. Nielsen, M.S.; Madsen, P.; Christensen, E.I.; Nykjaer, A.; Gliemann, J.; Kasper, D.; Pohlmann, R.; Petersen, C.M. The sortilin cytoplasmic tail conveys Golgi-endosome transport and binds the VHS domain of the GGA2 sorting protein. *EMBO J.* **2001**, *20*, 2180–2190. [[CrossRef](#)] [[PubMed](#)]
55. Collins, B.M.; Watson, P.J.; Owen, D.J. The structure of the GGA1-GAT domain reveals the molecular basis for ARF binding and membrane association of GGAs. *Dev. Cell* **2003**, *4*, 321–332. [[CrossRef](#)]
56. Gallusser, A.; Kirchhausen, T. The beta 1 and beta 2 subunits of the AP complexes are the clathrin coat assembly components. *EMBO J.* **1993**, *12*, 5237–5244. [[CrossRef](#)] [[PubMed](#)]

57. ter Haar, E.; Harrison, S.C.; Kirchhausen, T. Peptide-in-groove interactions link target proteins to the beta-propeller of clathrin. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 1096–1100. [[CrossRef](#)]
58. Simpson, F.; Peden, A.A.; Christopoulou, L.; Robinson, M.S. Characterization of the adaptor-related protein complex, AP-3. *J. Cell Biol.* **1997**, *137*, 835–845. [[CrossRef](#)]
59. Dell’Angelica, E.C.; Mullins, C.; Bonifacino, J.S. AP-4, a novel protein complex related to clathrin adaptors. *J. Biol. Chem.* **1999**, *274*, 7278–7285. [[CrossRef](#)]
60. Hirst, J.; Bright, N.A.; Rous, B.; Robinson, M.S. Characterization of a fourth adaptor-related protein complex. *Mol. Biol. Cell* **1999**, *10*, 2787–2802. [[CrossRef](#)]
61. Ohno, H.; Tomemori, T.; Nakatsu, F.; Okazaki, Y.; Aguilar, R.C.; Foelsch, H.; Mellman, I.; Saito, T.; Shirasawa, T.; Bonifacino, J.S. Mu1B, a novel adaptor medium chain expressed in polarized epithelial cells. *FEBS Lett.* **1999**, *449*, 215–220. [[CrossRef](#)]
62. Guo, X.; Mattera, R.; Ren, X.; Chen, Y.; Retamal, C.; Gonzalez, A.; Bonifacino, J.S. The adaptor protein-1  $\mu$ 1B subunit expands the repertoire of basolateral sorting signal recognition in epithelial cells. *Dev. Cell* **2013**, *27*, 353–366. [[CrossRef](#)] [[PubMed](#)]
63. Hille-Rehfeld, A. Mannose 6-phosphate receptors in sorting and transport of lysosomal enzymes. *BBA Rev. Biomembr.* **1995**, *1241*, 177–194. [[CrossRef](#)]
64. Meyer, C.; Zizioli, D.; Lausmann, S.; Eskelinen, E.L.; Hamann, J.; Saftig, P.; von Figura, K.; Schu, P. Mu1a-adaptin-deficient mice: Lethality, loss of AP-1 binding and rerouting of mannose 6-phosphate receptors. *EMBO J.* **2000**, *19*, 2193–2203. [[CrossRef](#)] [[PubMed](#)]
65. Hirst, J.; Borner, G.H.; Antrobus, R.; Peden, A.A.; Hodson, N.A.; Sahlender, D.A.; Robinson, M.S. Distinct and overlapping roles for AP-1 and GGAs revealed by the “knocksideways” system. *Curr. Biol.* **2012**, *22*, 1711–1716. [[CrossRef](#)] [[PubMed](#)]
66. Farias, G.G.; Cuitino, L.; Guo, X.; Ren, X.; Jarnik, M.; Mattera, R.; Bonifacino, J.S. Signal-mediated, AP-1/clathrin-dependent sorting of transmembrane receptors to the somatodendritic domain of hippocampal neurons. *Neuron* **2012**, *75*, 810–823. [[CrossRef](#)] [[PubMed](#)]
67. Gravotta, D.; Carvajal-Gonzalez, J.M.; Mattera, R.; Deborde, S.; Banfelder, J.R.; Bonifacino, J.S.; Rodriguez-Boulan, E. The clathrin adaptor AP-1A mediates basolateral polarity. *Dev. Cell* **2012**, *22*, 811–823. [[CrossRef](#)]
68. Zhang, H.; Kim, A.; Abraham, N.; Khan, L.A.; Hall, D.H.; Fleming, J.T.; Gobel, V. Clathrin and AP-1 regulate apical polarity and lumen formation during *C. elegans* tubulogenesis. *Development* **2012**, *139*, 2071–2083. [[CrossRef](#)]
69. Folsch, H.; Pypaert, M.; Schu, P.; Mellman, I. Distribution and function of AP-1 clathrin adaptor complexes in polarized epithelial cells. *J. Cell Biol.* **2001**, *152*, 595–606. [[CrossRef](#)]
70. Folsch, H.; Pypaert, M.; Maday, S.; Pelletier, L.; Mellman, I. The AP-1A and AP-1B clathrin adaptor complexes define biochemically and functionally distinct membrane domains. *J. Cell Biol.* **2003**, *163*, 351–362. [[CrossRef](#)]
71. Zizioli, D.; Meyer, C.; Guhde, G.; Saftig, P.; von Figura, K.; Schu, P. Early embryonic death of mice deficient in gamma-adaptin. *J. Biol. Chem.* **1999**, *274*, 5385–5390. [[CrossRef](#)] [[PubMed](#)]
72. Tarpey, P.S.; Stevens, C.; Teague, J.; Edkins, S.; O’Meara, S.; Avis, T.; Barthorpe, S.; Buck, G.; Butler, A.; Cole, J.; et al. Mutations in the gene encoding the Sigma 2 subunit of the adaptor protein 1 complex, AP1S2, cause x-linked mental retardation. *Am. J. Hum. Genet.* **2006**, *79*, 1119–1124. [[CrossRef](#)] [[PubMed](#)]
73. Saillour, Y.; Zanni, G.; Des Portes, V.; Heron, D.; Guibaud, L.; Iba-Zizen, M.T.; Pedespan, J.L.; Poirier, K.; Castelnau, L.; Julien, C.; et al. Mutations in the AP1S2 gene encoding the sigma 2 subunit of the adaptor protein 1 complex are associated with syndromic X-linked mental retardation with hydrocephalus and calcifications in basal ganglia. *J. Med. Genet.* **2007**, *44*, 739–744. [[CrossRef](#)] [[PubMed](#)]
74. Montpetit, A.; Cote, S.; Brustein, E.; Drouin, C.A.; Lapointe, L.; Boudreau, M.; Meloche, C.; Drouin, R.; Hudson, T.J.; Drapeau, P.; et al. Disruption of AP1S1, causing a novel neurocutaneous syndrome, perturbs development of the skin and spinal cord. *PLoS Genet.* **2008**, *4*, e1000296. [[CrossRef](#)] [[PubMed](#)]
75. Pevsner, J.; Volknandt, W.; Wong, B.R.; Scheller, R.H. Two rat homologs of clathrin-associated adaptor proteins. *Gene* **1994**, *146*, 279–283. [[CrossRef](#)]
76. Newman, L.S.; McKeever, M.O.; Okano, H.J.; Darnell, R.B. Beta- $\mu$ 1, a cerebellar degeneration antigen, is a neuron-specific vesicle coat protein. *Cell* **1995**, *82*, 773–783. [[CrossRef](#)]
77. Dell’Angelica, E.C.; Ohno, H.; Ooi, C.E.; Rabinovich, E.; Roche, K.W.; Bonifacino, J.S. AP-3: An adaptor-like protein complex with ubiquitous expression. *EMBO J.* **1997**, *16*, 917–928. [[CrossRef](#)] [[PubMed](#)]

78. Dell'Angelica, E.C.; Shotelersuk, V.; Aguilar, R.C.; Gahl, W.A.; Bonifacino, J.S. Altered trafficking of lysosomal proteins in hermansky-pudlak syndrome due to mutations in the beta 3A subunit of the AP-3 adaptor. *Mol. Cell* **1999**, *3*, 11–21. [[CrossRef](#)]
79. Kantheti, P.; Qiao, X.; Diaz, M.E.; Peden, A.A.; Meyer, G.E.; Carskadon, S.L.; Kapfhamer, D.; Sufalko, D.; Robinson, M.S.; Noebels, J.L.; et al. Mutation in AP-3 delta in the mocha mouse links endosomal transport to storage deficiency in platelets, melanosomes, and synaptic vesicles. *Neuron* **1998**, *21*, 111–122. [[CrossRef](#)]
80. Peden, A.A.; Oorschot, V.; Hesser, B.A.; Austin, C.D.; Scheller, R.H.; Klumperman, J. Localization of the AP-3 adaptor complex defines a novel endosomal exit site for lysosomal membrane proteins. *J. Cell Biol.* **2004**, *164*, 1065–1076. [[CrossRef](#)]
81. Abou Jamra, R.; Philippe, O.; Raas-Rothschild, A.; Eck, S.H.; Graf, E.; Buchert, R.; Borck, G.; Ekici, A.; Brockschmidt, F.F.; Nothen, M.M.; et al. Adaptor protein complex 4 deficiency causes severe autosomal-recessive intellectual disability, progressive spastic paraplegia, shy character, and short stature. *Am. J. Hum. Genet.* **2011**, *88*, 788–795. [[CrossRef](#)] [[PubMed](#)]
82. Bauer, P.; Leshinsky-Silver, E.; Blumkin, L.; Schlipf, N.; Schroder, C.; Schicks, J.; Lev, D.; Riess, O.; Lerman-Sagie, T.; Schols, L. Mutation in the AP4B1 gene cause hereditary spastic paraplegia type 47 (SPG47). *Neurogenetics* **2012**, *13*, 73–76. [[CrossRef](#)] [[PubMed](#)]
83. Abdollahpour, H.; Alawi, M.; Kortum, F.; Beckstette, M.; Seemanova, E.; Komarek, V.; Rosenberger, G.; Kutsche, K. An AP4B1 frameshift mutation in siblings with intellectual disability and spastic tetraplegia further delineates the AP-4 deficiency syndrome. *Eur. J. Hum. Genet.* **2015**, *23*, 256–259. [[CrossRef](#)] [[PubMed](#)]
84. Tuysuz, B.; Bilguvar, K.; Kocer, N.; Yalcinkaya, C.; Caglayan, O.; Gul, E.; Sahin, S.; Comu, S.; Gunel, M. Autosomal recessive spastic tetraplegia caused by AP4M1 and AP4B1 gene mutation: Expansion of the facial and neuroimaging features. *Am. J. Med. Genet. PART A* **2014**, *164*, 1677–1685. [[CrossRef](#)] [[PubMed](#)]
85. Accogli, A.; Hamdan, F.F.; Poulin, C.; Nassif, C.; Rouleau, G.A.; Michaud, J.L.; Srour, M. A novel homozygous AP4B1 mutation in two brothers with AP-4 deficiency syndrome and ocular anomalies. *Am. J. Med. Genet. PART A* **2018**, *176*, 985–991. [[CrossRef](#)]
86. Moreno-De-Luca, A.; Helmers, S.L.; Mao, H.; Burns, T.G.; Melton, A.M.; Schmidt, K.R.; Fernhoff, P.M.; Ledbetter, D.H.; Martin, C.L. Adaptor protein complex-4 (AP-4) deficiency causes a novel autosomal recessive cerebral palsy syndrome with microcephaly and intellectual disability. *J. Med. Genet.* **2011**, *48*, 141–144. [[CrossRef](#)]
87. Najmabadi, H.; Hu, H.; Garshasbi, M.; Zemojtel, T.; Abedini, S.S.; Chen, W.; Hosseini, M.; Behjati, F.; Haas, S.; Jamali, P.; et al. Deep sequencing reveals 50 novel genes for recessive cognitive disorders. *Nature* **2011**, *478*, 57–63. [[CrossRef](#)]
88. Raza, M.H.; Mattered, R.; Morell, R.; Sainz, E.; Rahn, R.; Gutierrez, J.; Paris, E.; Root, J.; Solomon, B.; Brewer, C.; et al. Association between rare variants in AP4E1, a component of intracellular trafficking, and persistent stuttering. *Am. J. Hum. Genet.* **2015**, *97*, 715–725. [[CrossRef](#)]
89. Verkerk, A.J.; Schot, R.; Dumee, B.; Schellekens, K.; Swagemakers, S.; Bertoli-Avella, A.M.; Lequin, M.H.; Dudink, J.; Govaert, P.; van Zwol, A.L.; et al. Mutation in the AP4M1 gene provides a model for neuroaxonal injury in cerebral palsy. *Am. J. Hum. Genet.* **2009**, *85*, 40–52. [[CrossRef](#)]
90. Hardies, K.; May, P.; Djemie, T.; Tarta-Arsene, O.; Deconinck, T.; Craiu, D.; Helbig, I.; Suls, A.; Balling, R.; Weckhuysen, S.; et al. Recessive loss-of-function mutations in ap4s1 cause mild fever-sensitive seizures, developmental delay, and spastic paraplegia through loss of AP-4 complex assembly. *Hum. Mol. Genet.* **2014**. [[CrossRef](#)]
91. Matsuda, S.; Miura, E.; Matsuda, K.; Kakegawa, W.; Kohda, K.; Watanabe, M.; Yuzaki, M. Accumulation of ampa receptors in autophagosomes in neuronal axons lacking adaptor protein AP-4. *Neuron* **2008**, *57*, 730–745. [[CrossRef](#)] [[PubMed](#)]
92. Yap, C.C.; Murate, M.; Kishigami, S.; Muto, Y.; Kishida, H.; Hashikawa, T.; Yano, R. Adaptor protein complex-4 (AP-4) is expressed in the central nervous system neurons and interacts with glutamate receptor delta2. *Mol. Cell. Neurosci.* **2003**, *24*, 283–295. [[CrossRef](#)]
93. Aguilar, R.C.; Boehm, M.; Gorshkova, I.; Crouch, R.J.; Tomita, K.; Saito, T.; Ohno, H.; Bonifacino, J.S. Signal-binding specificity of the mu4 subunit of the adaptor protein complex AP-4. *J. Biol. Chem.* **2001**, *276*, 13145–13152. [[CrossRef](#)] [[PubMed](#)]
94. Janvier, K.; Bonifacino, J.S. Role of the endocytic machinery in the sorting of lysosome-associated membrane proteins. *Mol. Biol. Cell* **2005**, *16*, 4231–4242. [[CrossRef](#)] [[PubMed](#)]

95. Simmen, T.; Honing, S.; Icking, A.; Tikkanen, R.; Hunziker, W. AP-4 binds basolateral signals and participates in basolateral sorting in epithelial mdck cells. *Nat. Cell Biol.* **2002**, *4*, 154–159. [[CrossRef](#)] [[PubMed](#)]
96. Borner, G.H.; Antrobus, R.; Hirst, J.; Bhumbra, G.S.; Kozik, P.; Jackson, L.P.; Sahlender, D.A.; Robinson, M.S. Multivariate proteomic profiling identifies novel accessory proteins of coated vesicles. *J. Cell Biol.* **2012**, *197*, 141–160. [[CrossRef](#)] [[PubMed](#)]
97. Burgos, P.V.; Mardones, G.A.; Rojas, A.L.; daSilva, L.L.; Prabhu, Y.; Hurley, J.H.; Bonifacino, J.S. Sorting of the alzheimer's disease amyloid precursor protein mediated by the AP-4 complex. *Dev. Cell* **2010**, *18*, 425–436. [[CrossRef](#)] [[PubMed](#)]
98. Tan, J.Z.A.; Gleeson, P.A. The *trans*-Golgi network is a major site for alpha-secretase processing of amyloid precursor protein in primary neurons. *J. Biol. Chem.* **2019**, *294*, 1618–1631. [[CrossRef](#)] [[PubMed](#)]
99. Mattera, R.; Park, S.Y.; De Pace, R.; Guardia, C.M.; Bonifacino, J.S. AP-4 mediates export of ATG9A from the *trans*-Golgi network to promote autophagosome formation. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, E10697–E10706. [[CrossRef](#)] [[PubMed](#)]
100. Davies, A.K.; Itzhak, D.N.; Edgar, J.R.; Archuleta, T.L.; Hirst, J.; Jackson, L.P.; Robinson, M.S.; Borner, G.H.H. AP-4 vesicles contribute to spatial control of autophagy via RUSC-dependent peripheral delivery of ATG9A. *Nat. Commun.* **2018**, *9*, 3958. [[CrossRef](#)] [[PubMed](#)]
101. De Pace, R.; Skirzewski, M.; Damme, M.; Mattera, R.; Mercurio, J.; Foster, A.M.; Cuitino, L.; Jarnik, M.; Hoffmann, V.; Morris, H.D.; et al. Altered distribution of ATG9A and accumulation of axonal aggregates in neurons from a mouse model of AP-4 deficiency syndrome. *PLoS Genet.* **2018**, *14*, e1007363. [[CrossRef](#)] [[PubMed](#)]
102. Govero, J.; Doray, B.; Bai, H.; Kornfeld, S. Analysis of gga null mice demonstrates a non-redundant role for mammalian GGA2 during development. *PLoS ONE* **2012**, *7*, e30184. [[CrossRef](#)] [[PubMed](#)]
103. Doray, B.; Govero, J.; Kornfeld, S. Impact of genetic background on neonatal lethality of GGA2 gene-trap mice. *G3* **2014**, *4*, 885–890. [[CrossRef](#)] [[PubMed](#)]
104. Doray, B.; Ghosh, P.; Griffith, J.; Geuze, H.J.; Kornfeld, S. Cooperation of GGAs and AP-1 in packaging MPRs at the *trans*-Golgi network. *Science* **2002**, *297*, 1700–1703. [[CrossRef](#)]
105. Toh, W.H.; Chia, P.Z.C.; Hossain, M.I.; Gleeson, P.A. GGA1 regulates signal-dependent sorting of BACE1 to recycling endosomes, which moderates Abeta production. *Mol. Biol. Cell* **2018**, *29*, 191–208. [[CrossRef](#)] [[PubMed](#)]
106. Uemura, T.; Kametaka, S.; Waguri, S. GGA2 interacts with EGFR cytoplasmic domain to stabilize the receptor expression and promote cell growth. *Sci. Rep.* **2018**, *8*, 1368. [[CrossRef](#)]
107. Puertollano, R.; Bonifacino, J.S. Interactions of GGA3 with the ubiquitin sorting machinery. *Nat. Cell Biol.* **2004**, *6*, 244–251. [[CrossRef](#)]
108. Tesco, G.; Koh, Y.H.; Kang, E.L.; Cameron, A.N.; Das, S.; Sena-Esteves, M.; Hiltunen, M.; Yang, S.H.; Zhong, Z.; Shen, Y.; et al. Depletion of GGA3 stabilizes BACE and enhances beta-secretase activity. *Neuron* **2007**, *54*, 721–737. [[CrossRef](#)]
109. Kang, E.L.; Cameron, A.N.; Piazza, F.; Walker, K.R.; Tesco, G. Ubiquitin regulates GGA3-mediated degradation of BACE1. *J. Biol. Chem.* **2010**, *285*, 24108–24119. [[CrossRef](#)]
110. Bonifacino, J.S.; Traub, L.M. Signals for sorting of transmembrane proteins to endosomes and lysosomes. *Annu. Rev. Biochem.* **2003**, *72*, 395–447. [[CrossRef](#)]
111. Park, S.Y.; Guo, X. Adaptor protein complexes and intracellular transport. *Biosci. Rep.* **2014**, *34*, 00123. [[CrossRef](#)] [[PubMed](#)]
112. Hofmann, M.W.; Honing, S.; Rodionov, D.; Dobberstein, B.; von Figura, K.; Bakke, O. The leucine-based sorting motifs in the cytoplasmic domain of the invariant chain are recognized by the clathrin adaptors AP1 and AP2 and their medium chains. *J. Biol. Chem.* **1999**, *274*, 36153–36158. [[CrossRef](#)] [[PubMed](#)]
113. Honing, S.; Sandoval, I.V.; von Figura, K. A di-leucine-based motif in the cytoplasmic tail of LIMP-II and tyrosinase mediates selective binding of AP-3. *EMBO J.* **1998**, *17*, 1304–1314. [[CrossRef](#)] [[PubMed](#)]
114. Le Borgne, R.; Alconada, A.; Bauer, U.; Hoflack, B. The mammalian AP-3 adaptor-like complex mediates the intracellular transport of lysosomal membrane glycoproteins. *J. Biol. Chem.* **1998**, *273*, 29451–29461. [[CrossRef](#)] [[PubMed](#)]
115. Owen, D.J.; Evans, P.R. A structural explanation for the recognition of tyrosine-based endocytotic signals. *Science* **1998**, *282*, 1327–1332. [[CrossRef](#)] [[PubMed](#)]

116. Mardones, G.A.; Burgos, P.V.; Lin, Y.; Kloer, D.P.; Magadan, J.G.; Hurley, J.H.; Bonifacino, J.S. Structural basis for the recognition of tyrosine-based sorting signals by the  $\mu$ 3A subunit of the AP-3 adaptor complex. *J. Biol. Chem.* **2013**, *288*, 9563–9571. [[CrossRef](#)] [[PubMed](#)]
117. Ohno, H.; Stewart, J.; Fournier, M.C.; Bosshart, H.; Rhee, I.; Miyatake, S.; Saito, T.; Gallusser, A.; Kirchhausen, T.; Bonifacino, J.S. Interaction of tyrosine-based sorting signals with clathrin-associated proteins. *Science* **1995**, *269*, 1872–1875. [[CrossRef](#)]
118. Honing, S.; Griffith, J.; Geuze, H.J.; Hunziker, W. The tyrosine-based lysosomal targeting signal in lamp-1 mediates sorting into Golgi-derived clathrin-coated vesicles. *EMBO J.* **1996**, *15*, 5230–5239. [[CrossRef](#)]
119. Salamero, J.; Le Borgne, R.; Saudrais, C.; Goud, B.; Hoflack, B. Expression of major histocompatibility complex class II molecules in hela cells promotes the recruitment of AP-1 Golgi-specific assembly proteins on golgi membranes. *J. Biol. Chem.* **1996**, *271*, 30318–30321. [[CrossRef](#)]
120. Prabhu, Y.; Burgos, P.V.; Schindler, C.; Farias, G.G.; Magadan, J.G.; Bonifacino, J.S. Adaptor protein 2-mediated endocytosis of the beta-secretase BACE1 is dispensable for amyloid precursor protein processing. *Mol. Biol. Cell* **2012**, *23*, 2339–2351. [[CrossRef](#)]
121. He, X.; Chang, W.P.; Koelsch, G.; Tang, J. Memapsin 2 (beta-secretase) cytosolic domain binds to the VHS domains of GGA1 and GGA2: Implications on the endocytosis mechanism of memapsin 2. *FEBS Lett.* **2002**, *524*, 183–187. [[CrossRef](#)]
122. He, X.; Li, F.; Chang, W.P.; Tang, J. GGA proteins mediate the recycling pathway of memapsin 2 (BACE). *J. Biol. Chem.* **2005**, *280*, 11696–11703. [[CrossRef](#)]
123. Navarro Negredo, P.; Edgar, J.R.; Wrobel, A.G.; Zaccai, N.R.; Antrobus, R.; Owen, D.J.; Robinson, M.S. Contribution of the clathrin adaptor AP-1 subunit micro1 to acidic cluster protein sorting. *J. Cell Biol.* **2017**, *216*, 2927–2943.
124. Stockli, J.; Honing, S.; Rohrer, J. The acidic cluster of the CK2 site of the cation-dependent mannose 6-phosphate receptor (CD-MPR) but not its phosphorylation is required for GGA1 and AP-1 binding. *J. Biol. Chem.* **2004**, *279*, 23542–23549. [[CrossRef](#)]
125. Teuchert, M.; Schafer, W.; Berghofer, S.; Hoflack, B.; Klenk, H.D.; Garten, W. Sorting of furin at the trans-Golgi network. Interaction of the cytoplasmic tail sorting signals with AP-1 Golgi-specific assembly proteins. *J. Biol. Chem.* **1999**, *274*, 8199–8207. [[CrossRef](#)]
126. Jia, X.; Singh, R.; Homann, S.; Yang, H.; Guatelli, J.; Xiong, Y. Structural basis of evasion of cellular adaptive immunity by HIV-1 Nef. *Nat. Struct. Mol. Biol.* **2012**, *19*, 701–706. [[CrossRef](#)]
127. Aguilar, R.C.; Ohno, H.; Roche, K.W.; Bonifacino, J.S. Functional domain mapping of the clathrin-associated adaptor medium chains  $\mu$ 1 and  $\mu$ 2. *J. Biol. Chem.* **1997**, *272*, 27160–27216. [[CrossRef](#)]
128. Ross, B.H.; Lin, Y.; Corales, E.A.; Burgos, P.V.; Mardones, G.A. Structural and functional characterization of cargo-binding sites on the  $\mu$ 4-subunit of adaptor protein complex 4. *PLoS ONE* **2014**, *9*, e88147. [[CrossRef](#)]
129. Chernomordik, L.V.; Kozlov, M.M. Protein-lipid interplay in fusion and fission of biological membranes. *Annu. Rev. Biochem.* **2003**, *72*, 175–207. [[CrossRef](#)]
130. Frolov, V.A.; Escalada, A.; Akimov, S.A.; Shnyrova, A.V. Geometry of membrane fission. *Chem. Phys. Lipids* **2015**, *185*, 129–140. [[CrossRef](#)]
131. Rosenthal, J.A.; Chen, H.; Slepnev, V.I.; Pellegrini, L.; Salcini, A.E.; Di Fiore, P.P.; De Camilli, P. The epsins define a family of proteins that interact with components of the clathrin coat and contain a new protein module. *J. Biol. Chem.* **1999**, *274*, 33959–33965. [[CrossRef](#)]
132. Legendre-Guillemain, V.; Wasiak, S.; Hussain, N.K.; Angers, A.; McPherson, P.S. ENTH/ANTH proteins and clathrin-mediated membrane budding. *J. Cell Sci.* **2004**, *117*, 9–18. [[CrossRef](#)]
133. Hirst, J.; Motley, A.; Harasaki, K.; Peak Chew, S.Y.; Robinson, M.S. EpsinR: An ENTH domain-containing protein that interacts with AP-1. *Mol. Biol. Cell* **2003**, *14*, 625–641. [[CrossRef](#)]
134. Archuleta, T.L.; Frazier, M.N.; Monken, A.E.; Kendall, A.K.; Harp, J.; McCoy, A.J.; Creanza, N.; Jackson, L.P. Structure and evolution of ENTH and VPS/ENTH-like domains in tepsin. *Traffic* **2017**, *18*, 590–603. [[CrossRef](#)]
135. Miller, S.E.; Collins, B.M.; McCoy, A.J.; Robinson, M.S.; Owen, D.J. A SNARE-adaptor interaction is a new mode of cargo recognition in clathrin-coated vesicles. *Nature* **2007**, *450*, 570–574. [[CrossRef](#)]
136. Ferguson, S.M.; De Camilli, P. Dynamin, a membrane-remodelling GTPase. *Nat. Rev. Mol. Cell Biol* **2012**, *13*, 75–88. [[CrossRef](#)]

137. Owen, D.J.; Vallis, Y.; Noble, M.E.; Hunter, J.B.; Dafforn, T.R.; Evans, P.R.; McMahon, H.T. A structural explanation for the binding of multiple ligands by the alpha-adaptin appendage domain. *Cell* **1999**, *97*, 805–815. [[CrossRef](#)]
138. Owen, D.J.; Vallis, Y.; Pearse, B.M.; McMahon, H.T.; Evans, P.R. The structure and function of the beta 2-adaptin appendage domain. *EMBO J.* **2000**, *19*, 4216–4227. [[CrossRef](#)]
139. Mills, I.G.; Praefcke, G.J.; Vallis, Y.; Peter, B.J.; Olesen, L.E.; Gallop, J.L.; Butler, P.J.; Evans, P.R.; McMahon, H.T. EpsinR: An AP1/clathrin interacting protein involved in vesicle trafficking. *J. Cell Biol.* **2003**, *160*, 213–222. [[CrossRef](#)]
140. Mattera, R.; Guardia, C.M.; Sidhu, S.S.; Bonifacino, J.S. Bivalent motif-ear interactions mediate the association of the accessory protein tepsin with the AP-4 adaptor complex. *J. Biol. Chem.* **2015**, *290*, 30736–43079. [[CrossRef](#)]
141. Frazier, M.N.; Davies, A.K.; Voehler, M.; Kendall, A.K.; Borner, G.H.; Chazin, W.J.; Robinson, M.S.; Jackson, L.P. Molecular basis for the interaction between AP4  $\beta$ 4 and its accessory protein, Tepsin. *Traffic* **2016**, *17*, 400–415. [[CrossRef](#)]
142. Itzhak, D.N.; Tyanova, S.; Cox, J.; Borner, G.H. Global, quantitative and dynamic mapping of protein subcellular localization. *Elife* **2016**, *5*, e16950. [[CrossRef](#)]
143. Itzhak, D.N.; Davies, C.; Tyanova, S.; Mishra, A.; Williamson, J.; Antrobus, R.; Cox, J.; Weekes, M.P.; Borner, G.H.H. A mass spectrometry-based approach for mapping protein subcellular localization reveals the spatial proteome of mouse primary neurons. *Cell Rep.* **2017**, *20*, 2706–2718. [[CrossRef](#)]
144. Hirst, J.; Itzhak, D.N.; Antrobus, R.; Borner, G.H.H.; Robinson, M.S. Role of the AP-5 adaptor protein complex in late endosome-to-Golgi retrieval. *PLoS Biol.* **2018**, *16*, e2004411. [[CrossRef](#)]
145. Theos, A.C.; Tenza, D.; Martina, J.A.; Hurbain, I.; Peden, A.A.; Sviderskaya, E.V.; Stewart, A.; Robinson, M.S.; Bennett, D.C.; Cutler, D.F.; et al. Functions of adaptor protein (AP)-3 and AP-1 in tyrosinase sorting from endosomes to melanosomes. *Mol. Biol. Cell* **2005**, *16*, 5356–5372. [[CrossRef](#)]
146. Boncompain, G.; Divoux, S.; Gareil, N.; de Forges, H.; Lescure, A.; Latreche, L.; Mercanti, V.; Jollivet, F.; Raposo, G.; Perez, F. Synchronization of secretory protein traffic in populations of cells. *Nat. Methods* **2012**, *9*, 493–498. [[CrossRef](#)]
147. Chen, Y.; Gershlick, D.C.; Park, S.Y.; Bonifacino, J.S. Segregation in the Golgi complex precedes export of endolysosomal proteins in distinct transport carriers. *J. Cell Biol.* **2017**, *216*, 4141–4151. [[CrossRef](#)]
148. Nakagawa, T.; Setou, M.; Seog, D.; Ogasawara, K.; Dohmae, N.; Takio, K.; Hirokawa, N. A novel motor, KIF13A, transports mannose-6-phosphate receptor to plasma membrane through direct interaction with ap-1 complex. *Cell* **2000**, *103*, 569–581. [[CrossRef](#)]
149. Schmidt, M.R.; Maritzen, T.; Kukhtina, V.; Higman, V.A.; Doglio, L.; Barak, N.N.; Strauss, H.; Oschkinat, H.; Dotti, C.G.; Haucke, V. Regulation of endosomal membrane traffic by a Gadkin/AP-1/kinesin KIF5 complex. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 15344–15349. [[CrossRef](#)]
150. Jia, X.; Weber, E.; Tokarev, A.; Lewinski, M.; Rizk, M.; Suarez, M.; Guatelli, J.; Xiong, Y. Structural basis of HIV-1 Vpu-mediated BST2 antagonism via hijacking of the clathrin adaptor protein complex 1. *Elife* **2014**, *3*, e02362. [[CrossRef](#)]

