

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Eapen, N;Borland, ML;Phillips, N;Kochar, A;Dalton, S;Cheek, JA;Gilhotra, Y;Neutze, J;Lyttle, MD;Donath, S;Crowe, L;Dalziel, SR;Oakley, E;Williams, A;Hearps, S;Bressan, S;Babl, FE

Title:

Neonatal head injuries: A prospective Paediatric Research in Emergency Departments International Collaborative cohort study

Date:

2020-05-01

Citation:

Eapen, N., Borland, M. L., Phillips, N., Kochar, A., Dalton, S., Cheek, J. A., Gilhotra, Y., Neutze, J., Lyttle, M. D., Donath, S., Crowe, L., Dalziel, S. R., Oakley, E., Williams, A., Hearps, S., Bressan, S. & Babl, F. E. (2020). Neonatal head injuries: A prospective Paediatric Research in Emergency Departments International Collaborative cohort study. *Journal of Paediatrics and Child Health*, 56 (5), pp.764-769. <https://doi.org/10.1111/jpc.14736>.

Persistent Link:

<https://hdl.handle.net/11343/286777>

Babl Franz (Orcid ID: 0000-0002-1107-2187)
Cheek John (Orcid ID: 0000-0002-3615-3821)

Neonatal Head Injuries - a prospective PREDICT cohort study
An Original Article

Nitaa Eapen, Meredith L. Borland, Natalie Phillips, Amit Kochar, Sarah Dalton, John A. Cheek, Yuri Gilhotra, Jocelyn Neutze, Mark D. Lyttle, Susan Donath, Louise Crowe, Stuart R Dalziel, Ed Oakley, Amanda Williams, Stephen Hearps, Silvia Bressan, Franz E Babl

Emergency Department, Royal Children's Hospital, 50 Flemington Rd, Parkville, VIC, 3052, Australia.

AUTHORS:

Nitaa Eapen Bsc^{1,2}
nitaa.eapen@mcri.edu.au

Meredith L. Borland MBBS^{3,4}
meredith.borland@health.wa.gov.au

Natalie Phillips MBBS⁵⁻⁷
natalie.phillips@merton-oxford.com

Amit Kochar MD⁸
amit.kochar@health.sa.gov.au

Sarah Dalton BMed⁹
sarah.dalton@health.nsw.gov.au

John A. Cheek MBBS^{1,2,10}
john.cheek@rch.org.au

Yuri Gilhotra MBBS⁵
yurigilhotra@me.com

Jocelyn Neutze MBChB¹¹
jocelyn.neutze@middlemore.co.nz

Mark D. Lyttle MBChB^{2,12,13}
mdlyttle77@gmail.com

Susan Donath MA^{2,14}
susan.donath@mcri.edu.au

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/jpc.14736](https://doi.org/10.1111/jpc.14736)

Louise Crowe PhD²
louise.crowe@mcri.edu.au

Stuart R Dalziel PhD^{15,16}
sdalziel@adhb.govt.nz

Ed Oakley MBBS^{1,2,14}
ed.oakley@rch.org.au

Amanda Williams²
Amanda.williams@mcri.edu.au

Stephen Hearps PGDipPsych²
stephen.hearps@mcri.edu.au

Silvia Bressan MD, PhD^{2,17}
silviabress@gmail.com

Franz E Babl MD^{1,2,14}
franz.babl@rch.org.au

¹ Royal Children's Hospital, Melbourne.
Address: Emergency Department, Royal Children's Hospital, 50 Flemington Rd, Parkville, VIC, 3052, Australia.

² Murdoch Childrens Research Institute, Melbourne.
Address: 50 Flemington Rd, Parkville, VIC, 3052, Australia.

³ Princess Margaret Hospital for Children, Perth.
Address: Emergency Department, Princess Margaret Hospital for Children, Roberts Rd, Subiaco, Perth, WA, 6008, Australia.

⁴ Schools of Paediatrics and Child Health and Primary, Aboriginal and Rural Healthcare, University of Western Australia.
Address: 35 Stirling Hwy, Crawley, WA, 6009, Australia.

⁵ Emergency department, Lady Cilento Children's Hospital, Brisbane.
Address: 501 Stanley St, South Brisbane, QLD, 4101, Australia.

⁶ Children's Health Research Centre, The University of Queensland, Medical Research Institute, Brisbane.
Address: 62 Graham St, South Brisbane, QLD, 4101, Australia

⁷ Faculty of Medicine, The University of Queensland, Brisbane.

Address: 20 Weightman St, Herston, QLD, 4006, Australia.

⁸ Women's & Children's Hospital, Adelaide.

Address: Emergency Department, Women's & Children's Hospital, 72 King William St, North Adelaide, SA, 5006, Australia.

⁹ The Children's Hospital at Westmead, Sydney.

Address: Emergency Department, The Children's Hospital at Westmead, 212 Hawkesbury Rd, Westmead, NSW, 2145, Australia.

¹⁰ Monash Medical Centre, Melbourne.

Address: Emergency Department, Monash Medical Centre, 246 Clayton Rd, Clayton, VIC, 3186, Australia.

¹¹ Kidzfirst Middlemore Hospital, Auckland.

Address: Emergency Department, Kidzfirst Middlemore Hospital, 100 Hospital Rd, Auckland, 2025, New Zealand.

¹² Bristol Royal Hospital for Children, Bristol.

Address: Emergency Department, Bristol Children's Hospital, Paul O'Gorman Building, Upper Maudlin St, Bristol, BS2 8BJ, United Kingdom.

¹³ Academic Department of Emergency Care, University of the West of England, Bristol.

Address: Blackberry Hill, Bristol, BS16 1XS, United Kingdom.

¹⁴ Department of Paediatrics, Faculty of Medicine, Dentistry and Health Sciences, University of Melbourne, Melbourne.

Address: Grattan St, Parkville, VIC, 3010, Australia.

¹⁵ Starship Children's Health, Auckland.

Address: Emergency Department, Starship Children's Health, 2 Park Rd, Grafton, Auckland, 1023, New Zealand.

¹⁶ Liggins Institute, University of Auckland, Auckland.

Address: 85 Park Ave, Grafton, Auckland, 1023, New Zealand.

¹⁷ Department of Women's and Child Health, University of Padova, Padova.

Address: Via Giustiniani3, 2 35128, Padova, PD, Italy.

CORRESPONDING AUTHOR:

Franz E Babl
Emergency Research
Murdoch Childrens Research Institute
Flemington Rd
Parkville
Victoria 3052
Australia
Phone 61 3 9345 6153
Fax 61 3 9345 600
Email franz.babl@rch.org.au

ABSTRACT

Aim: To characterise the causes, clinical characteristics and short-term outcomes of neonates who presented to paediatric emergency departments with head injury.

Methods: Secondary analysis of a prospective dataset of paediatric head injuries at 10 emergency departments in Australia and New Zealand. Patients without neuroimaging were followed up by telephone call. We extracted epidemiological information, clinical findings and outcomes in neonates (≤ 28 days).

Results: Of 20,137 children with head injuries, 93 (0.5%) occurred in neonates. These were mostly fall-related (75.2%), commonly from a caregiver's arms, or due to being accidentally struck by a person/object (20.4%). There were 3 cases of non-accidental head injuries (3.2%). Most neonates were asymptomatic (67.7%) and many had no findings on examination (47.3%). Most neonates had a GCS 15 (89.2%) or 14 (7.5%). 15.1% presented with vomiting and 5.4% were abnormally drowsy. None had experienced a loss of consciousness. The most common findings on examination were scalp haematoma (28.0%) and possible palpable skull fracture (6.5%). 8.6% underwent CT brain scan and 4.3% an ultrasound. Five of eight CTs (5.4% of neonates overall) showed traumatic brain injury (TBI) and 2 of 4 (2.2% overall) had TBI on ultrasound. 37% were admitted, 1 patient was intubated, none had neurosurgery or died.

Conclusions: Neonatal head injuries are rare with a mostly benign short term outcome and are appropriate for observation. However, non-accidental injuries need to be considered.

Keywords: Neonates, head injuries, emergency department

What is already known?

- Neonates presenting to emergency departments with head injuries are concerning for both clinicians and parents.
- They may have an increased anatomical vulnerability, yet information on out-of-hospital neonatal head injuries is limited.

What this paper adds?

- Most neonates presented asymptotically and many had no findings on examination.
- They mostly had benign short term outcomes and were generally appropriate for observation only.

INTRODUCTION

Children of all ages present to emergency departments (EDs) and acute care clinics with head injuries, one of the most concerning age groups are neonates. Their skulls are thin and pliable to facilitate passage through the birth canal^{1,2} and their brains less protected in head trauma^{3,4}. While minor trauma can result in intracranial injuries or skull fractures, it may be difficult for clinicians to assess deteriorating GCS in preverbal infants; intracranial injuries in children aged under two years can be asymptomatic^{5,6} and familiar signs of more severe head injury in older children and adults, like vomiting^{7,8,18}, may not be reliable. At the same time, clinicians will be particularly reluctant to use computed tomography (CT) scans in young children due to the increased risk of radiation induced cancer⁹⁻¹².

CT scan remains the preferred imaging modality for suspected intracranial injury¹³ requiring immediate intervention, as some areas of the brain are difficult to visualise with cranial ultrasound¹⁴. However, in mild neonatal HI, the significant risks of ionising radiation should be considered carefully. Magnetic resonance imaging (MRI) avoids ionising radiation but may be difficult to obtain in the acute setting and may require sedation¹⁴. While cranial ultrasound is a useful tool to evaluate brain abnormalities in neonates, it has limitations; cranial ultrasound is insensitive to detect parenchymal injuries and small acute subdural hematomas¹⁵ well as other intracranial injuries¹⁴

While a number of studies explore how head injuries in children less than 2 years of age differ from older children, particularly around the importance of scalp haematomas and vomiting^{6,16-18}, the literature on injuries in very young infants is limited. The mechanism of injury in young infants differs from older infants and often includes falls from caregiver's arms and from change tables^{19,20}. Non-accidental injuries also occur at higher rates in the early months of life^{21,22}. Neonatal specific papers focus on injuries sustained in hospital through falls in admitted infants^{23,24}.

We therefore set out to analyse the neonatal cohort from a large multicentre prospective study of head injuries in all childhood age groups. We extracted information on injury mechanism, severity and short term outcomes.

METHODS

Study design, setting and patients

We performed a secondary analysis for neonatal head injuries in a prospective multi-centre observational study which enrolled children presenting with head injuries of any severity to 10 paediatric emergency departments (EDs) in Australia and New Zealand between April 2011 and November 2014. All EDs are members of the Paediatric Research in Emergency Departments International Collaborative (PREDICT) research network²⁵. The study was approved by the ethics offices at all participating sites.

For the primary study we collected and analysed predictor and outcome variables for three head injury clinical decision rules (CDRs) for neuroimaging in head injured children^{18,26,27}. The following patients were excluded in the original study: trivial facial injury only, patient/family refusal to participate, referral from ED triage to an external provider (i.e. not seen in the ED), did not wait to be seen or neuroimaging done prior to the transfer to a study site.

The present study defined neonates as 28 days old or younger. Activity, place and mechanism of injury were identified by epidemiology codes assigned by research assistants (RAs) at each site from ED clinician documentation, medical records and the follow-up call. These epidemiology codes are based on Victorian government codes which are routinely collected in EDs in Victoria. We had collected these codes for patients across all sites; for the purposes of this study we added additional

activity codes to more accurately reflect the synopsis of the injury for the neonatal cohort if the existing Victorian government codes were not sufficiently detailed.

Study Procedures

Details of the study protocol are described in detail elsewhere^{28,29}. Patients were enrolled by the treating ED clinician who collected clinical data. RAs recorded ED and hospital management data after the visit and conducted telephone follow-up for patients who had not undergone neuroimaging. Data included clinical decision rule predictor variables, demographic and epidemiological information, and neuroimaging, admission and neurosurgery details.

To identify neonates in our population, we searched the database by age. We then manually reviewed all eligible case report forms (CRFs) to descriptively record the synopsis of the injury. Where available and necessary, we used medical records to augment the information from the CRFs.

Definitions

Glasgow Coma Scale

We used the Glasgow Coma Scale (GCS)³⁰ score as assigned by the ED clinician on their assessment, or if not available, GCS at triage²⁹.

Results of neuroimaging and neurosurgery

We used senior radiologist reports to determine the results of CT and ultrasound scans and operative reports for patients who underwent neurosurgery.

Traumatic brain injury (TBI) on CT or ultrasound was defined as intracranial haemorrhage or contusion, cerebral oedema, traumatic infarction, diffuse axonal injury, shearing injury, sigmoid sinus thrombosis, midline shift of intracranial contents or signs of brain herniation, diastasis of the skull, pneumocephalus or depressed skull fracture. In addition we included any type of skull fracture (depressed, non-depressed, basal, unknown type).

Neurosurgical intervention was defined as intracranial pressure monitoring, elevation of depressed skull fracture, ventriculostomy, haematoma evacuation, lobectomy, tissue debridement or dura repair.

Clinically important traumatic brain injury

Clinically-important TBI (ciTBI) was defined as a death from TBI, neurosurgery, intubation for more than 24 h for TBI, or hospital admission of 2 nights or more associated with TBI on CT¹⁸.

Statistical analysis

Data were entered into Epidata (The Epidata Association, Odense, Denmark), and later REDCap³¹, and analysed using Stata 13 (Statacorp, College Station, Texas, USA). Descriptive statistics were calculated with 95% confidence intervals (CI) where relevant.

RESULTS

We identified 93 neonates who presented with head injury (0.5% of the original cohort of 20,137 children). The mean age was 16.3 days; 59.1% were male. Most had a GCS of 15 (89.2%, 95%CI: 81.1-94.7) or 14 (7.5%, 95%CI: 3.1-14.9). Only 3.2% had a GCS of ≤ 13 , and none had a GCS of ≤ 8 . The most frequent signs and symptoms were vomiting (15.1%, 95%CI: 8.5-24.0) and abnormal drowsiness (5.4%, 95%CI: 1.8-12.1). None had loss of consciousness. The most common abnormalities on examination were scalp haematomas (28.0%, 95%CI: 19.1-38.2) and possible palpable skull fracture (6.5%, 95%CI: 2.4-13.5) (**Table 1**). Most were asymptomatic (67.7%); 47.3% were asymptomatic and had no findings on examination.

Most of the neonatal head injuries were fall related (75.2%) with a smaller percentage attributable to being struck by a person/object (20.4%). The most common mechanism was a fall from caregiver's arms (28.0%) followed by struck by inanimate stationary object (12.9%) and fall from couch (9.7%). Most injuries (75%) occurred at home in a bedroom or a living area (**Table 2**). Half presented between 2PM and 9PM; only 8.6% presented between midnight and 6 AM (**Figure 1**). There were three non-accidental head injuries confirmed by local multidisciplinary committees (3.2%).

Eleven of 93 neonates (11.8%) underwent imaging of the head which comprised of CT scan in eight neonates (8.6%), transfontanelle ultrasound in 4 (4.3%) neonates, MRI in two neonates (2.2%) and one X-ray (1.1). Two of the 11 neonates who had imaging of the head were asymptomatic and had no findings on examination. The presence of vomiting, abnormal drowsiness and irritability in the initial presentation was not correlated with neuroimaging. However, all six neonates who sustained palpable

skull fractures underwent neuroimaging and eight of the eleven neonates who underwent imaging of the head had a scalp haematoma. In the four neonates that underwent ultrasound, two showed TBI, which was confirmed by CT. In total, TBI on CT was observed in five of 11. Overall, one third (36.6%, 95%CI: 26.8-47.2) of neonates, including 13 asymptomatic infants with no examination findings, were admitted with a median length of stay of 1 day. Only one patient was intubated > 1 day. Four patients had ciTBI (4.3%, 95%CI: 1.2-10.6), as defined by admission $\geq$ 2 days. This included the one intubated patient. No neonates had neurosurgery and there were no deaths. (**Table 1, Table 3**).

Of the five TBI on CT patients, four had intracranial haemorrhage/contusion and skull fracture (of which two were depressed skull fractures). One had midline shift of intracranial contents or signs of brain herniation, diastasis of the skull and sigmoid sinus thrombosis. Of the three non-accidental head injuries, one neonate had no intracranial findings on neuroimaging, and two neonates had skull fractures and intracranial hemorrhages; one neonate had a ciTBI. (**Table 3**)

DISCUSSION

In this multicentre, prospective study we characterised the causes, clinical characteristics and outcomes of 93 neonates with head injuries presenting to EDs. The majority of neonatal head injuries presenting to Australasian EDs were surprisingly benign with two thirds discharged home, with two thirds asymptomatic and half having normal physical examinations. Only 11.8% had imaging of the head of any kind and in these children the incidence of TBI was high. Overall 4.3% had ciTBI which is 3 times higher than the percentage of ciTBI in the cohort of head injured children of all ages (1.4% ciTBI)²⁹. None of the neonates underwent neurosurgery or died. The risk of intracranial injury in young children is influenced by mechanism of injury (e.g. height of fall), size and location of scalp swelling and change in behaviour, among other factors⁶.

In line with previous research on head injuries in infants, most neonatal head injuries were caused by falls at home³² with the predominant mechanism being falls from caregiver's arms²⁰. Mulligan *et al.*³³ found that infants < 1 year dropped by caregivers were three times more likely to be admitted than infants from other fall types. A national Australian report³⁴ found that infants were most likely to be admitted when they were dropped by a caregiver and, consistent with our results, most head injuries in infants occurred at home. A smaller but significant proportion of head injuries were caused by

neonates being struck by object/person. Most neonates presented during the afternoon and early evening.

Of the four neonates with ciTBI, three had injuries from falls, including one who fell unrestrained from a pram. One neonate was accidentally struck by a person at home. All neonates with ciTBI had skull fractures, three of the four neonates had haematomas on head examination and two had intracranial haemorrhages. In contrast to our cohort of out of hospital head injuries Khan *et al.*²⁴ investigated in-hospital newborn head injuries, all due to falls. Seven of 24 received a skull radiograph followed by CT in six; two had a nondepressed linear parietal skull fracture and two had TBI with small subdural haemorrhages.

Three neonates had non-accidental trauma; 1 had no findings on neuroimaging, two of the three neonates had intracranial haemorrhages and skull fractures and of these two neonates, one had ciTBI. Non-accidental injuries must be considered in all neonates based on their anatomical vulnerability^{1,35,36} and susceptibility to serious head injuries caused by blunt impact and rotational movement of the brain within the cranial cavity^{3,37}. Non-accidental injuries or suspected non-accidental injuries are associated with a higher injury severity score and higher mortality compared to injuries sustained accidentally³⁸⁻⁴⁰. Additionally, TBI secondary to non-accidental trauma can require longer intensive care unit admission and higher mortality than TBI sustained by other mechanisms⁴⁰.

In our study, head injured neonates requiring neuroimaging most frequently underwent a CT or ultrasound. All neonates who had a palpable skull fracture underwent imaging of the head. The presence of vomiting, abnormal drowsiness and irritability did not appear to be correlated with imaging of the head. Two neonates who received neuroimaging were entirely asymptomatic and had no findings on examination. In this very young cohort, it is possible that clinician concern for ciTBI, variation in clinician practice and variation in the extent of paediatric emergency training may explain imaging of the head in the two asymptomatic neonates⁴¹ in the absence of supporting CDRs. At the same time, clinical assessment of neonates presenting with HI can be indeterminate and neuroimaging in these cases may be warranted¹³. Although the ciTBI rate in our cohort was high (4.3%), this was largely driven by the admission of neonates with TBI for two nights or more and may therefore represent increased concern by treating clinicians for potential sequelae in this young age group rather than more severe head injuries.

The gold standard imaging for paediatric TBI is CT but it carries risks of ionising radiation⁹⁻¹². Many clinical decision rules have been devised to help clinicians decide if a CT scan is required^{18,26-29, 42,43}. In addition to being able to identify skull fractures, in neonates the open fontanelle may allow the use of ultrasound to investigate if there is an intracranial injury. In our study four neonates underwent transfontanelle ultrasound with two showing TBI on ultrasound, which was confirmed by CT scan. As new ultrasound techniques emerge, it may increasingly complement CT scan as a safe (lack of radiation), cost-effective tool in the diagnosis and management of TBI⁴⁴ and skull fractures^{45,46}.

This study had some limitations. The coding of the injury mechanism depended on local research staff being able to extract relevant information from the medical records at the time of data collection. If not recorded these data could not be extracted. Our study was not designed to capture neonates who are admitted with medical issues unrelated to head injuries but are later found to have non-accidental head injuries. This may mean that we are underestimating the prevalence of non-accidental trauma. We did not collect information on why clinicians chose one imaging modality over another. Our data were focused on short term outcomes and we do not know the longer term outcome of neonatal head injuries.

Conclusions: Neonatal head injuries predominantly occurred after falls from caregiver's arms and have a mostly benign short term outcome that is appropriate for observation. However, non-accidental injuries remain an important cause of serious head injury in infants but were rare in our cohort

REFERENCES

1. Standring S. *Gray's Anatomy*. 40th ed. Churchill Livingstone, New York, 2008; 416-18
2. Pinto PS, Poretti A, Moeded A, Tekes A, Huisman TA. The unique features of traumatic brain injury in children. Review of the characteristics of the pediatric skull and brain, mechanisms of trauma, patterns of injury, complications and their imaging findings--part 1. *J. Neuroimaging*. 2012; **22**: e1-e17.
3. Case ME. Abusive head injuries in infants and young children. *Leg. Med. (Tokyo)*. 2007; **9**: 83-87.
4. Figaji AA. Anatomical and Physiological Differences between Children and Adults Relevant to Traumatic Brain Injury and the Implications for Clinical Assessment and Care. *Front. Neurol*. 2017; **8**: 685.
5. Greenes DS, Schultzman SA. Occult intracranial injury in infants. *Ann. Emerg. Med*. 1998; **32**: 680-6.
6. Schutzman SA, Barnes P, Duhaime AC et al. Evaluation and management of children younger than two years old with apparently minor head trauma: proposed guidelines. *Pediatrics*. 2001; **107**: 983-93.
7. Schnadower D, Vazquez H, Lee J, Dayan P, Roskind CG. Controversies in the evaluation and management of minor blunt head trauma in children. *Curr Opin Pediatr*. 2007; **19**: 258-64.
8. Willis AP, Latif SA, Chandratre S, Stanhope B, Johnson K. Not a NICE CT protocol for the acutely head injured child. *Clin Radiol*. 2008; **63**: 165-9.
9. Brenner DJ, Hall EJ. Computed tomography—an increasing source of radiation exposure. *N. Engl. J. Med*. 2007; **22**: 2277-84.
10. Pearce MS, Salotti JA, Little MP et al. Radiation exposure from CT scans in childhood and subsequent risk of leukaemia and brain tumours: a retrospective. *Lancet*. 2012; **380**: 499-505.
11. Miglioretti DL, Johnson E, Williams A et al. The use of computed tomography in pediatrics and the associated radiation exposure and estimated cancer risk. *JAMA Pediatr*. 2013; **167**: 700–07.
12. Mathews JD, Forsythe AV, Brady Z et al. Cancer risk in 680 000 people exposed to computed tomography scans in childhood or adolescence: data linkage study of 11 million Australians. *BMJ* 2013; **346**: 2360.
13. Ryan ME, Palasis S, Saigal G, et al. ACR Appropriateness Criteria head trauma--child. *J Am Coll Radiol*. 2014; **11**: 939-47.
14. van Wezel-Meijler G, Steggerda SJ, Leijser LM. Cranial Ultrasonography in Neonates: Roles and limitations. *Semin Perinatol*. 2010; **34**: 28-38.
15. Kemp AM, Jaspan T, Griffiths J, et al: Neuroimaging: What neuroradiological features distinguish abusive from non-abusive head trauma? A systematic review. *Arch Dis Child*. 2011; **96**: 1103-12.
16. Greenes DS, Schultzman SA. Clinical significance of scalp abnormalities in asymptomatic head-injured infants. *Pediatr. Emerg. Care*. 2001; **17**: 88-92.

17. Dayan PS, Holmes JF, Schutzman S et al. Traumatic Brain Injury Study Group of the Pediatric Emergency Care Applied Research Network (PECARN). Risk of traumatic brain injuries in children younger than 24 months with isolated scalp hematomas. *Ann. Emerg. Med.* 2014; **64**: 153-62
18. Kuppermann N, Holmes JF, Dayan PS et al, for the Pediatric Emergency Care Applied Research Network (PECARN). Identification of children at very low risk of clinically-important traumatic brain injuries after head trauma: a prospective cohort study. *Lancet.* 2009; **374**: 1160-70.
19. Crowe L, Babl F, Anderson V, Catroppa C. The epidemiology of paediatric head injuries: data from a referral centre in Victoria, Australia. *J. Paediatr. Child Health.* 2009; **45**: 346-50.
20. Agran PF, Anderson C, Winn D, Trent R, Walton-Haynes L, Thayer S. Rates of pediatric injuries by 3-month intervals for children 0 to 3 years of age. *Pediatrics.* 2003; **111**: e683-92.
21. Overpeck MD, Brenner RA, Trumble AC, Trifletti LB, Berendes HW. Risk factors for infant homicide in the United States. *N. Engl. J. Med.* 1998; **339**: 1211-1216.
22. Eisele JA, Kegler SR, Trent RB, Coronado VG. Nonfatal traumatic brain injury-related hospitalization in very young children-15 states, 1999. *J. Head Trauma Rehabil.* 2006; **21**: 537-43.
23. Graham CA, O'Toole SJ, Haddock G. Neonatal head injuries. *J. Accid. Emerg. Med.* 2000; **17**: 334-6
24. Khan DJ, Fisher PD, Hertzler DA. Variation in management of in-hospital newborn falls: a single-center experience. *J. Neurosurg. Pediatr.* 2017; **20**: 176-182.
25. Babl F, Borland M, Ngo P et al. Paediatric Research in Emergency Departments International Collaborative (PREDICT): first steps towards the development of an Australian and New Zealand research network. *Emerg. Med. Australas.* 2006; **18**: 143-47.
26. Osmond MH, Klassen TP, Wells GA et al. CATCH: a clinical decision rule for the use of computed tomography in children with minor head injury. *CMAJ.* 2010; **182**: 341-48.
27. Dunning J, Daly JP, Lomas JP et al. Derivation of the children's head injury algorithm for the prediction of important clinical events decision rule for head injury in children. *Arch. Dis. Child.* 2006; **91**: 885-91.
28. Babl FE, Lyttle MD, Bressan S et al. A prospective observational study to assess the diagnostic accuracy of clinical decision rules for children presenting to emergency departments after head injuries (protocol): the Australasian Paediatric Head Injury Rules Study (APHIRST). *BMC Pediatr.* 2014; **14**: 148.
29. Babl FE, Borland ML, Phillips N et al. Accuracy of PECARN, CATCH and CHALICE head injury decision rules in children: a prospective cohort study. *Lancet.* 2017; **389**: 2393-2402.
30. Reilly P, Simpson O, Sprod R, Thomas L. Assessing the conscious level in infants and young children: a pediatric version of Glasgow Coma Scale. *Childs Nervous System* 1998; **4**: 30-3.

31. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J. Biomed. Inform.* 2009; **42**: 377–81.
32. Warrington SA, Wright CM, ALSPAC Study Team. Accidents and resulting injuries in premobile infants: data from the ALSPAC study. *Arch. Dis. Child.* 2001; **85**: 104-7.
33. Mulligan CS, Adams S, Tzioumi D, Brown J. Injury from falls in infants under one year. *J. Paediatr. Child Health.* 2017; **53**: 754-760.
34. Pointer S. Hospitalised Injury in Children and Young People 2011–12. Canberra: Australian Institute of Health and Welfare, 2014. Injury Research and Statistics Series No.: 91. Cat. No.: INJCAT 167.
35. Greenley CS. Abusive head trauma: a review of the evidence base. *ALR. Am. J. Roentgenol.* 2015; **5**: 967-73.
36. Coats B, Margulies SS: Material properties of human infant skull and suture rates. *J. Neurotrauma.* 2006; **23**: 1222-1232.
37. Nadarasa J, Deck C, Meyer F, Willinger R, Raul JS. Update on injury mechanisms in abusive head trauma—shaken baby syndrome. *Pediatr. Radiol.* 2014; **44**: S565-S570.
38. Davies FC, Coats TJ, Fisher R, Lawrence T, Lecky FE. A profile of suspected child abuse as a subgroup of major trauma patients. *Emerg Med J.* 2015; **32**: 921-5.
39. Davies FC, Lecky FE, Fisher R, Frago-so-Iiguez M, Coats TJ. Major trauma from suspected child abuse: a profile of the patient pathway. *Emerg Med J.* 2017; **34**: 562-67.
40. Deans KJ, Minneci PC, Lowell W, Groner JL. Increased morbidity and mortality of TBI in victim of nonaccidental trauma. *J Trauma Acute Care Surg.* 2013; **75**: 157-60.
41. Stanley RM, Hoyle JD Jr, Dayan PS, et al. Emergency department practice variation in computed tomography use for children with minor blunt head trauma. *J Pediatr.* 2014; **165**: 1201-6.
42. Lyttle MD, Crowe L, Oakley E et al. Comparing CATCH, CHALICE and PECARN clinical decision rules for paediatric head injuries. *Emerg. Med. J.* 2012; **29**: 785–94.
43. Pickering A, Harnan S, Fitzgerald P, Pandor A, Goodacre S. Clinical decision rules for children with minor head injury: a systematic review. *Arch. Dis. Child.* 2011; **96**: 414–21.
44. Shin SS, Huisman TAGM, Hwang M. (2018) Ultrasound Imaging for Traumatic Brain Injury. *J. Ultrasound Med.* doi: 10.1002/jum.14547.
45. Choi JY, Lim YS, Jang JH, Park WB, Hyun SY, Cho JS. (2018) Accuracy of Bedside Ultrasound for the Diagnosis of Skull Fractures in Children Aged 0 to 4 Years. *Pediatr. Emerg. Care.* doi: 10.1097/PEC.0000000000001485.
46. Rabiner JE, Friedman LM, Khine H, Avner JR, Tsung JW. Accuracy of point-of-care ultrasound for diagnosis of skull fractures in children. *Pediatrics.* 2013; **131**: 1757-64.

Table 1: Demographics, signs and symptoms and outcomes of neonatal head injuries

	Total		
	n	%	(95% CI)
N	93		
Mean age [days], SD	16.3	8.1	(14.6-18.0)
Female	38	40.9	(30.7-51.5)
Signs:			
Vomiting	14	15.1	(8.5-24.0)
Abnormally drowsy	5	5.4	(1.8-12.1)
Irritable/agitated	3	3.2	(0.0-9.1)
Known of suspected LOC	0	0.0	
GCS:			
• 15	83	89.2	(81.1-94.7)
• 14	7	7.5	(3.1-14.9)
• 13	1	1.1	(0.0-5.8)
• 9-12	2	2.2	(0.0-7.5)
• 3-8	0	0.0	
Head Examination:			
Laceration	3	3.2	(0.7-9.1)
Haematoma	26	28.0	(19.1-38.2)
Possible palpable skull fracture	6	6.5	(2.4-13.5)
Tense fontanelle	2	2.2	(0.3-7.6)
Management and Outcome:			
Any neuroimaging	11	11.8	(6.1-20.2)
TBI on neuroimaging	5	5.4	(0.2-12.1)
CT scan	8	8.6	(3.8-16.2)
TBI on CT scan	5	5.4	(0.2-12.1)
Ultrasound	4	4.3	(1.2-10.6)
TBI on ultrasound	2	2.2	(0.3-7.6)
MRI	2	2.2	(0.3-7.6)
X-ray	1	1.1	(0.0-5.8)
ETT >1d	1	1.1	(0.0-5.8)
PICU admission	1	1.1	(0.0-5.8)
Admission	34	36.6	(26.8-47.2)
Median length of stay [days] (IQR)	1		(1.0-2.0)
Neurosurgery	0	0.0	
ciTBI	4	4.3	(1.2-10.6)
Mortality	0	0.0	(0.0-0.0)

LOC – loss of consciousness

GCS – Glasgow Coma Scale

CT – computed topography

TBI – traumatic brain injury

ETT – endotracheal tube

PICU –paediatric intensive care unit

IQR – interquartile range

ciTBI – clinically important traumatic brain injury

Table 2: Mechanism of neonatal head injuries


	n	%	%*
Total neonatal head injuries	93	100	
Activity Code:			
Fall related:	70	75.2	100.0
Fall from caregiver's arms	26	28.0	37.1
Fall from couch	9	9.7	12.9
Fall from pram	5	5.4	7.1
Car seat falling off table/stationary car	5	5.4	7.1
Fall from bed	4	4.3	5.7
Fall from bouncer	4	4.3	5.7
Fall from table	3	3.2	4.3
Fall from bassinet	3	3.2	4.3
Fall from chair	3	3.2	4.3
Fall from change table	2	2.2	2.9
Fall from parent's lap	2	2.2	2.9
Unknown fall related	4	4.3	5.7
Struck by:	19	20.4	100.0
Struck by inanimate stationary object	12	12.9	63.2
Struck by limb/body	4	4.3	21.1
Head strike with another child	3	3.2	15.8
Non accidental trauma (n=3)	3	3.2	100.0
Other: birth related trauma (n=1)	1	0.01	100.0
Location of injury:			
Home Area:	71	76.3	100.0
Living Area	31	33.3	43.7
Bedroom	18	19.4	25.4
Other Home Area	7	7.5	9.9
Stairs	4	4.3	5.6
Yard/Garden	4	4.3	5.6
Kitchen	3	3.2	4.2
Bathroom	2	2.2	2.8
Driveway	2	2.2	2.8
Other Areas:	22	23.7	100.0
Unknown	14.0	15.1	63.6
Public outdoor area	7.0	7.5	31.8
Restaurant/eating area	1.0	1.1	4.5

%* Proportion of specific head injury mechanism within each activity or location of injury code

Table 3: Traumatic brain injuries seen on neuroimaging

	Total
	n
Any traumatic brain injury	5
Intracranial haemorrhage or contusion	4
Any skull fracture	4
Depressed skull fracture	2
Midline shift of intracranial contents or signs of brain herniation	1
Diastasis of the skull	1
Sigmoid sinus thrombosis	1

Figure 1: Distribution of neonatal head injuries by time of injury



Neonatal Head Injuries - a prospective PREDICT cohort study
An Original Article

Nitaa Eapen, Meredith L. Borland, Natalie Phillips, Amit Kochar, Sarah Dalton, John A. Cheek, Yuri Gilhotra, Jocelyn Neutze, Mark D. Lyttle, Susan Donath, Louise Crowe, Stuart R Dalziel, Ed Oakley, Amanda Williams, Stephen Hearps, Silvia Bressan, Franz E Babl

Emergency Department, Royal Children's Hospital, 50 Flemington Rd, Parkville, VIC, 3052, Australia.

AUTHORS:

Nitaa Eapen BSc^{1,2}
nitaa.eapen@mcri.edu.au

Meredith L. Borland MBBS^{3,4}
meredith.borland@health.wa.gov.au

Natalie Phillips MBBS⁵⁻⁷
natalie.phillips@merton-oxford.com

Amit Kochar MD⁸
amit.kochar@health.sa.gov.au

Sarah Dalton BMed⁹
sarah.dalton@health.nsw.gov.au

John A. Cheek MBBS^{1,2,10}
john.cheek@rch.org.au

Yuri Gilhotra MBBS⁵
yurigilhotra@me.com

Jocelyn Neutze MBChB¹¹
jocelyn.neutze@middlemore.co.nz

Mark D. Lyttle MBChB^{2,12,13}
mdlyttle77@gmail.com

Susan Donath MA^{2,14}
susan.donath@mcri.edu.au

Louise Crowe PhD²
louise.crowe@mcri.edu.au

Stuart R Dalziel PhD^{15,16}
sdalziel@adhb.govt.nz

Ed Oakley MBBS^{1,2,14}
ed.oakley@rch.org.au

Amanda Williams²
Amanda.williams@mcri.edu.au

Stephen Hearps PGDipPsych²
stephen.hearps@mcri.edu.au

Silvia Bressan MD, PhD^{2,17}
silviabress@gmail.com

Franz E Babl MD^{1,2,14}
franz.babl@rch.org.au

¹ Royal Children's Hospital, Melbourne.
Address: Emergency Department, Royal Children's Hospital, 50 Flemington Rd, Parkville, VIC, 3052, Australia.

² Murdoch Childrens Research Institute, Melbourne.
Address: 50 Flemington Rd, Parkville, VIC, 3052, Australia.

³ Princess Margaret Hospital for Children, Perth.
Address: Emergency Department, Princess Margaret Hospital for Children, Roberts Rd, Subiaco, Perth, WA, 6008, Australia.

⁴ Schools of Paediatrics and Child Health and Primary, Aboriginal and Rural Healthcare, University of Western Australia.
Address: 35 Stirling Hwy, Crawley, WA, 6009, Australia.

⁵ Emergency department, Lady Cilento Children's Hospital, Brisbane.
Address: 501 Stanley St, South Brisbane, QLD, 4101, Australia.

⁶ Children's Health Research Centre, The University of Queensland, Medical Research Institute, Brisbane.
Address: 62 Graham St, South Brisbane, QLD, 4101, Australia

⁷ Faculty of Medicine, The University of Queensland, Brisbane.
Address: 20 Weightman St, Herston, QLD, 4006, Australia.

⁸ Women's & Children's Hospital, Adelaide.
Address: Emergency Department, Women's & Children's Hospital, 72 King William St, North Adelaide, SA, 5006, Australia.

⁹ The Children's Hospital at Westmead, Sydney.
Address: Emergency Department, The Children's Hospital at Westmead, 212 Hawkesbury Rd, Westmead, NSW, 2145, Australia.

¹⁰ Monash Medical Centre, Melbourne.

Address: Emergency Department, Monash Medical Centre, 246 Clayton Rd, Clayton, VIC, 3186, Australia.

¹¹ Kidzfirst Middlemore Hospital, Auckland.

Address: Emergency Department, Kidzfirst Middlemore Hospital, 100 Hospital Rd, Auckland, 2025, New Zealand.

¹² Bristol Royal Hospital for Children, Bristol.

Address: Emergency Department, Bristol Children's Hospital, Paul O'Gorman Building, Upper Maudlin St, Bristol, BS2 8BJ, United Kingdom.

¹³ Academic Department of Emergency Care, University of the West of England, Bristol.

Address: Blackberry Hill, Bristol, BS16 1XS, United Kingdom.

¹⁴ Department of Paediatrics, Faculty of Medicine, Dentistry and Health Sciences, University of Melbourne, Melbourne.

Address: Grattan St, Parkville, VIC, 3010, Australia.

¹⁵ Starship Children's Health, Auckland.

Address: Emergency Department, Starship Children's Health, 2 Park Rd, Grafton, Auckland, 1023, New Zealand.

¹⁶ Liggins Institute, University of Auckland, Auckland.

Address: 85 Park Ave, Grafton, Auckland, 1023, New Zealand.

¹⁷ Department of Women's and Child Health, University of Padova, Padova.

Address: Via Giustiniani3, 2 35128, Padova, PD, Italy.

CORRESPONDING AUTHOR:

Franz E Babl
Emergency Research
Murdoch Childrens Research Institute
Flemington Rd
Parkville
Victoria 3052
Australia
Phone 61 3 9345 6153
Fax 61 3 9345 600
Email franz.babl@rch.org.au