

DR. NIDHI MEDARA (Orcid ID : 0000-0002-5450-0647)

PROF. NEIL O'BRIEN-SIMPSON (Orcid ID : 0000-0001-8462-5603)

Article type : Original Article

## Peripheral neutrophil phenotypes during management of periodontitis

Nidhi Medara<sup>1</sup>, Jason C. Lenzo<sup>1,2</sup>, Katrina A. Walsh<sup>3</sup>, Eric C. Reynolds<sup>1,2\*</sup>, Neil M. O'Brien-Simpson<sup>1,2\*</sup> and Ivan B. Darby<sup>1\*</sup>

<sup>1</sup> Melbourne Dental School, The University of Melbourne, 720 Swanston Street, Carlton, VIC 3053, Australia

<sup>1,2</sup> Centre for Oral Health Research, The University of Melbourne, 720 Swanston Street, Carlton, VIC 3053, Australia

<sup>3</sup> Austin Hospital, 145 Studley Rd, Heidelberg, VIC 3084, Australia

\*Corresponding authors

Email:

N Medara – [nidhi.medara@unimelb.edu.au](mailto:nidhi.medara@unimelb.edu.au) (ORCID ID: 0000-0002-5450-0647)

JC Lenzo – [jlenzo@unimelb.edu.au](mailto:jlenzo@unimelb.edu.au)

KA Walsh – [kawalsh@unimelb.edu.au](mailto:kawalsh@unimelb.edu.au)

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/JRE.12793](https://doi.org/10.1111/JRE.12793)

This article is protected by copyright. All rights reserved

EC Reynolds – e.reynolds@unimelb.edu.au (ORCID ID: 0000-0002-6618-4856)

NM O'Brien-Simpson – neil.obs@unimelb.edu.au (ORCID ID: 0000-0001-8462-5603)

IB Darby – idarby@unimelb.edu.au (ORCID ID: 0000-0002-6457-5327)

## ABSTRACT

**Background and Objectives:** Neutrophils are emerging as a key player in periodontal pathogenesis. The surface expression of cellular markers enables functional phenotyping of neutrophils which have distinct roles in disease states. This study aimed to evaluate the effect of periodontal management on neutrophil phenotypes in peripheral blood in periodontitis patients over one year.

**Materials and methods:** Peripheral blood and the periodontal parameters, mean probing depth and percentage of sites with bleeding on probing (%BOP), were collected from 40 healthy controls and 54 periodontitis patients at baseline and 3-, 6- and 12-months post-treatment. Flow cytometry was used to identify CD11b<sup>+</sup>, CD16b<sup>+</sup>, CD62L<sup>-</sup> and CD66b<sup>+</sup> expression on neutrophils, neutrophil maturation stages as promyelocytes (CD11b<sup>-</sup>CD16b<sup>-</sup>), metamyelocytes (CD11b<sup>+</sup>CD16b<sup>-</sup>) and mature neutrophils (CD11b<sup>+</sup>CD16b<sup>+</sup>), and suppressive neutrophil phenotype as bands (CD16<sup>dim</sup>CD62L<sup>bright</sup>), normal neutrophils (CD16<sup>bright</sup>CD62L<sup>bright</sup>) and suppressive neutrophils (CD16<sup>bright</sup>CD62L<sup>dim</sup>).

**Results:** CD62L<sup>-</sup> expression decreased with treatment. No differences were observed in neutrophil maturation stages in health or disease upon treatment. Suppressive and normal neutrophils showed a reciprocal relationship, where suppressive neutrophils decreased with treatment and normal neutrophils increased with treatment. In addition, %BOP was associated with suppressive neutrophils.

**Conclusion:** This study demonstrates that management of periodontitis significantly modifies distinct neutrophil phenotypes in peripheral blood. Suppressive neutrophils may play a role in the pathogenesis of periodontitis. However, their exact role is unclear and requires further investigation.

## KEYWORDS

Periodontitis; Periodontal debridement; Granulocyte Precursor Cells; Suppressive Neutrophils

## INTRODUCTION

Neutrophils are a key player in the chronic inflammation seen in periodontitis. They are persistently recruited to the gingival crevice during inflammation, where they form a wall to block bacterial invasion into the underlying connective tissue<sup>1</sup>. Neutrophil extravasation through the post-capillary venules of the gingival plexus is tightly regulated through sequential expression of adhesion molecules. The initial stage of neutrophil adhesion from a freely circulating to a rolling state at reduced speeds is mediated by selectins, whereas integrins facilitate secondary 'firm' adhesion<sup>2</sup>. Once neutrophils reach the site of inflammation and are activated, they facilitate microbial clearance through a number of different processes including phagocytosis, generation of reactive oxygen species, release of granular contents, cytokine production and formation of neutrophil extracellular traps (NETs)<sup>3</sup>.

Previous studies have investigated CD11b, CD16b, CD62L and CD66b membrane markers in periodontitis to evaluate neutrophil adhesion and activation. CD11b non-covalently binds to CD18 to form the functional integrin heterodimer CD11b/CD18, and is involved in neutrophil adhesion and migration through the endothelium. Activation of neutrophils leads to a rapid increase in surface expression<sup>4,5</sup>, where CD11b/CD18 can also act as a complement receptor and transmembrane signalling adaptor<sup>6</sup>. CD62L (L-selectin) is part of the selectin family of cell adhesion molecules. It is expressed at high levels on resting circulating neutrophils<sup>7</sup>, but is shed rapidly during activation<sup>8</sup>. The same stimuli that cause up-regulation of CD11b expression can induce an equally rapid loss of CD62L during inflammation through proteolytic cleavage of the extracellular domain<sup>9</sup>. CD16b (FcγRIIIb) is the low affinity receptor for IgG1 and IgG3. CD16b plays a role in recognition and phagocytosis of IgG opsonised bacteria which triggers neutrophil activation, pro-inflammatory cascade and the production and release of anti-microbial compounds<sup>10</sup>. CD66b is a GPI-anchored glycoprotein of the carcinoembryonic antigen family. CD66b is involved in regulating CD11/CD18 function<sup>11-13</sup>, and CD66b cross-linking causes activation or priming in neutrophils and induces an increase in calcium and oxidative burst<sup>14</sup>, exocytosis of specific granules<sup>15</sup>, aggregate formation<sup>16</sup> and release of the pro-inflammatory cytokine IL-8<sup>17,18</sup>.

Neutrophils develop from progenitor cells in the bone marrow in several stages before exhibiting the characteristic segmented nuclear appearance seen in mature cells. Immature granulocytes are present rarely or in low numbers in health<sup>19</sup>, but their numbers increase in peripheral circulation during infection<sup>20,21</sup>, inflammation<sup>22,23</sup> or neoplastic processes<sup>24</sup>. Bacteria are the primary aetiological agents in the development of periodontitis and the differential expression of CD11b and CD16 may

be used to distinguish the various stages of neutrophil maturation in response to the microbial challenge in periodontitis<sup>25,26</sup>.

Neutrophils have previously been considered a homogenous population but mounting evidence suggests that distinct neutrophil subsets exist which have diverse roles in inflammation and cancer<sup>27</sup>. A subset of neutrophils with suppressive function can be identified based on the cell surface expression of CD16 and CD62L<sup>28</sup>. In addition to acute systemic inflammation, these CD16<sup>bright</sup>/CD62L<sup>dim</sup> suppressive neutrophils were also detected in patients with allergic rhinitis<sup>29</sup>, severe viral respiratory infections in infants<sup>30</sup>, head and neck squamous cell carcinoma<sup>31</sup> and other neoplasms<sup>32</sup>. Suppressing neutrophils display an activated phenotype with a hypersegmented nucleus, mediate T cell regulation in a CD11b manner<sup>28</sup>, form NETs, and migrate to tumour sites and perform anti-tumour functions such as inhibition of proliferation and induction of apoptosis in cancer cells<sup>31</sup>.

Previous studies that have evaluated CD11b, CD16b, CD62L and CD66b surface expression in peripheral neutrophils have only done so in cross-sectional studies and report conflicting results. Higher CD11b<sup>33</sup> expression and lower CD62L<sup>34,35</sup> and CD16<sup>36</sup> expression in periodontitis compared to health have been reported, whereas other studies reported no significant differences between health and disease<sup>37-41</sup>.

The aims of this study were three-fold. The first aim was to assess the longitudinal variation in the expression of the four adhesion and activation marker in periodontitis. Secondly, neutrophil maturation stage based on CD11b and CD16b expression was evaluated, where CD11b<sup>-</sup>CD16b<sup>-</sup> neutrophils were promyelocytes, CD11b<sup>+</sup>CD16b<sup>-</sup> neutrophils were metamyelocytes and CD11b<sup>+</sup>CD16b<sup>+</sup> neutrophils were mature neutrophils. Finally, the suppressive neutrophil phenotype was evaluated, where CD16<sup>bright</sup>CD62L<sup>bright</sup> neutrophils were phenotypically normal mature neutrophils, CD16<sup>dim</sup>CD62L<sup>bright</sup> neutrophils exhibited a banded nuclear morphology of cells recently released from bone marrow (bands) and CD16<sup>bright</sup>CD62L<sup>dim</sup> were suppressive neutrophils.

## MATERIALS AND METHODS

### Study population

The study was approved by the Human Ethics Sub-Committee at the University of Melbourne (1339812.3) and the Human Ethics Research Committee at Dental Health Services Victoria (279).

Sample size calculations were based on previously published work<sup>42</sup>. Informed written consent was obtained from each participant at the commencement of the study.

Fifty four subjects with periodontitis were recruited from the specialist periodontics clinic at The Royal Dental Hospital of Melbourne and Melbourne Dental Clinic, The University of Melbourne. The inclusion criteria for the periodontitis group were at least two non-adjacent sites per quadrant exhibiting probing depths (PD)  $\geq 5$ mm, excluding the third molars<sup>43</sup>. Subjects were  $>21$  years, had a minimum of 16 teeth (excluding third molars) and were systemically healthy with no periodontal treatment or antibiotic use within the preceding 6 months. Exclusion criteria included pregnancy or lactation and medical conditions affecting the progression of periodontitis (e.g. diabetes) or requiring pre-medication prior to treatment.

For the control group ( $n=40$ ), the inclusion criteria were a gender and age ( $\pm 5$  years) match to the periodontitis subject with no PD  $>4$ mm or percentage of sites with bleeding on probing (%BOP)  $>30\%$ . The remaining inclusion and exclusion criteria were the same as for the periodontitis subject.

The periodontal parameters, mean PD and %BOP, and peripheral blood were collected once from the healthy controls, and at baseline before treatment and 3-, 6- and 12-months during supportive periodontal therapy from the periodontitis group. Supportive periodontal therapy involved supra- and sub-gingival debridement every three months. PD and BOP were recorded at six sites around each tooth using a William's probe (Hu-Friedy Mfg. Co., Chicago, USA). PD was measured to the nearest millimeter from the base of the gingival sulcus to the free gingival margin. BOP was assessed visually following probing to the base of the sulcus and recorded as present or absent up to 30 seconds after probing. Smoking status was defined as current smoker or non-smoker at baseline.

### Neutrophil phenotyping

Peripheral blood was collected in potassium-EDTA tubes (BD Biosciences, NSW, Australia) and whole blood (20 $\mu$ L) was incubated for 30 minutes at 4 $^{\circ}$ C protected from light with pre-diluted fluorochrome-conjugated monoclonal antibodies against CD11b-PECy7 (3 $\mu$ L, clone-ICRF44), CD16b-PE (7 $\mu$ L, clone-CLBgran11.5), CD62L-APC (10 $\mu$ L, clone-DREG-56) and CD66b-FITC (15 $\mu$ L, clone-G10F5). All antibodies were purchased from BD Bioscience and the volumes used were optimised based on manufacturer's recommendation. After incubation, red blood cell lysis was performed by adding 1mL of cold MilliQ water (MilliporeSigma, MA, USA) and vigorously pipetting up and down for 30 seconds. After 30 seconds, 1mL of 2x phosphate buffered saline (PBS) was added to stop lysis. The cells were centrifuged at 800RCF for 5 minutes at 2 $^{\circ}$ C and the supernatant aspirated. The cells

were resuspended in 200µL of PBS and acquired by flow cytometry on BD LSRFortessa™ X-20 (BD Bioscience, CA, USA). Data analysis was performed on FlowJo software (v10, Tree Star Inc., OR, USA). The gating strategy to phenotype neutrophils is displayed in figure 1. CD62L is rapidly lost on activation, thus, neutrophils without CD62L represent an activated phenotype.

### Statistical analysis

Independent samples *t*-tests with unequal variances were used for comparison between health and periodontitis at each timepoint. Linear mixed models (LMMs) were used to determine changes in the mean of the outcome variable over timepoint. The base model for variation over timepoints included timepoint, age and sex as fixed effects and subject as random effect. Pairwise comparisons of estimated marginal means were used to assess significant differences. To assess the effect of periodontal management, mean PD and %BOP were added as fixed effects to the base model. Visual inspection of residual plots did not show any obvious deviations from normality. To assess difference in smoking status, independent samples *t*-tests with unequal variances were used for comparison between current smokers and non-smokers at each timepoint in health and periodontitis and smoking status was added as a fixed effect to the base LMMs. Spearman's correlations were used for associations with the periodontal parameters at each timepoint.

The level of significance was set at  $p \leq 0.05$  for all variables. Statistical analyses were performed in SPSS (v23, IBM SPSS Statistics for Windows, IBM Corp., NY, USA) and graphs were prepared using GraphPad Prism (v8, GraphPad Software, CA, USA).

## RESULTS

### Cohort demographics

The cohort demographics for healthy controls and periodontitis subjects are displayed in table 1. No significant differences were observed between healthy and periodontitis subjects at baseline with regards to age, sex or smoking status. The periodontal parameters, mean PD and %BOP, were significantly higher at all timepoints in periodontitis compared to health. With treatment, both parameters significantly decreased at all subsequent timepoints compared to baseline.

### CD11b, CD16b, CD62L and CD66b expression on neutrophils

Neutrophils and their expression of CD11b<sup>+</sup>, CD16b<sup>+</sup>, CD62L<sup>-</sup> and CD66b<sup>+</sup> are displayed in figure 2. Neutrophils were significantly higher at 3- and 6-months in periodontitis compared to health. CD62L<sup>-</sup> expression was significantly higher at 12-months compared to health, and with treatment, CD62L<sup>-</sup> expression significantly decreased at all subsequent timepoints compared to baseline. CD11b, CD16b and CD66b were all highly and constitutively expressed on neutrophils. There was a statistically significant increase in CD66b<sup>+</sup> expression at 6-months compared to baseline, but no differences were observed in CD11b<sup>+</sup> or CD16b<sup>+</sup> expression.

The CD16b antibody clone used in this study only bound to neutrophils expressing one of the two neutrophil antigen (NA) molecules. No statistical differences were observed between groups at all timepoints in the frequency of CD16b<sup>+</sup> expression (table 1) with regards to age, gender, mean PD or %BOP.

When using a combination of markers (figure 3), CD16b<sup>+</sup>CD66b<sup>+</sup>CD11b<sup>+</sup> identified 95% of neutrophils (mean±SD in health = 95.47±2.51 and baseline = 95.71±3.51). CD16b<sup>+</sup>CD66b<sup>+</sup>CD11b<sup>+</sup>CD62L<sup>-</sup> cells, which represent an activated neutrophil phenotype, were significantly decreased at 3- and 6-months in comparison to health and baseline.

Smokers had significantly higher neutrophil proportions at baseline (mean difference (*d*)=9.56%) and 6-months (*d*=6.69%) in comparison to non-smokers in periodontitis (supplementary table 1).

### Neutrophil maturation

Mature neutrophils were the most abundant cell type in peripheral blood, followed by metamyelocytes and promyelocytes (figure 4). These proportions of immature granulocytes and mature neutrophils remained constant in health and disease upon treatment.

### Suppressive neutrophils

When phenotyping neutrophils based on CD16b and CD62L expression, normal neutrophils were the largest subset followed by suppressive neutrophils then bands (figure 5A). The small proportion of bands remained relatively constant in health and disease (figure 5B). Age was a significant predictor for bands in the base model for timepoint (estimate of fixed effect (*b*) = 0.03, standard error (SE) = 0.01), and a 1 year increase in age was associated with 0.03% increase in bands. Smoking status also

had an effect on bands where smokers had significantly lower proportions compared to non-smokers at 6-months ( $d=0.31\%$ ).

Normal and suppressive neutrophils behaved in reciprocal ways (figure 5C, D), where normal neutrophils were significantly increased and suppressive neutrophils were significantly decreased at 3- and 6-months post-treatment compared to health and baseline.

#### Association with periodontal parameters

Mean PD and %BOP were correlated to various neutrophil subsets (table 2). In health, %BOP was significantly positively correlated with CD16b<sup>+</sup>CD66b<sup>+</sup>CD11b<sup>+</sup>CD62L<sup>-</sup> neutrophils ( $\rho=0.401$ ). At 3-months, %BOP was significantly positively correlated with CD16b<sup>+</sup>CD66b<sup>+</sup>CD11b<sup>+</sup> neutrophils ( $\rho=0.505$ ) and CD11b<sup>+</sup>CD16b<sup>+</sup> mature neutrophils ( $\rho=0.468$ ), and negatively correlated with promyelocytes ( $\rho=-0.547$ ). At 6-months, mean PD was significantly negatively correlated with total neutrophils ( $\rho=-0.379$ ), and at 12-months post-treatment, mean PD was significantly positively correlated with CD66b<sup>+</sup> expressing neutrophils ( $\rho=0.378$ ) and CD16<sup>bright</sup>CD62L<sup>bright</sup> normal neutrophils ( $\rho=0.514$ ).

When mean PD and %BOP were added as fixed effects to the base statistical model, %BOP ( $b=0.22$ ,  $SE=0.10$ ) was a significant predictor for suppressive neutrophils and a 1% increase in %BOP was associated with a 0.22% increase in suppressive neutrophils.

## DISCUSSION

Neutrophils are actively recruited to the gingival sulcus during periodontal inflammation. In normal conditions, 50% of peripheral neutrophils are marginating through post-capillary venules throughout the body<sup>44</sup>. Furthermore, neutrophil release from the bone marrow in response to proinflammatory stimuli is a rate-limiting step in the host response to pathogens<sup>45</sup>. Thus, a study of markers that facilitate neutrophil adhesion and activation will improve our understanding of the role of neutrophils in periodontitis.

CD62L or L-selectin is highly expressed on resting circulating neutrophils<sup>7</sup>, but is shed during activation<sup>8,9,46</sup>. CD62L<sup>-</sup> expression was not significantly different between health and baseline in this study, however, there was a significant decrease post-treatment at all timepoints compared to

baseline. The greatest decrease in CD62L<sup>-</sup> expressing cells was between baseline and 3-months. Likewise, mature activated CD16b<sup>+</sup>CD66b<sup>+</sup>CD11b<sup>+</sup>CD62L<sup>-</sup> cells decreased post-treatment compared to baseline. Previous studies that evaluated CD62L expression were cross-sectional and reported significantly lower expression in rapidly progressive periodontitis<sup>34</sup> and adult periodontitis<sup>35</sup> compared to healthy controls, while other studies observed no significant differences<sup>37,40,47</sup>. It is unclear at this stage as to what stimulus leads to the shedding of CD62L in peripheral blood. These neutrophils may have been activated due to systemic bacterial dissemination seen in periodontitis<sup>48,49,50</sup> or may represent a subset that is being actively recruited to periodontal sites. The latter scenario may be more likely as kinetic studies report that CD62L shedding is rapid and only 10-15% surface expression was found after 30 minutes<sup>51,52</sup>. It has been proposed that this shedding of CD62L may provide a rapid means for neutrophil de-adhesion necessary for detachment from endothelial cells before transigrations into sites of inflammation<sup>9</sup>.

Several CD16b clones are available for use in flow cytometry including ID3, REA589, 245514 and MM0272-5L11. The clone used in this study was CLBgran11.5 which binds with neutrophils expressing the NA1 molecule. CD16b monoclonal antibodies recognise the biallelic NA1/NA2 antigens, which differ in five nucleotides (nucleotide 141, 147, 227, 277 and 349) which leads to changes in four amino acids at positions 36, 65, 82 and 106, one of which is a silent mutation<sup>53,54</sup>. In this study, 64% of subjects in health and 58% in periodontitis at baseline were detected by the CD16b antibody representing the NA1NA1 homozygous forms. Although there were no differences in NA1NA1 expression in health and periodontitis groups, this effectively reduced the sample size for the study when CD16b was used in the analysis. CD16b is the low-affinity Fc receptor for immunoglobulin isotypes and is involved in adhesion and phagocytosis of opsonised bacteria. *In vitro* studies have reported that neutrophils homozygous for NA2 bind IgG3 less efficiently than NA1 homozygous neutrophils, and IgG1 or IgG3 opsonised particles are more efficiently ingested upon interaction with the NA1 allotype<sup>55,56</sup>. In periodontitis, a meta-analysis of fourteen studies reported that the NA2 phenotype was associated with an odd ratio of 2.01 for aggressive periodontitis compared to NA1, and the homozygous NA2NA2 phenotype was associated with an odd ratio of 1.40 for chronic periodontitis compared to the NA1NA1 and NA1NA2 phenotype<sup>57</sup>. Copy number variations of *FCGR3B* gene have also been reported to be associated with several other chronic inflammatory diseases, such as systemic lupus erythematosus<sup>58</sup>, rheumatoid arthritis<sup>59</sup> and immune-mediated glomerulonephritis<sup>60</sup>.

Neutrophils have traditionally been considered a homogenous population because of their short lifespan, restricted transcriptional activity and inability to reverse transigrate into peripheral blood.

However, recent studies demonstrate that they exhibit plasticity and heterogeneity that enables them to respond in a flexible manner in different disease states<sup>61-64</sup>. One such subset is the CD16<sup>bright</sup>CD62L<sup>dim</sup> suppressive neutrophil which displays a hypersegmented nuclear morphology and is involved in suppressing T cell proliferation via hydrogen peroxide release into the immunological synapse in a CD11b-dependent manner<sup>28</sup> and cell-cell contact through programmed death-ligand 1<sup>65</sup>. A recent study that used functional enrichment analysis showed that suppressive neutrophils were enriched in proteins involved in immune regulation<sup>66</sup> and interferon signalling<sup>65</sup>. In the present study, while there were no differences between health and baseline in periodontitis in both normal and suppressive neutrophils, these two subsets showed a reciprocal relationship upon treatment. Suppressive neutrophils significantly decreased with treatment at 3- and 6-months and were significantly associated with %BOP. This decrease with treatment may be associated with the decrease in inflammation in response to treatment. There is increased prevalence of systemic bacterial dissemination in periodontitis<sup>50</sup>, and the magnitude of bacteraemia is associated with the number of sites with BOP<sup>67</sup>. The regulatory role of suppressive neutrophils may be required to restrain excessive and potentially harmful T cell activation in peripheral blood as a result of this bacteraemia and to maintain compartmentalisation between peripheral circulation and gingival tissues. At the resolution of inflammation at 12-months post-treatment, CD16<sup>bright</sup>CD62L<sup>bright</sup> normal neutrophils were positively correlated with mean PD. Indeed, suppressive neutrophils showed lower bacterial containment<sup>68</sup> and may have immunoregulatory<sup>61</sup> rather than antibacterial functions. Further studies are needed to assess the functional role of suppressive neutrophils as well as their usefulness as a diagnostic or therapeutic marker in periodontitis.

Immature granulocytes were present at constant but low levels in health and during treatment in periodontitis. Promyelocytes and metamyelocytes have reduced effector functions compared to mature neutrophils<sup>19</sup> and their role in periodontitis is unclear. The chronic nature of periodontal disease and sampling well after establishment of plaque biofilm deposits may preclude detection of immature granulocyte release from bone marrow stores in this study. A previous study reported significantly more promyelocytes in early-onset periodontitis compared to health and suggested that there may be an abnormal haematopoietic mechanism involved in early-onset periodontitis pathogenesis<sup>36</sup>.

Smoking had an effect on neutrophil proportions where smokers had higher total neutrophil at baseline and 6-months compared to non-smokers. Higher neutrophil proportions in peripheral blood of smokers were also observed in other studies conducted in healthy subjects<sup>69-71</sup>. Smoking has several effects on neutrophil functions including impairment of migration and chemotaxis<sup>72</sup>,

phagocytosis<sup>73</sup> and respiratory burst<sup>74,75</sup>, and these effects maybe more evident in diseased gingival tissues. This study found no differences between smokers and non-smokers in CD62L<sup>-</sup> or CD11b expression which facilitates neutrophil entry into gingival tissues. However, this study only characterised current smokers and non-smokers, and previous history of smoking or the duration required to return to baseline after smoking cessation, and smoking quantity and history in current smokers may also affect expression of cell surface markers. Another limitation of this study is the small number of smokers in the cohort which did not allow for more robust statistical comparison to be made. Our results show that a larger prospective study with well-defined smoking categories are needed to expand and corroborate our results.

In this study as well as others<sup>37,39,76</sup>, no significant differences were observed in CD11b, CD16b and CD66b expressing neutrophils in health and disease in peripheral blood. However, others have reported significant differences in the expression of these markers in oral neutrophils in comparison with peripheral neutrophils<sup>77-80</sup>. Oral neutrophils may provide a more dynamic picture of cell subsets involved in periodontitis as these neutrophils are at the interface between periodontal tissues and the oral environment. Indeed, a previous study showed that neutrophils from sites with periodontal disease display increased number of gene changes compared to oral neutrophils in healthy periodontium<sup>81</sup>. Further studies that track the migration of activated and suppressive neutrophils from peripheral blood to the gingival tissues may provide a greater understanding of the role of these cells in periodontitis.

Several factors may affect extracellular surface marker expression on neutrophils. For example, CD62L expression was shown to be increased during active bone marrow release<sup>82</sup>, but neutrophil ageing<sup>83</sup> and several pharmaceuticals like non-steroidal anti-inflammatory drugs<sup>84</sup> and steroids<sup>85</sup> cause a loss of surface CD62L. Antigens such as CD10, CD11b, CD13, CD16 and CD66b have large intracellular storage pools and experimental manipulations can mobilise these from their intracellular location and increase their surface expression<sup>86,87</sup>. Furthermore, neutrophil apoptosis could increase surface expression of some antigens, such as CD13, and decrease others such as CD15 and CD16<sup>88</sup>.

In conclusion, this study found that CD11b, CD16b and CD66b were all highly expressed on neutrophils, whereas CD62L<sup>-</sup> was significantly decreased upon treatment. There was no variation in immature granulocyte proportions with treatment. Suppressive and normal neutrophils showed a reciprocal relationship upon treatment where suppressive neutrophils were decreased and normal neutrophils were increased at 3- and 6-months compared to baseline. To the best of our knowledge,

this is the first study that evaluated the longitudinal variation of these neutrophil phenotypes in peripheral blood in periodontitis.

## ACKNOWLEDGMENTS

The authors wish to acknowledge the Melbourne Cytometry Platform for equipment use and expertise. This study was supported by grants from the Australian National Health and Medical Research Council (APP1029878) and the Melbourne Dental School Higher Research Funds (2014). NM was supported by the Oral Health Cooperative Research Centre Top-up Scholarship.

## FIGURE & TABLE LEGENDS

### Table 1. Cohort demographics and periodontal parameters.

+ significantly different to health

‡ significantly different to baseline

§ significantly different to 3-months post-treatment

%BOP – percentage of sites exhibiting bleeding on probing, mm – millimetres, PD – probing depth,

SD – standard deviation

### Table 2. Correlations between periodontal parameters and neutrophil subsets.


Spearman's correlation coefficients ( $\rho$ ) between the periodontal parameters, mean probing depth (PD) and percentage of sites exhibiting bleeding on probing (%BOP), and various neutrophil subsets in health and at baseline, 3-, 6- and 12- months in periodontitis. Significant correlations ( $\rho \leq 0.05$ ) are represented in bold.

### Fig. 1. Gating strategy for phenotyping neutrophils.


Neutrophils **(A)** were identified based on forward scatter area (FSC-A) and side scatter area (SSC-A). Doublets were excluded using FSC-A and forward scatter height (FSC-H) followed by SSC-H and SSC-A. Single expression **(B)** of CD11b<sup>+</sup>, CD16b<sup>+</sup>, CD62L<sup>-</sup> and CD66b<sup>+</sup>, as well as a combination of markers **(C)** as CD16b<sup>+</sup>CD66b<sup>+</sup>CD11b<sup>+</sup> and CD16b<sup>+</sup>CD66b<sup>+</sup>CD11b<sup>+</sup>CD62L<sup>-</sup> were identified on neutrophils. Neutrophil maturation stages **(D)** were phenotyped as promyelocytes (CD11b<sup>-</sup>CD16b<sup>-</sup>), metamyelocytes (CD11b<sup>+</sup>CD16b<sup>-</sup>) and mature neutrophils (CD11b<sup>+</sup>CD16b<sup>+</sup>). Suppressive neutrophil

subsets (E) were phenotyped as bands (CD16<sup>dim</sup>CD62L<sup>bright</sup>), normal (CD16<sup>bright</sup>CD62L<sup>bright</sup>) or suppressive (CD16<sup>bright</sup>CD62L<sup>dim</sup>).


**Fig. 2. Surface expression of neutrophil markers.**

Neutrophils (A) (as a percentage of total cells) and CD11b<sup>+</sup> (B), CD16b<sup>+</sup> (C), CD62L<sup>-</sup> (D) and CD66b<sup>+</sup> (E) expressing cells (as a percentage of neutrophils) in health and periodontitis at baseline, 3-, 6-, and 12-months. Box and whisker plots represent the median, interquartile range and 10<sup>th</sup> – 90<sup>th</sup> percentile;  above the plots represent  $p \leq 0.05$ .


**Fig. 3. Combinations of neutrophil surface markers.**

CD16b<sup>+</sup>CD66b<sup>+</sup>CD11b<sup>+</sup> (A) and CD16b<sup>+</sup>CD66b<sup>+</sup>CD11b<sup>+</sup>CD62L<sup>-</sup> (B) expressing neutrophil subsets (as a percentage of neutrophils) in health and periodontitis at baseline, 3-, 6-, and 12-months. Box and whisker plots represent the median, interquartile range and 10<sup>th</sup> – 90<sup>th</sup> percentile;  above the plots represent  $p \leq 0.05$ .

**Fig. 4. Neutrophil maturation stages.**

Bar graph showing neutrophil maturation stages (A) and box and whisker plots showing promyelocytes (B), metamyelocytes (C) and mature neutrophils (D) (as a percentage of neutrophils) in health and periodontitis at baseline, 3-, 6-, and 12-months. Bars with error bars represent mean and standard deviation (SD); † all subsets are significantly different ( $p \leq 0.05$ ) to each other within the same timepoint. Box and whisker plots represent the median, interquartile range and 10<sup>th</sup> – 90<sup>th</sup> percentile;  above the plots represent  $p \leq 0.05$ .

**Fig. 5. Suppressive neutrophil subsets.**

Bar graph showing suppressive neutrophil subsets (A) and box and whisker plots showing bands (B), normal (C) and suppressive neutrophils (D) (as a percentage of neutrophils) in health and periodontitis at baseline, 3-, 6-, and 12-months. Bars with error bars represent mean and SD; † all subsets are significantly different ( $p \leq 0.05$ ) to each other within the same timepoint. Box and whisker plots represent the median, interquartile range and 10<sup>th</sup> – 90<sup>th</sup> percentile;  above the plots represent  $p \leq 0.05$ .

## REFERENCES

1. Ryder MI. Comparison of neutrophil functions in aggressive and chronic periodontitis. *Periodontol 2000*. 2010;53:124-137.
2. Kolaczowska E, Kubes P. Neutrophil recruitment and function in health and inflammation. *Nat Rev Immunol*. 2013;13(3):159-175.
3. Hirschfeld J. Neutrophil Subsets in Periodontal Health and Disease: A Mini Review. *Front Immunol*. 2020;10(3001).
4. Berger M, O'Shea J, Cross AS, et al. Human neutrophils increase expression of C3bi as well as C3b receptors upon activation. *J Clin Invest*. 1984;74(5):1566-1571.
5. Ginsberg MH, Du X, Plow EF. Inside-out integrin signalling. *Curr Opin Cell Biol*. 1992;4(5):766-771.
6. Anderson DC, Miller LJ, Schmalstieg FC, Rothlein R, Springer TA. Contributions of the Mac-1 glycoprotein family to adherence-dependent granulocyte functions: structure-function assessments employing subunit-specific monoclonal antibodies. *J Immunol*. 1986;137(1):15-27.
7. Butcher EC. Leukocyte-endothelial cell recognition: three (or more) steps to specificity and diversity. *Cell*. 1991;67(6):1033-1036.
8. Spertini O, Schleiffenbaum B, White-Owen C, Ruiz P, Jr., Tedder TF. ELISA for quantitation of L-selectin shed from leukocytes in vivo. *J Immunol Methods*. 1992;156(1):115-123.
9. Kishimoto TK, Jutila MA, Berg EL, Butcher EC. Neutrophil Mac-1 and MEL-14 adhesion proteins inversely regulated by chemotactic factors. *Science*. 1989;245(4923):1238-1241.
10. Gresham HD, Clement LT, Lehmeyer JE, Griffin FM, Jr., Volanakis JE. Stimulation of human neutrophil Fc receptor-mediated phagocytosis by a low molecular weight cytokine. *J Immunol*. 1986;137(3):868-875.
11. Kuijpers TW, van der Schoot CE, Hoogerwerf M, Roos D. Cross-linking of the carcinoembryonic antigen-like glycoproteins CD66 and CD67 induces neutrophil aggregation. *J Immunol*. 1993;151(9):4934-4940.
12. Stocks SC, Kerr MA, Haslett C, Dransfield I. CD66-dependent neutrophil activation: a possible mechanism for vascular selectin-mediated regulation of neutrophil adhesion. *J Leukoc Biol*. 1995;58(1):40-48.

13. Stocks S, Kerr MA. Stimulation of neutrophil adhesion by antibodies recognizing CD15 (Lex) and CD15-expressing carcinoembryonic antigen-related glycoprotein NCA-160. *Biochem J.* 1992;288(1):23-27.
14. Lund-Johansen F, Olweus J, Symington FW, et al. Activation of human monocytes and granulocytes by monoclonal antibodies to glycosylphosphatidylinositol-anchored antigens. *Eur J Immunol.* 1993;23(11):2782-2791.
15. Ducker TP, Skubitz KM. Subcellular localization of CD66, CD67, and NCA in human neutrophils. *J Leukoc Biol.* 1992;52(1):11-16.
16. Schmidt T, Zundorf J, Gruger T, et al. CD66b overexpression and homotypic aggregation of human peripheral blood neutrophils after activation by a gram-positive stimulus. *J Leukoc Biol.* 2012;91(5):791-802.
17. Schröder AK, Uciechowski P, Fleischer D, Rink L. Crosslinking of CD66b on Peripheral Blood Neutrophils Mediates the Release of Interleukin-8 from Intracellular Storage. *Hum Immunol.* 2006;67(9):676-682.
18. Beauchemin N, Draber P, Dveksler G, et al. Redefined nomenclature for members of the carcinoembryonic antigen family. *Exp Cell Res.* 1999;252(2):243-249.
19. Glasser L, Fiederlein RL. Functional differentiation of normal human neutrophils. *Blood.* 1987;69(3):937-944.
20. Nierhaus A, Klätte S, Linssen J, et al. Revisiting the white blood cell count: immature granulocytes count as a diagnostic marker to discriminate between SIRS and sepsis - a prospective, observational study. *BMC Immunol.* 2013;14(1):8.
21. Boettcher S, Gerosa RC, Radpour R, et al. Endothelial cells translate pathogen signals into G-CSF-driven emergency granulopoiesis. *Blood.* 2014;124(9):1393-1403.
22. Nagareddy PR, Murphy AJ, Stirzaker RA, et al. Hyperglycemia promotes myelopoiesis and impairs the resolution of atherosclerosis. *Cell Metab.* 2013;17(5):695-708.
23. Soehnlein O, Steffens S, Hidalgo A, Weber C. Neutrophils as protagonists and targets in chronic inflammation. *Nature Reviews Immunology.* 2017;17:248.
24. Trellakis S, Farjah H, Bruderek K, et al. Peripheral Blood Neutrophil Granulocytes from Patients with Head and Neck Squamous Cell Carcinoma Functionally Differ from Their Counterparts in Healthy Donors. *Int J Immunopathol Pharmacol.* 2011;24(3):683-693.
25. Fujimoto H, Sakata T, Hamaguchi Y, et al. Flow cytometric method for enumeration and classification of reactive immature granulocyte populations. *Cytometry.* 2000;42(6):371-378.
26. Terstappen L, Safford M, Loken MR. Flow cytometric analysis of human bone marrow. III. Neutrophil maturation. *Leukemia.* 1990;4(9):657-663.

27. Rodriguez FM, Novak ITC. What about the neutrophils phenotypes? *Hematol Med Oncol*. 2017;2(3):1-6.
28. Pillay J, Kamp VM, Van Hoffen E, et al. A subset of neutrophils in human systemic inflammation inhibits T cell responses through Mac-1. *J Clin Invest*. 2012;122(1):327-336.
29. Arebro J, Ekstedt S, Hjalmarsson E, Winqvist O, Kumlien Georén S, Cardell L-O. A possible role for neutrophils in allergic rhinitis revealed after cellular subclassification. *Sci Rep*. 2017;7:43568.
30. Cortjens B, Ingelse SA, Calis JC, et al. Neutrophil subset responses in infants with severe viral respiratory infection. *Clin Immunol*. 2017;176:100-106.
31. Millrud CR, Kågedal Å, Kumlien Georén S, et al. NET-producing CD16high CD62Ldim neutrophils migrate to tumor sites and predict improved survival in patients with HNSCC. *Int J Cancer*. 2017;140(11):2557-2567.
32. Hao S, Andersen M, Yu H. Detection of immune suppressive neutrophils in peripheral blood samples of cancer patients. *Am J Blood Res*. 2013;3(3):239-245.
33. Agarwal S, Suzuki JB, Piesco NP, Aichelmann-Reidy MB. Neutrophil function in juvenile periodontitis: induction of adherence. *Oral Microbiol Immunol*. 1994;9(5):262-271.
34. Macey MG, McCarthy DA, Howells GL, Curtis MA, King G, Newland AC. Multiparameter flow cytometric analysis of polymorphonuclear leucocytes in whole blood from patients with adult rapidly progressive periodontitis reveals low expression of the adhesion molecule L-selectin (Cd62L). *Cytometry*. 1998;34(3):152-158.
35. Gainet J, Dang PM, Chollet-Martin S, et al. Neutrophil dysfunctions, IL-8, and soluble L-selectin plasma levels in rapidly progressive versus adult and localized juvenile periodontitis: variations according to disease severity and microbial flora. *J Immunol*. 1999;163(9):5013-5019.
36. Nemoto E, Nakamura M, Shoji S, Horiuchi H. Circulating promyelocytes and low levels of CD16 expression on polymorphonuclear leukocytes accompany early-onset periodontitis. *Infect Immun*. 1997;65(9):3906-3912.
37. Asman B, Gustafsson A, Bergstrom K. Gingival crevicular neutrophils: membrane molecules do not distinguish between periodontitis and gingivitis. *J Clin Periodontol*. 1997;24(12):927-931.
38. Watanabe K, Hagen KL, Ramakrishnan V, Andersen BR. Kinetics of CD11b expression on neutrophils isolated from subjects with healthy gingivae and patients with advanced periodontitis. *J Periodontal Res*. 1993;28(2):137-144.

39. Watanabe K, Blew B, Scherer M, et al. CD11b mRNA expression in neutrophils isolated from peripheral blood and gingival crevicular fluid. *J Clin Periodontol.* 1997;24(11):814-822.
40. Pietruska M, Zak J, Pietruski J, Wysocka J. Expressions of selected adhesion molecules on peripheral blood leukocytes in patients with aggressive periodontitis. *Arch Immunol Ther Exp (Warsz).* 2005;53(3):266-271.
41. Leino L, Hurttia H, Sorvajärvi K, Sewon L. Increased respiratory burst activity is associated with normal expression of IgG-Fc-receptors and complement receptors in peripheral neutrophils from patients with juvenile periodontitis. *J Periodontol Res.* 1994;29(3):179-184.
42. Byrne S, Dashper S, Darby I, Adams G, Hoffmann B, Reynolds E. Progression of chronic periodontitis can be predicted by the levels of Porphyromonas gingivalis and Treponema denticola in subgingival plaque. *Oral Microbiol Immunol.* 2009;24(6):469-477.
43. Darby IB, Mooney J, Kinane DF. Changes in subgingival microflora and humoral immune response following periodontal therapy. *J Clin Periodontol.* 2001;28(8):796-805.
44. Summers C, Rankin SM, Condliffe AM, Singh N, Peters AM, Chilvers ER. Neutrophil kinetics in health and disease. *Trends Immunol.* 2010;31(8):318-324.
45. Sima C, Glogauer M. Neutrophil Dysfunction and Host Susceptibility to Periodontal Inflammation: Current State of Knowledge. *Current Oral Health Reports.* 2014;1(2):95-103.
46. Griffin JD, Spertini O, Ernst TJ, et al. Granulocyte-macrophage colony-stimulating factor and other cytokines regulate surface expression of the leukocyte adhesion molecule-1 on human neutrophils, monocytes, and their precursors. *J Immunol.* 1990;145(2):576-584.
47. Krugluger W, Nell A, Solar P, Matejka M, Boltz-Nitulescu G. Influence of sE-selectin and L-selectin on the regulation of cell migration during chronic periodontitis. *J Periodontol Res.* 1995;30(3):198-203.
48. Reyes L, Herrera D, Kozarov E, Rolda S, Progulske-Fox A. Periodontal bacterial invasion and infection: contribution to atherosclerotic pathology. *J Periodontol.* 2013;84(4 Suppl):S30-50.
49. Nakajima M, Arimatsu K, Kato T, et al. Oral Administration of P. gingivalis Induces Dysbiosis of Gut Microbiota and Impaired Barrier Function Leading to Dissemination of Enterobacteria to the Liver. *PLoS One.* 2015;10(7):e0134234.
50. Tomás I, Diz P, Tobías A, Scully C, Donos N. Periodontal health status and bacteraemia from daily oral activities: Systematic review/meta-analysis. *J Clin Periodontol.* 2012;39(3):213-228.
51. Monteseirin J, Chacon P, Vega A, et al. L-selectin expression on neutrophils from allergic patients. *Clin Exp Allergy.* 2005;35(9):1204-1213.

52. Ohsaka A, Saionji K, Sato N, Mori T, Ishimoto K, Inamatsu T. Granulocyte colony-stimulating factor down-regulates the surface expression of the human leucocyte adhesion molecule-1 on human neutrophils in vitro and in vivo. *Br J Haematol.* 1993;84(4):574-580.
53. Kuwano ST, Bordin JO, Chiba AK, et al. Allelic polymorphisms of human Fcγ receptor IIa and Fcγ receptor IIIb among distinct groups in Brazil. *Transfusion (Paris).* 2000;40(11):1388-1392.
54. Flesch B, Bauer F, Neppert J. Rapid typing of the human Fc gamma receptor IIA polymorphism by polymerase chain reaction amplification with allele-specific primers. *Transfusion (Paris).* 1998;38(2):174-176.
55. Bredius RG, Fijen CA, De Haas M, et al. Role of neutrophil Fc gamma RIIa (CD32) and Fc gamma RIIIb (CD16) polymorphic forms in phagocytosis of human IgG1- and IgG3-opsonized bacteria and erythrocytes. *Immunology.* 1994;83(4):624-630.
56. Salmon JE, Edberg JC, Brogle NL, Kimberly RP. Allelic polymorphisms of human Fc gamma receptor IIA and Fc gamma receptor IIIB. Independent mechanisms for differences in human phagocyte function. *J Clin Invest.* 1992;89(4):1274-1281.
57. Dimou NL, Nikolopoulos GK, Hamodrakas SJ, Bagos PG. Fcγ receptor polymorphisms and their association with periodontal disease: a meta-analysis. *J Clin Periodontol.* 2010;37(3):255-265.
58. Willcocks LC, Lyons PA, Clatworthy MR, et al. Copy number of FCGR3B, which is associated with systemic lupus erythematosus, correlates with protein expression and immune complex uptake. *J Exp Med.* 2008;205(7):1573-1582.
59. McKinney C, Fanciulli M, Merriman ME, et al. Association of variation in Fcγ receptor 3B gene copy number with rheumatoid arthritis in Caucasian samples. *Ann Rheum Dis.* 2010;69(9):1711-1716.
60. Aitman TJ, Dong R, Vyse TJ, et al. Copy number polymorphism in Fcgr3 predisposes to glomerulonephritis in rats and humans. *Nature.* 2006;439(7078):851-855.
61. Tak T, Wijten P, Heeres M, et al. Human CD62Ldim neutrophils identified as a separate subset by proteome profiling and in vivo pulse-chase labeling. *Blood.* 2017;129(26):3476-3485.
62. Hong CW. Current understanding in neutrophil differentiation and heterogeneity. *Immune Netw.* 2017;17(5):298-306.
63. Hellebrekers P, Hietbrink F, Vrisekoop N, Leenen LPH, Koenderman L. Neutrophil functional heterogeneity: Identification of competitive phagocytosis. *Front Immunol.* 2017;8(NOV):1498.

64. Silvestre-Roig C, Hidalgo A, Soehnlein O. Neutrophil heterogeneity: Implications for homeostasis and pathogenesis. *Blood*. 2016;127(18):2173-2181.
65. de Kleijn S, Langereis JD, Leentjens J, et al. IFN- $\gamma$ -Stimulated Neutrophils Suppress Lymphocyte Proliferation through Expression of PD-L1. *PLoS One*. 2013;8(8):e72249.
66. Tak T, Wijten P, Heeres M, et al. Human CD62L(dim) neutrophils identified as a separate subset by proteome profiling and in vivo pulse-chase labeling. *Blood*. 2017;129(26):3476-3485.
67. Förner L, Larsen T, Kilian M, Holmstrup P. Incidence of bacteremia after chewing, tooth brushing and scaling in individuals with periodontal inflammation. *J Clin Periodontol*. 2006;33(6):401-407.
68. Leliefeld PHC, Pillay J, Vrisekoop N, Heeres M, Tak T, Kox M. Differential antibacterial control by neutrophil subsets. *Blood Adv*. 2018;2:1344-1355.
69. Tulgar YK, Cakar S, Tulgar S, Dalkilic O, Cakiroglu B, Uyanik BS. The effect of smoking on neutrophil/lymphocyte and platelet/lymphocyte ratio and platelet indices: a retrospective study. *Eur Rev Med Pharmacol Sci*. 2016;20(14):3112-3118.
70. Pedersen KM, Colak Y, Ellervik C, Hasselbalch HC, Bojesen SE, Nordestgaard BG. Smoking and Increased White and Red Blood Cells. *Arterioscler Thromb Vasc Biol*. 2019;39(5):965-977.
71. Schwartz J, Weiss ST. Cigarette smoking and peripheral blood leukocyte differentials. *Ann Epidemiol*. 1994;4(3):236-242.
72. Ryder MI, Wu TC, Kallaos SS, Hyun W. Alterations of neutrophil f-actin kinetics by tobacco smoke: implications for periodontal diseases. *J Periodontal Res*. 2002;37(4):286-292.
73. MacFarlane GD, Herzberg MC, Wolff LF, Hardie NA. Refractory periodontitis associated with abnormal polymorphonuclear leukocyte phagocytosis and cigarette smoking. *J Periodontol*. 1992;63(11):908-913.
74. Drost EM, Selby C, Lannan S, Lowe GD, MacNee W. Changes in neutrophil deformability following in vitro smoke exposure: mechanism and protection. *Am J Respir Cell Mol Biol*. 1992;6(3):287-295.
75. Palmer RM, Wilson RF, Hasan AS, Scott DA. Mechanisms of action of environmental factors--tobacco smoking. *J Clin Periodontol*. 2005;32 Suppl 6:180-195.
76. Lakschevitz FS, Hassanpour S, Rubin A, Fine N, Sun C, Glogauer M. Identification of neutrophil surface marker changes in health and inflammation using high-throughput screening flow cytometry. *Exp Cell Res*. 2016;342(2):200-209.
77. Watanabe K, Hagen KL, Andersen BR. CD11b expression on neutrophils in human crevicular fluid collected from clinically healthy gingivae. *J Periodontal Res*. 1991;26(2):91-96.

78. Mouynet P, Picot C, Nicolas P, et al. Ex vivo studies of polymorphonuclear neutrophils from patients with early-onset periodontitis: (III). CR3 and LFA-1 expression by peripheral blood and gingival crevicular polymorphonuclear neutrophils. *J Clin Periodontol*. 1995;22(2):110-117.
79. Rijkschroeff P, Jansen IDC, van der Weijden FA, Keijser BJF, Loos BG, Nicu EA. Oral polymorphonuclear neutrophil characteristics in relation to oral health: a cross-sectional, observational clinical study. *International Journal Of Oral Science*. 2016;8:191.
80. Wellappuli NC, Fine N, Lawrence HP, Goldberg M, Tenenbaum HC, Glogauer M. Oral and Blood Neutrophil Activation States during Experimental Gingivitis. *JDR Clin Trans Res*. 2018;3(1):65-75.
81. Lakschevitz FS, Aboodi GM, Glogauer M. Oral neutrophil transcriptome changes result in a pro-survival phenotype in periodontal diseases. *PLoS One*. 2013;8(7):e68983.
82. Van Eeden S, Miyagashima R, Haley L, Hogg JC. L-selectin expression increases on peripheral blood polymorphonuclear leukocytes during active marrow release. *Am J Respir Crit Care Med*. 1995;151(2):500-507.
83. Van Eeden SF, Bicknell S, Walker B, Hogg JC. Polymorphonuclear leukocytes L-selectin expression decreases as they age in circulation. *American Journal of Physiology-Heart and Circulatory Physiology*. 1997;272(1):H401-H408.
84. Díaz-González F, González-Alvaro I, Campanero MR, et al. Prevention of in vitro neutrophil-endothelial attachment through shedding of L-selectin by nonsteroidal antiinflammatory drugs. *The Journal of clinical investigation*. 1995;95(4):1756-1765.
85. Burton J, Kehrl M, Kapil S, Horst R. Regulation of L-selectin and CD18 on bovine neutrophils by glucocorticoids: effects of cortisol and dexamethasone. *J Leukoc Biol*. 1995;57:317 - 325.
86. Razak K, Newland AC. Induction of CD13 expression on fresh myeloid leukaemia: correlation of CD13 expression with aminopeptidase-N activity. *Leuk Res*. 1992;16(6):625-630.
87. Kuijpers T, Tool A, Van der Schoot C, et al. Membrane surface antigen expression on neutrophils: a reappraisal of the use of surface markers for neutrophil activation. *Blood*. 1991;78(4):1105-1111.
88. Hart SP, Ross JA, Ross K, Haslett C, Dransfield I. Molecular characterization of the surface of apoptotic neutrophils: implications for functional downregulation and recognition by phagocytes. *Cell Death Differ*. 2000;7(5):493-503.

**Table 1. Cohort demographics and periodontal parameters.**

	Health (n = 40)	Periodontitis			
		Baseline (n = 54)	3-months (n = 46)	6-months (n = 44)	12-months (n = 37)
Sex (n, % male)	14 (35%)	20 (37%)			
Age at baseline (years, mean ± SD)	49.30 ± 10.62	53.28 ± 11.44			
Smokers (n, %smokers)	6 (15.0%)	15 (27.8%)	12 (26.1%)	10 (22.7%)	7 (18.9%)
CD16b present (n, % present)	25 (62.5%)	31 (57.4%)	25 (54.3%)	23 (52.3%)	19 (51.4%)
<b>Periodontal parameters</b>					
Mean PD (mm, mean ± SD)	2.80 ± 0.28	3.72 ± 0.74†	3.01 ± 0.51†‡	2.92 ± 0.55†‡	2.80 ± 0.44†‡§
%BOP (%mean ± SD)	10.86 ± 7.72	51.61 ± 25.57†	28.35 ± 15.17†‡	25.82 ± 18.84†‡	24.66 ± 18.05†‡

† significantly different to health

‡ significantly different to baseline

§ significantly different to 3-months post-treatment

%BOP – percentage of sites exhibiting bleeding on probing, mm – millimetres, PD – probing depth,

SD – standard deviation

**Table 2. Correlations between periodontal parameters and neutrophil subsets.**

Neutrophil phenotypes	Health		Periodontitis							
			Baseline		3-months		6-months		12-months	
	Mean PD	%BOP	Mean PD	%BOP	Mean PD	%BOP	Mean PD	%BOP	Mean PD	%BOP
Total Neutrophils	0.068	0.243	-0.073	-0.226	0.039	-0.050	<b>-0.379</b>	-0.343	-0.149	0.068
CD11b <sup>+</sup> neutrophils	-0.077	-0.066	-0.033	0.131	-0.052	0.212	-0.296	-0.223	0.193	0.315
CD16b <sup>+</sup> neutrophils	0.136	-0.063	0.142	-0.034	0.091	0.445	0.208	0.226	0.179	-0.086
CD62L <sup>-</sup> neutrophils	0.037	0.200	-0.161	-0.139	-0.176	-0.321	0.197	0.171	0.002	-0.187
CD66b <sup>+</sup> neutrophils	0.228	0.104	0.213	0.264	0.174	0.063	0.157	0.241	<b>0.378</b>	0.344
CD16b <sup>+</sup> CD66b <sup>+</sup> CD11b <sup>+</sup> neutrophils	0.094	-0.051	0.170	-0.057	0.143	<b>0.505</b>	0.205	0.290	0.217	-0.006
CD16b <sup>+</sup> CD66b <sup>+</sup> CD11b <sup>+</sup> CD62L <sup>-</sup> neutrophils	0.209	<b>0.401</b>	-0.185	-0.264	0.172	-0.172	0.461	0.168	-0.325	-0.086
Promyelocytes (CD11b <sup>+</sup> CD16b <sup>-</sup> )	0.020	-0.311	-0.162	0.127	-0.229	<b>-0.547</b>	0.081	0.194	-0.266	-0.113
Metamyelocytes (CD11b <sup>+</sup> CD16b <sup>-</sup> )	-0.010	0.218	-0.292	-0.133	-0.036	-0.372	-0.272	-0.393	-0.223	0.000
Mature neutrophils (CD11b <sup>+</sup> CD16b <sup>+</sup> )	0.124	-0.045	0.175	0.042	0.081	<b>0.468</b>	0.205	0.279	0.232	0.017
Bands (CD16 <sup>dim</sup> CD62L <sup>bright</sup> )	-0.032	-0.255	-0.148	-0.112	-0.325	-0.198	-0.172	-0.388	-0.260	-0.308
Normal neutrophils (CD16 <sup>bright</sup> CD62L <sup>bright</sup> )	-0.116	-0.238	0.098	0.095	-0.018	0.328	-0.316	-0.112	<b>0.514</b>	0.412
Suppressive neutrophils (CD16 <sup>bright</sup> CD62L <sup>dim</sup> )	0.169	0.266	-0.098	-0.120	0.022	-0.084	0.446	0.171	-0.360	-0.236

Spearman's correlation coefficients ( $\rho$ ) between the periodontal parameters, mean probing depth (PD) and percentage of sites exhibiting bleeding on probing (%BOP), and various neutrophil subsets in health and at baseline, 3-, 6- and 12- months in periodontitis. Significant correlations ( $p \leq 0.05$ ) are represented in bold.




Fig. 1

jre\_12793\_f1.tif




Fig. 2

jre\_12793\_f2.tif




Fig. 3

jre\_12793\_f3.tif




Fig. 4

jre\_12793\_f4.tif




Fig. 5

jre\_12793\_f5.tif