



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Fakhri, O;Hartley, CA;Devlin, JM;Browning, GF;Noormohammadi, AH;Lee, SW

Title:

Development and application of high-resolution melting analysis for the classification of infectious laryngotracheitis virus strains and detection of recombinant progeny

Date:

2019-02-01

Citation:

Fakhri, O., Hartley, C. A., Devlin, J. M., Browning, G. F., Noormohammadi, A. H. & Lee, S. W. (2019). Development and application of high-resolution melting analysis for the classification of infectious laryngotracheitis virus strains and detection of recombinant progeny. *Archives of Virology*, 164 (2), pp.427-438. <https://doi.org/10.1007/s00705-018-4086-1>.

Persistent Link:

<https://hdl.handle.net/11343/253127>

License:

CC BY



# Development and application of high-resolution melting analysis for the classification of infectious laryngotracheitis virus strains and detection of recombinant progeny

Omid Fakhri<sup>1</sup> · Carol A. Hartley<sup>1</sup> · Joanne M. Devlin<sup>1</sup> · Glenn F. Browning<sup>1</sup> · Amir H. Noormohammadi<sup>2</sup> · Sang-Won Lee<sup>1,3</sup>

Received: 9 July 2018 / Accepted: 15 October 2018 / Published online: 12 November 2018

© The Author(s) 2018

## Abstract

Live attenuated vaccines against infectious laryngotracheitis virus (ILTV) are widely used in the poultry industry to control disease and help prevent economic losses. Molecular epidemiological studies of currently circulating strains of ILTV within poultry flocks in Australia have demonstrated the presence of highly virulent viruses generated by genomic recombination events between vaccine strains. In this study, high-resolution melting (HRM) analysis was used to develop a tool to classify ILTV isolates and to investigate ILTV recombination. The assay was applied to plaque-purified progeny viruses generated after co-infection of chicken embryo kidney (CEK) monolayers with the A20 and Serva ILT vaccine strains and also to viruses isolated from field samples. The results showed that the HRM analysis is a suitable tool for the classification of ILTV isolates and can be used to detect recombination between ILTV vaccine strains *in vitro*. This method can be used to classify a broad range of ILTV strains to facilitate the classification and genotyping of ILTV and help to further understand recombination in these viruses.

## Introduction

Infectious laryngotracheitis (ILT), caused by infectious laryngotracheitis virus (ILTV, species *Gallid alphaherpesvirus 1*), has an impact on the poultry industry worldwide [1]. Live attenuated vaccines are widely available and are often administered in the early life of commercial birds, followed by one or more rounds of revaccination in order to reach

the desired level of immunity [2, 3]. Although the level of immunity developed by this vaccination procedure is adequate to protect the chickens from developing clinical disease (respiratory signs followed by reduced weight gain and egg production), the challenge virus is often able to replicate in the vaccinated birds [2, 4].

All of the ILTV strains form a single serotype [5], yet have shown a notable level of genotypic variation in different studies (0.1–0.8% nucleotide sequence difference) [6]. A number of different methods have been published to classify these viruses by genotype [7–10], including polymerase chain reaction restriction fragment length polymorphism (PCR-RFLP) [9–13], targeted sequencing [7, 8], and strain-specific fluorescent probe hydrolysis [14]. In all methods, multiple regions of the ~150-kbp genome are analyzed in order to more precisely differentiate genotypes. These methods have been used to classify ILT viruses to facilitate epidemiological studies and clinical diagnosis [15].

Intraspecific genomic recombination is considered an important driver in the evolution of ILTV and other herpesviruses [16] that have low synonymous nucleotide sequence substitution rates [17]. Genomic recombination requires coinfection with two or more strains, where replication of challenge virus in vaccinated chickens and reactivation of

Handling Editor: Scott Schmid.

**Electronic supplementary material** The online version of this article (<https://doi.org/10.1007/s00705-018-4086-1>) contains supplementary material, which is available to authorized users.

✉ Omid Fakhri  
omid.fakhri@unimelb.edu.au

<sup>1</sup> Asia-Pacific Centre for Animal Health, Faculty of Veterinary and Agricultural Sciences, The University of Melbourne, Parkville, VIC, Australia

<sup>2</sup> Asia-Pacific Centre for Animal Health, Faculty of Veterinary and Agricultural Sciences, The University of Melbourne, Werribee, VIC, Australia

<sup>3</sup> College of Veterinary Medicine, Konkuk University, Seoul, Republic of Korea

latent ILTV infection (either vaccine or field strains) can provide an environment that allows genomic recombination [18]. Intraspecific genomic recombination between vaccine strains of ILTV has been reported to occur in the field [18] and has been demonstrated for field strains under laboratory conditions [14, 18–20]. However, the recombination between modified live vaccine strains has not been studied in experimental settings.

High-resolution melting (HRM) analysis is a simple, rapid and cost-efficient method for genotyping of multiple single-nucleotide polymorphisms (SNPs) [21]. This method has been used in several pathogen identification procedures [22], as well as mutation scanning and diagnosis of genetic disorders in human genome research [23–25]. The extensive applications of this method demonstrate its wide potential to be used as an adjunct to current detection methods.

This study aims to develop an accurate and cost-effective HRM analysis that can be applied to the classification of known and emerging ILTV strains, and to study genomic recombination of ILTV vaccine strains under controlled *in vitro* conditions.

## Materials and methods

### Viruses

The eight ILTV isolates used in this study are listed in Table 1. The viruses (other than the commercial vaccine viruses) were plaque purified, propagated on primary CEK cells and re-typed by PCR-RFLP [9] prior to testing. In order to increase the titre of commercial vaccines, each vaccine was passaged twice on the chorioallantoic membranes (CAM) of specific-pathogen-free (SPF) embryonated eggs (Australian SPF Services, Woodend, Australia) using artificial air-sac inoculation [26]. The infected CAMs were extracted and thoroughly minced using a mortar and pestle.

**Table 1** Viruses used in this study

Isolate	PCR-RFLP class [9]	GenBank accession number	Reference
NOBILIS® ILT (Serva)	7	HQ630064	MSD Animal Health
Poultvac® Laryngo A20	1	JN596963	Zoetis
ACC78	8	JN804826	[18]
Class 9	9	JN804827	[18]
Class 10	10	KR822401	[15]
SA2	1	JN596962	[6]
CSW-1	4	JX646899	[19]
V1-99	2	JX646898	[19]

The homogenates were allowed to sediment for 5 minutes at room temperature to let the larger fragments settle before the supernatants were collected. The harvested viruses were cultured on CEK monolayers. Viruses were harvested at 48 hours after infection of monolayers, and aliquots were stored at -80 °C.

### Cell culture

CEK cells were prepared using kidneys harvested from 18-day-old chick embryos as described previously [27]. Monolayers of CEK cells were maintained in Dulbecco's modified Eagle medium (DMEM) containing 1% v/v foetal bovine serum (FBS), and 100 µg of ampicillin and 10 µg of amphotericin B per mL. All *in vitro* and *in ovo* experiments were performed on CEK cells or CAM from embryonated SPF hen eggs (Australian SPF Services, Woodend, Australia).

### Coinfection of CEK monolayers

Primary CEK monolayers were infected with a mixture of Serva and A20 ILTV strains in 6-well tissue culture plates. After 1 hour of incubation at 37 °C, the inoculum was removed and 3 mL of fresh medium was added. The total multiplicity of infection (MOI) for the coinfection studies was 10. The progeny virus populations were collected by scraping the cells into culture supernatant 48 hours post infection when an extensive cytopathic effect was observed. Samples collected from this experiment were stored at -80 °C prior to virus isolation and plaque purification.

### Virus isolation and plaque purification

Ten-fold dilutions of progeny viruses from an *in vitro* coinfection experiment were prepared and inoculated onto CEK monolayers in 6-well plates as described previously [14]. Following 1 hour of incubation at 37 °C, the inoculum was removed and the monolayers were overlaid with methylcellulose semi-solid medium containing 1% w/v methylcellulose in DMEM, with 1% v/v FBS, and 50 µg of ampicillin, 50 µg of gentamicin and 5 µg of amphotericin B per mL. After 48 to 72 hours of incubation at 37 °C in a humidified atmosphere of 5% v/v CO<sub>2</sub>, 20 well-isolated plaques were picked using a micropipette guided under an inverted light microscope. Picked plaques were resuspended in 500 µL of DMEM, and each plaque was purified through three rounds of plaque purification before a final propagation step on CEK cells in 24-well plates. One freeze/thaw cycle was performed between rounds of plaque purification.

## DNA extraction

Nucleic acid extraction was performed using a PureLink<sup>®</sup> Pro 96 viral RNA/DNA purification kit (Invitrogen, USA) and a QIAxtractor automated vacuum system. Purified DNA from 200- $\mu$ L samples was eluted in 200  $\mu$ L of pyrogen-free water (Milli-Q<sup>®</sup> Integral system, Darmstadt, Germany) and stored at -20 °C until testing. Pyrogen-free water, rather than the elution buffer supplied with the extraction kit, was used to avoid introducing any variation in HRM patterns.

## HRM-based multiple-SNP ILTV genotyping assay

### Assay conditions

HRM assays were developed to be used as a method of classification and genotyping ILTV. Following the alignment of full genome sequences of the virus isolates (Table 1), 662 SNPs were identified. These potential target regions were further analyzed *in silico* using uMelt software [28] in order to identify the regions with a detectable melting temperature ( $T_m$ ) gap in their PCR products. Ultimately, six regions were selected based on their even distribution across the genome (in order to increase the chance of detecting recombinants), the ability to measure differences in their  $T_m$  values, and their ability to produce unique patterns for known/tested ILTV strains. Targeted sites included both transition and transversion mutations and resulted in either synonymous or non-synonymous changes (Table 2 and 3). The primers used for this assay are listed in Table 2. A unique single-nucleotide polymorphism (SNP) pattern was predicted for each strain based on the SNPs present in six genomic regions (Fig. 1A). These regions were also amplified and sequenced (Big Dye Terminator v3.1, Life Technologies) to confirm the presence of the targeted SNPs in these viruses.

For the HRM assay, all reactions were prepared using a QIAgility<sup>®</sup> automated system (QIAGEN). Reaction mixtures (20  $\mu$ L) were prepared containing 2 mM MgCl<sub>2</sub>, 200  $\mu$ M each dNTP, 400 nM each forward and reverse primer, 8  $\mu$ M SYTO<sup>®</sup> 9 (Thermo Fisher Scientific), 0.04 U of GoTaq<sup>®</sup> Flexi DNA Polymerase (Promega, Wisconsin, USA) per mL, Colorless GoTaq<sup>®</sup> Flexi Buffer (Promega, Wisconsin, USA), and 2  $\mu$ L of DNA template. PCR amplification and the HRM assay were performed on a Rotor-Gene Q 2plex HRM instrument (QIAGEN Inc., MD, USA) in 100-tube rings (Rotor-Disc 100; QIAGEN, Australia).

The thermal cycling profile included 45 cycles of a denaturation step at 95 °C for 30 seconds, an annealing step at a primer-specific temperature (listed in Table 2) for 30 seconds, and 30 seconds of extension at 72 °C. A final extension step at 72 °C for 5 minutes was included before the melting step. The amplified products were incubated at 95 °C for 2 minutes before a renaturation step at 60 °C for 2

minutes. The HRM was performed in the temperature range of 70–95 °C with fluorescence acquisition at every 0.3 °C and 2-second ramps. Every reaction was performed in triplicate and in three independent experiments unless stated otherwise. Controls in each run included a no-template control and the appropriate reference-virus-positive controls (Table 1). In order to generate a reference for identification of possible mixed populations that were not plaque purified as expected, a 1:1 mixture of the two parent strains was also included in the experimental setup to identify samples containing more than one variant type.

### Principal component analysis of the HRM output

The HRM data were processed using Rotor-Gene Q Series software (version 2.1.0 Build 9) and Rotor-Gene ScreenClust HRM software (version 1.10.1.3) [29] in supervised mode. Principal component analysis (PCA) was used to better differentiate the HRM profiles based on the sequence variations in the parent viruses (A20 and Serva). The ScreenClust software normalizes the pre- and post-melting fluorescence levels, generates residual plots, identifies two or three principal components (PC) in the normalized data, defines clusters based on the control samples, assigns unknown/test samples into these defined clusters, and reports the probabilities and typicalities of unknown samples belonging to known clusters based on posterior class probabilities [29]. When all of the unknown samples are classified, the covariance of each cluster is determined (represented by ellipsoids in Fig. 2b). The mean of the reference samples forms the center of the cluster. The test samples are classified into each cluster with posterior probabilities and typicality scores (Supplementary Table 1). These were determined for pure populations of each of the parent viruses as well as for the control samples containing the 1:1 premixed template of each parent virus. The PCA plots of each parent and the mixed populations showed distinct and non-overlapping clusters that made it possible to clearly differentiate each population (each of the two parents and the mixed population). A range of recent studies have used the ScreenClust software for further analysis alongside traditional HRM analysis in higher-throughput settings [30–34].

### *In silico* analysis for classification of published ILTV strains

The full genome sequences of five additional ILTV strains were collected from GenBank (Table 4) to enable *in silico* analyses and to examine the potential for a wider application of the HRM-based genotyping to ILTV strains, including strains from other geographical regions. Using the methods previously established with the Australian ILTV isolates, a series of *in silico* amplifications was performed in Geneious 11.1.4 [35] on multiple regions of five ILTV strains that

**Table 2** List of primers and selected target SNPs tested on Australian ILTV isolates

Target gene	Primer set (5'-3')	Annealing temp. (°C)	Amplicon length	$T_m$ gap (°C) experimental [predicted <sup>a</sup> ]	No. of SNPs in amplicon	Type of SNP (NS/S) <sup>b</sup>	SNPs									
							Serva	A20	CL8	CL9	CL10	SA2	CSW-1	V1-99		
UL52	F: GGTCGGTTTCTAGCTGTGG	49	130	0.4 [0.8]	2	TS (S)	A	G	A	A	A	G	G	G		
	R: TTGCTCCCTTAAAGTAGGTACTTT						T	G	T	T	T	G	G			
UL27	F: CATGGTTGGTCCCTGG	50	186	0.4 [0.6]	2	TS (NS)	T	C	C	C	C	C	T	T		
	R: CGCCAACTGATTGTGGCA						A	G	G	G	G	G	G			
UL36	F: AGATGTTTTAGTTCTGTGGGC	50	120	1.4 [1.3]	3	TS (NS)	T	C	T	C	T	C	T	T		
	R: GGCTGCACTCCTCCAAGATC						T	C	T	C	T	C	T	C	T	C
UL8	F: ATTGATGGAAGAGGAATGAAGCA	55	100	0.8 [1.2]	3	TS (NS)	T	C	T	C	T	C	T	T		
	R: TTGTTTTCCGGGGTGTAAAGG						C	A	C	A	C	A	C	C	A	C
IR	F: GTCAACGGGGCTAGATCATT	50	100	0.6 [0.9]	2	TS (n/a)	T	C	T	C	C	C	T	T		
	R: ATTACTTCTGGTTGCGTGCTC						T	A	T	T	A	A	T	T		
US4	F: TTCGTGTTCCGGGTGGATAG	54	125	0.5 [0.5]	3	TV (NS)	C	A	C	C	A	A	A	A		
	R: TCGAAATCAGGTATCCCGGC						T	G	T	T	G	G	G	G		
ORFB	F: CAGTGACGTTAATTTACCGGGA	50	71	0.6 [1.1]	1	TS (NS)	T	C	T	T	T	T	T	T		
	R: CAGCTCGAGAAATGCAGCG						C	T	C	C	T	T	C	T		
US7	F: ACTGGAATCACGTCTCCGC	50	76	1.5 [1.7]	3	TS (NS)	A	A	A	A	A	A	G	A		
	R: GAATTGTAGCTTCGGGGCGA						A	G	A	A	A	G	G	G		
							T	T	T	T	T	T	C	T		

<sup>a</sup> Predicted  $T_m$  gap values were calculated using uMelt Batch software<sup>b</sup> TS transition, TV transversion, S synonymous, NS non-synonymous

**Table 3** List of primers and selected target SNPs for the classification of US isolates

Target gene	Primer set (5'-3')	Amplicon length	Predicted <sup>a</sup> $T_m$ gap (°C)	No. of SNPs in amplicon	Type of SNP (NS/S) <sup>b</sup>	SNPs						
						Laryngo Vac	TCO IVAX	CEO TRVX	LT Blen	1874C5	1874C5	1874C5
UL54	F: TGTTCTGGCCGGTCTATTG	62	1	1	TS (NS)	T	T	C	C	C	T	
	R: TTGTGTATCCCGGACCCAGAG											
UL10	F: AAACCCCTGCTTGGGACTAA	57	1.2	1	TS (NS)	T	C	T	T	T	C	
	R: CTGGTGTGATATCACTGGCCT											
ICP4-2	F: ATCGTCGTTGTCGGTCTTCC	69	0.9	1	TS (NS)	T	T	T	C	T	T	
	R: CACGTAGTAATGGACAGGCCGA											
ICP4	F: CGGAGCACTTGCCGGTAC	126	1.2	3	TS (S)	T	T	T	T	T	C	
	R: GTTGGCCGGGAGATTCTTGGG											

<sup>a</sup> Predicted  $T_m$  gap values were calculated using uMelt Batch software

<sup>b</sup> TS transition, TV transversion, S synonymous, NS non-synonymous

<sup>c</sup> In these 4-character HRM combination patterns, the letters represent the sequence type of UL54, UL10, ICP4 and a second region in ICP4 genes, respectively. The viruses with melting characteristics similar to those of the Laryngo Vac strain in each region are indicated by an "A" and otherwise by a "B"

originated from the USA (Table 3) in order to obtain strain-specific PCR products. Four new sets of primers (Table 3) were designed using Primer3 software (v. 2.3.4) [36]. The products of *in silico* PCRs were imported to uMelt Batch software (v. 2.2) [28] to predict the melting characteristics of each product.

## Results

### Differentiation of closely related ILTV genotypes by HRM analysis of SNPs


An HRM panel (Table 2) targeting SNPs across six sites (UL52, UL27, UL36, UL8, IR and US4) was applied to A20 and Serva vaccine strains and the well-characterized recombinant class 8, 9 and 10 ILTV strains (Table 1). A schematic view of the SNP pattern is shown in Figure 1A. The melt curve analysis from each of these reactions (Fig. 1B) on A20 and Serva was consistent with that predicted by the uMelt software, particularly in regards to the melting temperature separation predicted for each of the sites. This HRM analysis of six genomic regions enabled differentiation of each of the vaccine and recombinant field strains of ILTV (Fig. 1). The results obtained between assays were reproducible and showed 0.02 to 0.03% coefficients of variation between runs.

### Differentiation of parent from recombinant progeny viruses after recombination between Serva and A20 ILTV in coinfecting primary cell cultures

The HRM panel targeting the same six sites (UL52, UL27, UL36, UL8, IR and US4) was applied to DNA extracted from plaque-purified viruses generated after a 1:1 coinfection of CEK monolayers with the A20 and Serva ILTV vaccine strains (Fig. 2). Each tested region was classified as Serva-like or A20-like. Four isolates out of 20 tested samples were detected as mixed population of viruses and excluded from further analysis. The remaining 16 isolates were classified as either recombinant viruses or parent viruses (Fig. 2). In total, six different patterns of recombinants were detected, with no genotype(s) dominating amongst the recombinant progeny. Equal proportions of parent viruses and recombinant viruses were detected among the 20 progeny viruses tested (Fig. 2A).

### Adaptation of the HRM panel for genotyping a broader range of ILTV isolates

Analysis of the six genomic regions across A20 and Serva genomes by HRM has shown the ability of this technology to build a picture of recombination across the ILTV genome, and this system was adapted to explore the potential for use




**Fig. 1** HRM analysis of five fully sequenced ILTV isolates. (a) A schematic view of six target genomic markers with distinct melting characteristics showing the distribution of Serva-like (closed circles) and A20-like (open circles) SNPs in Serva, A20, class 8, 9 and 10 ILTV strains. (b) HRM plots showing the differentiation of Serva (red) and A20 (blue), along with the melt curves for class 8 (green),

class 9 (orange) and class 10 (purple) ILTV strains by the six target genomic markers (UL52, UL27, UL36, UL8, IR and US4). Each sample was clustered into SNP identities that were either Serva-like (closed circles) or A20-like (open circles). The plots were generated by Rotor-Gene Q software. Each curve represents one replicate

in the genotypic classification of other ILTV isolates. To do so, the genomes of three more Australian strains (one vaccine [SA2] and two field strains [class 2 and 4]) were retrieved from GenBank and compared to the Australian ILTV isolates used previously in this study (Table 1). A panel of four SNPs was identified that would differentiate the three vaccine strains used in Australia (SA2, A20 and

Serva) and five Australian genotypes (Table 1). This panel (Fig. 3A) includes two previously characterized regions (UL27 and UL36, Table 2, Fig. 1 and 2) and two new regions (ORFB and US7, Table 2, Fig. 3) predicted by uMelt to have a  $T_m$  gap of  $>1$  °C. When this new panel of four sets of HRM primers (ORFB, UL27, UL36 and US7) was applied to extracted DNA from these viruses (Table 1), the actual

**Fig. 2** HRM analysis of parent and progeny viruses generated two days after infection of CEK cells with a 1:1 mixture of Serva and A20 ILTV. (a) A schematic representation of SNP patterns in parent and progeny viruses is shown, along with (b) PCA cluster analysis and (c) HRM plots of the assays for each of the six genomic regions. (a) A schematic view of genotypes of plaque-purified parent and progeny viruses after coinfection. The closed circles represent the Serva-like SNPs, and open circles indicate A20-like SNPs. Each virus is genotyped as A20, Serva, or recombinant with a unique genotype code number (1 to 62) for each recombination pattern. (b) PCA cluster plots and (c) HRM plots show triplicate Serva (red) and A20 (blue) ILTV parent strains as references. A 1:1 mixture of Serva and A20 ILTV (green) are shown to identify mixed samples. Plaque-purified progeny viruses (grey squares/curves) are clustered into Serva-origin (red oval) SNPs, A20-origin SNPs (blue oval) or mixed populations (green oval) on PCA plots. Clusters are highlighted by ovals that represent the covariance of the classified samples in a two-PC-dimension plot. The samples that plot outside the ovals are classified into the closest cluster and receive a lower typicality score



**Table 4** Viruses used in *in silico* classification of US isolates

Isolate	PCR-RFLP class [10, 37]	GenBank accession number	Reference
Laryngo Vac	IV	JQ083494	[38]
TCO IVAX	II	JN580312	[39]
CEO TRVX	IV	JN580313	[39]
LT Blen	IV	JQ083493	[38]
1874C5	VI	JN542533	[37]

HRM findings were used to generate a four-character combination code based on the melting characteristics of each of the target regions. Using these four sites, the genotype of the five distinct Australian field strains and the three vaccine strains used in Australia could be differentiated from each other (Fig. 3). Moreover, the SA2 and A20 vaccine strains could be differentiated from each other in one region (ORF B) despite their close relationship (sharing 99.9% nucleotide sequence identity) and both being classified by PCR-RFLP as class 1 viruses.

### ***In silico* exploration of HRM-based genotyping to ILTV strains from other countries**

Given the consistency between the predicted and experimentally observed melting temperatures for HRM analysis, this method could be broadened as a means of genotyping viruses from different geographical regions. To illustrate this, similar *in silico* analysis of five US strains (Table 4) has identified four regions that could be used to differentiate US field and vaccine isolates (Table 3), suggesting the potential for the simple application of this technology to characterize a broad range of ILTV isolates across the world.



## **Discussion**

In this study, we aimed to develop a reliable cost- and time-efficient method that can be applied to the classification and genotyping of ILTV strains and the identification of ILTV recombinant viruses. The majority of existing ILTV classification systems are based on identifying the sequence variations in representative regions of the genome that are first amplified by PCR. Sequence variation is identified in these regions by a range of methodologies including RFLP [9, 10], Sanger sequencing [7, 8] or fluorescently labelled TaqMan<sup>®</sup> probes [14, 40, 41]. Both the number of regions being analyzed and the method of analysis will influence the precision of the classification and differentiation of ILTV isolates. Although PCR-RFLP results are relatively easy to interpret, other studies have suggested the use of Sanger

**Fig. 3** The genotyping of eight Australian ILTV field and vaccine isolates using HRM analysis. (a) A schematic view of a panel of four SNPs with distinct melting characteristics showing the distribution of Serva-like (closed circles), A20-like (open circles) and other SNPs (closed triangle) used to differentiate Australian field strains in Serva, A20, class 2, 4, 8, 9 and 10 ILTV strains and (b) HRM plots comparing four target genomic markers (ORFB, UL27, UL36 and US7) in a panel of viruses including Serva (red) and A20 (blue) to SA2 (brown), class 2 (V1-99) (light blue), class 4 (CSW-1) (light green), class 8 (dark green), class 9 (pink) and class 10 (orange) ILTV isolates. Each sample was clustered into either Serva-like (closed circles), A20-like (open circles) or other (triangles) SNP identities. The plots were generated by Rotor-Gene Q software. Each curve represents one replicate

sequencing of one, two [7, 42–45] or multiple [8] region(s) of ILTV genome as an alternative to RFLP.

The HRM analysis used in the current study could differentiate field and vaccine strains of ILTV and classify them into distinct genotypes that had previously been determined by PCR-RFLP [9] and full-genome sequencing [6, 18, 19]. Australian ILTV classes 3, 4, 5 and 6 were not examined in this study, as no genomic sequences were available for these viruses to validate the results. The results of HRM analysis were reproducible and were in accordance with other methods; however, the successful use of this method relies on the precision of techniques used during sample preparation, DNA isolation, amplification and melting, and post-reaction analysis steps, as described in detail by Slomka *et al.* [21]. In order to overcome the variations introduced by different isolation methods and sample composition, a single type of DNA isolation kit was used for both test samples and the samples used as reference strains in this study. The commercial vaccine strains were passaged once in CEK cells so that the composition of the reference strains and test samples would be similar. Automated systems were used for extraction and PCR preparation to maintain the uniformity of samples and reactions. The sizes of amplicons for each target region was also kept short (<200 bp) where possible in order to obtain a higher  $T_m$  gap between variant types. The uMelt melting curve prediction software provided a reliable guide for selecting target SNPs based on the melting peaks. The predicted melting temperatures were close to experimental melting peaks with only slight differences (zero to 0.4 °C) in some regions. Therefore, when designing the second and third set of primers, the regions with a  $T_m$  gap of  $\geq 1$  °C were preferred. In order to use this system to classify viruses submitted from outbreaks of disease in the field it is likely that viruses from any clinical samples would also require processing (isolation and DNA extraction) using uniform techniques. It is conceivable that future assays could be developed and optimized for direct use on specific clinical samples (e.g., tracheal tissue or conjunctival swabs) after extensive evaluations using DNA extracted directly from clinical specimens. This would have time and



cost advantages and represents an important area of future work. The direct use of the HRM assay on clinical samples has been successfully applied to several pathogens in the past [46–49]. The potential of HRM in identifying mixed specimens has been reported to be primarily dependent on the quantity and proportion of the target DNAs in the mixture [50]. Current detection or genotyping methods for ILTV have not been able to rapidly identify the presence of mixed virus populations in laboratory or field samples. The potential of this method to identify mixed infections in these settings requires further exploration.

The HRM system developed in this study could also be used successfully to detect recombinant viral progeny. This represents the first time that HRM methods have been used to detect recombination in herpesviruses. Previous studies have shown that genomic recombination between Serva and A20 ILTV strains has occurred in the natural host and has resulted in field recombinants with higher levels of virulence [18]. The results of the current study showed that these two vaccine strains can recombine in coinfecting (MOI = 10) CEK cells and generate a diverse pattern of recombinants. After this coinfection, recombinants accounted for 50% (8/16) of progeny viruses, and the highest proportion of an individual genotype (6/16) had an SNP profile identical to that of the A20 parent strain. The proportion of recombinants in the current study is consistent with the greater opportunity for coinfection of cells at this high MOI so that recombination can occur. These findings are consistent with findings from an *in vivo* coinfection study in SPF chickens using a mixture of CSW-1 and V1-99 ILTV strains delivered intratracheally, where 46.5% (in birds co-inoculated with 10<sup>3</sup> PFU of each virus) to 79.2% (in birds co-inoculated with 10<sup>4</sup> PFU of each virus) of isolated progeny viruses were identified as recombinants.

In conclusion, the HRM assay developed in this study was successfully applied as a high-throughput screening tool for recombinant detection and could be used to successfully classify different strains of ILTV. In these applications the assay was shown to be accurate, time-efficient and cost-efficient. The HRM genotyping method promises to be an important tool to facilitate the study of ILTV epidemiology and recombination in both experimental and field settings.

**Acknowledgements** We gratefully acknowledge Nino Ficorilli and Cynthia Brown for their technical assistance.

**Funding** SWL is funded by an Australian Research Council Discovery Early Career Researcher award. OF is supported by a Melbourne International Research Scholarship from The University of Melbourne.

## Compliance with ethical standards

**Conflict of interest** The authors declare that they have no conflict of interest.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

1. Bagust T, Jones R, Guy J (2000) Avian infectious laryngotracheitis. *Revue Sci Tech-Office Int Epizoot* 19(2):483–488
2. García M, Spatz S, Guy JS (2013) Infectious laryngotracheitis. In: Swayne D, Glisson J, McDougald L, Nolan L, Suarez D (eds) *Diseases of poultry*, 13th edn. Wiley-Blackwell, Iowa, pp 161–179
3. Vagnozzi A, Zavala G, Riblet SM, Mundt A, García M (2012) Protection induced by commercially available live-attenuated and recombinant viral vector vaccines against infectious laryngotracheitis virus in broiler chickens. *Avian Pathol* 41(1):21–31
4. Coppo MJ, Noormohammadi AH, Hartley CA, Gilkerson JR, Browning GF, Devlin JM (2011) Comparative *in vivo* safety and efficacy of a glycoprotein G-deficient candidate vaccine strain of infectious laryngotracheitis virus delivered via eye drop. *Avian Pathol* 40(4):411–417
5. Cover M, Benton W (1958) The biological variation of the infectious laryngotracheitis virus. *Avian Dis* 2(4):375–383
6. Lee SW, Devlin JM, Markham JF, Noormohammadi AH, Browning GF, Ficorilli NP, Hartley CA, Markham PF (2011) Comparative analysis of the complete genome sequences of two Australian origin live attenuated vaccines of infectious laryngotracheitis virus. *Vaccine* 29(52):9583–9587. <https://doi.org/10.1016/j.vaccine.2011.10.055>
7. Chacon JL, Ferreira AJ (2009) Differentiation of field isolates and vaccine strains of infectious laryngotracheitis virus by DNA sequencing. *Vaccine* 27(48):6731–6738. <https://doi.org/10.1016/j.vaccine.2009.08.083>
8. Choi EJ, La TM, Choi IS, Song CS, Park SY, Lee JB, Lee SW (2016) Genotyping of infectious laryngotracheitis virus using allelic variations from multiple genomic regions. *Avian Pathol* 45(4):443–449. <https://doi.org/10.1080/03079457.2016.1155692>
9. Kirkpatrick NC, Mahmoudian A, O'Rourke D, Noormohammadi AH (2006) Differentiation of infectious laryngotracheitis virus isolates by restriction fragment length polymorphic analysis of polymerase chain reaction products amplified from multiple genes. *Avian Dis* 50(1):28–34. <https://doi.org/10.1637/7414-072205R.1>
10. Oldoni I, Garcia M (2007) Characterization of infectious laryngotracheitis virus isolates from the US by polymerase chain reaction and restriction fragment length polymorphism of multiple genome regions. *Avian Pathol* 36(2):167–176. <https://doi.org/10.1080/03079450701216654>
11. Andreasen JR Jr, Glisson JR, Villegas P (1990) Differentiation of vaccine strains and Georgia field isolates of infectious laryngotracheitis virus by their restriction endonuclease fragment patterns. *Avian Dis* 34(3):646–656
12. Guy JS, Barnes HJ, Munger LL, Rose L (1989) Restriction endonuclease analysis of infectious laryngotracheitis viruses: comparison of modified-live vaccine viruses and North Carolina field isolates. *Avian Dis* 33(2):316–323
13. Keeler CL Jr, Hazel JW, Hastings JE, Rosenberger JK (1993) Restriction endonuclease analysis of Delmarva field isolates of infectious laryngotracheitis virus. *Avian Dis* 37(2):418–426
14. Loncoman CA, Hartley CA, Coppo MJ, Vaz PK, Diaz-Mendez A, Browning GF, Lee SW, Devlin JM (2017) Development and

- application of a TaqMan single nucleotide polymorphism genotyping assay to study infectious laryngotracheitis virus recombination in the natural host. *PLoS One* 12(3):e0174590. <https://doi.org/10.1371/journal.pone.0174590>
15. Agnew-Crumpton R, Vaz PK, Devlin JM, O'Rourke D, Blacker-Smith HP, Konsak-Ilievski B, Hartley CA, Noormohammadi AH (2016) Spread of the newly emerging infectious laryngotracheitis viruses in Australia. *Infect Genet Evol* 43:67–73
  16. Thiry E, Meurens F, Muylkens B, McVoy M, Gogev S, Thiry J, Vanderplasschen A, Epstein A, Keil G, Schynts F (2005) Recombination in alphaherpesviruses. *Rev Med Virol* 15(2):89–103. <https://doi.org/10.1002/rmv.451>
  17. Sakaoka H, Kurita K, Iida Y, Takada S, Umene K, Kim YT, Ren CS, Nahmias AJ (1994) Quantitative analysis of genomic polymorphism of herpes simplex virus type 1 strains from six countries: studies of molecular evolution and molecular epidemiology of the virus. *J Gen Virol* 75(Pt 3):513–527. <https://doi.org/10.1099/0022-1317-75-3-513>
  18. Lee S-W, Markham PF, Coppo MJ, Legione AR, Markham JF, Noormohammadi AH, Browning GF, Ficorilli N, Hartley CA, Devlin JM (2012) Attenuated vaccines can recombine to form virulent field viruses. *Science* 337(6091):188
  19. Lee SW, Devlin JM, Markham JF, Noormohammadi AH, Browning GF, Ficorilli NP, Hartley CA, Markham PF (2013) Phylogenetic and molecular epidemiological studies reveal evidence of multiple past recombination events between infectious laryngotracheitis viruses. *PLoS One* 8(2):e55121. <https://doi.org/10.1371/journal.pone.0055121>
  20. Loncoman CA, Hartley CA, Coppo MJC, Vaz PK, Diaz-Mendez, Browning GF, Garcia M, Spatz S, Devlin JM (2017) Genetic diversity of infectious laryngotracheitis virus during in vivo coinfection parallels viral replication and arises from recombination hot spots within the genome. *Appl Environ Microbiol*. <https://doi.org/10.1128/aem.01532-17>
  21. Slomka M, Sobalska-Kwapis M, Wachulec M, Bartosz G, Strapagiel D (2017) High resolution melting (HRM) for high-throughput genotyping—limitations and caveats in practical case studies. *Int J Mol Sci*. <https://doi.org/10.3390/ijms18112316>
  22. Tong SY, Giffard PM (2012) Microbiological applications of high-resolution melting analysis. *J Clin Microbiol* 50(11):3418–3421
  23. Er T-K, Chang J-G (2012) High-resolution melting: applications in genetic disorders. *Clin Chim Acta* 414:197–201
  24. Erali M, Voelkerding KV, Wittwer CT (2008) High resolution melting applications for clinical laboratory medicine. *Exp Mol Pathol* 85(1):50–58
  25. Montgomery JL, Sanford LN, Wittwer CT (2010) High-resolution DNA melting analysis in clinical research and diagnostics. *Expert Rev Mol Diagn* 10(2):219–240. <https://doi.org/10.1586/erm.09.84>
  26. Gentry R (1963) Cultivation of infectious laryngotracheitis virus in embryonated chicken eggs by various routes of inoculation. *Avian Dis* 7(1):31–37
  27. Lukert P (1965) Comparative sensitivities of embryonating chicken's eggs and primary chicken embryo kidney and liver cell cultures to infectious bronchitis virus. *Avian Dis* 9(2):308–316
  28. Dwight Z, Palais R, Wittwer CT (2011) uMELT: prediction of high-resolution melting curves and dynamic melting profiles of PCR products in a rich web application. *Bioinformatics* 27(7):1019–1020
  29. Reja V, Kwok A, Stone G, Yang L, Missel A, Menzel C, Bassam B (2010) ScreenClust: Advanced statistical software for supervised and unsupervised high resolution melting (HRM) analysis. *Methods* 50(4):S10–S14
  30. Grando D, Said MM, Mayall BC, Gurtler V (2012) High resolution melt analysis to track infections due to ribotype 027 *Clostridium difficile*. *J Microbiol Methods* 89(2):87–94
  31. Gurtler V, Grando D, Mayall BC, Wang J, Ghaly-Derias S (2012) A novel method for simultaneous *Enterococcus* species identification/genotyping and van genotyping by high resolution melt analysis. *J Microbiol Methods* 90(3):167–181
  32. Naze F, Desvars A, Picardeau M, Bourhy P, Michault A (2015) Use of a new high resolution melting method for genotyping pathogenic *Leptospira* spp. *PLoS One* 10(7):e0127430
  33. Roth AL, Hanson ND (2013) Rapid detection and statistical differentiation of KPC gene variants in Gram-negative pathogens by use of high-resolution melting and ScreenClust analyses. *J Clin Microbiol* 51(1):61–65
  34. Winterhagen P, Wunsche JN (2016) Single nucleotide polymorphism analysis reveals heterogeneity within a seedling tree population of a polyembryonic mango cultivar. *Genome* 59(5):319–325. <https://doi.org/10.1139/gen-2015-0216>
  35. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C (2012) Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28(12):1647–1649
  36. Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, Rozen SG (2012) Primer3—new capabilities and interfaces. *Nucleic Acids Res* 40(15):e115–e115
  37. Spatz SJ, Volkening JD, Keeler CL, Kutish GF, Riblet SM, Boettger CM, Clark KF, Zsak L, Afonso CL, Mundt ES, Rock DL, Garcia M (2012) Comparative full genome analysis of four infectious laryngotracheitis virus (Gallid herpesvirus-1) virulent isolates from the United States. *Virus Genes* 44(2):273–285. <https://doi.org/10.1007/s11262-011-0696-3>
  38. Chandra YG, Lee J, Kong BW (2012) Genome sequence comparison of two United States live attenuated vaccines of infectious laryngotracheitis virus (ILTV). *Virus Genes* 44(3):470–474. <https://doi.org/10.1007/s11262-012-0728-7>
  39. Garcia M, Volkening J, Riblet S, Spatz S (2013) Genomic sequence analysis of the United States infectious laryngotracheitis vaccine strains chicken embryo origin (CEO) and tissue culture origin (TCO). *Virology* 440(1):64–74. <https://doi.org/10.1016/j.virol.2013.02.007>
  40. Callison S, Riblet S, Oldoni I, Sun S, Zavala G, Williams S, Resurreccion R, Spackman E, Garcia M (2007) Development and validation of a real-time Taqman® PCR assay for the detection and quantitation of infectious laryngotracheitis virus in poultry. *J Virol Methods* 139(1):31–38
  41. Shil NK, Legione AR, Markham PF, Noormohammadi AH, Devlin JM (2014) Development and validation of TaqMan real-time polymerase chain reaction assays for the quantitative and differential detection of wild-type infectious Laryngotracheitis Viruses from a glycoprotein G-deficient candidate vaccine strain. *Avian Dis* 59(1):7–13
  42. Creelan JL, Calvert VM, Graham DA, McCullough SJ (2006) Rapid detection and characterization from field cases of infectious laryngotracheitis virus by real-time polymerase chain reaction and restriction fragment length polymorphism. *Avian Pathol* 35(02):173–179
  43. Han MG, Kim SJ (2001) Analysis of Korean strains of infectious laryngotracheitis virus by nucleotide sequences and restriction fragment length polymorphism. *Vet Microbiol* 83(4):321–331
  44. Ojkic D, Swinton J, Vallieres M, Martin E, Shapiro J, Sanei B, Binnington B (2006) Characterization of field isolates of infectious laryngotracheitis virus from Ontario. *Avian Pathol* 35(4):286–292
  45. Shehata AA, Halami MY, Sultan HH, El-Razik AGA, Vahlenkamp TW (2013) Chicken embryo origin-like strains are responsible for infectious laryngotracheitis virus outbreaks in Egyptian cross-bred broiler chickens. *Virus Genes* 46(3):423–430

46. Zhu L, Konsak BM, Olaogun OM, Agnew-Crumpton R, Kanci A, Marenda MS, Browning GF, Noormohammadi AH (2017) Identification of a new genetic marker in *Mycoplasma synoviae* vaccine strain MS-H and development of a strategy using polymerase chain reaction and high-resolution melting curve analysis for differentiating MS-H from field strains. *Vet Microbiol* 210:49–55. <https://doi.org/10.1016/j.vetmic.2017.08.021>
47. Shahid MA, Markham PF, Marenda MS, Agnew-Crumpton R, Noormohammadi AH (2014) High-resolution melting-curve analysis of *obg* gene to differentiate the temperature-sensitive *Mycoplasma synoviae* vaccine strain MS-H from non-temperature-sensitive strains. *PLoS One* 9(3):e92215. <https://doi.org/10.1371/journal.pone.0092215>
48. Robertson T, Bibby S, O'Rourke D, Belfiore T, Agnew-Crumpton R, Noormohammadi A (2010) Identification of chlamydial species in crocodiles and chickens by PCR-HRM curve analysis. *Vet Microbiol* 145(3–4):373–379
49. Patterson JL, Lynch M, Anderson GA, Noormohammadi AH, Legione A, Gilkerson JR, Devlin JM (2015) The prevalence and clinical significance of *Chlamydia* infection in island and mainland populations of Victorian koalas (*Phascolarctos cinereus*). *J Wildl Dis* 51(2):309–317
50. Ghorashi SA, Kanci A, Noormohammadi AH (2015) Evaluation of the capacity of PCR and high-resolution melt curve analysis for identification of mixed infection with *Mycoplasma gallisepticum* Strains. *PLoS One* 10(5):e0126824