



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Greenwood, CJ;Letcher, P;Hutchinson, DM;Macdonald, JA;Grigg, J;Boden, JM;Toumbourou, JW;Olsson, CA

Title:

Pre-pandemic predictors of parental substance use during COVID-19

Date:

2025-01-01

Citation:

Greenwood, C. J., Letcher, P., Hutchinson, D. M., Macdonald, J. A., Grigg, J., Boden, J. M., Toumbourou, J. W. & Olsson, C. A. (2025). Pre-pandemic predictors of parental substance use during COVID-19. *Frontiers in Child and Adolescent Psychiatry*, 4, pp.1587146-. <https://doi.org/10.3389/frcha.2025.1587146>.

Persistent Link:

<https://hdl.handle.net/11343/367703>

License:

[CC BY](#)



OPEN ACCESS

EDITED BY

Tjhin Wiguna,
University of Indonesia, Indonesia

REVIEWED BY

Antonella Pomè,
Heinrich Heine University of Düsseldorf,
Germany
Prafulla Shriyan,
Public Health Foundation of India, India

*CORRESPONDENCE

Christopher J. Greenwood
✉ christopher.greenwood@deakin.edu.au

RECEIVED 04 March 2025

ACCEPTED 08 September 2025

PUBLISHED 07 October 2025

CITATION

Greenwood CJ, Letcher P, Hutchinson DM,
Macdonald JA, Grigg J, Boden JM,
Toumbourou JW and Olsson CA (2025) Pre-
pandemic predictors of parental substance
use during COVID-19.
Front. Child Adolesc. Psychiatry 4:1587146.
doi: 10.3389/frcha.2025.1587146

COPYRIGHT

© 2025 Greenwood, Letcher, Hutchinson,
Macdonald, Grigg, Boden, Toumbourou and
Olsson. This is an open-access article
distributed under the terms of the [Creative
Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The
use, distribution or reproduction in other
forums is permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original publication in
this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Pre-pandemic predictors of parental substance use during COVID-19

Christopher J. Greenwood^{1,2,3*}, Primrose Letcher^{1,2,3},
Delyse M. Hutchinson^{1,2,3,4}, Jacqui A. Macdonald^{1,2,3},
Jasmin Grigg^{5,6}, Joseph M. Boden⁷, John W. Toumbourou^{1,2} and
Craig A. Olsson^{1,2,3}

¹SEED Lifespan Strategic Research Centre, Faculty of Health, School of Psychology, Deakin University, Geelong, VIC, Australia, ²Centre for Adolescent Health, Murdoch Children's Research Institute, Melbourne, VIC, Australia, ³Department of Paediatrics, Royal Children's Hospital, University of Melbourne, Melbourne, VIC, Australia, ⁴National Drug and Alcohol Research Centre, Faculty of Medicine, University of New South Wales, Sydney, NSW, Australia, ⁵Turning Point, Eastern Health, Melbourne, VIC, Australia, ⁶Monash Addiction Research Centre, Eastern Health Clinical School, Monash University, Melbourne, VIC, Australia, ⁷Department of Psychological Medicine, University of Otago Christchurch, Christchurch, New Zealand

Aims: To examine pre-pandemic predictors of parent substance use during COVID-19 in Australia, where some of the longest periods of public health restrictions in the world occurred.

Methods: We used data from the Australian Temperament Project Generation 3 Study on 560 parents (59% female) who completed COVID-19 specific surveys (2020/2021), including assessment of alcohol, tobacco, and illicit substance use. Comprehensive pre-pandemic assessments were conducted during the postpartum period when offspring turned 1-year of age (2012–2018), including 33 indicators spanning parent and child factors (individual, relational, and contextual).

Results: During the pandemic, 39% of parents reported drinking alcohol 3-to-4 days per week or more, 12% used tobacco, and 6% used illicit substances. In Least Absolute Shrinkage and Selection Operator (LASSO) logistic regression models, a variety of pre-pandemic predictors [k] were identified across alcohol (AUC = 0.72, k = 2; OR = 0.92–2.03), tobacco (AUC = 0.96, k = 10; OR = 0.61–4.21), and illicit substance use (AUC = 0.78, k = 2; OR = 1.44–1.60). The strongest predictors were pre-pandemic use of the same substance (OR = 1.60–4.21). While few other predictors were identified for alcohol and illicit substance use, several family characteristics predicted tobacco use.

Conclusions: Results indicated that parents engaging in alcohol, tobacco and illicit drug use in our cohort reported strong continuity of use from before, to during the COVID-19 pandemic in Australia. This highlights the importance of public health initiatives that provide accessible substance use support and treatment options to parents during periods of public health emergencies. Further, enriched population interventions targeted across socioeconomic and family contexts may be important in identifying risk, particularly for tobacco use.

KEYWORDS

families, substance use, COVID-19, machine learning, cohort study

1 Introduction

The COVID-19 pandemic has had a significant impact on population health and wellbeing worldwide. In Australia, strict social distancing and isolation measures were implemented across 2020–2021 to slow the rate of infection (1). While these measures achieved their direct aim of containing virus transmission, they also provoked a cascade of population-wide outcomes including changes to patterns of alcohol, tobacco, and illicit substance use. For example, during the COVID-19 period in Australia, there was an increase in alcohol-induced deaths, coupled with a decrease in drug-induced deaths (2). More broadly, patterns of substance use varied across populations, pointing to considerable heterogeneity in risk for substance use (3, 4) and highlighting the “need to recognise the existence of disadvantage in our response to crises” (5).

One of the hardest hit sectors of society were families with young children, where pandemic-related restrictions radically reduced access to childcare and education services, and introduced new and often complex challenges balancing working from home, with childcare and/or home-schooling demands (6, 7). In this context, potential changes in substance use in the family environment during COVID-19 have received growing attention because of the unique risks that substance use can pose to parents and children (8, 9) including substance-induced intimate partner violence and child maltreatment (10, 11).

Systematic reviews on the COVID-19 context have identified parenting and caregiving responsibilities to be associated with increased substance use, although evidence has primarily focused on alcohol use (3, 4). Further, findings from one Australian sample of parents identified the importance of demographic (e.g., gender, single parent household) and individual (e.g., chronic physical conditions) factors in parental tobacco and alcohol use (12, 13). Nevertheless, further examination of a broad range of factors related to parental substance use during the pandemic, including substances other than alcohol and tobacco, is warranted.

Further to this, to accurately identify parents at risk in future public health emergencies (e.g., disasters and disease outbreaks), comprehensive studies of risks occurring prior to such emergencies, are needed, spanning individual, relational, and contextual domains, relevant to both the parent and child(ren). Given there is likely a large set of potential predictors of risk, one method to identifying the most salient risks for a given outcome is to utilise machine learning approaches. Machine learning methods can also be used to maximise prediction (as opposed to understanding causal explanations). Least Absolute Shrinkage and Selection Operator (LASSO) regression models (14) are one such approach which extend upon traditional regression models. Notably, such models (and their predictor coefficients) are not interpreted causally, but rather with the aim of identifying which parents are likely to be at risk in public health emergencies.

In this study we aimed to examine the extent to which pre-pandemic predictors (parent and child, individual, relational, contextual), that were prospectively measured when offspring were 1-year old (2012–2018), were associated with parent

substance use (alcohol, tobacco, and illicit substances) during COVID-19 (mid-2020 and late-2021). We employed machine learning methods, specifically LASSO (Least Absolute Shrinkage and Selection Operator) penalised regression models, to identify the most predictive subset of indicators.

2 Methods

2.1 Participants

Participants were from the Australian Temperament Project (ATP), a 16-wave longitudinal study tracking the psychosocial development of young people from infancy to adulthood. In 1983, 67 Local Government Areas (LGAs) in the state of Victoria were randomly selected based on advice from the Australian Bureau of Statistics, to provide a representative sample of Victorian LGAs (15). Parents (Generation 1; G1) of every 4–8-month-old infant (Generation 2; G2) who visited one of the selected infant welfare centres between 22nd April and 6th May (1983) were invited to participate. Approximately 3,000 questionnaires were administered. The baseline sample consisted of 2,443 infants aged 4–8 months. Subsequently, families have participated via mail surveys approximately every 2 years until children were 19–20 years of age, and every 4 years thereafter (16).

The ATP Generation 3 (ATPG3) commenced in 2012, with recruitment of infant offspring (Generation 3; G3) of G2 participants and their partners. Identification of pregnancies occurred via participant contact every six months between 2012 and 2018 when participants were aged 29–35 years, representing the peak period of first births in Australia. Telephone or web-based surveys were conducted with parents at the third trimester of pregnancy, and at 8 weeks and 1-year postpartum, for which the 1-year postpartum data is used in the current study. Further information on sampling and procedures is available elsewhere (17) and a flowchart of the ATPG3 sample recruitment is presented in the supplementary material (17; [Supplementary Figure S1](#)). ATP Generation 3 Study protocols have been approved by the Royal Children’s Hospital Human Research Ethics Committee (34185). Prior ATP waves were approved by the relevant committee at the time.

G2 participants in the ATPG3 study with one or more children were invited to complete a brief online survey about the impacts of the COVID-19 pandemic in: (1) 2020 between May and September, during the height of COVID-19 lockdowns ($n = 515$ parents; 60% female; 73% of the existing ATPG3 parent sample); and (2) 2021 between October and December ($n = 488$ parents; 59% female; 69% of existing ATPG3 parent sample). To assess bias due to attrition, ATPG3 participants were compared on characteristics collected at baseline (1983, at 4–8 months), including G2 sex, difficult temperament, and behaviour problems, as well as G1 education and country of birth. Compared to the full ATP sample, those who were screened for the ATPG3 study had marginally lower rates of G1s born overseas and G1s with less education. Those who participated in either of the COVID-19 surveys were representative of the ATPG3 sample. A total of 574

TABLE 1 Descriptive statistics for postpartum (1-year) predictor variables.

Measures	n	%	M	SD	95% CI	n
Infant (G3) characteristics						
Individual						
Female	295	53%			(49, 57)	560
Birth weight			3.4	0.5	(3.4, 3.5)	549
Gestational age			39.1	1.7	(39.0, 39.2)	552
Number of siblings			1.0	0.9	(0.9, 1.1)	560
Health			3.7	0.6	(3.6, 3.7)	552
Behaviour problems			7.3	4.3	(6.9, 7.7)	550
Behaviour competencies			15.8	2.8	(15.5, 16.0)	550
Approach			13.7	4.6	(13.3, 14.1)	549
Cooperation			18.7	4.3	(18.4, 19.1)	549
Persistence			13.1	3.5	(12.8, 13.4)	550
Rhythmicity			9.4	2.9	(9.2, 9.6)	551
Distractibility			16.6	2.6	(16.4, 16.8)	549
Reactivity			29.3	4.5	(28.9, 29.7)	550
Relational						
Family care hours			4.6	10.3	(3.7, 5.5)	552
Non-family care hours			4.8	9.8	(4.0, 5.6)	552
Parent (G2) characteristics						
Individual						
Female	330	59%			(55, 63)	560
Depression			1.2	1.9	(1.0, 1.4)	490
Anxiety			0.7	1.5	(0.6, 0.9)	490
Stress			3.6	3.1	(3.3, 3.8)	490
Alcohol use (≥ 3 –4 days per week)	98	20%			(17, 24)	489
Tobacco use (≥ 1 –2 day per week)	61	13%			(10, 16)	488
Illicit use (≥ 1 –2 day per week)	24	5%			(3, 7)	489
Relational						
Parenting self-efficacy			3.2	0.8	(3.1, 3.3)	489
Parental warmth			4.4	0.5	(4.3, 4.4)	490
Parental hostility			1.8	0.5	(1.8, 1.9)	490
Parental anxiety			3.0	0.6	(3.0, 3.1)	490
Social support			4.4	0.5	(4.4, 4.4)	366
Maternal-infant bonding [†]			80.9	6.4	(80.2, 81.6)	329
Paternal-infant bonding [†]			76.2	7.6	(75.0, 77.4)	157
Contextual						
Stressful life events			0.4	0.7	(0.3, 0.4)	491
Married/de facto	469	86%			(83, 89)	543
Paid employment	447	80%			(76, 83)	560
Education			5.5	1.9	(5.3, 5.6)	478
Financial stress			0.9	0.9	(0.8, 1.0)	549

[†]Scored separately for mothers and fathers.

parents completed either COVID-19 assessment (59% female, 82% of the existing ATPG3 parent sample). To be included in the current sample, participants needed to have further provided data at the 1-year postpartum survey. The resulting sample included 560 G2 parents (59% female).

2.2 Measures

2.2.1 Substance use during the COVID-19 pandemic

Frequency of alcohol (item: “Alcohol”), tobacco (item: “Cigarettes or other tobacco?”), cannabis [item: “Cannabis (e.g.,

marijuana joint, blunt, pipe, bong)”, and other illicit substances (item: “Other drugs including opiates, heroin, or narcotics cocaine, crack, amphetamine, methamphetamine, hallucinogens, or ecstasy”) in the past two weeks were assessed at both COVID-19 waves. Response options were: 1 = “Not at all”, 2 = “1 or 2 days a week”, 3 = “3 or 4 days a week”, 4 = “Almost every day”. Responses to cannabis and other illicit substances were combined to indicate any illicit use. Binary variables were derived to indicate potentially harmful levels of use. For alcohol, derived binary variables indicated drinking around half the week or more (i.e., reporting at least “3 or 4 days a week”). For tobacco and illicit substance use derived binary variables indicated any use (i.e., reporting at least “1 or 2 days a week”). Given the relatively low prevalence of substance use (particularly tobacco and illicit substance use) and consistent pattern at each COVID-19 timepoint [alcohol: 32% (t1), 29% (t2); tobacco: 7% (t3), 8% (t2); illicit substance use: 3% (t1), 2% (t2)], variables were combined across COVID-19 timepoints.

2.2.2 Pre-COVID-19 postpartum (1-year) predictors

A summary of all pre-COVID-19 postpartum (1-year) predictor variables is presented in Table 1 (see Supplementary Table S1 for further details). Measures included: G3 individual factors (i.e., sex, birth weight, gestational age, number of siblings, health, behaviour problems and competencies, temperament) and relational factors (i.e., hours in care with family members and non-family members); G2 individual factors (i.e., sex, mental health, substance use), relational factors (i.e., parenting style, social support, parent-infant bonding), and contextual factors (i.e., stressful life events, marital status, employment status, education level, financial situation). G2 participant responses were used for all measures, however, for measures related to the G3 and those reflecting the G2 context, partner responses supplemented missing participant responses.

2.3 Statistical analysis

All analyses were conducted in Stata 17 (18). Missing data within the analysis sample ranged from 0% to 35%. Multiple imputation was used to handle missing data. Twenty complete datasets were imputed, based on a multivariate normal model (19). Binary variables were imputed as continuous variables and then back transformed with adaptive rounding following imputation (20). This approach to imputation has been shown to perform similarly to imputation models using chained equations (19). Estimates were obtained by pooling results across the 20 imputed datasets using Rubin’s rules (21).

First, descriptive statistics (mean/SD, *n*%) were provided for all variables included in the current study using the non-imputed data. Next, predictors [*k* = number of predictors] were identified using Least Absolute Shrinkage and Selection Operator (LASSO) penalised logistic regression models, for which each outcome was regressed onto the 33 predictors, in multivariable models. In comparison to traditional regression methods, the LASSO models

shrink the size of coefficients by applying a penalty factor and, in turn, retain the most important predictor variables (i.e., coefficients greater than zero) (14). Ten-fold cross-validation was used to select the penalty strength which minimises the cross-validation function (i.e., to identify the best fitting model). Predictive performance was assessed using the area under the Receiver Operating Characteristic (ROC) Curve (AUC; ≥ 0.7 is acceptable, ≥ 0.8 is excellent, and ≥ 0.9 is outstanding) (22). Given that the LASSO model has not been embedded within Stata's imputation commands, models were run on each imputed dataset and then coefficients were averaged if a predictor was retained in at least 80% of imputed datasets (with AUC values averaged across all datasets). Traditional logistic regressions models were used to estimate the association between each pre-COVID-19 postpartum predictor and substance use during the COVID-19 pandemic (adjusting for the time since the 1-year survey). Of note, while confidence intervals are estimated from traditional regression models, no confidence intervals are estimated as part of the LASSO model – these models aim to maximise prediction, not quantify the uncertainty of individual parameter estimates. All models included adjustment for time since the 1-year survey and the start of the COVID-19 surveys (12 May 2020) to account for variation in the timing between assessments. All continuous variables were z-scored prior to analysis.

3 Results

3.1 Descriptive information

During the COVID-19 pandemic, 39% (95% CI = 35, 43) of parents reported using alcohol at least 3–4 days per week [79% (95% CI = 76, 83) used at least 1 day per week], 12% (95% CI = 9, 15) reported using any tobacco, and 6% (95% CI = 4, 8) reported use any illicit substance use. Patterns of use were similar or somewhat higher in males [alcohol: 46% (95% CI = 39, 52); tobacco: 14% (95% CI = 10, 19); illicit substance use: 7% (95% CI = 4, 11)] relative to females [alcohol: 34% (95% CI = 29, 39); tobacco: 10% (95% CI = 7, 14); illicit substance use: 6% (95% CI = 4, 9)] participants. A descriptive summary of all postpartum predictor variables is presented in [Supplementary Table S1](#).

3.2 Postpartum predictors of substance use during the COVID-19 pandemic

Associations between pre-pandemic, postpartum predictors and substance use during the COVID-19 pandemic are presented in [Table 2](#) for LASSO and traditional logistic regression models.

For alcohol use during the pandemic, LASSO models (AUC=0.72) identified two predictors (k=2): prior alcohol use (OR=2.03) and parent-infant bonding (OR=0.92). These associations were supported by evidence from traditional regression models: prior alcohol use (OR=7.97, 95% CI = 4.77, 13.32) and parent-infant bonding (OR = 0.77, 95% CI = 0.64, 0.93).

For tobacco use during the pandemic, LASSO models (AUC=0.96) identified ten (k=10) predictors including G3 behaviour competencies (OR = 1.73), distractibility (OR = 0.87), reactivity (OR = 1.29), and hours at in family (OR = 0.85) and non-family care (OR = 0.88), and G2 stress (OR = 0.81), tobacco use (OR = 4.21), marital status (OR = 0.61), employment (OR = 0.86), and education (OR = 0.78). Five of these predictors were also evidenced in traditional regression models: behaviour competencies (OR = 1.49, 95% CI = 1.11, 1.98), tobacco use (OR = 69.16, 95% CI = 29.61, 161.55), marital status (OR = 0.29, 95% CI = 0.15, 0.54), paid employment (OR = 0.48, 95% CI = 0.27, 0.85), and education level (OR = 0.56, 95% CI = 0.42, 0.75).

For illicit substance use during the pandemic, LASSO models (AUC=0.78) identified two predictors (k=2): postpartum tobacco use (OR = 1.44) and illicit substance use (OR = 1.60). These were also identified in traditional regression models: tobacco use (OR = 8.22, 95% CI = 3.63, 18.60) and illicit substance use (OR = 18.97, 95% CI = 7.41, 48.57).

4 Discussion

Here we draw on prospective data to examine pre-pandemic predictors of parent substance use during the COVID-19 pandemic in Australia, which experienced some of the longest periods of public health restrictions in the world throughout 2020 and 2021. Most prominently, pre-pandemic substance use was identified as the strongest predictor for all substance types during COVID-19 (alcohol, tobacco and illicit substance use); a finding which is consistent with prior literature that has identified strong continuities in substance use behaviour into parenthood (23).

Further, pre-pandemic parent-infant bonding and tobacco use were identified in LASSO models as the only other predictors of parental alcohol and illicit drug use, respectively, during the pandemic. The association with parent-infant bonding is consistent with previously observed patterns of poorer quality parent-infant bonding among parents who report substance use (24). One explanation is that infants of parents with stronger bonds have better behavioural and developmental outcomes (25), which is linked to reduced parenting stress (26) and lower likelihood of parental alcohol consumption. Further to this, we note that the association between pre-pandemic tobacco use and COVID-19 illicit drug use is in line with the high prevalence of tobacco co-use identified among people using illicit substances (27).

A more complex predictive model emerged for tobacco use during the pandemic, with family factors spanning both child (individual and relational) and parent (individual and contextual) characteristics emerging as important predictors of tobacco use. Associations were consistent across several contextual factors (e.g., marital status, employment, and education level). Current findings support previous work identifying socioeconomic disadvantage as a key predictor of tobacco use (28) and reflect patterns of risk identified in parents assessed during the pandemic (12). While population-wide strategies such as media campaigns, tax increases, and smoke-free legislation have been effective in reducing smoking across socioeconomic groups (29), current findings suggest that

TABLE 2 Penalised and traditional logistic regression models examining relationships between COVID-19 substance use and pre-COVID-19 postpartum (1-year) predictors (N = 560).

Measures	Alcohol use			Tobacco use			Illicit use		
	OR _L	OR	95% CI	OR _L	OR	95% CI	OR _L	OR	95% CI
Infant (G3) characteristics									
Individual									
Female	–	0.73	(0.52, 1.03)	–	1.05	(0.62, 1.78)	–	1.31	(0.64, 2.65)
Birth weight	–	0.99	(0.83, 1.18)	–	0.70	(0.54, 0.91)	–	0.85	(0.61, 1.20)
Gestational age	–	1.06	(0.90, 1.27)	–	0.83	(0.65, 1.07)	–	1.20	(0.83, 1.75)
Number of siblings	–	0.83	(0.70, 0.99)	–	0.98	(0.75, 1.28)	–	0.97	(0.69, 1.38)
Health	–	0.99	(0.84, 1.18)	–	0.97	(0.74, 1.26)	–	1.15	(0.78, 1.69)
Behaviour problems	–	1.09	(0.92, 1.30)	–	1.14	(0.88, 1.48)	–	1.20	(0.86, 1.67)
Behaviour competencies	–	0.89	(0.75, 1.06)	1.73	1.49	(1.11, 1.98)	–	1.28	(0.89, 1.85)
Approach	–	0.91	(0.76, 1.08)	–	0.78	(0.59, 1.03)	–	1.06	(0.75, 1.50)
Cooperation	–	1.14	(0.96, 1.36)	–	1.00	(0.77, 1.31)	–	0.95	(0.67, 1.35)
Persistence	–	1.06	(0.90, 1.26)	–	0.87	(0.66, 1.14)	–	0.84	(0.58, 1.20)
Rhythmicity	–	0.96	(0.81, 1.14)	–	1.05	(0.81, 1.36)	–	1.10	(0.78, 1.54)
Distractibility	–	0.96	(0.80, 1.14)	0.87	0.78	(0.60, 1.02)	–	0.92	(0.65, 1.31)
Reactivity	–	0.96	(0.81, 1.14)	1.29	1.21	(0.92, 1.58)	–	1.07	(0.76, 1.51)
Relational									
Family care hours	–	0.89	(0.74, 1.08)	0.85	0.72	(0.51, 1.03)	–	0.78	(0.49, 1.25)
Non-family care hours	–	1.02	(0.86, 1.21)	0.88	0.93	(0.70, 1.24)	–	1.08	(0.78, 1.49)
Parent (G2) characteristics									
Individual									
Female	–	0.61	(0.43, 0.86)	–	0.60	(0.35, 1.02)	–	0.83	(0.41, 1.67)
Depression	–	0.93	(0.77, 1.13)	–	1.03	(0.79, 1.33)	–	1.22	(0.91, 1.65)
Anxiety	–	1.07	(0.88, 1.30)	–	0.93	(0.70, 1.24)	–	1.00	(0.70, 1.45)
Stress	–	1.13	(0.94, 1.35)	0.81	0.75	(0.55, 1.03)	–	1.01	(0.69, 1.46)
Alcohol use (≥3/4days/week)	2.03	7.97	(4.77, 13.32)	–	1.52	(0.76, 3.06)	–	1.93	(0.86, 4.31)
Tobacco use (any)	–	1.29	(0.76, 2.19)	4.21	69.16	(29.61, 161.55)	1.44	8.22	(3.63, 18.60)
Illicit use (any)	–	2.41	(1.05, 5.51)	–	5.79	(2.15, 15.59)	1.60	18.97	(7.41, 48.57)
Relational									
Parenting self-efficacy	–	1.05	(0.87, 1.27)	–	0.97	(0.71, 1.31)	–	1.17	(0.79, 1.73)
Parental warmth	–	0.88	(0.73, 1.05)	–	1.19	(0.87, 1.62)	–	0.77	(0.55, 1.09)
Parental hostility	–	1.15	(0.96, 1.38)	–	0.93	(0.70, 1.25)	–	1.19	(0.83, 1.72)
Parental anxiety	–	1.07	(0.89, 1.29)	–	1.01	(0.75, 1.35)	–	0.87	(0.59, 1.28)
Social support	–	0.97	(0.77, 1.24)	–	1.06	(0.74, 1.52)	–	1.04	(0.67, 1.61)
Parent-infant bonding	0.92	0.77	(0.64, 0.93)	–	1.23	(0.88, 1.71)	–	0.83	(0.58, 1.20)
Contextual									
Stressful life events	–	0.85	(0.71, 1.03)	–	1.04	(0.79, 1.39)	–	1.18	(0.85, 1.63)
Married/de facto	–	0.76	(0.46, 1.25)	0.61	0.29	(0.15, 0.54)	–	0.44	(0.19, 0.98)
Paid employment	–	1.20	(0.78, 1.85)	0.86	0.48	(0.27, 0.85)	–	0.64	(0.29, 1.40)
Education	–	1.05	(0.88, 1.27)	0.78	0.56	(0.42, 0.75)	–	0.92	(0.63, 1.36)
Financial stress	–	1.07	(0.90, 1.27)	–	1.36	(1.06, 1.73)	–	1.02	(0.73, 1.43)

OR_L, odds ratio from LASSO model; traditional regression coefficients at $p < 0.05$ bolded.

parents impacted by greater disadvantage would benefit from more targeted support around smoking cessation (e.g., social media targeting based on geographic location).

Further, a range of family characteristics, spanning child and parent factors, were identified in risk profiles for tobacco use during the pandemic. Need a summary sentence here. Interestingly, while the direction of the reactivity finding supports prior risk associations between parental smoking and negative child outcomes (30), several associations were in unexpected directions (e.g., a risk association for behaviour competencies and a protective association for parent stress).

Given the non-causal modelling approach in the current study (i.e., predictive LASSO models) further research is required to clarify these relationships. The current findings also suggest that the identification of predictors of tobacco use behaviours requires a multifaceted approach.

4.1 Implications

The strong prediction of pandemic substance use by pre-pandemic patterns of substance use highlights the need for

treatment services and early intervention to prevent use from becoming entrenched. The COVID-19 pandemic has profoundly changed the way in which people access healthcare. In Australia, despite high rates of substance use (31), rates of treatment uptake remain low and delays in treatment-seeking are substantial (32, 33). Telehealth provides a promising strategy to increase earlier engagement with alcohol and substance use treatment, by providing a format of treatment delivery that offers a sense of privacy and anonymity, which can help to overcome treatment barriers related to fear of shame and stigma, as well as structural barriers such as constraints related to travel time and need for childcare. Such models of treatment may be filling a gap in service provision for women who often perform significant childcare duties and who may be less able to access traditional in-person services due to childcare responsibilities (34, 35). Online treatment/app interventions could be another option.

For parents of young families, as well as those planning families, routine prenatal visits provide an opportunity for alcohol and drug screening, provision of information on substance use risks and supports, and where suitable, referral and intervention, which may in turn increase engagement with alcohol and other drug support services (as well as reduce risk of harm to infants, e.g., FASD, family violence) (36). However, it also important to recognise that, for many people, prenatal substance use behaviours have a history of continuity from adolescence and into the young adult period, well before conception (23). This suggests that universal monitoring and intervention efforts, extending earlier in the life course, are likely to be beneficial in disrupting future patterns of continuity into the parenting period.

4.2 Limitations

In mature cohort studies there are important sources of selection, measurement, and confounding bias that require consideration. Although multiple imputation was used to minimize missing data bias, it is likely that there was some selective attrition of the most vulnerable individuals, such as those with high levels of substance use. Further, parents who had a child prior to the recruitment period (parents aged 29–35 years) are excluded, unless they also went on to have another child in this period. Additionally, as all measures are self-reported, there is a risk of shared method variance and social desirability bias. To mitigate these issues, one option could be to incorporate data from additional sources, such as family members or peers, or employ approaches like the Randomized Response Technique (37) into future research methods. Further, substance use outcomes were defined based only on frequency, and therefore do not capture aspects of substance use related to quantity or disorder, which may have differential predictive profiles. The strength of associations may also be impacted by the variation in time (i.e., less than one year up to nine years) between completing the 1-year postpartum survey and the COVID-19 survey, although accounted for analytically. Further,

the relatively small sample size may have meant that the study was underpowered to detect weaker associations of interest (that might be expected across long periods of time), as indicated by the imprecise confidence intervals. As such, the pattern of results are likely to be conservative in nature. Finally, while the current aims were not causal in nature, future work could examine causal pathways to determine appropriate intervention opportunities.

5 Conclusions

In our study, the most important predictor of parental substance use during the COVID-19 pandemic was prior substance use, pointing to the importance of enriching support to those with histories of substance use. Given the strong continuity in substance use behaviours, public health policy should consider improved access to substance use support and treatment services, particularly in times of public health emergencies. However, to comprehensively prepare for future public health emergencies, continued efforts are needed to identify and protect against broader social inequalities and vulnerabilities. This was evident in the more complex predictive models that emerged for parental tobacco use. Together, findings suggest that beyond patterns of substance use continuity, approaches to targeting interventions, particularly for tobacco use, likely requires efforts which consider a range of individual, relational and contextual factors.

Data availability statement

The datasets presented in this article are not readily available because Ethics approvals for this study do not permit these potentially re-identifiable participant data to be made publicly available. Enquires about collaboration are possible through our institutional data access protocol: <https://lifecourse.melbournechildrens.com/data-access/>. The current institutional body responsible for ethical approval is The Royal Children's Hospital Human Research Ethics Committee. Requests to access the datasets should be directed to Craig Olsson, craig.olsson@deakin.edu.au.

Ethics statement

The studies involving humans were approved by Royal Children's Hospital Human Research Ethics Committee. The studies were conducted in accordance with the local legislation and institutional requirements. Written informed consent was obtained from the minor(s)' legal guardian/next of kin for the publication of any potentially identifiable images or data included in this article.

Author contributions

CG: Conceptualization, Formal analysis, Writing – original draft, Writing – review & editing. PL: Conceptualization, Funding acquisition, Writing – original draft, Writing – review & editing. DH: Conceptualization, Funding acquisition, Writing – original draft, Writing – review & editing. JM: Conceptualization, Funding acquisition, Writing – original draft, Writing – review & editing. JG: Conceptualization, Writing – original draft, Writing – review & editing. JB: Conceptualization, Writing – original draft, Writing – review & editing. JT: Conceptualization, Funding acquisition, Supervision, Writing – original draft, Writing – review & editing. CO: Conceptualization, Funding acquisition, Supervision, Writing – original draft, Writing – review & editing.

Funding

The author(s) declare that financial support was received for the research and/or publication of this article. Data collection for the ATP study was supported primarily through Australian grants from the Melbourne Royal Children's Hospital Research Foundation, National Health and Medical Research Council, Australian Research Council, and the Australian Institute of Family Studies. Funding for this work was supported by grants from the Australian Research Council [DP130101459; DP160103160; DP180102447; DP200100571] and the National Health and Medical Research Council of Australia [APP1082406]. CAO and DMH were supported by fellowships from the National Health and Medical Research Council (Investigator grants APP1175086 and APP1197488, respectively).

Acknowledgments

The ATP study is located at The Royal Children's Hospital Melbourne and is a collaboration between Deakin University, The University of Melbourne, the Australian Institute of Family Studies, The University of New South Wales, The University of Otago (New Zealand), and the Royal Children's Hospital (further information available at <https://www.melbournechildrens.com/atp/>). The views expressed in this article are those of the authors and may

References

1. Stobart A, Duckett S. Australia's response to COVID-19. *Health Econ Policy Law*. (2022) 17(1):95–106. doi: 10.1017/S1744133121000244
2. Australian Bureau of Statistics. Causes of Death, Australia [Internet]. 2021. Available online at: <https://www.abs.gov.au/statistics/health/causes-death/causes-death-australia/2021> (Accessed January 01, 2023).
3. Roberts A, Rogers J, Mason R, Siriwardena AN, Hogue T, Whitley GA, et al. Alcohol and other substance use during the COVID-19 pandemic: a systematic review. *Drug Alcohol Depend*. (2021) 229:1–35. doi: 10.1016/j.drugalcdep.2021.109150
4. Schmidt RA, Genois R, Jin J, Vigo D, Rehm J, Rush B. The early impact of COVID-19 on the incidence, prevalence, and severity of alcohol use and other drugs:

not reflect those of their organizational affiliations, nor of other collaborating individuals or organizations. We acknowledge all collaborators who have contributed to the ATP, especially Professors Ann Sanson, Margot Prior, Frank Oberklaid, and Dr Diana Smart. We would also like to sincerely thank the participating families for their time and invaluable contribution to the study.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Generative AI statement

The author(s) declare that no Generative AI was used in the creation of this manuscript.

Any alternative text (alt text) provided alongside figures in this article has been generated by Frontiers with the support of artificial intelligence and reasonable efforts have been made to ensure accuracy, including review by the authors wherever possible. If you identify any issues, please contact us.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/frcha.2025.1587146/full#supplementary-material>

a systematic review. *Drug Alcohol Depend*. (2021) 228:1–17. doi: 10.1016/j.drugalcdep.2021.109065

5. Shergold P, Broadbent J, Marshall I, Varghese P. *Fault Lines: An Independent Review in Australia's Response to COVID-19*. Melbourne, VIC: John and Myriam Wylie Foundation, Minderoo Foundation and the Paul Ramsay Foundation (2022).

6. Verlenden J, Pampati S, Rasberry CN, Liddon N, Hertz M, Kilmer G, et al. Association of children's mode of school instruction with child and parent experiences and well-being during the COVID-19 pandemic — COVID experiences survey, United States, October 8–November 13, 2020. *MMWR Recomm Rep*. (2021) 70(11):369–76. doi: 10.15585/mmwr.mm7011a1

7. Power K. The COVID-19 pandemic has increased the care burden of women and families. *Sustain Sci Pract Policy*. (2020) 16(1):67–73. doi: 10.1080/15487733.2020.1776561
8. Hutchinson DM, Mattick RP, Braunstein D, Maloney E, Wilson J. *The Impact of Alcohol use Disorders on Family Life: A Review of the Empirical Literature*. Sydney, NSW: National Drug and Alcohol Research Centre, University of New South Wales (2014).
9. Kuppens S, Moore SC, Gross V, Lowthian E, Siddaway AP. The enduring effects of parental alcohol, tobacco, and drug use on child well-being: a multilevel meta-analysis. *Dev Psychopathol*. (2020) 32(2):765–78. doi: 10.1017/S0954579419000749
10. Choenni V, Hammink A, van de Mheen D. Association between substance use and the perpetration of family violence in industrialized countries: a systematic review. *Trauma Violence Abuse*. (2017) 18:37–50. doi: 10.1177/1524838015589253
11. Laslett AM, Room R, Dietze P, Ferris J. Alcohol's involvement in recurrent child abuse and neglect cases. *Addiction*. (2012) 107(10):1786–93. doi: 10.1111/j.1360-0443.2012.03917.x
12. Westrupp EM, Bennett C, Berkowitz T, Youssef GJ, Toumbourou JW, Tucker R, et al. Child, parent, and family mental health and functioning in Australia during COVID-19: comparison to pre-pandemic data. *Eur Child Adolesc Psychiatry*. (2023) 32(2):317–30. doi: 10.1007/s00787-021-01861-z
13. Greenwood CJ, Fuller-Tyszkiewicz M, Hutchinson DM, Macdonald JA, Bereznicki HGK, Youssef GJ, et al. Alcohol use among Australian parents during the COVID-19 pandemic—April-2020 to May 2021. *Addict Behav*. (2023) 138:1–7. doi: 10.1016/j.addbeh.2022.107561
14. Tibshirani R. Regression shrinkage and selection via the lasso. *J R Stat Soc Ser B Methodol*. (1996) 58(1):267–88. doi: 10.1111/j.2517-6161.1996.tb02080.x
15. Sanson AV, Prior M, Oberklaid F. Normative data on temperament in Australian infants. *Aust J Psychol*. (1985) 37(2):185–95. doi: 10.1080/00049538508256397
16. Vassallo S, Sanson A. *The Australian Temperament Project: The First 30 Years*. Melbourne: Australian Institute of Family Studies (2013).
17. Olsson CA, Letcher P, Greenwood CJ, McIntosh JE, Barker S, Olsson CM, et al. The Australian temperament project generation 3 study: a population-based multigenerational prospective cohort study of socioemotional health and development. *BMJ Open*. (2022) 12(9):e061854. doi: 10.1136/bmjopen-2022-061854
18. StataCorp. *Stata Statistical Software: Release 17*. College Station, TX: StataCorp LLC (2021).
19. Lee KJ, Carlin JB. Multiple imputation for missing data: fully conditional specification versus multivariate normal imputation. *Am J Epidemiol*. (2010) 171(5):624–32. doi: 10.1093/aje/kwp425
20. Bernaards CA, Belin TR, Schafer JL. Robustness of a multivariate normal approximation for imputation of incomplete binary data. *Stat Med*. (2007) 26(6):1368–82. doi: 10.1002/sim.2619
21. Rubin DB. *Multiple Imputation for Nonresponse in Surveys*. Hoboken, NJ: Wiley Series in Probability and Statistics (1987).
22. Hosmer DW, Lemeshow S. *Applied Logistic Regression [Internet]*. 2nd ed. New York, NY: John Wiley & Sons, Inc. (2000). doi: 10.1002/0471722146
23. Thomson KC, Greenwood CJ, Letcher P, Spry EA, Macdonald JA, McAnally HM, et al. Continuities in maternal substance use from early adolescence to parenthood: findings from the intergenerational cohort consortium. *Psychol Med*. (2023) 53(5):2136–45. doi: 10.1017/S0033291721003925
24. Quinlivan JA, Evans SF. Impact of domestic violence and drug abuse in pregnancy on maternal attachment and infant temperament in teenage mothers in the setting of best clinical practice. *Arch Womens Ment Health*. (2005) 8(3):191–9. doi: 10.1007/s00737-005-0079-7
25. le Bas GA, Youssef GJ, Macdonald JA, Rossen L, Teague SJ, Kothe EJ, et al. The role of antenatal and postnatal maternal bonding in infant development: a systematic review and meta-analysis. *Social Development*. (2020) 29:3–20. doi: 10.1111/sode.12392
26. de Cock ESA, Henrichs J, Klimstra TA, Janneke A, Vreeswijk CMJM, Meeus WHJ, et al. Longitudinal associations between parental bonding, parenting stress, and executive functioning in toddlerhood. *J Child Fam Stud*. (2017) 26(6):1723–33. doi: 10.1007/s10826-017-0679-7
27. Australian Institute of Health and Welfare. *National Drug Strategy Household Survey 2022–2023*. Canberra: AIHW Canberra (2024).
28. Hiscock R, Bauld L, Amos A, Fidler JA, Munafo M. Socioeconomic status and smoking: a review. *Ann N Y Acad Sci*. (2012) 1248:107–23. doi: 10.1111/j.1749-6632.2011.06202.x
29. Germain D, Durkin S, Scollo M, Wakefield M. The long-term decline of adult tobacco use in Victoria: changes in smoking initiation and quitting over a quarter of a century of tobacco control. *Aust N Z J Public Health*. (2012) 36(1):17–23. doi: 10.1111/j.1753-6405.2011.00805.x
30. Button TMM, Maughan B, McGuffin P. The relationship of maternal smoking to psychological problems in the offspring. *Early Hum Dev*. (2007) 83(11):727–32. doi: 10.1016/j.earlhumdev.2007.07.006
31. Australian Institute of Health and Welfare. *National Drug Strategy Household Survey 2019*. Canberra: AIHW Canberra (2020).
32. Australian Institute of Health and Welfare. *Alcohol and Other Drug Treatment Services in Australia: 2016–17 [Internet]*. Canberra: AIHW Canberra (2018). Available online at: www.aihw.gov.au
33. Reavley NJ, Cvetkovski S, Jorm AF, Lubman DI. Help-seeking for substance use, anxiety and affective disorders among young people: results from the 2007 Australian national survey of mental health and wellbeing. *Aust N Z J Psychiatry*. (2010) 44(8):729–35. doi: 10.3109/00048671003705458
34. Garde EL, Manning V, Lubman DI. Characteristics of clients currently accessing a national online alcohol and drug counselling service. *Australas Psychiatry*. (2017) 25(3):250–3. doi: 10.1177/1039856216689623
35. Grigg J, Volpe I, Tyler J, Hall K, McPherson B, Lubman DI, et al. Ready2Change: preliminary effectiveness of a telephone-delivered intervention program for alcohol, methamphetamine and cannabis use problems. *Drug Alcohol Rev*. (2022) 41(2):517–27. doi: 10.1111/dar.13363
36. Wright TE, Terplan M, Ondersma SJ, Boyce C, Yonkers K, Chang G, et al. The role of screening, brief intervention, and referral to treatment in the perinatal period. *Am J Obstet Gynecol*. (2016) 215(5):539–47. doi: 10.1016/j.ajog.2016.06.038
37. Blair G, Imai K, Zhou YY. Design and analysis of the randomized response technique. *J Am Stat Assoc*. (2015) 110:1304–19. doi: 10.1080/01621459.2015.1050028