



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Rogerson, SJ

Title:

Management of malaria in pregnancy

Date:

2017-09-01

Citation:

Rogerson, S. J. (2017). Management of malaria in pregnancy. *Indian Journal of Medical Research*, 146 (September), pp.328-333. https://doi.org/10.4103/ijmr.IJMR_1304_17.

Persistent Link:

<https://hdl.handle.net/11343/283710>

License:

[CC BY-NC-SA](#)

Title Management of malaria in pregnancy

Professor Stephen J Rogerson, FRACP
Department of Medicine at the Doherty Institute,
The University of Melbourne,
792 Elizabeth Street,
Melbourne VIC 3000,
Australia

E sroger@unimelb.edu.au

M +61439336694

Abstract

Pregnant women are especially susceptible to malaria infection. Without existing immunity, severe malaria can develop requiring emergency treatment, and pregnancy loss is common. In semi-immune women, consequences of malaria for the mother include anaemia, while stillbirth, premature delivery and fetal growth restriction affect the developing fetus. Preventive measures include insecticide treated nets and (in some African settings) intermittent preventive treatment.

Prompt management of maternal infection is key, using parenteral artemisinins for severe malaria, and artemisinin combination treatments (ACTs) in second and third trimesters of pregnancy. ACTs may soon also be recommended as an alternative to quinine as treatment in first trimester of pregnancy. Monitoring the safety of antimalarials and understanding their pharmacokinetics is particularly important in pregnancy with the altered maternal physiology and the risks to the developing fetus.

As increasing numbers of countries embrace malaria elimination as a goal, the special needs of the vulnerable group of pregnant women and their babies should not be overlooked.

Keywords artemisinin; diagnosis; plasmodium; pregnant; treatment;

Introduction

The global malaria burden has declined in recent years, but over 40% of the world's population remain at risk of infection, and over 400,000 people die ¹. In India over 90% of the population live in malaria transmission areas, with two thirds of infections caused by *Plasmodium falciparum* and one third by *P. vivax* and an estimate of 13 million cases and 24,000 deaths each year ¹.

Pregnant women are particularly susceptible to malaria infection ², and this susceptibility is attributed to immunological changes occurring in pregnancy, and to the unique predilection of a subset of *P. falciparum* parasites to sequester in the maternal blood spaces of the placenta (reviewed in ³). This placental malaria infection helps the parasite avoid clearance by the immune system and especially filtration by the spleen. Placental falciparum parasites express a protein on the red cell surface called VAR2CSA, which sticks to the placental receptor CSA ⁴. Antibodies to VAR2CSA have been associated with protection from placental malaria and adverse birth outcomes ⁵ and a VAR2CSA-based vaccine is now in early clinical trials (NCT02647489 and NCT02658253, clinicaltrials.gov), with the aim of preventing pregnancy malaria.

Placental sequestration is not a major feature of *P. vivax* infection ⁶, but the infection is common in pregnancy and has been associated with adverse pregnancy outcomes including decreased birth weight and maternal anaemia (reviewed in ⁷), although a recent study found relatively low infection prevalence and modest associations with morbidity in five countries including India ⁸.

In high transmission areas, malaria in pregnancy is commonest in first-time mothers and prevalence and densities of parasitaemia both decline over subsequent pregnancies. By contrast in low transmission areas all pregnancies are equally at risk for *P. falciparum*, and probably *P. vivax* infection ⁹.

Consequences of malaria in pregnancy

The consequences vary with transmission intensity. When transmission is high, maternal anaemia is common, and infant low birth weight due to fetal growth restriction and/or premature delivery is frequent ². In low transmission areas, when non-immune pregnant women become infected, malaria infection may become severe and life-threatening, requiring emergency treatment ². Maternal complications include acute lung injury, severe hypoglycaemia, and coma, while pregnancy loss due to miscarriage or stillbirth is also frequent. Malaria may be an under-recognised cause of maternal death ¹⁰.

Diagnosis

Microscopy

Microscopy of stained blood smears remains widely used to monitor prevalence of malaria. For point-of-care testing, Rapid Diagnostic Tests (RDT) are very effective for the diagnosis of symptomatic malaria infection, which tends to be accompanied by high parasitaemia. They are less effective as screening tools, being unable to reliably detect low-density infections which are common ¹¹. New, high-sensitivity RDTs will soon be evaluated in pregnant women.

Polymerase chain reaction

Polymerase chain reaction (PCR) is highly sensitive, with quantitative PCR able to detect very low density malaria infection¹². However, a specialised laboratory with trained staff is required, and assays are relatively time-consuming. LAMP (Loop-mediated isothermal amplification) has similar sensitivity to PCR but is more rapid and robust, and potentially applicable at point of care^{13,14}. Both are presently reserved for research settings.

We have a strong understanding of the relationship between infection detected by microscopy and adverse pregnancy outcomes², but the significance of sub-patent infection detected by PCR but not by microscopy and/or RDT is not yet clear. Sub-patent infection has been associated with lower mean haemoglobin, and with low birth weight in primigravidae and premature delivery in multigravidae¹⁵, but other studies do not find similar associations^{16,17}. A pooled analysis currently in preparation should more clearly determine whether these low-density infections affect pregnancy outcomes.

Placental histology

Histological examination of placental tissue at delivery is a sensitive tool for detection of active or past malaria infection. Past infection is detected as the malaria pigment, haemozoin, most commonly in fibrin deposits. Active infection can be accompanied by leukocyte infiltrates, principally monocytes, termed intervillitis, particularly in first time mothers with little pregnancy-associated malaria immunity. In this group, it is strongly associated with low birth weight and maternal anaemia¹⁸.

Management of malaria in pregnancy

Prevention

The World Health Organization (WHO) recommends a three-pronged strategy for control of malaria in pregnancy in Africa including case management (prompt treatment with highly effective drug), use of insecticide treated nets (ITNs), and intermittent preventive treatment (IPTp), the administration of a full treatment course of an effective antimalarial at regular antenatal visits, usually a month apart.

Intermittent Preventive Treatment in pregnancy

Currently IPTp is only recommended in Africa, using sulphadoxine-pyrimethamine (SP). A meta-analysis revealed that three doses of IPTp were superior to two¹⁹, and a multisite study showed that IPTp remains effective against low birth weight and anaemia, even when resistance to SP is high, which is common in Eastern and Southern Africa²⁰. In recent years, mefloquine and a combination of azithromycin and chloroquine have been evaluated for prevention of malaria in pregnancy. Mefloquine decreased parasite prevalence at delivery, but was poorly tolerated²¹, and the trial of azithromycin and chloroquine was stopped due to futility (i.e. inability to show a clear benefit over the comparator, SP); it, too, was poorly tolerated²².

Recent studies^{23,24} suggested that dihydroartemisinin-piperaquine (DHA-PQ) may be a suitable replacement for SP. It has also been studied in Indonesia as

IPTp or IST, and larger trials presently in progress should provide a definitive answer as to whether it is a suitable replacement for SP in Africa. We presently lack evidence regarding the benefit of IPTp in lower transmission countries (such as India). Challenges for SP IPTp include lack of evidence regarding its effectiveness against *P. vivax*, and low levels of uptake ²⁵.

An alternative to IPTp is intermittent screening and treatment (IST). Using this approach, at each antenatal visit an RDT is performed, and women testing positive receive an effective antimalarial, most commonly an ACT. In part because of the low sensitivity of current RDTs, IST was not more effective than IPTp in preventing malaria and its complications in several studies in Africa (REFS ²⁶), and it has not been endorsed by WHO for wider use (MPAC). A trial of IST has recently concluded in India, while high-sensitivity RDTs as mentioned above hold promise to overcome some of the weaknesses of the IST approach. A recent review summarised trials of IPTp and IST over the last five years ²⁷.

Insecticide treated nets and indoor residual spraying

Evidence primarily from trials in Africa shows that ITNs can prevent malaria and decrease low birth weight and other adverse pregnancy outcomes. There is limited data from Asia, where mosquito vectors and their biting behaviour are different (refs), and where *P. vivax* is prevalent. Nevertheless, ITNs are a relatively cheap and probably effective tool for malaria prevention in pregnancy in India. Indoor residual spraying is used in parts of Africa and Asia, but there is little pregnancy specific data on its impact.

Severe malaria in pregnancy

When a pregnant woman presents with severe malaria, the first priority is to save her life. The recommended treatment for severe malaria at any time in pregnancy is with parenteral artesunate ²⁸. The SEAQUAMAT trial showed that this is superior to parenteral quinine in Asian adults ²⁹. Once the woman recovers, treatment can be continued with appropriate oral medication, as discussed below. Pregnant women are particularly prone to hypoglycaemia, and blood sugar levels should be closely monitored.

Uncomplicated malaria in pregnancy

Falciparum malaria in first trimester

Currently quinine and clindamycin is the recommended treatment for women in the first trimester of pregnancy ²⁸. In many places, clindamycin is unavailable, and quinine monotherapy is prescribed. Side effects of the seven-day quinine regime, such as tinnitus or fullness in the ears, result in poor compliance, and the risk of recrudescence.

Artemisinin combination treatments (ACTs) are a logical alternative to quinine, but establishing their safety in the first trimester has been challenging, especially because in animal studies the drugs show embryo toxicity ³⁰. Pharmacovigilance, the monitoring of drug safety, is particularly important in pregnancy where both mother and developing fetus are at risk. A recent meta-analysis investigated the safety of ACTs in early pregnancy. Compared to quinine treatment, ACTs were not associated with increased risk of miscarriage, stillbirth, or embryo toxicity ³¹.

This analysis has informed the decision of WHO's Malaria Policy Advisory Committee to endorse use of ACTs in first trimester³², although this is not yet recommended in the treatment guidelines²⁸.

Second and third trimester

A recent large trial compared four different ACTs, namely artemether-lumefantrine, amodiaquine-artesunate, mefloquine-artesunate, or DHA-PQ, for treatment of uncomplicated malaria in the second and third trimesters of pregnancy, in 3428 women³³. All drugs were highly effective (95% or more), with artemether-lumefantrine being associated with the fewest adverse events, while DHA-PQ was associated with the greatest post-exposure prophylaxis. This evidence strongly supports the use of ACTs as first line treatment for malaria in second and third trimester²⁸. In Papua, Indonesia both *P. falciparum* and *P. vivax* were resistant to previous first line drugs, and introduction of DHA-PQ was associated with substantial declines in maternal malaria and anaemia, congenital malaria and low birth weight³⁴.

Vivax malaria

Chloroquine remains an effective treatment for vivax malaria in most of the world, although resistance that emerged in Papua New Guinea and Indonesia is spreading. ACTs (apart from artesunate-SP) are also highly effective for treatment of vivax malaria, and parenteral artesunate is highly effective in severe vivax disease²⁸. An increasing number of countries recommend ACTs for treatment of all forms of uncomplicated malaria³⁵.

Primaquine is contraindicated in pregnancy because of the risk of severe haemolysis in G6PD deficient individuals. Pregnant women may require suppressive treatment, usually with chloroquine, until delivery. Small amounts of primaquine enter breast milk, so G6PD testing of the infant is recommended in lactating women before they are prescribed primaquine²⁸.

Pharmacokinetics of antimalarials in pregnancy

Over recent years, studies have investigated the pharmacokinetics of antimalarials in pregnant women, with the aim of determining whether dose adjustment is required. Many of these studies are reviewed in³¹. For artemether-lumefantrine some evidence points towards a need to extend treatment duration³¹ but at present no dose adjustments have been recommended²⁸. There is no clear evidence for dose adjustments for other drugs in pregnancy at this time.

Special groups: HIV-infected women

HIV infection increases the risk of malaria in pregnancy³⁶. HIV infected women who are prescribed cotrimoxazole to decrease opportunistic infections should not receive IPTp (where otherwise recommended) because of the increased risk of severe skin reactions such as Stevens-Johnson syndrome. Mefloquine decreased parasite prevalence in HIV-infected women, but appeared to increase the risk of mother-to-child transmission of HIV³⁷. DHA-PQ is currently being evaluated as an alternative for IPTp. An additional consideration is the risk of drug interactions between antimalarials and antiretrovirals³⁸, which may require dose adjustment or even avoidance of certain drugs. To date, little is

known about the impact of antiretroviral therapy or immune system reconstitution on the interactions between malaria and HIV.

Access

An optimal approach to control and prevention of malaria in pregnancy requires close cooperation between malaria control and reproductive health programmes. Training to improve knowledge of both health service staff and private sector suppliers of antimalarials regarding optimal case management in pregnancy may be required. Diagnostic tools for pregnancy malaria and drugs recommended for treatment of pregnant women need to be available at all levels of the health system.

Future directions

Malaria elimination

Countries across the Asia-Pacific including India have committed to malaria elimination by 2030. There are implications for management of malaria in pregnant women, and for campaigns such as mass drug administration. Currently DHA-PQ is the most widely used drug for MDA, but it does not yet have an established safety record in early pregnancy, so women who are possibly in early stages of pregnancy should be excluded from such campaigns. Where pregnant women are excluded from MDA campaigns, a significant part of the remaining burden may be found in these women ³⁹.

Pregnant women may be a useful sentinel population to monitor malaria changes ⁴⁰. The great majority of women attend ANC at least once, so they are readily accessed, and parasite prevalence can be easily monitored. An analysis suggests these measures correlate well with a more established measure, the parasite prevalence in children under 5 ⁴¹.

Conclusion

Pregnant women are uniquely susceptible to malaria. Optimal malaria prevention varies with transmission; in higher transmission areas ITNs have demonstrated benefits. Whether preventive treatment approaches such as IPTp or IST will have a place outside Africa will depend on results of studies presently in progress. In lower transmission settings, women may lack malaria immunity and are at risk of developing severe, potentially fatal disease or losing their babies to miscarriage or stillbirth; they require immediate diagnosis and treatment. ACTs are recommended in most circumstances, although quinine remains first choice in first trimester of pregnancy. The approach to treatment must be tailored according to pregnancy trimester and clinical severity of malaria.

Acknowledgements:

SJR's work is supported by the National Health and Medical Research Council of Australia. He thanks Siddhartha Mohanty for helpful review of the manuscript.

References

1. World Health Organization. World Malaria Report 2016. Geneva: World Health Organization, 2016.

2. Desai M, ter Kuile FO, Nosten F, McGready R, Asamo K, Brabin B, et al. Epidemiology and burden of malaria in pregnancy. *The Lancet infectious diseases* 2007; **7**(2): 93-104.
3. Rogerson SJ, Desai M, Mayor A, Sicuri E, S.M. T, van Eijk AM. Burden, pathology and costs of malaria in pregnancy: new developments for an old problem. *The Lancet infectious diseases* 2017; **in press**.
4. Salanti A, Dahlback M, Turner L, Nielsen MA, Barfod L, Magistrado P, et al. Evidence for the involvement of VAR2CSA in pregnancy-associated malaria. *J Exp Med* 2004; **200**(9): 1197-203.
5. Ndam NT, Denoed-Ndam L, Doritchamou J, Viwami F, Salanti A, Nielsen MA, et al. Protective Antibodies against Placental Malaria and Poor Outcomes during Pregnancy, Benin. *Emerg Infect Dis* 2015; **21**(5): 813-23.
6. Mayor A, Bardaji A, Felger I, King CL, Cistero P, Dobano C, et al. Placental infection with Plasmodium vivax: a histopathological and molecular study. *The Journal of infectious diseases* 2012; **206**(12): 1904-10.
7. Rijken MJ, McGready R, Boel ME, Poespoprodjo R, Singh N, Syafruddin D, et al. Malaria in pregnancy in the Asia-Pacific region. *The Lancet infectious diseases* 2012; **12**(1): 75-88.
8. Bardaji A, Martinez-Espinosa FE, Arevalo-Herrera M, Padilla N, Kochar S, Ome-Kaius M, et al. Burden and impact of Plasmodium vivax in pregnancy: A multi-centre prospective observational study. *PLoS Negl Trop Dis* 2017; **11**(6): e0005606.
9. Nosten F, Rogerson SJ, Beeson JG, McGready R, Mutabingwa TK, Brabin B. Malaria in pregnancy and the endemicity spectrum: what can we learn? *Trends Parasitol* 2004; **20**(9): 425-32.
10. Menendez C, Romagosa C, Ismail MR, Carrilho C, Saute F, Osman N, et al. An autopsy study of maternal mortality in Mozambique: the contribution of infectious diseases. *PLoS Med* 2008; **5**(2): e44.
11. Umbers AJ, Unger HW, Rosanas-Urgell A, Wangnapi RA, Kattenberg JH, Jally S, et al. Accuracy of an HRP-2/panLDH rapid diagnostic test to detect peripheral and placental Plasmodium falciparum infection in Papua New Guinean women with anaemia or suspected malaria. *Malaria journal* 2015; **14**: 412.
12. Britton S, Cheng Q, McCarthy JS. Novel molecular diagnostic tools for malaria elimination: a review of options from the point of view of high-throughput and applicability in resource limited settings. *Malaria journal* 2016; **15**: 88.
13. Prah M, Jagannathan P, McIntyre TI, Auma A, Farrington L, Wamala S, et al. Timing of in utero malaria exposure influences fetal CD4 T cell regulatory versus effector differentiation. *Malaria journal* 2016; **15**(1): 497.
14. Tegegne B, Getie S, Lemma W, Mohon AN, Pillai DR. Performance of loop-mediated isothermal amplification (LAMP) for the diagnosis of malaria among malaria suspected pregnant women in Northwest Ethiopia. *Malaria journal* 2017; **16**(1): 34.
15. Cottrell G, Moussiliou A, Luty AJ, Cot M, Fievet N, Massougboji A, et al. Submicroscopic Plasmodium falciparum Infections Are Associated With Maternal Anemia, Premature Births, and Low Birth Weight. *Clin Infect Dis* 2015; **60**(10): 1481-8.

16. Taylor SM, Madanitsa M, Thwai KL, Khairallah C, Kalilani-Phiri L, van Eijk AM, et al. Minimal impact by antenatal subpatent *P. falciparum* infections on delivery outcomes in Malawian women: a cohort study. *The Journal of infectious diseases* 2017.
17. Williams JE, Cairns M, Njie F, Laryea Quaye S, Awine T, Oduro A, et al. The Performance of a Rapid Diagnostic Test in Detecting Malaria Infection in Pregnant Women and the Impact of Missed Infections. *Clin Infect Dis* 2016; **62**(7): 837-44.
18. Menendez C, Ordi J, Ismail MR, Ventura PJ, Aponte JJ, Kahigwa E, et al. The impact of placental malaria on gestational age and birth weight. *J Infect Dis* 2000; **181**: 1740-5.
19. Kayentao K, Garner P, van Eijk AM, Naidoo I, Roper C, Mulokozi A, et al. Intermittent preventive therapy for malaria during pregnancy using 2 vs 3 or more doses of sulfadoxine-pyrimethamine and risk of low birth weight in Africa: systematic review and meta-analysis. *JAMA* 2013; **309**(6): 594-604.
20. Desai M, Gutman J, Taylor SM, Wiegand RE, Khairallah C, Kayentao K, et al. Impact of Sulfadoxine-Pyrimethamine Resistance on Effectiveness of Intermittent Preventive Therapy for Malaria in Pregnancy at Clearing Infections and Preventing Low Birth Weight. *Clin Infect Dis* 2016; **62**(3): 323-33.
21. Gonzalez R, Mombo-Ngoma G, Ouedraogo S, Kakolwa MA, Abdulla S, Accrombessi M, et al. Intermittent preventive treatment of malaria in pregnancy with mefloquine in HIV-negative women: a multicentre randomized controlled trial. *PLoS Med* 2014; **11**(9): e1001733.
22. Kimani J, Phiri K, Kamiza S, Duparc S, Ayoub A, Rojo R, et al. Efficacy and Safety of Azithromycin-Chloroquine versus Sulfadoxine-Pyrimethamine for Intermittent Preventive Treatment of Plasmodium falciparum Malaria Infection in Pregnant Women in Africa: An Open-Label, Randomized Trial. *PloS one* 2016; **11**(6): e0157045.
23. Desai M, Gutman J, L'Lanziva A, Otieno K, Juma E, Kariuki S, et al. Intermittent screening and treatment or intermittent preventive treatment with dihydroartemisinin-piperaquine versus intermittent preventive treatment with sulfadoxine-pyrimethamine for the control of malaria during pregnancy in western Kenya: an open-label, three-group, randomised controlled superiority trial. *Lancet* 2015; **386**(10012): 2507-19.
24. Kakuru A, Jagannathan P, Muhindo MK, Natureeba P, Awori P, Nakalembe M, et al. Dihydroartemisinin-Piperaquine for the Prevention of Malaria in Pregnancy. *N Engl J Med* 2016; **374**(10): 928-39.
25. Agarwal K, Alonso P, Chico RM, Coleman J, Dellicour S, Hill J, et al. Global Call to Action to scale-up coverage of intermittent preventive treatment of malaria in pregnancy: seminar report. *Malaria journal* 2015; **14**: 206.
26. Madanitsa M, Kalilani L, Mwapasa V, van Eijk AM, Khairallah C, Ali D, et al. Scheduled Intermittent Screening with Rapid Diagnostic Tests and Treatment with Dihydroartemisinin-Piperaquine versus Intermittent Preventive Therapy with Sulfadoxine-Pyrimethamine for Malaria in Pregnancy in Malawi: An Open-Label Randomized Controlled Trial. *PLoS Med* 2016; **13**(9): e1002124.
27. Rogerson SJ, Unger HW. Prevention and control of malaria in pregnancy - new threats, new opportunities? *Expert Rev Anti Infect Ther* 2017; **15**(4): 361-75.
28. World Health Organization. Guidelines for the treatment of malaria. Geneva: WHO, 2015.

29. Dondorp A, Nosten F, Stepniewska K, Day N, White N, South East Asian Quinine Artesunate Malaria Trial g. Artesunate versus quinine for treatment of severe falciparum malaria: a randomised trial. *Lancet* 2005; **366**(9487): 717-25.
30. Clark RL. Animal Embryotoxicity Studies of Key Non-Artemisinin Antimalarials and Use in Women in the First Trimester. *Birth Defects Res* 2017.
31. D'Alessandro U, Hill J, Tarning J, Pell C, Webster J, Gutman J, et al. Treatment of uncomplicated malaria and complicated malaria during pregnancy; . *Lancet Infectious Diseases* 2017; **in press**.
32. Malaria Policy Advisory Committee. Malaria Policy Advisory Committee to the WHO: conclusions and recommendations of eighth biannual meeting (September 2015). *Malaria journal* 2016; **15**.
33. Pregact Steering Group, Pekyi D, Ampromfi AA, Tinto H, Traore-Coulibaly M, Tahita MC, et al. Four Artemisinin-Based Treatments in African Pregnant Women with Malaria. *N Engl J Med* 2016; **374**(10): 913-27.
34. Poespoprodjo JR, Fobia W, Kenangalem E, Lampah DA, Sugiarto P, Tjitra E, et al. Treatment policy change to dihydroartemisinin-piperaquine contributes to the reduction of adverse maternal and pregnancy outcomes. *Malaria journal* 2015; **14**: 272.
35. Grigg MJ, William T, Menon J, Dhanaraj P, Barber BE, Wilkes CS, et al. Artesunate-mefloquine versus chloroquine for treatment of uncomplicated Plasmodium knowlesi malaria in Malaysia (ACT KNOW): an open-label, randomised controlled trial. *The Lancet infectious diseases* 2016; **16**(2): 180-8.
36. ter Kuile FO, Parise ME, Verhoeff FH, Udhayakumar V, Newman RD, van Eijk AM, et al. The burden of co-infection with human immunodeficiency virus type 1 and malaria in pregnant women in sub-saharan Africa. *The American journal of tropical medicine and hygiene* 2004; **71**(2 Suppl): 41-54.
37. Gonzalez R, Desai M, Macete E, Ouma P, Kakolwa MA, Abdulla S, et al. Intermittent preventive treatment of malaria in pregnancy with mefloquine in HIV-infected women receiving cotrimoxazole prophylaxis: a multicenter randomized placebo-controlled trial. *PLoS Med* 2014; **11**(9): e1001735.
38. Russo G, Paganotti GM, Soeria-Atmadja S, Haverkamp M, Ramogola-Masire D, Vullo V, et al. Pharmacogenetics of non-nucleoside reverse transcriptase inhibitors (NNRTIs) in resource-limited settings: Influence on antiretroviral therapy response and concomitant anti-tubercular, antimalarial and contraceptive treatments. *Infect Genet Evol* 2016; **37**: 192-207.
39. Goncalves BP, Walker PG, Cairns M, Tiono AB, Bousema T, Drakeley C. Pregnant Women: An Overlooked Asset to Plasmodium falciparum Malaria Elimination Campaigns? *Trends Parasitol* 2017; **33**(7): 510-8.
40. Ataide R, Mayor A, Rogerson SJ. Malaria, primigravidae, and antibodies: knowledge gained and future perspectives. *Trends Parasitol* 2014; **30**(2): 85-94.
41. van Eijk AM, Hill J, Noor AM, Snow RW, ter Kuile FO. Prevalence of malaria infection in pregnant women compared with children for tracking malaria transmission in sub-Saharan Africa: a systematic review and meta-analysis. *Lancet Glob Health* 2015; **3**(10): e617-28.