

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Wang, S;Wang, CX;Zhang, N;Xiang, YT;Yang, Y;Shi, YZ;Deng, YM;Zhu, MF;Liu, F;Yu, P;Ungvari, GS;Ng, CH

Title:

The association between post-stroke depression, aphasia, and physical independence in stroke patients at 3-month follow-up

Date:

2018-08-20

Citation:

Wang, S., Wang, C. X., Zhang, N., Xiang, Y. T., Yang, Y., Shi, Y. Z., Deng, Y. M., Zhu, M. F., Liu, F., Yu, P., Ungvari, G. S. & Ng, C. H. (2018). The association between post-stroke depression, aphasia, and physical independence in stroke patients at 3-month follow-up. *Frontiers in Psychiatry*, 9 (AUG), <https://doi.org/10.3389/fpsy.2018.00374>.

Persistent Link:

<https://hdl.handle.net/11343/271110>

License:

CC BY



# The Association Between Post-stroke Depression, Aphasia, and Physical Independence in Stroke Patients at 3-Month Follow-Up

Shuo Wang<sup>1,2,3,4,5</sup>, Chun-Xue Wang<sup>1,2,3,4,5,6\*</sup>, Ning Zhang<sup>1,2,3,4,5</sup>, Yu-Tao Xiang<sup>7</sup>, Yang Yang<sup>1,2,3,4,5</sup>, Yu-Zhi Shi<sup>1,3</sup>, Yi-Ming Deng<sup>3,4,8</sup>, Mei-Fang Zhu<sup>1,2,3,4,5</sup>, Fei Liu<sup>1,2,3,4,5</sup>, Ping Yu<sup>1,2,3,4,5</sup>, Gabor S. Ungvari<sup>9</sup> and Chee H. Ng<sup>10</sup>

<sup>1</sup> Department of Neurology, Beijing Tian Tan Hospital, Capital Medical University, Beijing, China, <sup>2</sup> Department of Neuropsychiatry and Behavioral Neurology and Clinical Psychology, Beijing Tian Tan Hospital, Capital Medical University, Beijing, China, <sup>3</sup> China National Clinical Research Center for Neurological Diseases, Beijing, China, <sup>4</sup> Center of Stroke, Beijing Institute for Brain Disorders, Beijing, China, <sup>5</sup> Beijing Key Laboratory of Translational Medicine for Cerebrovascular Disease, Beijing, China, <sup>6</sup> Department of Clinical Psychology, Capital Medical University, Beijing, China, <sup>7</sup> Unit of Psychiatry, Faculty of Health Sciences, University of Macau, Taipa, Macau, <sup>8</sup> Department of Interventional Neuroradiology, Beijing Tian Tan Hospital, Capital Medical University, Beijing, China, <sup>9</sup> University of Notre Dame Australia, Marian Centre and Graylands Hospital, Perth, WA, Australia, <sup>10</sup> Department of Psychiatry, University of Melbourne, Melbourne, VIC, Australia

## OPEN ACCESS

### Edited by:

Gianluca Serafini,  
Ospedale San Martino (IRCCS), Italy

### Reviewed by:

Jeffrey Guina,  
Wright State University, United States  
Paul David Barrows,  
University of Nottingham,  
United Kingdom

### \*Correspondence:

Chun-Xue Wang  
snowsen@126.com

### Specialty section:

This article was submitted to  
Mood and Anxiety Disorders,  
a section of the journal  
Frontiers in Psychiatry

Received: 08 May 2018

Accepted: 25 July 2018

Published: 20 August 2018

### Citation:

Wang S, Wang C-X, Zhang N,  
Xiang Y-T, Yang Y, Shi Y-Z, Deng Y-M,  
Zhu M-F, Liu F, Yu P, Ungvari GS and  
Ng CH (2018) The Association  
Between Post-stroke Depression,  
Aphasia, and Physical Independence  
in Stroke Patients at 3-Month  
Follow-Up. *Front. Psychiatry* 9:374.  
doi: 10.3389/fpsy.2018.00374

**Objective:** Few studies have examined the association between post-stroke depression (PSD), aphasia, and physical independence in Chinese patients. This study investigated the above association in stroke patients in China at 3-month follow-up.

**Methods:** Altogether 270 patients within 14 days after ischemic stroke were recruited and followed up at 3 months. PSD, aphasia, and physical functional status were measured using the Stroke Aphasia Depression Questionnaire (SADQ), Western Aphasia Battery (WAB), and modified Rankin Scale (mRS), respectively. Patients with mRS total score >2 were considered as having “physical dependence.”

**Results:** Out of 248 patients at 3-month follow up, 119 (48%) were rated as having physical dependence. Multiple logistic regression analyses revealed that female ( $p = 0.04$ ; OR = 2.2; 95% CI: 1.0–5.1), more severe stroke at admission ( $p < 0.01$ ; OR = 1.4; 95% CI: 1.3–1.5), and more severe PSD at 3 months ( $p = 0.01$ ; OR = 1.05; 95% CI: 1.01–1.1) were independently associated with physical dependence at 3 months.

**Conclusions:** Greater PSD and stroke severity were independently associated with physical dependence at 3 months after stroke. Aphasia was also associated with physical dependence but the relationship was not significant. Early and effective depression screening, treatment and stroke rehabilitation appear to be important to improve the physical outcome and reduce the burden of stroke survivors.

**Keywords:** aphasia, ischemic stroke, depression, PSD, physical independence

## INTRODUCTION

Post-stroke depression (PSD) is one of the most common psychiatric comorbidities in stroke survivors, with a prevalence ranging from 20 to 65% (1–4). PSD is significantly associated with poor treatment adherence and increased risk of disability, mortality, stroke recurrence, and poor quality of life (5, 6). Aphasia occurs in about a third of ischemic stroke patients and is associated with impaired activity of daily life (7, 8) and higher risk of PSD (9, 10). Poor functional outcome after stroke could result in significant personal distress and family burden (11). Understanding the association between PSD, aphasia and physical independence is hence important to develop comprehensive treatment strategies for stroke survivors.

In China, the lifetime prevalence of stroke was 2.08% (95% CI, 2.02–2.13%) (12) in 2017, which translates to ~2.9 million stroke patients. An Italy study found PSD was the only significant factor related to functional recovery from discharge to 3-month follow-up after stroke but not aphasia (13). To the best of our knowledge, there are no studies that have investigated the independent association between PSD, aphasia, and physical independence in stroke survivors in China. This study thus aimed to examine the association between PSD, aphasia, and physical independence in Chinese stroke patients at 3-month follow-up.

## METHODS

### Participants and Study Setting

This prospective cohort study was conducted between April 2014 and October 2015 in the Stroke Centre of Beijing Tiantan Hospital. A total of 320 patients were consecutively screened if they fulfilled the following criteria: (1) aged 18 years or older; (2) had an acute ischemic stroke within 14 days according to the WHO diagnostic criteria (14) confirmed by computed tomography (CT) or magnetic resonance imaging (MRI); and (3) had the ability to provide informed consent and complete the assessment. Exclusion criteria included: (1) history of language impairment; (2) drug and alcohol abuse and severe psychiatric disorders; (3) other major medical conditions, such as Parkinson's disease; and (4) severe cognitive deficit defined by the Mini Mental State Examination (MMSE) total score <18 (15, 16). The first three were excluded based on the self-reported pre-stroke histories by unstructured interviews and the last was made at the start of the study. The study protocol was approved by the ethics committee of Beijing Tiantan Hospital, Capital Medical University. All participants provided written informed consent.

### Measurement Instruments and Evaluation

Assessment was conducted at baseline and 3 months after index ischemic stroke. Patients' socio-demographic and clinical characteristics at baseline were recorded via a review of electronic medical records and confirmed by a clinical interview conducted by trained research neurologists. Severity and type of stroke was assessed with the National Institutes of Health Stroke Scale (NIHSS) (17–19) and the Trial of Org 10172 Acute Stroke Treatment (TOAST) (20, 21).

Physical independence and degree of handicap was evaluated using the modified Ranking Scale (mRS) at 3 months, with mRS total score >2 indicating physical dependence (22, 23). As most depression scales cannot be used in stroke patients with aphasia due to their impaired communication ability, the severity of depressive symptoms at  $14 \pm 2$  days and 3 months after index stroke was measured using the 21-item Stroke Aphasic Depression Questionnaire (SADQ) (24, 25) with total score ranges from 0 to 63,  $\geq 19$  indicating the presence of PSD,  $\geq 22$  indicating moderate depressive symptoms and  $\geq 26$  indicating major depressive symptoms. The SADQ relies on external observation of emotional behavior by nursing staff and family members in recent time. The hospital version (SADQ-H) focus on the recent week. The severity and type of aphasia was evaluated with the Aphasia Quotient (AQ) derived from the Western Aphasia Battery (WAB), with AQ < 93.8 indicating the presence of aphasia (26). The language assessment was also conducted at 3-month follow-up with much missing data on AQ score, so the data was not analyzed.

### Statistical Analysis

Data were analyzed with SPSS 21.0 (SPSS Inc., Chicago, IL, USA). Socio-demographic and clinical variables were compared between physical independence and dependence groups using Chi-square test, independent sample *t*-test and Mann-Whitney *U*-test, as appropriate. Independent correlates of the physical dependence at 3 months were examined using multivariate logistic regression analysis with the "enter" method. The outcome at 3 months was the dependent variable, while variables that significantly differed between both groups in the univariate analyses were entered as independent variables. Significance was set at 0.05 (two-tailed).

## RESULTS

Out of 320 patients with ischemic stroke who were consecutively screened, 270 fulfilled the study entry criteria and participated in the study, giving a participation rate of 84.4%. At baseline, 160 (59.3%) patients had aphasic symptom with the incidence of PSD was 47.5% compared with 29.1% in non-aphasiac patients ( $p < 0.01$ ). At the 3-month assessment, 22 patients dropped out due to lack of interest, moving house or other unknown reasons. Of the 248 patients who completed the 3-month assessment, 119 (48.0%) had physical dependence.

**Table 1** shows the socio-demographic and clinical characteristics of the whole sample and separately by outcome. Patients with physical dependence were less likely to be married, and have large-artery atherosclerosis (LAA) TOAST type and pulmonary infection, but more likely to be female, have aphasia at baseline and more severe depressive symptoms at both baseline and 3 months after stroke. In addition, they had higher NIHSS scores at baseline, and higher SADQ scores at baseline and 3 months (all *p*-values < 0.05).

The independent correlates of physical dependence are shown in **Tables 2, 3**. Due to collinearity between SADQ-H (Hospital version using during hospitalization) and SADQ score, two multivariate logistic regression analyses were performed with

**TABLE 1** | Comparison of demographic and clinical variables between physical independence and dependence groups.

|                                                | Total sample<br>( <i>n</i> = 248) |      | Physical independence<br>( <i>n</i> = 129) |      | Physical dependence<br>( <i>n</i> = 119) |      | Statistics |     |                  |
|------------------------------------------------|-----------------------------------|------|--------------------------------------------|------|------------------------------------------|------|------------|-----|------------------|
|                                                | <i>N</i>                          | %    | <i>N</i>                                   | %    | <i>N</i>                                 | %    | $\chi^2$   | df  | <i>p</i>         |
| Female                                         | 72                                | 29.0 | 26                                         | 20.2 | 46                                       | 38.7 | 10.2       | 1   | <b>0.001</b>     |
| Higher education (high middle school or above) | 88                                | 35.5 | 49                                         | 38.0 | 39                                       | 32.8 | 0.7        | 1   | 0.39             |
| Being married                                  | 229                               | 92.3 | 124                                        | 96.1 | 105                                      | 88.2 | 5.4        | 1   | <b>0.02</b>      |
| Living with others                             | 235                               | 94.8 | 123                                        | 95.3 | 112                                      | 94.1 | 0.1        | 1   | 0.66             |
| Diabetes                                       | 61                                | 24.6 | 29                                         | 22.5 | 32                                       | 26.9 | 0.6        | 1   | 0.42             |
| Hypertension                                   | 156                               | 62.9 | 87                                         | 67.4 | 69                                       | 58.0 | 2.3        | 1   | 0.12             |
| Hyperlipidemia                                 | 33                                | 13.5 | 19                                         | 14.7 | 14                                       | 11.8 | 0.4        | 1   | 0.49             |
| Cardiac disease                                | 21                                | 8.5  | 10                                         | 7.8  | 11                                       | 9.2  | 0.1        | 1   | 0.67             |
| Stroke history                                 | 58                                | 23.4 | 27                                         | 20.9 | 31                                       | 26.1 | 0.9        | 1   | 0.34             |
| TOAST type (LAA)                               | 209                               | 84.3 | 115                                        | 89.1 | 94                                       | 79.0 | 4.8        | 1   | <b>0.02</b>      |
| Aphasia at baseline                            | 148                               | 59.7 | 66                                         | 51.2 | 82                                       | 68.9 | 8.1        | 1   | <b>0.004</b>     |
| Depressive symptoms at baseline                | 100                               | 40.3 | 39                                         | 30.2 | 61                                       | 51.3 | 11.3       | 1   | <b>0.001</b>     |
| Moderate/Major depressive symptoms at baseline | 54                                | 21.8 | 15                                         | 11.6 | 39                                       | 32.8 | 16.2       | 1   | <b>&lt;0.001</b> |
| Depressive symptoms at 3 months                | 85                                | 34.7 | 26                                         | 20.2 | 59                                       | 50.9 | 25.4       | 1   | <b>&lt;0.001</b> |
| Moderate/Major depressive symptoms at 3 months | 48                                | 19.6 | 8                                          | 6.2  | 40                                       | 34.5 | 31.0       | 1   | <b>&lt;0.001</b> |
| Pulmonary infection at baseline                | 36                                | 14.5 | 13                                         | 10.1 | 23                                       | 9.3  | 4.2        | 1   | <b>0.03</b>      |
| Upper gastrointestinal hemorrhage at baseline  | 8                                 | 3.2  | 3                                          | 2.3  | 5                                        | 4.4  | 0.2        | 1   | 0.63             |
| Cardiac diseases at baseline                   | 12                                | 4.8  | 5                                          | 3.9  | 7                                        | 5.9  | 0.5        | 1   | 0.46             |
| Urinary infection at baseline                  | 11                                | 4.4  | 3                                          | 2.3  | 8                                        | 6.7  | 2.8        | 1   | 0.09             |
| Deep venous thrombosis at baseline             | 20                                | 8.1  | 5                                          | 3.9  | 15                                       | 12.6 | 6.3        | 1   | <b>0.01</b>      |
|                                                | Mean                              | SD   | Mean                                       | SD   | Mean                                     | SD   | T/Z        | df  | <i>p</i>         |
| Age (years)                                    | 57.3                              | 14.2 | 57.3                                       | 14.0 | 57.3                                     | 14.6 | 0.01       | 246 | 0.9              |
| Stroke duration from onset to admission        | 6.1                               | 6.1  | 6.3                                        | 6.1  | 5.9                                      | 6.0  | -1.2       | __a | 0.22             |
| NIHSS at baseline                              | 8.3                               | 5.3  | 5.1                                        | 3.7  | 11.8                                     | 4.5  | -10.1      | __a | <b>&lt;0.001</b> |
| SADQ-H at baseline                             | 14.5                              | 7.9  | 12.5                                       | 7.2  | 16.8                                     | 8.0  | -4.2       | __a | <b>&lt;0.001</b> |
| SADQ at 3 months                               | 12.8                              | 8.0  | 9.7                                        | 7.1  | 16.2                                     | 7.6  | -6.3       | __a | <b>&lt;0.001</b> |

Values *p* < 0.05 are bolded; a, Mann-Whitney U-test; LAA, Large-Artery Atherosclerosis; NIHSS, National Institutes of Health Stroke Scale; SADQ-H, Stroke Aphasic Depression Questionnaire (Hospital version); SADQ, Stroke Aphasic Depression Questionnaire; TOAST, Trial of Org 10172 in Acute Stroke Treatment.

SADQ-H and SADQ separately. Finally, female, NIHSS score at admission and SADQ score at 3 months were independently associated with physical dependence at 3-month follow up (adjusted  $R^2 = 0.410$  in **Table 2** and adjusted  $R^2 = 0.423$  in **Table 3**).

## DISCUSSION

Generally about a third of patients with ischemic stroke have poor functional outcome at different follow-up time points (27, 28). In this study, the prevalence of physical dependence was 48% using the mRS cut-off score, which is similar to the 5-year prevalence of poor functional outcome (45%) in another study with the same measure (29). However, any direct comparison should be done with caution due to the different assessment measures, follow-up time points, and demographic characteristics.

A report of the American Heart Association/American Stroke Association in 2017 and meta-analyses found that approximately a third of stroke patients develop PSD at any point after stroke (30–32). A large-scale prospective cohort study in China

involving over 2,000 stroke patients found that the cumulative incidence of PSD at 1 year after the index stroke was 42% (4). Depression was found predictive of worse functional outcome in an updated meta-analysis about the impact of depression on stroke outcome (33). While up to 62–70% in aphasic patients with stroke were diagnosed as major depression according to DSM-III-R criteria at 3 months and 1 year after stroke (9, 10, 34) which was higher than our results. The reason may due to the different scales, population and time points. Aphasia have been considered for the potential association with PSD with inconsistent conclusion (35, 36). We found that the incidence of PSD in stroke patients with aphasia and non-aphasia was significantly different indicating aphasia may be a risk factor of the development of PSD.

In this study, the incidence of PSD was 34.7% at 3 months after stroke, which was independently associated with physical dependence. Similar findings were also reported previously (37–39). Depressive symptoms after stroke could result in behavioral and biological abnormalities, such as poor treatment adherence and dysregulation in autonomic system activation, which in turn, could lead to physical dependence (11, 37). However,

**TABLE 2 |** Independent correlates of physical dependence at 3 months<sup>#</sup>.

| Variables                          | Ischemic stroke patients |      |          |
|------------------------------------|--------------------------|------|----------|
|                                    | P                        | OR   | 95%CI    |
| Feale                              | <b>0.04</b>              | 2.2  | 1.0,5.1  |
| Being married                      | 0.22                     | 0.3  | 0.08,1.7 |
| TOAST type (LAA)                   | 0.75                     | 1.1  | 0.4,3.5  |
| Aphasia at baseline                | 0.32                     | 1.4  | 0.6,2.9  |
| SADQ-H score at baseline           | 0.72                     | 1.08 | 0.9,1.05 |
| Pulmonary infection at baseline    | 0.74                     | 1.1  | 0.4,3.4  |
| Deep venous thrombosis at baseline | 0.27                     | 2.4  | 0.5,11.5 |
| NIHSS at admission                 | <b>&lt;0.01</b>          | 1.4  | 1.3,1.5  |

Values  $p < 0.05$  are bolded; a, Mann-Whitney U-test; LAA, Large-Artery Atherosclerosis; NIHSS, National Institutes of Health Stroke Scale; SADQ-H, Stroke Aphasic Depression Questionnaire (Hospital version); TOAST, Trial of Org 10172 in Acute Stroke Treatment.

<sup>#</sup>The severity of depressive symptoms was tested by SADQ-H at baseline and SADQ at 3 months. There was collinearity between SADQ-H at baseline and SADQ at 3 months, therefore they were not entered in the analysis concurrently.

**TABLE 3 |** Independent correlates of physical dependence at 3 months<sup>#</sup>.

| Variables                          | Ischemic stroke patients |      |          |
|------------------------------------|--------------------------|------|----------|
|                                    | P                        | OR   | 95%CI    |
| Female                             | 0.08                     | 2.0  | 0.9,4.8  |
| Being married                      | 0.26                     | 0.4  | 0.08,1.9 |
| TOAST type (LAA)                   | 0.80                     | 1.1  | 0.3,3.4  |
| Aphasia at baseline                | 0.53                     | 1.2  | 0.6,2.5  |
| SADQ score at 3 months             | <b>0.01</b>              | 1.05 | 1.01,1.1 |
| Pulmonary infection at baseline    | 0.69                     | 1.2  | 0.4,3.6  |
| Deep venous thrombosis at baseline | 0.30                     | 2.2  | 0.4,10.3 |
| NIHSS at admission                 | <b>&lt;0.01</b>          | 1.4  | 1.2,1.5  |

Values  $p < 0.05$  are bolded; a, Mann-Whitney U test; LAA, Large-Artery Atherosclerosis; NIHSS, National Institutes of Health Stroke Scale; SADQ, Stroke Aphasic Depression Questionnaire; TOAST, Trial of Org 10172 in Acute Stroke Treatment. <sup>#</sup>The severity of depressive symptoms was tested by SADQ-H at baseline and SADQ at 3 months. There was collinearity between SADQ-H at baseline and SADQ at 3 months, therefore they were not entered in the analysis concurrently.

since more than half of the patients suffered from aphasia in this sample, the use of the SADQ required the input of nursing staff and/or family members. Therefore, we could not exclude the possibility that nursing staff and family members were unable to distinguish between insomnia, irritability, poor appetite, anxiety, and depressive symptoms, which would cause bias in the incidence of PSD to an uncertain extent.

In this study, aphasia was measured using the AQ that covered complete aphasia, motor aphasia, sensory aphasia, transcortical mixed aphasia, transcortical motor aphasia, transcortical sensory aphasia, anomic aphasia, and conduction aphasia. Aphasia may lead to communication or comprehension difficulties, social avoidance and decreased attention, which is associated with physical dependence in stroke survivors (40). However, this finding was only confirmed in the univariate, but not in the multivariate analyses in this study. It is speculated that the association between aphasia and poor physical independence was moderated by other variables, such as depressive symptoms and severity of stroke. In addition, traditional Confucian culture

favors family support and inter-dependence, particularly for family members with illness. For example, 94.8% of patients in this study were living with others. Thus, the strong family support may have offset the association between aphasia and physical dependence.

The association between demographic characteristics and physical dependence in stroke survivors have been inconsistent (41, 42). In this study, only female was independently associated with physical dependence, which is supported by previous findings (43, 44). The gender difference in physical independence in stroke survivors may be related to menstrual cycles, neuro-endocrine regulation and more frequent physical comorbidities, such as diabetes, atrial fibrillation, and coronary heart disease in women with stroke (45). As expected, stroke severity as measured by the NIHSS was positively associated with physical dependence, which is consistent with previous findings (27, 29).

There are several methodological limitations to this study. First, this was a single-center study with relatively small sample size, therefore the findings could not be generalized to all stroke patients in China. Second, depressive symptoms were measured using the SADQ based on the observation by nursing staff or family members. There may be a gap between observer-rated and self-reported measures of depression, although there are a number of self-reported measures specific for aphasia such as the Visual Analogue Mood Scales (VAMS) (46), Visual Analogue Self-Esteem Scales (VASES) (47), Disc Intensity Scale Circles (DISCS) (48), and Dynamic Visual Analogue Mood Scales (D-VAMS) (49). We chose SADQ from the perspective of relatively short items, easy operation and short time. Third, some important variables related to physical dependence, such as the use of medication, treatment adherence, the size and lesions of infarcts and the missing data about aphasia at 3-month, were not evaluated in the 3 months assessment.

In conclusion, physical dependence at 3-month follow up was common in Chinese stroke patients, which was associated with gender, greater PSD and stroke severity. Aphasia was also associated with physical dependence but the relationship was not significant. Our findings call for early and effective depression screening, treatment, and stroke rehabilitation to improve physical outcome and reduce the burden of stroke survivors in China.

## AUTHOR CONTRIBUTIONS

SW and C-XW: study design. SW, NZ, YY, Y-ZS, Y-MD, M-FZ, FL, PY, and Y-TX: collection, analysis and interpretation of data. SW, C-XW, and Y-TX: drafting of the manuscript. GU, CN, and Y-TX: critical revision of the manuscript. All coauthors approval of the final version for publication.

## FUNDING

This study was funded by the Ministry of Science and Technology and the Ministry of Health of the People's Republic of China. Individual grants include the National Key Research & Development Program of China (No. 2016YFC1301720), Beijing

Brain Research (Z161100000216131), the Beijing Municipal Science and Technology Commission (Z151100004015127), and the Build High Level Technology Talents of Health System in Beijing (No.2015-3-038).

## REFERENCES

- Kotila M, Numminen H, Waltimo O, Kaste M. Post-stroke depression and functional recovery in a population-based stroke register. The Finnstroke study. *Eur J Neurol.* (1999) 6:309–12. doi: 10.1046/j.1468-1331.1999.630309.x
- Naess H, Nyland HI, Thomassen L, Aarseth J, Myhr KM. Mild depression in young adults with cerebral infarction at long-term follow-up: a population-based study. *Eur J Neurol.* (2005) 12:194–8. doi: 10.1111/j.1468-1331.2004.00937.x
- Robinson RG, Jorge RE. Post-stroke depression: a review. *Am J Psychiatry* (2016) 173:221–31. doi: 10.1176/appi.ajp.2015.15030363
- Zhang N, Wang CX, Wang AX, Bai Y, Zhou Y, Wang YL, et al. Time course of depression and one-year prognosis of patients with stroke in mainland China. *CNS Neurosci Ther.* (2012) 18:475–81. doi: 10.1111/j.1755-5949.2012.00312.x
- House A, Knapp P, Bamford J, Vail A. Mortality at 12 and 24 months after stroke may be associated with depressive symptoms at 1 month. *Stroke* (2001) 32:696–701. doi: 10.1161/01.STR.32.3.696
- Morris PL, Robinson RG, Andrzejewski P, Samuels J, Price TR. Association of depression with 10-year poststroke mortality. *Am J Psychiatry* (1993) 150:124–9.
- Engelter ST, Gostynski M, Papa S, Frei M, Born C, Ajdacic-Gross V, et al. Epidemiology of aphasia attributable to first ischemic stroke: incidence, severity, fluency, etiology, and thrombolysis. *Stroke* (2006) 37:1379–84. doi: 10.1161/01.STR.0000221815.64093.8c
- Thomas SA, Lincoln NB. Predictors of emotional distress after stroke. *Stroke* (2008) 39:1240–5. doi: 10.1161/STROKEAHA.107.498279
- De Ryck A, Franssen E, Brouns R, Geurden M, Peij D, Mariën P, et al. Poststroke depression and its multifactorial nature: results from a prospective longitudinal study. *J Neurol Sci.* (2014) 347:159–66. doi: 10.1016/j.jns.2014.09.038
- Shehata GA, El Mistikawi T, Al Sayed KR, Hassan HS. The effect of aphasia upon personality traits, depression and anxiety among stroke patients. *J Affect Disord.* (2015) 172:312–4. doi: 10.1016/j.jad.2014.10.027
- Dhamoon MS, McClure LA, White CL, Lakshminarayan K, Benavente OR, Elkind MS. Long-term disability after lacunar stroke: secondary prevention of small subcortical strokes. *Neurology* (2015) 84:1002–8. doi: 10.1212/WNL.0000000000001331
- Li Q, Wu H, Yue W, Dai Q, Liang H, Bian H, et al. Prevalence of stroke and vascular risk factors in China: a nationwide community-based Study. *Sci Rep.* (2017) 7:6402. doi: 10.1038/s41598-017-06691-1
- Nannetti L, Paci M, Pasquini J, Lombardi B, Taiti PG. Motor and functional recovery in patients with post-stroke depression. *Disabil Rehabil.* (2005) 27:170–5. doi: 10.1080/09638280400009378
- Kunitz SC, Gross CR, Heyman A, Kase CS, Mohr JP, Price TR, et al. The pilot stroke data bank: definition, design, and data. *Stroke* (1984) 15:740–6. doi: 10.1161/01.STR.15.4.740
- Folstein MF, Folstein SE, McHugh PR. “Mini-mental state”. A practical method for grading the cognitive state of patients for the clinician. *J Psychiatric Res.* (1975) 12:189–98. doi: 10.1016/0022-3956(75)90026-6
- Shen M, Jiamei M. Cognitive scale in healthy Chinese elderly. *Chin J Gerontol.* (1991) 11:203–7.
- Brott T, Adams HP, Olinger CP, Marle JR, Barsan WG, Biller J, et al. Measurements of acute cerebral infarction: a clinical examination scale. *Stroke* (1989) 20:864–70. doi: 10.1161/01.STR.20.7.864
- Lyden P, Brott T, Tilley B, Welch KM, Mascha EJ, Levine S, et al. Improved reliability of the NIH Stroke Scale using video training. NINDS TPA Stroke Study Group. *Stroke* (1994) 25:2220–6.
- Sun TK, Chiu SC, Yeh SH, Chang KC. Assessing reliability and validity of the Chinese version of the stroke scale: scale development. *Int J Nurs Stud.* (2006) 43:457–63. doi: 10.1016/j.ijnurstu.2005.07.004
- Adams HP, Bendixen BH, Kappelle LJ, Biller J, Love BB, Gordon DL, et al. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of Org 10172 in Acute Stroke Treatment. *Stroke* (1993) 24:35–41.
- Zhou Heng LJ, Wang Yong-jun J. The reliability of ischemic stroke subtype classification using the TOAST criteria. *Chin J Intern Med.* (2005) 44:825–7. doi: 10.3760/j.issn:0578-1426.2005.11.011
- American College of Sports Medicine, Chodzko-Zajko WJ, Proctor DN, Fiatarone Singh MA, Minson CT, Nigg CR, et al. American College of Sports Medicine position stand. Exercise and physical activity for older adults. *Med Sci Sports Exerc.* (2009) 41:1510–30. doi: 10.1249/MSS.0b013e3181a0c95c
- Uyttenboogaart M, Stewart RE, Vroomen, P. C, De Keyser J, Luijckx, GJ. Optimizing cutoff scores for the Barthel index and the modified Rankin scale for defining outcome in acute stroke trials. *Stroke* (2005) 36:1984–7. doi: 10.1161/01.STR.0000177872.87960.61
- Chang Y, Jing X, Junping W. Validation of Chinese version of Stroke Aphasic Depression Questionnaire (SADQ). *Chin J Clin Psychol.* (2006) 14:230–2. doi: 10.3969/j.issn.1005-3611.2006.03.004
- Sutcliffe LM, Lincoln NB. The assessment of depression in aphasic stroke patients: the development of the stroke aphasic depression questionnaire. *Clin Rehabil.* (1998) 12:506–13. doi: 10.1191/026921598672167702
- Shewan CM, Kertesz A. Reliability and validity characteristics of the Western Aphasia Battery (WAB). *J Speech Hear Disord.* (1980) 45:308–24.
- Hankey GJ. Long-term outcome after ischaemic stroke/transient ischaemic attack. *Cerebrovasc Dis.* (2003) 16 (Suppl. 1):14–9. doi: 10.1159/000069936
- Ji R, Du W, Shen H, Pan Y, Wang P, Liu G, et al. Web-based tool for dynamic functional outcome after acute ischemic stroke and comparison with existing models. *BMC Neurol.* (2014) 14:214. doi: 10.1186/s12883-014-0214-z
- Yang Y, Shi YZ, Zhang N, Wang S, Ungvari GS, Ng CH, et al. The disability rate of 5-year post-stroke and its correlation factors: a national survey in China. *PLoS ONE* (2016) 11:e0165341. doi: 10.1371/journal.pone.0165341
- Hackett ML, Pickles K. Part I: frequency of depression after stroke: an updated systematic review and meta-analysis of observational studies. *Int J Stroke* (2014) 9:1017–25. doi: 10.1111/ijs.12357
- Hackett ML, Yapa C, Parag V, Anderson CS. Frequency of depression after stroke: a systematic review of observational studies. *Stroke* (2005) 36:1330–40. doi: 10.1161/01.STR.0000165928.19135.35
- Towfighi A. Poststroke depression: a scientific statement for healthcare professionals from the American heart association/American stroke association. *Stroke* (2017) 48:e30–43. doi: 10.1161/STR.0000000000000113
- Kutlubaev MA, Hackett ML. Part II: predictors of depression after stroke and impact of depression on stroke outcome: an updated systematic review of observational studies. *Int J Stroke* (2014) 9:1026–36. doi: 10.1111/ijs.12356
- Kauhanen ML, Korpelainen JT, Hiltunen P, Määttä R, Mononen H, Brusin E, et al. Aphasia, depression, and non-verbal cognitive impairment in ischaemic stroke. *Cerebrovasc Dis.* (2000) 10:455–61. doi: 10.1159/000016107
- Sienkiewicz-Jaroszy H, Milewska D, Bochynska A, Chelminiak A, Dworek N, Kasprzyk K, et al. Predictors of depressive symptoms in patients with stroke - a three-month follow-up. *Neurol Neurochir Pol.* (2010) 44:13–20. doi: 10.1016/S0028-3843(14)60402-3
- Starkstein SE, Robinson RG. Affective disorders and cerebral vascular disease. *Br J Psychiatry* (1989) 154:170–82. doi: 10.1192/bjp.154.2.170
- Ayerbe L, Ayis S, Crichton S, Wolfe CD, Rudd AG. The long-term outcomes of depression up to 10 years after stroke; the South London Stroke Register. *J Neurol Neurosurg Psychiatry* (2014) 85:514–21. doi: 10.1136/jnnp-2013-306448

## ACKNOWLEDGMENTS

The authors would like to thank all of the participating colleagues, patients, and their families.

38. Robinson RG. *The Clinical Neuropsychiatry of Stroke*. Cambridge: Cambridge University Press (2006).
39. Shi YZ, Xiang YT, Yang Y, Zhang N, Wang S, Ungvari GS, et al. Depression after minor stroke: the association with disability and quality of life - a 1-year follow-up study. *Int J Geriatr Psychiatry* (2016) 31:421–7. doi: 10.1002/gps.4353
40. El Hachoui H, Lingsma HF, Van De Sandt-Koenderman, MW, Dippel DW, Koudstaal PJ, Visch-Brink EG. Long-term prognosis of aphasia after stroke. *J Neurol Neurosurg Psychiatry* (2013) 84:310–5. doi: 10.1136/jnnp-2012-302596
41. Fahey M, Crayton E, Wolfe C, Douiri A. Clinical prediction models for mortality and functional outcome following ischemic stroke: a systematic review and meta-analysis. *PLoS ONE* (2018) 13:e0185402. doi: 10.1371/journal.pone.0185402
42. Rempe DA. Predicting outcomes after transient ischemic attack and stroke. *Cerebrovasc Dis.* (2014) 20:412–28. doi: 10.1212/01.CON.0000446110.97667.58
43. Glader EL, Stegmayr B, Norrving B, Terént A, Hulter-Asberg K, Wester, PO, et al. Sex differences in management and outcome after stroke: a Swedish national perspective. *Stroke* (2003) 34:1970–5. doi: 10.1161/01.STR.0000083534.81284.C5
44. Liu X, Lv Y, Wang B, Zhao G, Yan Y, Xu D. Prediction of functional outcome of ischemic stroke patients in northwest China. *Clin Neurol Neurosurg.* (2007) 109:571–7. doi: 10.1016/j.clineuro.2007.05.008
45. Wang Z, Li J, Wang C, Yao X, Zhao X, Wang Y, et al. Gender differences in 1-year clinical characteristics and outcomes after stroke: results from the China National Stroke Registry. *PLoS ONE* (2013) 8:e56459. doi: 10.1371/journal.pone.0056459
46. Stern RA, Arruda JE, Hooper CR, Wolfner GD, Morey CE. Visual analogue mood scales to measure internal mood state in neurologically impaired patients: description and initial validity evidence. *Aphasiology* (1997) 11:59–71. doi: 10.1080/02687039708248455
47. Brumfitt SM, Sheeran P. The development and validation of the Visual Analogue Self-Esteem Scale (VASES). *Br J Clin Psychol.* (1999) 38 (Pt 4):387–400.
48. Turner-Stokes L. (2005). The Depression Intensity Scale Circles (DISCs): a first evaluation of a simple assessment tool for depression in the context of brain injury. *J Neurol Neurosurg Psychiatry* 76:1273–8. doi: 10.1136/jnnp.2004.050096
49. Barrows PD, Thomas SA. Assessment of mood in aphasia following stroke: validation of the Dynamic Visual Analogue Mood Scales (D-VAMS). *Clin Rehabil.* (2018) 32:94–102. doi: 10.1177/0269215517714590

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Wang, Wang, Zhang, Xiang, Yang, Shi, Deng, Zhu, Liu, Yu, Ungvari and Ng. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.