



Minerva Access is the Institutional Repository of The University of Melbourne

**Author/s:**

Nguyen, HTH;Ong, L;Beaucher, E;Madec, M-N;Kentish, SE;Gras, SL;Lopez, C

**Title:**

Buffalo milk fat globules and their biological membrane: in situ structural investigations

**Date:**

2015-01-01

**Citation:**

Nguyen, H. T. H., Ong, L., Beaucher, E., Madec, M. -N., Kentish, S. E., Gras, S. L. & Lopez, C. (2015). Buffalo milk fat globules and their biological membrane: in situ structural investigations. *Food Research International*, 67, pp.35-43. <https://doi.org/10.1016/j.foodres.2014.10.022>.

**Persistent Link:**

<https://hdl.handle.net/11343/238440>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

**Buffalo milk fat globules and their biological membrane: *in situ* structural investigations**

Hanh T.H. Nguyen <sup>a,b,c</sup>, Lydia Ong <sup>a,b,c</sup>, Eric Beaucher <sup>d,e</sup>, Marie-Noëlle Madec <sup>d,e</sup>, Sandra E. Kentish <sup>a,c</sup>, Sally L. Gras <sup>a,b,c</sup>, Christelle Lopez <sup>d,e,\*</sup>

<sup>a</sup> Department of Chemical and Biomolecular Engineering, The University of Melbourne, Parkville, Vic 3010, Australia.

<sup>b</sup> The Bio21 Molecular Science and Biotechnology Institute, The University of Melbourne, Parkville, Vic 3010, Australia.

<sup>c</sup> The ARC Dairy Innovation Hub, The University of Melbourne, Parkville, Vic 3010, Australia.

<sup>d</sup> INRA, UMR 1253, Science and Technology of Milk and Egg, F-35000 Rennes, France.

<sup>e</sup> Agrocampus Ouest, UMR 1253, Science and Technology of Milk and Egg, F-35000 Rennes, France.

\* Corresponding author  
E-mail address: [Christelle.Lopez@rennes.inra.fr](mailto:Christelle.Lopez@rennes.inra.fr) (C. Lopez)  
Tel.: +33 2 23 48 56 17  
Fax: +33 2 23 48 53 50

**Abbreviations:** MFGM: milk fat globule membrane; Lo phase: liquid ordered phase; CLSM: confocal laser scanning microscopy; DIC: differential interference contrast; TAG: triacylglycerol; SM: sphingomyelin; PC: phosphatidylcholine

25 **ABSTRACT**

26 Milk fat globules and their surrounding biological membrane (the MFGM) are not well  
27 understood despite the importance of these milk components in human nutrition and the role  
28 of fat globules in determining the properties of dairy products. The objectives of this study  
29 were to investigate these unique colloidal assemblies and the microstructure of the MFGM in  
30 buffalo milk, which is the second largest global source of dairy products. *In-situ* structural  
31 investigations were performed at room temperature using confocal microscopy with multiple  
32 fluorescent probes (Nile Red, Rh-DOPE, the lectin WGA-488). Microscopic observations  
33 showed cytoplasmic crescents around fat globules and the heterogeneous distribution of  
34 glycosylated molecules and polar lipids with the occurrence of lipid domains. The lipid  
35 domains in the buffalo MFGM appear to form by the segregation of lipids with a high phase  
36 transition temperature (e.g. sphingomyelin and saturated phosphatidylcholine molecular  
37 species) and cholesterol resulting in a gel phase or a Lo phase forming circular domains. The  
38 structure of the buffalo MFGM results from a non-random mixing of components, consistent  
39 with observations for other species. Structural heterogeneities of the MFGM could affect the  
40 processability of buffalo fat globules and the bioavailability of milk lipids.

41

42 **Keywords:** milk fat globule membrane, phospholipid, sphingomyelin, high phase transition  
43 temperature lipid, lipid domain

44

## 45 **1. Introduction**

46 Milk is an exceptionally complex biological fluid used for the manufacture of a wide range of  
47 dairy products. Numerous studies have focused on cow's milk although milks from other  
48 animal species such buffalos, ewes, goats and camels are essential to the human diet in  
49 various parts of the world. Buffalo milk represents the second largest volume of milk  
50 produced globally after cow's milk with more than 97 million tonnes produced each year  
51 (FAOSTAT, 2012). Buffalo milk is also one of the richest milks from a compositional point  
52 of view (Ménard et al., 2010). Fat constitutes the main fraction of buffalo milk, with the  
53 almost twice the fat content of bovine milk (7.4-8.8% w/w vs. 3.6-4.7% w/w) (Solh, Staines,  
54 Honda, & Limley, 2007; Varricchio, Di Francia, Masucci, Romano, & Proto, 2007) and this  
55 fat is responsible for the high energetic and nutritive value of buffalo milk. Despite the  
56 nutritional value of milk fat and the influence of fat on dairy product properties, information  
57 about buffalo milk fat is scarce.

58 All milk fat is thought to be dispersed in colloidal assemblies called milk fat globules. The  
59 core of the milk fat globule is mainly composed of triacylglycerols (TAG, esters of fatty acids  
60 and glycerol; 98% of milk lipids). This core is surrounded by a biological membrane called  
61 the milk fat globule membrane (MFGM). This MFGM contains many bioactive compounds  
62 which are involved with several biological functions and health benefits, such as neonatal gut  
63 maturation, antibacterial infection, inhibition of colon development and lowering cholesterol  
64 absorption (Dewettinck et al., 2008; Lopez, 2011). Previous studies have demonstrated that  
65 buffalo milk fat globules are significantly larger in size (5.0  $\mu\text{m}$  vs. 3.5  $\mu\text{m}$ ) and have a higher  
66 absolute zeta potential ( $|-11.0 \text{ mV}|$  vs.  $|- 9.4 \text{ mV}|$ ) compared to bovine milk fat globules  
67 (Ménard et al., 2010). Few authors have characterized the buffalo MFGM (Abou-Dawood,  
68 Moussaa, El-Demerdash, & Ahmed, 1988; D'Ambrosio et al., 2008; Ménard et al., 2010).

69 This biological membrane is thought to be comprised of three layers, containing mainly of  
70 glycerophospholipids (phosphatidylcholine, PC; phosphatidylethanolamine, PE;  
71 phosphatidylinositol, PI; phosphatidylserine, PS), sphingolipids (mainly sphingomyelin, SM),  
72 cholesterol and proteins (Keenan & Patton, 1995; Ménard et al., 2010). Polar lipids, which are  
73 mainly located in the MFGM, account for about 2.6 mg/g fat and about 190 mg/L of buffalo  
74 milk (Ménard et al., 2010), which is 28% higher than in bovine milk (Ménard et al., 2010).  
75 Also, buffalo milk contains less cholesterol than bovine milk (7.0-10.2 mg/100mL vs. 10.5-  
76 19.8 mg/100 mL) (Strzalkowska, Jozwik, Baghnicka, Krzyzewski, & Horbanczuk, 2009;  
77 Talpur, Memom, & Bhangar, 2007; Zotos & Bampidis, 2014). The average content of  
78 membrane protein was reported to be less in buffalo MFGM compared to bovine milk, either  
79 when calculated as a percentage of fat (4.2% vs. 4.9%) or as a proportion of the total MFGM  
80 material (29.5% vs. 33.9%) (Abou-Dawood et al., 1988). A recent proteomic study identified  
81 50 proteins within buffalo MFGM, with the major proteins being as xanthine  
82 dehydrogenase/oxidase, butyrophilin, adipophilin, lactadherin and mucin, similar to bovine  
83 MFGM proteins (D'Ambrosio et al., 2008; Fong, Norris, & MacGibbon, 2007). Both the  
84 composition and the structure of the MFGM result from the mechanisms of secretion of fat  
85 globules from the epithelial cells of the mammary gland (Heid & Keenan, 2005). The  
86 compositional differences reported to date suggest there may be further underlying structural  
87 differences in the MFGM in buffalo milk and possible differences occurring during the *in vivo*  
88 secretion of these globules during milk production.

89 A number of studies performed using confocal microscopy have revealed heterogeneities in  
90 the organization of the MFGM (Evers et al., 2008), in the localization of membrane proteins  
91 (Lopez, Madec, & Jimenez-Flores, 2010; Lopez & Ménard, 2011) and the presence of lipid  
92 domains in the membranes of bovine (Gallier, Gragson, Jimenez-Flores, & Everett, 2010;  
93 Lopez et al., 2011; Lopez et al., 2010) and human milk (Lopez & Ménard, 2011; Zou et al.,

94 2012). This heterogeneous distribution has been attributed to the phase separation of polar  
95 lipids according to their temperature of phase transition ( $T_m$ ). The exogenous phospholipid  
96 fluorescently head-labelled with rhodamine (i.e. Rh-DOPE; PE containing 2 oleic acids with  
97 very low phase transition temperature) preferentially partitions in the fluid  $L_d$  phase  
98 containing unsaturated polar lipids (i.e. containing one or two long-chain unsaturated fatty  
99 acids) (Lopez et al., 2010). Hence, areas of the MFGM not stained by Rh-DOPE may  
100 correspond to gel phase domains rich in polar lipids with a high  $T_m$ . Polar lipids not stained  
101 by Rh-DOPE could adopt a second possible phase consisting of liquid ordered ( $L_o$ ) phase  
102 domains composed of high  $T_m$  lipids and cholesterol. These high  $T_m$  lipids include SM,  
103 which accounts for about 25% of polar lipids in the MFGM (Lopez, 2011) and contains long  
104 chain saturated fatty acids ( $T_m = 35\text{ }^\circ\text{C}$ ; Malmsten, Bergentahl, Nyberg, and Odham (1994)).  
105 Saturated PC are also high  $T_m$  lipids found in the MFGM (e.g. dipalmitoyl  
106 phosphatidylcholine, DPPC,  $T_m = 41.7\text{ }^\circ\text{C}$ ; Benesch and McElhaney (2014)). The  
107 surrounding membrane stained by Rh-DOPE may correspond to the fluid liquid disordered  
108 ( $L_d$ ) phase of the unsaturated glycerophospholipids (PC, PE, PI, PS) which have a low  $T_m$   
109 and are responsible for the fluidity of the biomembranes (Gallier et al., 2010; Lopez et al.,  
110 2010). A recent study performed by atomic force microscopy with a model membrane showed  
111 that milk SM segregates from the fluid  $L_d$  phase of unsaturated polar lipids to form domains  
112 in the gel phase (Guyomarc'h et al., 2014). This study also showed that cholesterol has a large  
113 effect on the domains formed by milk SM; the shapes and nanomechanical properties of the  
114 domains formed differ in the gel phase in the absence of cholesterol and in the  $L_o$  phase when  
115 cholesterol is present. The domains also protrude further from the fluid phase of the  
116 membrane in the absence of cholesterol (Guyomarc'h et al., 2014).

117 The lipid domains recently revealed in the MFGM could potentially be responsible for several  
118 biological processes (e.g. digestion of milk fat globules, interaction with pathogens and

119 viruses in the gut) and cellular functions. Also, the lipid domains present in outer bilayer of  
120 the MFGM could be involved in functional properties of milk fat globules, important for dairy  
121 applications.

122 The objective of this study was to perform a structural analysis of fat globules and their  
123 biological membrane, *in situ* in buffalo milk. The microstructure was investigated using the  
124 combination of optical microscopy with differential interferential contrast together with  
125 confocal laser scanning microscopy (CLSM) with adapted fluorescent dyes able to label  
126 TAG, total proteins, polar lipids and glycosylated molecules located in the MFGM.

127

## 128 **2. Materials and methods**

129

### 130 **2.1. Milk samples**

131 The buffalo milks used in this study were a mixture of the individual milks produced by 30  
132 buffaloes of the Mediterranean breed *Bubalus bubalis* and collected from evening and  
133 morning milking from Coopérative de Bufflonnes (Mauris, Cantal region, France). NaN<sub>3</sub>  
134 (0.02% w/v) was added to the milk to prevent the growth of bacteria. Milk samples were  
135 stored at ambient temperature before fat globule size measurements and CLSM experiments.  
136 The content of fat in the milk was determined using the Gerber method method (IDF, 2008).

### 137 **2.2 Fat globule size measurements**

138 **2.3** The fat globule size distributions were determined by laser light scattering, using a  
139 Mastersizer 2000 (Malvern Instruments, Malvern, U.K.) equipped with a He/Ne laser ( $\lambda =$   
140 633 nm) and an electroluminescent diode ( $\lambda = 466$  nm). The refractive indexes of milk fat

141 were set at 1.46 (at 466 nm) and 1.458 (at 633 nm) and the refractive index was set at 1.33  
142 for water. The absorption coefficient used was 0.0001. The experiments were performed  
143 at room temperature. Aliquots of approximately 70  $\mu$ L of whole milk were introduced into  
144 the measurement cell of the apparatus, which contained 100 ml of water, in order to reach  
145 10% obscuration (optimal conditions for particle-size measurements with this apparatus).  
146 A solution of 35 mM EDTA/NaOH pH 7.0 buffer (>98% disodium salt dihydrat, Prolabo,  
147 Fortenay-sous-Bois, France) was added to the measurement cell to disrupt the casein  
148 micelles. All analyses were performed in triplicate with three replicate milk samples  
149 collected from different days, giving the total of nine measurements. The specific surface  
150 area and volume-weighted average diameter  $d_{43}$ , defined as  $\frac{\sum n_i d_i^4}{\sum n_i d_i^3}$ , where  $n_i$  is the  
151 number of fat globules of diameter  $d_i$ , were calculated by the software. ***Microstructural***  
152 ***analysis***

153 Microstructural analysis was performed using an inverted microscope NIKON Eclipse-  
154 TE2000-C1si (NIKON, Champigny sur Marne, France). Confocal laser scanning microscopy  
155 (CLSM) was performed using an argon laser operating at an excitation wavelength of 488 nm  
156 with emission detected between 500 nm and 530 nm, a He-Ne laser operating at 543nm  
157 wavelength excitation with emission detected between 565 nm and 615 nm and a diode  
158 operating at 633 nm, detected with a long pass filter > 650 nm. The observations were  
159 performed using a  $\times 100$  (numerical aperture NA 1.4) oil immersion objective.

160 The staining protocols followed previously described methods (Lopez et al., 2010; Lopez &  
161 Ménard, 2011). Briefly, Nile Red (5H-Benzo  $\alpha$ -phenoxazine-5-one, 9-diethylamino, supplied  
162 by Sigma-Aldrich, St. Louis, USA) was prepared at a concentration of 42  $\mu$ g/mL in  
163 propanediol and used to stain the triacylglycerol core of the fat globules. Fast Green FCF  
164 (Sigma-Aldrich, St. Louis, USA) was prepared at a concentration of 10 mg/mL in water and

165 used to stain proteins. N-(Lissamine rhodamine B sulfonyl) dioleoylphosphatidyl  
166 ethanolamine (Rh-DOPE, Avanti polar lipids Inc., Birmingham, England) was provided at a  
167 concentration of 1 mg/mL in chloroform and used to label the phospholipids in the membrane  
168 surrounding buffalo milk fat globules. Wheat germ agglutinin Alexa fluor 488 (WGA488,  
169 Invitrogen, Cergy Pontoise, France) was prepared at a concentration of 1 mg/mL in phosphate  
170 saline buffer and used to label the glycosylated molecules in the membrane, i.e. carbohydrate  
171 moieties containing N-acetylglucosamine and N-acetyl neuraminic acid (sialic acid) residues.

172 The concentrations of the fluorescent dyes were adapted to the specific composition of  
173 buffalo milk. For observation of the fat globules, 100  $\mu$ L of Nile Red and 10  $\mu$ L of Fast Green  
174 FCF were added to 1 mL of buffalo milk samples. For observation of the MFGM polar lipids,  
175 40  $\mu$ L of the Rh-DOPE solution was placed in a glass vial and the chloroform was evaporated  
176 under nitrogen, to avoid the possible artefacts caused by this organic solvent. Then 1 mL of  
177 milk sample was introduced in the vial. For dual staining of the MFGM polar lipids and  
178 carbohydrate moieties, 10  $\mu$ L of the WGA-488 solution was also added into the vial  
179 containing Rh-DOPE. Then, 1 mL of milk sample was added to the vial. The stained milk  
180 samples were kept at room temperature for at least 1 h prior to observation by CLSM. The  
181 microstructural analyses were performed at room temperature ( $19 \pm 1$  °C)

182 Optical microscopy using differential interference contrast (DIC, also called Nomarski,  
183 Cogswell and Sheppard (1992)) was also used to characterize buffalo milk fat globules. DIC  
184 images were superimposed on the fluorescent emission recorded in the CLSM images. The  
185 two dimensional images had a resolution of 512 $\times$ 512 pixels and the pixel scale values were  
186 converted into micrometers using a scaling factor. In the multiple labeled samples, different  
187 colors were used to locate the fluorescent probes (e.g. red color for phospholipids stained by

188 Rh-DOPE, green color for the carbohydrate moieties stained by WGA-488 and grey levels for  
189 DIC images).

### 190 **3. Results and discussion**

#### 191 *3.1. Size distribution of buffalo milk fat globules*

192 The buffalo milks used in this study contained  $74.1 \pm 5.8$  g fat per kg milk, in agreement with  
193 previous studies (Ahmad et al., 2008; Ménard et al., 2010; Varricchio et al., 2007). The size  
194 distribution of the fat globules, determined using light scattering, ranged from 0.4  $\mu\text{m}$  to 15.8  
195  $\mu\text{m}$ , with a mean volume-weighted diameter of  $5.0 \pm 0.4$   $\mu\text{m}$  (Figure 1A). The surface of fat  
196 globules covered by the MFGM was  $2.0 \pm 0.2$   $\text{m}^2/\text{g}$  fat, which corresponded to about 148  $\text{m}^2$   
197 per kg milk. The biological membrane surrounding fat globules represents a large interface  
198 between the TAG core of fat globules and the aqueous environment, confirming the  
199 importance of its characterization. These size parameters are consistent with previous reports  
200 (Ahmad et al., 2008; Ménard et al., 2010).

201 The fat globules, observed by the combination of differential interference contrast (DIC) light  
202 microscopy and CLSM after staining with Nile Red, were spherical in shape and dispersed in  
203 the aqueous phase of milk containing proteins (Figure 1B and D). A rough surface could be  
204 observed on some fat globules (indicated by arrows in Figure 1C) and may arise from surface  
205 heterogeneities or partial crystallisation of the TAG core. The overlay of DIC and fluorescent  
206 CLSM images showed that the triacylglycerols are located in the core of fat globules (Figure  
207 1D), similar to previous observations for bovine and human milk (Lopez et al., 2010; Lopez  
208 & Ménard, 2011; Zou et al., 2012).

### 209 ***3.2 Cytoplasmic remnants attached to buffalo milk fat globules***

210 Cytoplasmic remnants (CR) from lactating cells were observed to be connected to some  
211 buffalo fat globules of various sizes, using optical microscopy and fluorescence microscopy  
212 (Figure 2). The volume of CR varied from one globule to another and their thickness was 1  
213  $\mu\text{m}$  up to 5  $\mu\text{m}$ , as determined in the fluorescence images (Figure 2). The CR can exceed the  
214 buffalo milk fat globule volume (Figure 2 D). The morphology of CR can be compared to a  
215 crescent attached to fat globules, consistent with another name for these remnants of  
216 cytoplasmic crescents. The majority of the buffalo fat globules did not contain CR and were  
217 enveloped compactly by the MFGM (Figure 2 D and E).

218 CR occur when the mechanisms involved in the closure of the apical plasma membrane  
219 behind the projecting lipid droplet lead to their secretion with a piece of cytoplasm remaining  
220 attached to fat globule in milk (Heid & Keenan, 2005). The occurrence of the CR has also  
221 been reported in previous CLSM studies of human milk (Evers et al., 2008; Lopez & Ménard,  
222 2011) but CR have not been observed in bovine milk (Lopez et al., 2010) when the same  
223 CLSM methods and fluorescent probes have been applied. This is probably due to variations  
224 in CR concentration between the milk of different species, arising from differences in the  
225 membrane secretion process, which is poorly understood (Robenek et al., 2005). CR were  
226 observed for 7.2% of the fat globules within human milk but for less than 1% of the fat  
227 globules in bovine milk, quantified using microscopic observation with the assistance of a  
228 counting chamber slide (Huston & Patton, 1990). In goat milk, between 1% and 5% of milk  
229 fat globules had CR (Wooding, Peaker, & Linzell, 1970). The current study is the first to  
230 report the presence of CR in buffalo milk but the number of CR was not quantified, which  
231 may be worthy of further investigation.

232 Previous authors who observed CR in human milk employed Rh-DOPE or DiIC<sub>18</sub>(3)-DS to  
233 label the MFGM (Evers et al., 2008; Lopez & Ménard, 2011). DiIC<sub>18</sub>(3)-DS is a lipophilic  
234 fluorescent probe that can penetrate the membrane by inserting its two acyl chains into the  
235 interior of the phospholipid membrane (Evers et al., 2008). The labelling of the CR by Rh-  
236 DOPE or DiIC<sub>18</sub>(3)-DS suggests that these remnants are rich in phospholipid, consistent with  
237 their origin as fragments of the cytoplasm or apical plasma membrane (Heid & Keenan,  
238 2005). The size of CR has been observed to range from a thin sliver to a size exceeding the fat  
239 globule core (Heid & Keenan, 2005). CR contain all organelles and intracellular membrane  
240 materials (except for nuclei) of the lactating cells (Heid & Keenan, 2005; Wooding et al.,  
241 1970) and thus, these CR may be a good source of enzymes, micronutrients and growth  
242 factors necessary for newborn development (Huston & Patton, 1990). RNA has also been  
243 successfully isolated from the CR of human milk (Lemay et al., 2013; Maningat et al., 2009),  
244 suggesting CR could impact on protein expression in host cells, similar to other RNA  
245 containing membrane vesicles found in milk and other body fluids (Valadi et al., 2007; Zhou  
246 et al., 2012). CR could similarly be examined as potential markers of milk quality.

### 247 ***3.3 Heterogeneous distribution of polar lipids in buffalo MFGM***

248 The fluorescent dye Rh-DOPE was used to assess the distribution of polar lipids in the  
249 MFGM surrounding buffalo fat globules, *in situ* in buffalo milk at  $19 \pm 1$  °C, as already  
250 performed for bovine and human MFGM (Gallier et al., 2010; Lopez et al., 2010; Lopez &  
251 Ménard, 2011; Zou et al., 2012).

252 Rh-DOPE integrated the buffalo MFGM and appeared fluorescent at the periphery of the fat  
253 globules viewed as an equatorial section by CLSM (Figure 3, Figure 4 A and C). The interior  
254 of buffalo milk fat globules, which is mainly composed of triacylglycerols, as shown in  
255 Figure 1, was not labelled by Rh-DOPE (Figure 3 and Figure 4 C). In some cases, the entire

256 periphery of fat globules was fluorescent, showing the homogeneous distribution of the  
257 fluorescent dye, while in other cases, both fluorescent and non-fluorescent areas were visible,  
258 indicating the partial labelling of the MFGM by the exogenous phospholipid Rh-DOPE. Three  
259 dimensional observations (i.e. recording of thin optical sections at different z-depths of milk  
260 samples) allowed the characterisation of non-fluorescently-labelled domains at the surface of  
261 milk fat globules composed by the MFGM (noted Z-depth 1 in Figure 4A) and of fluorescent  
262 rings seen in an equatorial cross section of the fat globule (noted Z-depth 2 in Figure 4B).  
263 These non-fluorescent areas were mostly circular in shape with a size of approximately 0.5 to  
264 2  $\mu\text{m}$  and a number generally ranging from zero to nine in one side of fat globule (Figures 4).  
265 Differences in size and shape of the non-fluorescent areas, however, were also observed  
266 (Figure 4). For example, Figure 4 shows fat globules with circular non-fluorescent areas of  
267 size ranging from very small (Figure 4 B) to very large (Figure 4 E), while fat globules with  
268 irregular shaped non-fluorescent regions have also been characterized (Figure 4 F, G, H). The  
269 irregular shapes could result from the connection of several circular domains at the surface of  
270 fat globules or to different physical states of the lipids in the domains. Non-fluorescent  
271 domains not stained by the Rh-DOPE can diffuse in the plane of the membrane as a function  
272 of time, as observed at the surface of a fat globule (highlighted by arrows in Figure 5). This  
273 result shows that the MFGM is a highly dynamic system, as previously reported for the  
274 bovine MFGM (Lopez et al., 2010).

### 275 ***3.4. Simultaneous localisation of polar lipids, glycoproteins and glycolipids in buffalo*** 276 ***MFGM***

277 The organisation of lipids and the localisation of the MFGM glycoproteins and glycolipids, *in*  
278 *situ* around buffalo milk fat globules, was investigated by CLSM with the simultaneous  
279 labelling of phospholipids with Rh-DOPE that stain the Ld phase and glycosylated molecules

280 that are specifically probed using the fluorescently labelled lectin wheat germ agglutinin  
281 (WGA-488). DIC optical microscopy in combination with CLSM permitted the observation  
282 of fat globules and the localisation of the components in the MFGM surrounding fat globules.

283 Figure 6 shows the heterogeneous lateral organisation of lipids and glycoproteins in the  
284 MFGM. Figure 6 A focuses on the top view of a fat globule (noted Z-depth 1) of about 10  $\mu\text{m}$   
285 diameter, while Figure 6 B-D shows an equatorial cross section of the same fat globule (at a  
286 higher Z-depth, noted Z-depth 2), with TAG in the core and the MFGM at the periphery. The  
287 lectin stain WGA-488 is reported to bind to N-acetylglucosamine and N-acetylneuraminic  
288 acid residues (Evers et al., 2008; Lopez et al., 2010), including the glycolipids such as  
289 gangliosides and the main glycoproteins of the MFGM such as butyrophilin, mucin and  
290 lactadherin (Evers et al., 2008; Heid & Keenan, 2005; Lopez et al., 2010). The distribution of  
291 these glycosylated molecules is non-homogeneous occurring in patches or networks in some  
292 areas of the MFGM of buffalo milk fat globules (Figure 6 and 7).

293 The co-localisation of polar lipids and glycosylated molecules reveals that most of the non-  
294 fluorescent areas observed when the MFGM was labelled with Rh-DOPE were also not  
295 fluorescent after labelling with WGA-488 (Figure 6). This is an additional argument in favour  
296 of the interpretation of the domains in which Rh-DOPE did not integrate, as being domains  
297 rich in SM and DPPC (i.e. high  $T_m$  lipids that are not glycosylated). Some exceptions were  
298 observed, however, where areas that were not stained by Rh-DOPE but stained by WGA-488  
299 (areas indicated by arrows in Figure 7), indicating that some glycoproteins and/or glycolipids  
300 are located in the high  $T_m$  lipid rich domains. Also, some glycosylated molecules were  
301 observed at the periphery of the domains not labelled by Rh-DOPE (Figure 8). These results  
302 suggest that the glycosylated molecules are mostly present in the fluid phase of the MFGM  
303 but they can also participate in the high  $T_m$  lipid domains at times. Figure 7 E shows that

304 glycosylated molecules protrude in the aqueous phase surrounding buffalo milk fat globules,  
305 and contribute in the formation of a glycocalyx.

306 The heterogeneous distribution of glycosylated molecules has also been reported in  
307 previous studies for bovine, sheep and human milk (Evers et al., 2008; Gallier et al., 2010;  
308 Lopez et al., 2011; Lopez et al., 2010). Non-fluorescent regions have been observed within  
309 human and bovine MFGM when dual-staining with Rh-DOPE and WGA-488 has been  
310 applied (Gallier et al., 2010; Lopez et al., 2011; Lopez et al., 2010; Lopez & Ménard, 2011)  
311 and these non-fluorescent areas were attributed to SM-rich domains. Non-fluorescent areas  
312 were also observed within the bovine, sheep and human MFGM when the milk was single  
313 stained with WGA488 or when dual stained with WGA488 and the lipophilic probe FMN-64  
314 (Evers et al., 2008), although these studies interpreted these non-fluorescent areas as the  
315 regions where the MFGM is absent. The lateral organisation of polar lipids and proteins in the  
316 MFGM could be involved in specific functions (including the mechanisms of lipid digestion  
317 or protection of the neonates from infections), which currently remain unknown and require  
318 further investigation.

### 319 ***3.5. The MFGM: a non-random mixing of components***

320 We observed that the MFGM is a non-random mixture of components characterised by the  
321 lateral separation of compositionally distinct lipid phases. The occurrence of both fluorescent  
322 and non-fluorescent areas observed with Rh-DOPE staining was interpreted as a  
323 heterogeneous distribution of the polar lipids within the buffalo MFGM. Phase separation  
324 would result in preferential partition of select lipids in the coexisting phases as a function of  
325 their chemical and physical properties. Such heterogeneities in the partitioning of Rh-DOPE  
326 stained lipids have been previously observed in the human and bovine MFGM (Lopez et al.,  
327 2011; Lopez et al., 2010; Lopez & Ménard, 2011; Zou et al., 2012).

328 The non-fluorescent areas characterised at room temperature for the buffalo MFGM could  
329 correspond to the preferential tight packing of polar lipids with a high gel to liquid crystalline  
330 phase transition temperature (high  $T_m$ ), leading to their lateral segregation from very low  $T_m$   
331 polar lipids in the liquid phase of the MFGM. These high  $T_m$  polar lipids consist of long-  
332 chain saturated fatty acids (i.e. C16:0, P; C18:0, S; and fatty acids with a number of carbon >  
333 20). According to the fatty acid composition of polar lipids in the MFGM (Sanchez-Juanes,  
334 Alonso, Jancada, & Hueso, 2009) and the preferential concentration of PC and SM in the  
335 outer layer of the MFGM (Deeth, 1997), these polar lipids could be SM (the main fatty acids  
336 being C16:0, C20:0, C22:0, C23:0, C24:0; Lopez (2011)) and PC (i.e. DPPC, PSPC or SSPC).  
337 Under the conditions examined ( $T=19\text{ }^\circ\text{C}$ ), these milk SM and saturated PC molecular species  
338 are expected to be in the gel phase, as they have a  $T_m > 19\text{ }^\circ\text{C}$ . Such high  $T_m$  polar lipids can  
339 also segregate together with cholesterol in the plane of the biological membranes to form  
340 ordered lipid domains in the  $L_o$  phase (Brown & London, 1998). The role played by  
341 cholesterol in the MFGM is important, since a minimum molar percentage of cholesterol is  
342 necessary to form the  $L_o$  phase with high  $T_m$  lipids, as reported in phase diagrams built with  
343 controlled lipid compositions (Veatch & Keller, 2005). The coexistence of two ordered phases  
344 (i.e. the  $L_o$  phase domains including high  $T_m$  lipids and cholesterol and the gel phase  
345 domains composed only by high  $T_m$  lipids) with one fluid phase composed by unsaturated  
346 polar lipids may occur, as already discussed for ternary model systems (Veatch & Keller,  
347 2005). The ordered phases (gel and  $L_o$  phases) are immiscible with the MFGM fluid phase  
348 and form micron-scale domains, permitting their observation by confocal microscopy. The  
349 circular shape of the domains observed in the buffalo MFGM at room temperature is  
350 consistent with the formation of  $L_o$  phase domains, since polar lipids organised in a gel phase  
351 are semi-crystallised and form angles (Brown & London, 1998; Morales-Pennington et al.,  
352 2011). Non-circular domains present in the outer bilayer of MFGM have been previously

353 reported at room temperature for human milk (Lopez & Ménard, 2011) but have not been  
354 observed for the bovine MFGM (Lopez et al., 2010). This difference in the morphology of the  
355 lipid domains observed in the MFGM from various species could possibly be due to the  
356 differences in the composition of the polar lipids such as the concentration and individual  
357 lipid species with high  $T_m$  (for example SM and DPPC) and the relative proportion of low  
358  $T_m$  lipids, high  $T_m$  lipids and cholesterol (Veatch & Keller, 2005). The relative proportion of  
359 SM, which is assumed to be the major component of the  $L_o$  phase domains (Gallier et al.,  
360 2010; Lopez et al., 2010), is lower in the buffalo MFGM compared to the bovine MFGM  
361 (24.8% vs. 26.9% of polar lipid) (Ménard et al., 2010). It has been reported that the  
362 morphology of the SM-rich domains and the composition of the bovine MFGM could be  
363 linked to fat globule size (Gallier et al., 2010; Lopez et al., 2011). Although some studies  
364 have reported that small bovine fat globules appear to have fewer domains compared to large  
365 fat globules (Gallier et al., 2010), others did not find any significant differences arising from  
366 the characterisation of domains in small fat globules (diameter < 2  $\mu\text{m}$ ) (Lopez et al., 2011).  
367 The lipid composition of the outer bilayer of the MFGM could be considered as a ternary  
368 system consisting of high  $T_m$  polar lipids (e.g. SM and saturated PC), cholesterol and low  $T_m$   
369 polar lipids (e.g. unsaturated glycerophospholipids) with differences in membrane structure  
370 and dynamics arising from compositional differences between species.

371

372 The organisation of lipids in the MFGM can govern the functional properties of this  
373 biomembrane, affect the localisation of membrane-associated proteins and the chemical  
374 reactions that occur in the MFGM or that are mediated by this membrane. Lipid phase  
375 separation could also influence the binding of molecules to the MFGM, such as bacteria and  
376 viruses (Lopez, 2011). The MFGM plays a key role in the physical stability of fat globules in  
377 milk including protection against coalescence and aggregation. Buffalo milk fat globules are

378 important in the preparation of dairy products such as yoghurt and cheese. Hence, the lipid  
379 domains formed in the MFGM, could be involved in a number of functions, warranting  
380 further investigation of these microstructural features.

381

## 382 **4. Conclusion**

383 Buffalo fat globules and their biological membrane are poorly characterised despite the  
384 importance of buffalo milk to human nutrition and dairy processing. This study investigated  
385 the microstructure of the buffalo MFGM improving our knowledge of this dynamic and  
386 complex biological membrane. We showed that the MFGM is heterogeneous with respect to  
387 lipid, cholesterol and protein assemblies. Confocal experiments revealed that polar lipids  
388 segregated in the outer bilayer of the MFGM into two or more liquid phases (e.g. gel, Lo and  
389 Ld phases) that can coexist and have different physical properties, with gel and Ld phases  
390 being most probable. Glycoproteins and glycolipids were mainly present in the Ld fluid phase  
391 but also existed in the lipid domains (gel or Lo phases) at times. Whether or not the lipid  
392 domains in buffalo MFGM are lipid rafts remains unknown and would be worthy of further  
393 study.

394

## 395 **Acknowledgements**

396 Joëlle Léonil, the head of INRA STLO (Rennes, France), is acknowledged for hosting Hanh  
397 Nguyen (PhD student). The authors thank Joël Guillemin (Coopérative de Bufflonnes, Maurs,  
398 France) for providing buffalo milks. The authors also acknowledge The Australian  
399 Government including The ARC Dairy Innovation Hub, The Rural Industries Research and  
400 Development Cooperation (RIRDC), The University of Melbourne, The Bio21 Molecular

401 Science and Biotechnology Institute, The Particulate Fluids Processing Centre (PFPC) and  
402 The Clive Pratt Family for financial support.

403

404 All authors have approved the final article.

405

## 406 **References**

407 Abou-Dawood, A. E., Moussaa, A. E., El-Demerdash, O., & Ahmed, R. B. (1988).  
408 Comparative study on fat globule membrane of buffalo, cow, goat and sheep raw milk.  
409 *Annals of Agricultural Science*, 26(4), 2527-2536.

410 Ahmad, S., Gaucher, I., Rousseau, F., Beaucher, E., Piot, M., Grongnet, J. F., & Gaucheron,  
411 F. (2008). Effects of acidification on physico-chemical characteristics of buffalo milk:  
412 A comparison with cow's milk. *Food Chemistry*, 106(1), 11-17.

413 Benesch, M. G. K., & McElhaney, R. N. (2014). A comparative calorimetric study of the  
414 effects of cholesterol and the plant sterols campesterol and brassicasterol on the  
415 thermotropic phase behavior of dipalmitoylphosphatidylcholine bilayer membranes.  
416 *Biochimica et Biophysica Acta (BBA) - Biomembranes*, 1838(7), 1941-1949.

417 Brown, D. A., & London, E. (1998). Functions of lipid rafts in biological membranes. *Annual*  
418 *Review of Cell and Developmental Biology*, 14, 111-136.

419 Cogswell, C. J., & Sheppard, C. J. R. (1992). Confocal differential interference contrast (DIC)  
420 microscopy: Including a theoretical analysis of conventional and confocal DIC  
421 imaging. *Journal of Microscopy*, 165(1), 81-101.

422 D'Ambrosio, C., Arena, S., Salzano, A. M., Renzone, G., Ledda, L., & Scaloni, A. (2008). A  
423 proteomic characterization of water buffalo milk fractions describing PTM of major  
424 species and the identification of minor components involved in nutrient delivery and  
425 defense against pathogens. *Proteomics*, 8, 3657-3666.

426 Deeth, H. C. (1997). The role of phospholipids in the stability of milk fat globules. *Australian*  
427 *Journal of Dairy Technology*, 52(1), 44-46.

428 Dewettinck, K., Rombaut, R., Thienpont, N., Le, T. T., Messens, K., & Camp, J. V. (2008).  
429 Nutritional and technological aspects of milk fat globule membrane material.  
430 *International Dairy Journal*, 18, 436-457.

431 Evers, J. M., Haverkamp, R. G., Holroyd, S. E., Jameson, G. B., Mackenzie, D. D. S., &  
432 McCarthy, O. J. (2008). Heterogeneity of milk fat globule membrane structure and  
433 composition as observed using fluorescence microscopy techniques. *International*  
434 *Dairy Journal*, 18(12), 1081-1089.

- 435 FAOSTAT. (2012). Production quantity of livestock primary 2012. Food and Agriculture  
436 Organisation of the United Nations - Statistics Division. Retrieved 23/10/2014 from  
437 <http://faostat3.fao.org/download/Q/QL/E>.
- 438 Fong, B. Y., Norris, C. S., & MacGibbon, A. K. H. (2007). Protein and lipid composition of  
439 bovine milk-fat-globule membrane. *International Dairy Journal*, 17, 275-288.
- 440 Gallier, S., Gragson, D., Jimenez-Flores, R., & Everett, D. (2010). Using confocal laser  
441 scanning microscopy to probe the milk fat globule membrane and associated proteins.  
442 *Journal of Agricultural and Food Chemistry*, 58(7), 4250-4257.
- 443 Guyomarc'h, F., Zou, S., Chen, M., Milhiet, P. E., Godefroy, C., Vié, V., & Lopez, C. (2014).  
444 Milk sphingomyelin domains in biomimetic membranes and the role of cholesterol:  
445 morphology and nanomechanical properties investigated using AFM and force  
446 spectroscopy. *Langmuir*, 30(22), 6516-6524.
- 447 Heid, H. W., & Keenan, T. W. (2005). Intracellular origin and secretion of milk fat globules.  
448 *European Journal of Cell Biology*, 84(2-3), 245-258.
- 449 Huston, G. E., & Patton, S. (1990). Factors related to the formation of cytoplasmic crescents  
450 on milk fat globules. *Journal of Dairy Science*, 73(8), 2061-2066.
- 451 IDF. (2008). *Milk - Determination of fat content (Gerber method)*. (International standard  
452 ISO-IDF 2446:2008). Brussels, Belgium: International Dairy Federation.
- 453 Keenan, T. W., & Patton, S. (1995). The structure of milk: implications for sample and  
454 storage. A. The milk lipid globule membrane. In R. G. Jensen (Ed.), *Handbook of milk*  
455 *composition* (pp. 5-49). San Diego, CA, USA: Academic Press.
- 456 Lemay, D. G., Ballard, O. A., Hughes, M. A., Morrow, A. L., Horseman, N. D., & Nommsen-  
457 Rivers, L. A. (2013). RNA sequencing of the human milk fat layer transcriptome  
458 reveals distinct gene expression profiles at three stages of lactation. *Plosone*, 8(7). doi:  
459 10.1371/journal.pone.0067531. Retrieved 25/09/2014 from  
460 <http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0067531>.
- 461 Lopez, C. (2011). Milk fat globules enveloped by their biological membrane: Unique  
462 colloidal assemblies with a specific composition and structure. *Current Opinion in*  
463 *Colloid & Interface Science*, 16(5), 391-404.
- 464 Lopez, C., Briard-Bion, V., Menard, O., Beaucher, E., Rousseau, F., Fauquant, J., . . . Robert,  
465 B. (2011). Fat globules selected from whole milk according to their size: Different  
466 compositions and structure of the biomembrane, revealing sphingomyelin-rich  
467 domains. *Food Chemistry*, 125(2), 355-368.
- 468 Lopez, C., Madec, M. N., & Jimenez-Flores, R. (2010). Lipid rafts in the bovine milk fat  
469 globule membrane revealed by the lateral segregation of phospholipids and  
470 heterogeneous distribution of glycoproteins. *Food Chemistry*, 120(1), 22-33.
- 471 Lopez, C., & Ménard, O. (2011). Human milk fat globules: Polar lipid composition and in situ  
472 structural investigations revealing the heterogeneous distribution of proteins and the  
473 lateral segregation of sphingomyelin in the biological membrane. *Colloids and*  
474 *Surfaces B: Biointerfaces*, 83(1), 29-41.

- 475 Malmsten, M., Bergentahl, B., Nyberg, L., & Odham, G. (1994). Sphingomyelin from milk -  
 476 characterisation of liquid crystalline, liposome and emulsion properties. *Journal of the*  
 477 *American Oil Chemists Society*, 91(9), 1021-1026.
- 478 Maningat, P. D., Sen, P., Rijnkels, M., Sunehag, A. L., Hadsell, D. L., Bray, M., & Haymond,  
 479 M. W. (2009). Gene expression in the human mammary epithelium during lactation:  
 480 the milk fat globule transcriptome. *Physiological Genomics*, 37(1), 12-22.
- 481 Ménard, O., Ahmad, S., Rousseau, F., Briard-Bion, V., Gaucheron, F., & Lopez, C. (2010).  
 482 Buffalo vs. cow milk fat globules: Size distribution, zeta-potential, compositions in  
 483 total fatty acids and in polar lipids from the milk fat globule membrane. *Food*  
 484 *Chemistry*, 120(2), 544-551.
- 485 Morales-Pennington, N. F., Wu, J., Farkas, E. R., Goh, S. L., Konyakhina, T. M., Zheng, J.  
 486 Y., . . . Feigenson, G. W. (2011). GUV preparation and imaging: minimizing artifacts.  
 487 *Biochimica et Biophysica Acta*, 1798(7), 1324-1332.
- 488 Robenek, H., Robenek, J. M., Buers, I., Lorkowski, S., Hofnagel, O., Troyer, D., & Severs, N.  
 489 J. (2005). Lipid droplets gain PAT family proteins by interaction with specialised  
 490 plasma membrane domains. *The Journal of Biological Chemistry*, 280, 26330-26338.
- 491 Sanchez-Juanes, F., Alonso, J. M., Jancada, L., & Hueso, P. (2009). Distribution and fatty  
 492 acid content of phospholipids from bovine milk and bovine milk fat globule  
 493 membranes. *International Dairy Journal*, 19, 273-278.
- 494 Solh, V. A., Staines, V., Honda, S., & Limley, H. A. (2007). Measurement of milk color and  
 495 composition: Effect of dietary intervention on western Australian Holstein-Friesian  
 496 cow's milk quality. *Journal of Food Science*, 72(8), 560-566.
- 497 Strzalkowska, N., Jozwik, A., Baghnicka, E., Krzyzewski, J., & Horbanczuk, J. O. (2009).  
 498 Studies upon genetic and environmental factors affecting the cholesterol content of  
 499 cow milk. I. Relationship between the polymorphic form of beta-lactoglobulin,  
 500 somatic cell count, cow age and stage of lactation and cholesterol content of milk.  
 501 *Animal Science Papers and Reports*, 27(2), 95-103.
- 502 Talpur, F. N., Memom, N. N., & Bhangar, M. I. (2007). Comparison of fatty acid and  
 503 cholesterol content of Pakistani water buffalo breeds. *Pakistan Journal of Analytical*  
 504 *and Environmental Chemistry*, 8, 15-20.
- 505 Valadi, H., Ekstrom, K., Bossios, A., Sjostrand, M., Lee, J. J., & Lotvall, J. O. (2007).  
 506 Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of  
 507 genetic exchange between cells. *Nature Cell Biology*, 9(6), 654-659.
- 508 Varricchio, M. L., Di Francia, A., Masucci, F., Romano, R., & Proto, V. (2007). Fatty acid  
 509 composition of Mediterranean buffalo milk fat. *Italian Journal of Animal Science*, 6,  
 510 509-511.
- 511 Veatch, S. L., & Keller, S. L. (2005). Seeing spots: Complex phase behavior in simple  
 512 membranes. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*,  
 513 1746(3), 172-185.

- 514 Wooding, F. B. P., Peaker, M., & Linzell, J. L. (1970). Theories of milk secretion : evidence  
515 from the electron microscopic examination of milk. *Nature*, 226(5247), 762-764.
- 516 Zhou, Q., Li, M., Wang, X., Li, Q., Wang, T., Zhu, Q., . . . Li, X. (2012). Immune-related  
517 microRNAs are abundant in breast milk exosomes. *International Journal of Biological*  
518 *Sciences*, 8(1), 118-123.
- 519 Zotos, A., & Bampidis, V. A. (2014). Milk fat quality of Greek buffalo (*Bubalus bubalis*).  
520 *Journal of Food Composition and Analysis*, 33(2), 181-186.
- 521 Zou, X. Q., Guo, Z., Huang, J. H., Jin, Q. Z., Cheong, L. Z., Wang, X. G., & Xu, X. B.  
522 (2012). Human milk fat globules from different stages of lactation: a lipid composition  
523 analysis and microstructure characterisation. *Journal of Agricultural and Food*  
524 *Chemistry*, 60, 7158-7167.

525

526

527

528

## 529 **FIGURE CAPTION**

530

531 **Figure 1:** Size distribution of buffalo milk fat globules determined using light scattering (A)  
532 in combination with microscopy (B-D). (B) CLSM image with Nile Red labeled fat globule  
533 cores appearing red and Fast Green FCF labeled protein appearing green, (C) Image taken  
534 with differential interferential contrast and (D) overlay of images A and B. The scale bars are  
535 10  $\mu\text{m}$  in length. Arrows indicate fat globules with a rough surface.

536

537 **Figure 2:** Microscopy images showing the presence of cytoplasmic remnants (CR) attached  
538 to buffalo milk fat globules. (A) DIC image, (B) CLSM image of fat globules stained with  
539 Rh-DOPE, (C) overlay of images A and B, (D and E) CLSM images with Rh-DOPE. Images  
540 were captured using a 100x objective. The scale bars are indicated in the images. Thick  
541 arrows indicate cytoplasmic remnant, thin arrows show the non-fluorescent domain in the  
542 biological membrane surrounding fat globules. TAG core = triacylglycerol core of milk fat  
543 globules.

544

545 **Figure 3:** Microscopy images showing an equatorial cross-section of fat globules and the  
546 heterogeneous distribution of polar lipids within the buffalo MFGM surrounding the milk fat

547 globules. **(A)** CLSM image after labelling with Rh-DOPE (red colour), **(B)** overlay of DIC  
548 optical microscopy image and CLSM image. The non-fluorescent areas (dark areas) observed  
549 around the fat globules correspond to the lateral segregation of high phase transition  
550 temperature polar lipids in domains (indicated by the white arrows). Images were captured at  
551 room temperature ( $19 \pm 1^\circ\text{C}$ ) using a 100x objective. The scale bars are indicated in the  
552 figures.

553

554 **Figure 4:** Microscopy images showing the heterogeneous distribution of polar lipids within  
555 the buffalo MFGM. (A, B, C) CLSM images after labelling with Rh-DOPE (red colour), (D-  
556 H) overlay of DIC optical microscopy image and CLSM image. The non-fluorescent areas  
557 (dark areas) observed around the fat globules correspond to the lateral segregation of high  
558 phase transition temperature polar lipids in domains (indicated by the white arrows). Images  
559 were captured at room temperature ( $19 \pm 1^\circ\text{C}$ ) using a 100x objective. The scale bars are  
560 indicated in the figures.

561

562 **Figure 5:** CLSM images taken at room temperature showing the diffusion of a non-  
563 fluorescent lipid domain within the MFGM as a function of time (A)  $t = 0$  sec, (B)  $t = 30$  sec.  
564 The polar lipids within the MFGM were stained using Rh-DOPE (red colour). The scale bars  
565 are indicated in the figures.

566

567 **Figure 6:** Microscopy images showing the heterogeneous distribution of polar lipids and  
568 glycosylated molecules (glycoproteins and glycolipids) in the buffalo milk fat globule  
569 membrane characterised at different z-depths. In-situ observations performed by the  
570 combination of DIC optical microscopy and CLSM, using Rh-DOPE to label the  
571 phospholipids (red colour) and the lectin WGA-488 to stain the glycosylated molecules (green  
572 colour). **(A)** overlay of DIC and fluorescence (Rh-DOPE and WGA) at z-depth 1  
573 corresponding to the polar region of fat globules, **(B)** overlay of DIC and fluorescence (Rh-  
574 DOPE and WGA-488) at z-depth 2 corresponding to the equatorial section of fat globules, **(C)**  
575 overlay of DIC and CLSM image with WGA-488, **(D)** overlay of DIC and CLSM image with  
576 Rh-DOPE. Images were captured using a 100x objective. The scale bars are indicated in the  
577 figures. White arrows correspond to non-fluorescent domains, green arrows correspond to  
578 glycosylated molecules located at the periphery or within a domain in which Rh-DOPE did  
579 not integrate.

580

581 **Figure 7:** Microscopy images showing the heterogeneous distribution of polar lipids and  
582 glycosylated molecules (glycoproteins and glycolipids) in the buffalo milk fat globule  
583 membrane. In-situ observations performed by the combination of DIC optical microscopy  
584 and CLSM, using Rh-DOPE to label the phospholipids (red colour) and the lectin WGA-488  
585 to stain the glycosylated molecules (green colour). **(A)** DIC image, **(B)** overlay of DIC and  
586 fluorescence with Rh-DOPE, **(C)** overlay of DIC image and fluorescence with WGA-488, **(D)**  
587 overlay of DIC image with Rh-DOPE and WGA-488, **(E)** equatorial cross section of the same  
588 fat globule showing the overlay of DIC image with Rh-DOPE and WGA-488. Images were  
589 captured using a 100x objective. The scale bars are indicated in the figures. White arrows  
590 correspond to non-fluorescent domains, green arrows correspond to glycosylated molecules  
591 located at the periphery or within a domain in which Rh-DOPE did not integrate.

592

593

594 **Figure 8:** Microscopy images showing the heterogeneous distribution of polar lipids and  
595 glycosylated molecules (glycoproteins and glycolipids) in the buffalo milk fat globule  
596 membrane. In-situ observations performed by the combination of DIC optical microscopy  
597 and CLSM, using Rh-DOPE to label the phospholipids (red colour) and the lectin WGA-488  
598 to stain the glycosylated molecules (green colour). **(A)** overlay of DIC image with Rh-DOPE  
599 and WGA-488, **(B)** overlay of DIC image with fluorescence with WGA-488, **(C)** overlay of  
600 DIC image and fluorescence with Rh-DOPE, **(D)** DIC image. Images were captured using a  
601 100x objective. The scale bars are indicated in the figures. White arrows correspond to  
602 glycosylated molecules located at the periphery or within a domain in which Rh-DOPE did  
603 not integrate.

604

605

606


607

## 608 FIGURES

609

610 **Figure 1**

611




612

613

614 **Figure 2**

615




616

617

618

619 **Figure 3**

620




621

622

623

624 **Figure 4**

625




626

627


628

629 **Figure 5**




633 **Figure 6**

634



644 **Figure 7**

645




646

647

648

649 **Figure 8**

650



651