

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Ong, WL;Foroudi, F;Evans, S;Millar, J

Title:

Androgen deprivation therapy use with post-prostatectomy radiotherapy in the Prostate Cancer Outcomes Registry Victoria

Date:

2019-02-01

Citation:

Ong, W. L., Foroudi, F., Evans, S. & Millar, J. (2019). Androgen deprivation therapy use with post-prostatectomy radiotherapy in the Prostate Cancer Outcomes Registry Victoria. *Journal of Medical Imaging and Radiation Oncology*, 63 (1), pp.124-130. <https://doi.org/10.1111/1754-9485.12818>.

Persistent Link:

<https://hdl.handle.net/11343/284590>

Androgen deprivation therapy (ADT) use with post-prostatectomy radiotherapy (PPRT) in the Prostate Cancer Outcomes Registry Victoria (PCOR-Vic)

Wee Loon Ong^{1,2,3,4}, Farshad Foroudi¹, Sue Evans³, Jeremy Millar^{5,6}

¹Department of Radiation Oncology, Olivia Newton-John Cancer and Wellness Centre, Austin Health

²Department of Radiation Oncology, Peter MacCallum Cancer Centre

³Department of Epidemiology and Preventive Medicine, Monash University

⁴Health and Biomedical Informatics Centre, University of Melbourne

⁵Alfred Health Radiation Oncology Services

⁶Central Clinical School, Monash University

Abstract word count: 250 words

Manuscript word count: 1683 words

Number of tables: 2

Number of figures: 1

Running title: ADT use with post-prostatectomy RT

Keywords: prostate cancer, androgen deprivation, post-prostatectomy, radiotherapy, registry

Corresponding Author:

Dr Wee Loon Ong
Department of Radiation Oncology
Olivia Newton-John Cancer and Wellness Centre
Austin Health
145 Studley Road
Heidelberg 3084
E: weeloonong@cantab.net
P: 03 9496 2800

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/1754-9485.12818](https://doi.org/10.1111/1754-9485.12818)

This article is protected by copyright. All rights reserved

DR. WEE LOON ONG (Orcid ID : 0000-0001-6657-7193)

A/PROF. FARSHAD FOROUDI (Orcid ID : 0000-0001-8387-0965)

Article type : Radiation Oncology Original Article

Androgen deprivation therapy (ADT) use with post-prostatectomy radiotherapy (PPRT) in the Prostate Cancer Outcomes Registry Victoria (PCOR-Vic)

ABSTRACT

Introduction: The aim of this study is to evaluate the use of androgen deprivation therapy (ADT) with post-prostatectomy radiotherapy (PPRT) in a population-based cohort of Australian men.

Methods: This is a prospective cohort of men with localised prostate cancer captured in the Prostate Cancer Outcomes Registry Victoria (PCOR-Vic), who received PPRT between January 2010 and December 2015. The primary outcome was ADT use with PPRT. Multivariate logistic regressions were used to identify patient, tumour and institutional factors influencing ADT use.

Results: 485 men were included in this study – 115 (24%) had pT2 disease, 231 (48%) pT3a, 134 (28%) pT3b and 5 (1%) pT4. Eighteen (4%) men had ISUP grade 1 disease, 139 (29%) ISUP grade 2, 170 (35%) ISUP grade 3, and 158 (33%) ISUP grade 4/5, while 267 (64%) men had positive surgical margins. Median time from prostatectomy to PPRT was 8.1 months (IQR=5.3-13.9). Sixty-six (14%) patients had ADT with PPRT. In multivariate analyses, men who had increased age (OR=1.06; 95%CI=1.01-1.11), seminal vesicle involvement (OR=3.81; 95%CI=1.63-8.91) and underwent treatment in regional centres (OR=2.17; 95%CI=1.08-4.33) were more likely to have ADT with PPRT.

Conclusion: We reported that 14% of men treated with PPRT received ADT in a population-based cohort of Australian men, which was less than half of the proportion of ADT use with PPRT in the US. It will be of interest to evaluate the uptake of ADT with PPRT in the coming

years following recent publications of level 1 evidence confirming overall survival benefits of ADT with PPRT.

INTRODUCTION

A significant number of men with localised prostate cancer treated with radical prostatectomy (RP) will develop local and biochemical recurrence (1, 2), and the risk could be as high as 70% for men with high-risk features such as extra-prostatic extension or seminal vesicle invasion (3, 4). Adjuvant or early salvage post-prostatectomy radiotherapy (PPRT), has been shown to be associated with oncological and overall survival benefits in prospective randomised controlled trials (5-7), but men with aggressive prostate cancer are still at risk of disease progression and distant metastases despite PPRT. Recent long-term follow-up results of RTOG-9601 and GETUG-AFU16, have provided level 1 evidence supporting the use of androgen deprivation therapy (ADT) with PPRT to further improve disease control and overall survival (8, 9). In this study, we aimed to evaluate the pattern of ADT use with PPRT in an Australian population-based cohort of men with prostate cancer, and to evaluate the factors associated with ADT use with PPRT.

METHODS

Study population: The study comprised men with prostate cancer from the Prostate Cancer Outcome Registry Victoria (PCOR-Vic). Detailed patient recruitment and data collection methodology for PCOR-Vic has been previously described (10). For this study, we only included men with localised prostate cancer who had RP as their primary treatment and subsequently had PPRT between January 2010 and December 2015.

Primary outcomes and covariates: The primary outcome of interest was ADT use with PPRT, as a binary outcome. There was no information on type or duration of ADT given the nature of data collection in PCOR-Vic. Covariates included in the analyses were: age at PPRT, year of PPRT, PSA level at prostate cancer diagnosis, International Society of Urological Pathology (ISUP) grade group (11), pathological stage, RP surgical margin status, and PPRT treatment institutions, which were classified into public or private, and metropolitan or regional.

Statistical analyses: Differences in covariates between men who received ADT vs. no ADT were analysed using Pearson's chi-squared test. Univariate and multivariate logistic regression analyses were used to estimate the effect of each factor associated with likelihood of ADT use with PPRT. Covariates with $P < 0.1$ in univariate analyses were

included in the multivariate analyses. A two-sided P-value of <0.05 was considered to indicate statistical significance. All statistical analyses were performed using STATA/IC 13 (STATA Corp, College Station, TX, USA).

RESULTS

A total of 485 men treated with RP and PPRT were included in this study. Baseline characteristics of the study population are summarised in Table 1. Of the patients included in this study, 115 (24%) had organ-confined disease (pT2), 231 (48%) had extracapsular extension (pT3a), 134 (28%) had seminal vesicle invasion (pT3b), and 5 (1%) had bladder/rectum invasion (pT4). Prostate cancer was classified based on ISUP grade group – 18 (4%), 139 (29%), 170 (35%), and 158 (33%) had ISUP grade group 1, 2, 3, and 4/5 respectively. 312 (64%) had positive surgical margins on RP. The median time from RP to RT was 8 months (IQR: 5.3-13.9 months).

There were 66 (14%) patients who had ADT with PPRT. Men who had ADT were older (mean=65.7) compared to men who did not have ADT (mean=63.4) ($P=0.006$). Men with higher ISUP grade group were more likely to have ADT ($P<0.001$) – 11%, 3%, 12% and 25% of men with ISUP Grade 1, 2, 3 and 4/5 respectively (Figure-1A). Men with non-organ confined disease were more likely to have ADT ($P<0.001$) – 7% pT2, 10% pT3a, 25% pT3b, and 20% pT4 had ADT (Figure-1B). There was no statistically significant difference in ADT use between men with positive (14%) vs. negative (12%) surgical margins ($P=0.5$) (Figure-1C). There was also no difference in ADT use in men who had PPRT at different intervals post-RP ($P=0.8$) (Figure-1D). Men treated in regional centres (23%) were more likely to have ADT compared to those treated in metropolitan centres (12%) ($P=0.01$) (Figure-1E), but there was no significant difference in ADT use in men treated in public (12%) vs. private centres (18%) ($P=0.1$) (Figure-1F).

In multivariate analyses, patient age, pathological stage, and treatment centres were independently associated with ADT use with PPRT. For every year increase in age, there was a relative 6% (95%CI=1.01-1.11, $P=0.02$) increase in likelihood of ADT use with PPRT. Men with seminal vesicle involvement were 3.8 (95%CI=1.6-8.9, $P=0.002$) times more likely to have ADT compared to men with organ-confined disease. Men treated in regional centres were 2.2 times (95%CI=1.1-4.3, $P=0.03$) more likely to have ADT with PPRT.

DISCUSSION

The establishment of PCOR-Vic has provided an excellent platform for risk-adjusted evaluation of patterns of care for men with prostate cancer in Victoria (12). Over the years, we have reported the pattern of RT practice as part of a multimodality treatment for men with prostate cancer using the extensively collected data in PCOR-Vic, including the pattern of adjuvant RT use following RP (13), ADT use with definitive prostate RT (14, 15), and brachytherapy-based prostate RT (16). Using the same database, we evaluated the pattern of ADT use in the PPRT setting in this study, and reported that 14% of men treated with PPRT between 2010 and 2015 in Victoria received ADT. This is less than half of the proportion of men who received ADT with PPRT (32%) in the US National Cancer Database (NCDB) between 2004 and 2012 (17).

We did not observe any increase in uptake of ADT use with PPRT over the study period from 2010 to 2015. This is despite an initial report of RTOG-9601 that was presented at the ASTRO meeting in 2010, which showed significant benefits of ADT use with PPRT, in terms of PSA progression-free survival, with a median follow-up of 7-years (18). It is likely that clinicians at large are still awaiting the final publication of the study prior to adoption of the evidence into clinical practice. With the recent publication of the RTOG-9601, which showed overall survival benefits of ADT use with PPRT, it will be of interest to evaluate the adoption of this practice in the coming years. Nonetheless, we also observed low uptake of ADT use in the definitive prostate RT setting in Victoria (14), despite ample level 1 evidence confirming overall survival benefits of ADT with definitive prostate RT since early 2010 (19-22), suggesting that in general there is a lag in adoption of evidence into radiation oncology practice.

The pattern of low uptake of ADT with PPRT was also observed throughout Australia and New Zealand. In a survey conducted by the RANZCR Faculty of Radiation Oncology Genitourinary Group (FROGG) in 2012, radiation oncologists with an interest in uro-oncology were asked to complete a survey regarding treatment recommendation for a 54 year-old man post-RP, who had Gleason 4+4 disease with bilateral seminal vesicle involvement (pT3b), positive surgical margins, negative node status (N0), and detectable PSA (0.4ng/mL) 8 weeks post-RP (23). Of the 46 radiation oncologists who responded, only one-third recommended ADT in conjunction with PPRT.

It is important to bear in mind that there is significant prostate cancer heterogeneity in men treated with PPRT, and ADT use may not be appropriate and necessary for every one of them (24). In the exploratory sub-group analyses in the RTOG-9601 study, Shipley et al reported that overall survival benefits of ADT was most evident in men with high-risk

features such as positive surgical margin, pre-PPRT PSA >0.7ng/mL, and Gleason 7 and above (8). In our study, however, the only high-risk feature associated with the use of ADT with PPRT in multivariate analyses was pathological stage. Also, older patients were more likely to have ADT. Given the wide spectrum of side effects associated with ADT, ranging from risk of osteoporosis and cardiovascular events, to metabolic changes and sexual dysfunction, as well as cognitive impairment and Alzheimer's disease (25-27), we postulated that younger patients were less likely to be willing to accept these side effects, and hence the lower use of ADT with PPRT in younger patients.

Apart from patient and tumour factors, we also observed that institution factors were significant determinants of ADT use – patients treated in regional centres were more likely to receive ADT with PPRT, after adjusting for other covariates. This pattern of higher proportion of ADT use in regional centres was also observed in our earlier study in the definitive prostate RT setting (14). While we postulated that patient preference may again come into play, such that men in regional centres were more willing to accept the side effects of ADT compared to men in metropolitan centres, we could not discount the possibility of preferential ADT prescription by clinicians i.e. clinicians in regional centres were possibly more likely to prescribe ADT with PPRT. This could be due to metropolitan clinicians' lack of awareness of published level 1 evidence, or their belief that the side effects outweigh the survival benefits of ADT. We do not have information to confirm or refute any of these hypotheses, and this offers potential areas for future research in order to identify the barriers and facilitators for adoption of ADT use with PPRT, from both patients' and clinicians' perspective.

There are several other limitations in the current study, which are inherent limitations of PCOR-Vic database. These include the lack of information on RT dose and fractionations as well as type and duration of ADT use. One of the possible avenues to overcome these limitations in future study using the PCOR-Vic data is through data-linkage with other population-based administrative database such as the Victorian Radiotherapy Minimum Data Set, which captures detailed RT information in the state of Victoria, as well as the Pharmaceutical Benefit Scheme, which will provide details on type and duration of ADT.

In summary, we report the contemporary patterns of ADT use with PPRT in a population-based cohort of Australian men with prostate cancer, with only 14% receiving ADT with PPRT. It will be of great interest to evaluate the pattern of uptake of ADT with PPRT in the coming years, following the publications of the RTOG 9601 and GETUG-AFU16 findings. With long-term follow-up, it will also be interesting to evaluate if the pattern of ADT use with

PPRT in a population-based real-life clinical setting translates into overall survival differences as observed in the trials.

ACKNOWLEDGEMENT

PCOR-Vic is funded by the Movember Foundation. SME receives funding from the Monash Partners Academic Health Science Centre Clinical Research Fellowship. We acknowledge the assistance of data collectors, clinicians, and hospitals contributing to the PCOR-Vic.

REFERENCES

1. Pound CR, Partin AW, Eisenberger MA, Chan DW, Pearson JD, Walsh PC. Natural history of progression after PSA elevation following radical prostatectomy. JAMA. 1999;281(17):1591-7.
2. Roehl KA, Han M, Ramos CG, Antenor JA, Catalona WJ. Cancer progression and survival rates following anatomical radical retropubic prostatectomy in 3,478 consecutive patients: long-term results. J Urol. 2004;172(3):910-4.
3. Carver BS, Bianco FJ, Jr., Scardino PT, Eastham JA. Long-term outcome following radical prostatectomy in men with clinical stage T3 prostate cancer. J Urol. 2006;176(2):564-8.
4. Nguyen CT, Reuther AM, Stephenson AJ, Klein EA, Jones JS. The specific definition of high risk prostate cancer has minimal impact on biochemical relapse-free survival. J Urol. 2009;181(1):75-80.
5. Bolla M, van Poppel H, Collette L, van Cangh P, Vekemans K, Da Pozzo L, et al. Postoperative radiotherapy after radical prostatectomy: a randomised controlled trial (EORTC trial 22911). Lancet. 2005;366(9485):572-8.
6. Thompson IM, Jr., Tangen CM, Paradelo J, Lucia MS, Miller G, Troyer D, et al. Adjuvant radiotherapy for pathologically advanced prostate cancer: a randomized clinical trial. JAMA. 2006;296(19):2329-35.
7. Wiegel T, Bottke D, Steiner U, Siegmann A, Golz R, Storkel S, et al. Phase III postoperative adjuvant radiotherapy after radical prostatectomy compared with radical prostatectomy alone in pT3 prostate cancer with postoperative undetectable prostate-specific antigen: ARO 96-02/AUO AP 09/95. J Clin Oncol. 2009;27(18):2924-30.

- 216 8. Shipley WU, Seiferheld W, Lukka HR, Major PP, Heney NM, Grignon DJ, et al.
217 Radiation with or without Antiandrogen Therapy in Recurrent Prostate Cancer. *N Engl J Med*.
218 2017;376(5):417-28.
- 219 9. Carrie C, Hasbini A, de Laroche G, Richaud P, Guerif S, Latorzeff I, et al. Salvage
220 radiotherapy with or without short-term hormone therapy for rising prostate-specific
221 antigen concentration after radical prostatectomy (GETUG-AFU 16): a randomised,
222 multicentre, open-label phase 3 trial. *Lancet Oncol*. 2016;17(6):747-56.
- 223 10. Evans SM, Millar JL, Davis ID, Murphy DG, Bolton DM, Giles GG, et al. Patterns of
224 care for men diagnosed with prostate cancer in Victoria from 2008 to 2011. *Med J Aust*.
225 2013;198(10):540-5.
- 226 11. Gordetsky J, Epstein J. Grading of prostatic adenocarcinoma: current state and
227 prognostic implications. *Diagn Pathol*. 2016;11:25.
- 228 12. Evans SM, Millar JL, Wood JM, Davis ID, Bolton D, Giles GG, et al. The Prostate
229 Cancer Registry: monitoring patterns and quality of care for men diagnosed with prostate
230 cancer. *BJU Int*. 2013;111(4 Pt B):E158-66.
- 231 13. Daniels CP, Millar JL, Spelman T, Sengupta S, Evans SM. Predictors and rate of
232 adjuvant radiation therapy following radical prostatectomy: A report from the Prostate
233 Cancer Registry. *J Med Imaging Radiat Oncol*. 2016;60(2):247-54.
- 234 14. Ong WL, Foroudi F, Evans S, Millar J. Large institutional variations in use of androgen
235 deprivation therapy with definitive radiotherapy in a population-based cohort of men with
236 intermediate- and high-risk prostate cancer. *BJU Int*. 2017;120 Suppl 3:35-42.
- 237 15. Ong WL, Evans S, Millar J. In Regard to Yang et al. *Int J Radiat Oncol Biol Phys*.
238 2018;100(5):1294-5.
- 239 16. Ong WL, Evans SM, Millar JL. Under-utilisation of high-dose-rate brachytherapy
240 boost in men with intermediate-high risk prostate cancer treated with external beam
241 radiotherapy. *J Med Imaging Radiat Oncol*. 2018; 62(2): 256-261.
- 242 17. Yang DD, Muralidhar V, Mahal BA, Nezoslosky MD, Labe SA, Vastola ME, et al. Low
243 rates of androgen deprivation therapy use with salvage radiation therapy in patients with
244 prostate cancer after radical prostatectomy. *Urol Oncol*. 2017;35(9):542 e25- e32.
- 245 18. Shipley WU, Hunt D, Lukka H, Major PP, Heney NM, Grignon D, et al. Initial report of
246 RTOG 9601: a phase III trial in prostate cancer: anti-androgen therapy with bicalutamide
247 during and after radiation therapy improves freedom from progression and reduces the

- incidence of metastatic disease in patients following radical prostatectomy with pT2–3, N0 disease and elevated PSA levels. *Int J Radiat Oncol Biol Phys*. 2010;78(suppl. 3):S27.
19. Pilepich MV, Winter K, Lawton CA, Krisch RE, Wolkov HB, Movsas B, et al. Androgen suppression adjuvant to definitive radiotherapy in prostate carcinoma--long-term results of phase III RTOG 85-31. *Int J Radiat Oncol Biol Phys* 2005;61(5):1285-90.
20. Bolla M, Van Tienhoven G, Warde P, Dubois JB, Mirimanoff RO, Storme G, et al. External irradiation with or without long-term androgen suppression for prostate cancer with high metastatic risk: 10-year results of an EORTC randomised study. *Lancet Oncol*. 2010;11(11):1066-73.
21. Denham JW, Steigler A, Lamb DS, Joseph D, Turner S, Matthews J, et al. Short-term neoadjuvant androgen deprivation and radiotherapy for locally advanced prostate cancer: 10-year data from the TROG 96.01 randomised trial. *Lancet Oncol*. 2011;12(5):451-9.
22. Jones CU, Hunt D, McGowan DG, Amin MB, Chetner MP, Bruner DW, et al. Radiotherapy and short-term androgen deprivation for localized prostate cancer. *N Engl J Med*. 2011;365(2):107-18.
23. Lehman M, Sidhom M, Kneebone AB, Hayden AJ, Martin JM, Christie D, et al. FROGG high-risk prostate cancer workshop: patterns of practice and literature review. Part II post-radical prostatectomy. *J Med Imaging Radiat Oncol*. 2014;58(3):392-400.
24. Lee WR. Invited commentary on GETUG-AFU 16. *Transl Androl Urol*. 2016;5(6):958-60.
25. Rhee H, Gunter JH, Heathcote P, Ho K, Stricker P, Corcoran NM, et al. Adverse effects of androgen-deprivation therapy in prostate cancer and their management. *BJU Int*. 2015;115 Suppl 5:3-13.
26. Keating NL, O'Malley AJ, Smith MR. Diabetes and cardiovascular disease during androgen deprivation therapy for prostate cancer. *J Clin Oncol*. 2006;24(27):4448-56.
27. Nead KT, Gaskin G, Chester C, Swisher-McClure S, Dudley JT, Leeper NJ, et al. Androgen Deprivation Therapy and Future Alzheimer's Disease Risk. *J Clin Oncol*. 2016;34(6):566-71.

Table -1 | Baseline characteristics of men who had ADT vs. no ADT

	No ADT	ADT	All	P-value
	N=419 (86%)	N=66 (14%)	N=485	
Age at PPRT (year), mean (SD)	63.4 (6.5)	65.7 (5.8)	63.7 (6.5)	0.006
Serum PSA at diagnosis (ng/L), median (IQR)	7.2 (5.5-10.2)	8.2 (5.8-13)	7.2 (5.6-10.8)	0.07
<10	306 (73%)	38 (58%)	334 (71%)	0.08
10-20	75 (17%)	18 (27%)	93 (19%)	
>20	31 (7%)	8 (12%)	39 (8%)	
Unknown	7 (2%)	2 (3%)	9 (2%)	
ISUP Grade , n (%)				
1 (GS 3+3)	16 (4%)	2 (3%)	18 (4%)	<0.001
2 (GS 3+4)	133 (32%)	4 (6%)	139 (29%)	
3 (GS4+3)	150 (36%)	20 (30%)	170 (35%)	
4/5 (GS 8/9/10)	118 (28%)	40 (61%)	158 (33%)	
Pathological stage, n (%)				
pT2	107 (26%)	8 (12%)	115 (24%)	<0.001
pT3a	207 (49%)	24 (36%)	231 (48%)	
pT3b	101 (24%)	33 (50%)	134 (28%)	
pT4	4 (1%)	1 (2%)	5 (1%)	
Positive margin, n (%)				
No	152 (36%)	21 (32%)	173 (36%)	0.5
Yes	267 (64%)	45 (68%)	312 (64%)	
Time from RP to PPRT (month), median (IQR)	8.3 (5.3-14.1)	7.5 (5.7-10.3)	8.1 (5.3-13.9)	0.2
<6 month	134 (32%)	24 (36%)	158 (33%)	0.7
6-12 month	155 (37%)	26 (39%)	181 (37%)	
12-24 month	98 (23%)	12 (18%)	110 (23%)	
>24 month	32 (8%)	4 (6%)	36 (7%)	
RT centre type, n (%)				
Public	322 (77%)	45 (68%)	367 (76%)	0.1
Private	97 (23%)	21 (32%)	118 (24%)	
RT centre location, n (%)				

Metropolitan	370 (88%)	51 (77%)	421 (87%)	0.01
Regional	49 (12%)	15 (23%)	64 (13%)	
Year of PPRT, n (%)				
2010	38 (9%)	5 (8%)	43 (9%)	0.9
2011	60 (14%)	7 (11%)	67 (14%)	
2012	95 (23%)	18 (27%)	113 (23%)	
2013	80 (19%)	12 (18%)	92 (19%)	
2014	77 (18%)	12 (18%)	89 (18%)	
2015	69 (16%)	12 (18%)	81 (17%)	

Table-2 | Univariate and multivariate logistic regressions of factors associated with ADT use with PPRT

	Univariate		Multivariate	
	OR (95%CI)	P-value	OR (95%CI)	P-value
Age at PPRT	1.06 (1.02-1.11)	0.006	1.06 (1.01-1.11)	0.02
Serum PSA at diagnosis				
<10	1.0 (Ref.)		1.0 (Ref.)	
10-20	1.93 (1.04-3.57)	0.04	1.42 (0.73-2.74)	0.3
>20	2.08 (0.89-4.85)	0.09	1.51 (0.61-3.74)	0.4
ISUP Grade				
1	1.0 (Ref.)		-	-
2	0.24 (0.04-1.49)	0.1	-	-
3	1.07 (0.23-4.99)	0.9	-	-
4/5	2.71 (0.60-12.31)	0.1	-	-
Pathological stage				
pT2	1.0 (Ref.)		1.0 (Ref.)	
pT3a	1.55 (0.67-3.57)	0.3	1.36 (0.58-3.20)	0.5
pT3b	4.37 (1.93-9.91)	<0.001	3.81 (1.63-8.91)	0.002
pT4	3.34 (0.33-33.55)	0.3	4.29 (0.39-47.16)	0.2
Positive margin				
No	1.0 (Ref.)		-	-
Yes	1.22 (0.70-2.12)	0.5	-	-
Time from RP to RT				
<6 months	1.0 (Ref.)			
6-12 months	0.94 (0.51-1.71)	0.8	-	-

12-24 months	0.68 (0.33-1.43)	0.3	-	-
>24 months	0.70 (0.23-2.15)	0.5	-	-
RT centre type				
Public	1.0 (Ref.)		-	-
Private	0.65 (0.37-1.14)	0.1	-	-
RT centre location				
Metropolitan	1.0 (Ref.)		1.0 (Ref.)	
Regional	2.22 (1.16-4.25)	0.02	2.17 (1.08-4.33)	0.03
Year of RT				
2010	1.0 (Ref.)		-	-
2011	0.89 (0.26-3.00)	0.8	-	-
2012	1.44 (0.50-4.16)	0.5	-	-
2013	1.14 (0.37-3.47)	0.8	-	-
2014	1.18 (0.39-3.61)	0.8	-	-
2015	1.32 (0.43-4.03)	0.6	-	-

Figure 1 | Proportion of men who received ADT with PPRT, stratified by ISUP grading (1A), pathological stage (1B), surgical margin (1C), interval between RP and PPRT (1D), metropolitan-regional centres (1E), and public-private centres (1F)

