



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Badurdeen, S;Galinsky, R;Roberts, CT;Crossley, KJ;Zahra, VA;Thiel, A;Pham, Y;Davis, PG;Hooper, SB;Polglase, GR;Camm, EJ

Title:

Rapid oxygen titration following cardiopulmonary resuscitation mitigates cerebral overperfusion and striatal mitochondrial dysfunction in asphyxiated newborn lambs

Date:

2025-04-01

Citation:

Badurdeen, S., Galinsky, R., Roberts, C. T., Crossley, K. J., Zahra, V. A., Thiel, A., Pham, Y., Davis, P. G., Hooper, S. B., Polglase, G. R. & Camm, E. J. (2025). Rapid oxygen titration following cardiopulmonary resuscitation mitigates cerebral overperfusion and striatal mitochondrial dysfunction in asphyxiated newborn lambs. *Journal of Cerebral Blood Flow and Metabolism*, 45 (4), pp.630-642. <https://doi.org/10.1177/0271678X241302738>.

Persistent Link:

<https://hdl.handle.net/11343/359303>

License:

CC BY

Rapid oxygen titration following cardiopulmonary resuscitation mitigates cerebral overperfusion and striatal mitochondrial dysfunction in asphyxiated newborn lambs

Journal of Cerebral Blood Flow & Metabolism

0(0) 1–13

© The Author(s) 2024





Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/0271678X241302738

journals.sagepub.com/home/jcbfm



Shiraz Badurdeen^{1,2,3} , Robert Galinsky¹ ,
Calum T Roberts^{1,4}, Kelly J Crossley¹, Valerie A Zahra¹,
Alison Thiel¹, Yen Pham¹, Peter G Davis³, Stuart B Hooper^{1,6},
Graeme R Polglase^{1,5,*} and Emily J Camm^{1,6,*}

Abstract

Asphyxiated neonates must have oxygenation rapidly restored to limit ongoing hypoxic-ischemic injury. However, the effects of transient hyperoxia after return of spontaneous circulation (ROSC) are poorly understood. We randomly allocated acutely asphyxiated, near-term lambs to cardiopulmonary resuscitation in 100% oxygen (“standard oxygen”, $n = 8$) or air ($n = 7$) until 5 minutes after ROSC, or to resuscitation in 100% oxygen immediately weaned to air upon ROSC (“rapid-wean”, $n = 7$). From 5 minutes post-ROSC, oxygen was titrated to target preductal oxygen saturation between 90–95%. Cerebral tissue oxygenation was transiently but markedly elevated following ROSC in the standard oxygen group compared to the air and rapid-wean groups. The air group had a delayed rise in cerebral tissue oxygenation from 5 minutes after ROSC coincident with up-titration of oxygen. These alterations in oxygen kinetics corresponded with similar overshoots in cerebral perfusion (pressure and flow), indicating a physiological mechanism. Transient cerebral tissue hyperoxia in the standard oxygen and air groups resulted in significant alterations in mitochondrial respiration and dynamics, relative to the rapid-wean group. Overall, rapid-wean of oxygen following ROSC preserved striatal mitochondrial respiratory function and reduced the expression of genes involved in free radical generation and apoptosis, suggesting a potential therapeutic strategy to limit cerebral reperfusion injury.

Keywords

Infant, hypoxia, reperfusion injury, metabolism, hypoxia-ischemia

Received 24 June 2024; Revised 31 October 2024; Accepted 6 November 2024

Introduction

Cardiopulmonary resuscitation (CPR) of neonates with severe perinatal asphyxia currently focuses on rapid restoration of the circulation to limit ongoing hypoxic-ischemic brain injury.^{1,2} Given the challenges in conducting adequately powered trials of CPR in neonates, international recommendations rely on transitional animal models to guide practice.¹ Current

³Department of Obstetrics, Gynaecology and Newborn Health, University of Melbourne, Parkville VIC, Australia

⁴Monash Newborn, Monash Children’s Hospital, Clayton, Australia

⁵Department of Paediatrics, Monash University, Clayton, Australia

⁶Department of Obstetrics and Gynaecology, Monash University, Clayton, Australia

*These authors contributed equally to this work.

Corresponding authors:

Shiraz Badurdeen, The Ritchie Centre, Hudson Institute of Medical Research, Clayton VIC 3168, Australia.

Email: shiraz.badurdeen@unimelb.edu.au

Emily J Camm, The Ritchie Centre, Hudson Institute of Medical Research, Clayton VIC 3168, Australia.

Email: emily.camm@hudson.org.au

¹The Ritchie Centre, Hudson Institute of Medical Research, Clayton, Australia

²Melbourne Children’s Global Health, Murdoch Children’s Research Institute, Parkville, Australia

guidelines recommend increasing the fraction of inspired oxygen (FiO_2) to 100% if chest compressions are required to restore myocardial contractility.^{2,3} Following return of spontaneous circulation (ROSC), the optimal strategy for oxygen provision is not known. In practice, oxygen is typically titrated to mimic oxygen saturation (SpO_2) levels of neonates with normal perinatal transition.^{2,4} However, due to the rapid cerebral overperfusion that occurs immediately after ROSC, this strategy results in global cerebral oxygen delivery that is markedly in excess of oxygen consumption in the minutes following ROSC which can lead to oxidative stress and disruption of cerebral metabolism.^{5–11}

The effect of transient excess oxygen delivery immediately after ROSC on region-specific cerebral tissue oxygenation and cellular function is unknown. Acute perinatal asphyxia predominantly affects the deep grey nuclei (striatum) and the perirolandic cortex.¹² Following asphyxia and re-perfusion, alterations in mitochondrial bioenergetics and mitochondrial-derived reactive oxygen species (ROS) result in oxidative damage to mitochondrial proteins, membranes and DNA, impairing their ability to synthesize ATP, ultimately leading to cell death and secondary brain injury.^{13,14} Transient hyperoxia may therefore exacerbate ROS production, further impairing mitochondrial function and contributing to secondary energy failure that is characteristic of hypoxic-ischemic encephalopathy.¹⁵ Strategies to limit hypoxia while avoiding hyperoxia-mediated oxidative stress in the reperfusion phase have not been well studied.^{6,16}

Among neonates with mild perinatal asphyxia requiring respiratory support alone, commencing mask ventilation in 100% oxygen versus air increases the risk of mortality.¹⁷ In neonates needing CPR therefore, the more severe initial insult may render pathways leading to secondary reperfusion injury particularly susceptible to even transient hyperoxic exposure. Here, we hypothesized that in acutely asphyxiated newborn lambs, a brief delay in weaning supplemental oxygen leads to cerebral tissue hyperoxia and mitochondrial dysfunction, which can be mitigated by rapidly weaning oxygen to air immediately after ROSC.

Materials and methods

All procedures were approved by the Monash University Animal Ethics Committee (MMCA-2020-04; 24/6/2020) and conducted in accordance with the National Health and Medical Research Council of Australia code for the care and use of animals for scientific purposes. This study is reported according to the ARRIVE guidelines.¹⁸ Additional methods are provided in the Supplementary Materials.

Instrumentation

Border-Leicester sheep fetuses at 139 ± 2 days' gestation (mean \pm SD; term ~ 148 days) were intubated and instrumented after exteriorization of the head and chest via laparotomy. Cerebral oxygenation parameters were measured using fibre-optic oxygen probes placed in the right cerebral striatum (caudate nucleus) and right common carotid artery. Carotid blood flow is known to strongly correlate with cerebral perfusion.¹⁹ Regional cortical tissue oxygen saturation (crSO_2) was measured over the left frontal cortex using near-infrared spectroscopy (NIRS).

Asphyxia and management after birth

Asphyxia was induced by cord clamping while withholding ventilation. Immediately upon terminal asphyxia, defined by asystole, mean arterial blood pressure of ~ 0 mmHg and $\text{pH} < 7.0$, ventilation was initiated in air with pressures of 30/5 cmH₂O at 60 ventilations per minute (NeoPuff, Fisher & Paykel). One minute later, chest compressions were initiated at a 3:1 ratio with ventilation and FiO_2 delivered according to one of 3 strategies that were randomly assigned prior to resuscitation using a web-based random sequence generator (www.random.org/lists):

1. Standard oxygen (100% oxygen from the commencement of chest compressions until 5 minutes after ROSC), $n = 8$
2. Air (21% oxygen until 5 minutes after ROSC), $n = 7$
3. Rapid-wean (100% oxygen from the commencement of chest compressions until onset of ROSC, upon which oxygen was immediately weaned to 21% until 5 minutes after ROSC), $n = 7$.

Epinephrine (1:10,000, 0.2 ml/kg) was administered 1 minute after initiating chest compressions. ROSC was defined as continuous unsupported cardiac output with diastolic blood pressure > 20 mmHg. Lambs then received ventilator-driven breaths at 30/5 cmH₂O for 10 minutes followed by volume-guarantee mode (7 mL/kg). From 5 minutes post-ROSC, FiO_2 was adjusted in all groups to target pre-ductal arterial oxygen saturation of 90–95%. Lambs were sedated from 10 minutes after ROSC (Alfaxalone intravenous 5 mg/kg/hour in 5% dextrose; Jurox) to prevent spontaneous breathing.

Data acquisition and analysis

Continuous recordings of physiological parameters were made at 1 kHz. Carotid arterial blood gases were taken before induction of asphyxia, at terminal asphyxia, during CPR, at ROSC, and at regular intervals until 60 minutes after ROSC. All analyses were conducted blind to group allocation.

Tissue collection and analysis

At 1 hour after ROSC, lambs were euthanized and tissue samples taken from the left striatum (matching the level where the oxygen probe was placed) and left cerebral cortex (at the level of the ansate sulcus, equivalent to the perirolandic cortex in humans). Mitochondrial respiratory rates and bioenergetics were measured using respirometry, western blotting and quantitative PCR (q-PCR), as previously described.^{20,21}

Statistical analysis

From our previous work, 7–8 lambs per group were able to demonstrate differences in mitochondrial oxidative phosphorylation (OXPHOS).²¹ Baseline fetal characteristics, mitochondrial respirometry and western blot data were compared using a one-way analysis of variance, t-test or Mann-Whitney U test, as appropriate. Comparisons of physiological data between study groups were done using two-way repeated measures analysis of variance (ANOVA), or mixed effects models when there was missing data. The Tukey test was used to compare group means at each timepoint (Supplement) and calculate p-values and 95% confidence intervals adjusted for multiple comparisons. Quantitative PCR data was expressed as relative change ($2^{-\Delta\Delta CT}$) were log-transformed to maintain normal distribution, and all statistical analysis and

conducted on log-transformed $2^{-\Delta\Delta CT}$ data. Volcano plot data visualizations present comparisons of standard care versus rapid-wean or 21% oxygen as fold changes. P values were adjusted for multiple comparisons using the Benjamini-Hochberg method. All analyses were performed using PRISM version 10 for Mac OS X (GraphPad Software, La Jolla California USA).

Results

Baseline physiological comparisons

One lamb in the rapid-wean group died shortly after ROSC due to delayed recognition of a mal-positioned endotracheal tube and was excluded from all analyses. Baseline characteristics and blood gas parameters of the remaining 22 lambs were similar between groups except for fetal pO₂, which was lowest in the standard oxygen group (Table 1).

Time to achieve ROSC was similar between lambs receiving CPR in 100% oxygen versus air

The mean time to achieve ROSC after commencing ventilation was similar between lambs receiving CPR in 100% oxygen (standard and rapid-wean groups, n = 15, 185 s, 95% confidence interval [CI] 173–196 s) compared to the 21% oxygen group (n = 7, 191 s, 95%

Table 1. Cohort characteristics.

	Standard oxygen n = 8	Air n = 7	Rapid-wean n = 7
Weight (kg) ^a	4.94 (0.55)	5.16 (0.71)	4.76 (0.85)
Sex (n female/total)	5/8	5/7	4/7
Single:Twins (n:n)	2:6	2:5	1:6
Fetal (baseline) ^a			
pH	7.27 (0.03)	7.27 (0.04)	7.28 (0.06)
PaO ₂ (mmHg)	17.4 (4.2)	23.3 (2.9)	22.0 (5.4)
PCO ₂ (mmHg)	58.7 (3.8)	59.3 (3.1)	56.1 (8.3)
Base excess	-0.36 (2.4)	-1.14 (3.2)	-0.74 (1.8)
Lactate	4.0 (0.8)	2.9 (1.3)	3.9 (0.8)
End asphyxia ^a			
pH	6.91 (0.05)	6.92 (0.04)	6.90 (0.05)
PaO ₂ (mmHg)	1.9 (2.2)	1.79 (2.2)	2.3 (2.3)
PCO ₂ (mmHg)	112.2 (14.4)	111.7 (15.0)	111.4 (14.1)
Base excess	-10.4 (3.5)	-9.9 (1.3)	-11.4 (1.5)
Lactate	9.2 (1.6)	8.2 (2.2)	9.5 (1.7)
60 min post-ROSC ^a			
pH	7.25 (0.06)	7.20 (0.12)	7.23 (0.07)
PaO ₂ (mmHg)	61.1 (12.9)	58.0 (17.7)	63.9 (5.0)
PCO ₂ (mmHg)	39.7 (2.5)	44.5 (8.8)	42.7 (4.1)
Base excess	-9.8 (2.5)	-11.0 (4.3)	-9.6 (3.2)
Lactate	8.4 (1.3)	8.6 (2.2)	8.2 (1.0)

^aValues are mean (SD). PaO₂: partial pressure of arterial oxygen; PCO₂: partial pressure of arterial carbon dioxide.

CI 174–207 s). All lambs achieved ROSC following a single dose of epinephrine.

Rapid-wean of FiO_2 dampened the rise in cerebral tissue oxygenation following ROSC

During CPR, arterial oxygen saturation, striatal pO_2 and $crSO_2$ remained low despite the use of 100%

oxygen in the standard and rapid-wean groups (Figure 1(b) to (e)). Upon ROSC, lambs in the air group had the lowest arterial oxygen saturation and $crSO_2$ (Figure 1(b) and (e)). In the standard oxygen group, arterial oxygen saturation, carotid arterial pO_2 , striatal tissue pO_2 and $crSO_2$ were markedly elevated within 1–3 min post-ROSC (Figure 1(b) to (e)). In comparison, the fast rise in oxygenation was




Figure 1. Fraction of inspired oxygen (FiO_2 , a) and oxygenation parameters (b–f) for lambs in the standard oxygen group (“Standard”, orange), rapid-wean group (“Rapid”, blue), and air group (“21%”, black). The standard oxygen group had a marked rise in carotid arterial oxygen saturation (b) and pO_2 (c), as well as striatal tissue pO_2 (d) and regional cerebral oxygen saturation (e) from the onset of ROSC that was mitigated by rapidly weaning FiO_2 to air. The air group had a slow initial rise in oxygenation parameters followed by a secondary rise once FiO_2 was increased at 5 min after ROSC. Regional cerebral fractional tissue oxygen extraction (f) was similar between groups. Error bars represent 95% confidence intervals. Asterisks in red and black represent statistically significant differences between the rapid-wean versus standard oxygen group, and rapid-wean versus air groups, respectively. Comparisons between the standard oxygen and air group are not shown for clarity ($n = 7–8$ /group). *Adjusted $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ from Tukey test following two-way repeated measures analysis of variance or mixed effects models (when there was missing data).

dampened in the rapid-wean group- for example, mean carotid arterial pO_2 at 3 min post-ROSC was 56 mmHg vs 298 mmHg (adjusted $p = 0.0007$, Tukey test); striatal pO_2 at 6 min post-ROSC was 29 mmHg vs 104 mmHg (adjusted $p = 0.008$, Tukey test). The air group had the slowest initial rise in oxygenation parameters.

From 5 minutes post-ROSC, all lambs had FiO_2 titrated to target pre-ductal arterial oxygen saturation of 90–95% as per protocol. In the standard oxygen group, FiO_2 was steadily weaned (Figure 1(a)). All lambs in the air group had FiO_2 increased (range 25–50%) and 4/7 lambs in the rapid-wean group had FiO_2 increased (range 30–50%). In the air group, this corresponded with a distinct secondary rise in striatal pO_2 and $crSO_2$ that was not apparent in the rapid-wean group, although differences did not meet statistical significance (Figures 1(d) and (e)). From approximately 12 minutes post-ROSC, oxygen parameters were similar between groups.

Regional cortical tissue oxygen saturation (Figure 1(e)) was significantly higher between 30 s–5 min post-ROSC in the standard oxygen versus the air group (adjusted $p < 0.05$, Tukey test). From 5 min post-ROSC, the secondary rise in $crSO_2$ that followed up-titration of FiO_2 in the air group resulted in both $crSO_2$ and cerebral fractional tissue oxygen extraction values that were similar to the standard oxygen group (Figure 1(e) and (f)). Correspondingly, peak $crSO_2$ was $\geq 85\%$ in 7/8 lambs and 6/7 lambs in the standard oxygen and air groups respectively. In contrast, 2/7 lambs in the rapid-wean group had peak $crSO_2 \geq 85\%$.

The post-ROSC overshoot in pulmonary and carotid blood flow matched cerebral oxygen kinetics and was least prominent in the rapid-wean group

Carotid arterial pO_2 was significantly higher in the standard care group at 3 and 6 minutes after ROSC compared to the rapid wean and air groups (Figure 2(a)). Arterial pCO_2 was similar between groups at all timepoints (Figure 2(b)). Pulmonary artery blood flow was higher in the standard care group for the first 5 minutes (by 21.3% vs. Rapid Wean and 21.0% vs. Air; $p_{\text{group} \times \text{time}} < 0.001$ from two-way ANOVA; Figure 2(c)) but did not reach significance at individual time points (adjusted $p > 0.05$, Tukey test). Lambs in the air group had a delayed secondary rise in pulmonary artery blood flow following the increase in FiO_2 from 5 min after ROSC, which was not apparent in the rapid-wean group.

Common carotid arterial pressures were similar between groups (Figure 2(f)). However, the post-ROSC overshoot in common carotid blood flow (Figure 2(d)) was lowest in the rapid-wean group, with a significantly lower common carotid blood flow

than the standard care (from 20 s – 10 min; adjusted $p < 0.05$, Tukey test) and air (from 5:40 min – 10 min; adjusted $p < 0.05$, Tukey test) groups. Lambs in the standard care group had a rapid rise in both carotid arterial pressure and blood flow consistent with cerebral overperfusion, which was markedly reduced in the air and rapid-wean groups during the first 10 minutes. Taken together, the post-ROSC overshoot in pulmonary and carotid blood flows was least prominent in the rapid-wean group and indicates that the reduction to cerebral oxygenation is driven by this reduction in cerebral perfusion.

Transient hyperoxia had region-specific effects on mitochondrial respiration and bioenergetics

Both the standard oxygen and air groups had significantly decreased striatal mitochondrial Complex I- (CI), combined CI&CII, and CIV-driven respiration, and reduced maximal electron transfer system capacity relative to the rapid-wean group (all $p < 0.05$, Tukey test, Figure 3(a)). CII-driven respiration in the standard group was significantly lower than in the rapid-wean group ($p < 0.05$, Tukey test). No effect of oxygen exposure on the remaining outcome measures were observed, or when data was expressed relative to ETS capacity (all $p > 0.05$, Tukey test, Figure 5(a)). In the cortex (Figures 4 and 5(b)), only LEAK respiration relative to ETS capacity was reduced in the standard and air groups relative to the rapid-wean group ($p < 0.05$, Tukey test).

Striatal abundance of CII was increased in both the air and rapid-wean groups relative to the standard oxygen group ($p < 0.05$, Tukey test Figure 3(b)). The abundances of the remaining complexes were not altered in the striatum or cortex ($p > 0.05$, Figures 3(b) and 4). In the cortex, the abundance of citrate synthase, a proxy marker for mitochondrial content, and 75-kDa glucose-regulated protein (GRP75), a mitochondrial stress protein, were increased and decreased, respectively in the rapid-wean group relative to the standard care and air groups (both $p < 0.05$, Tukey test, Figure 4).

Relative to the standard oxygen group, rapid-wean resulted in significant downregulation of multiple genes in the striatum (adjusted $p < 0.05$, Benjamini-Hochberg correction for multiple comparisons, Figure 3(c)), including genes that encode for NADPH oxidase enzymes which are crucial in the generation of free radicals (*NADPH Oxidase 1* and *2*, *Nox1* and *Nox2*), as well as those involved in mitochondrial biogenesis (*Tfam*, *Opal*, *Drp1*, *Mfn1*, *PGC-1 α*), apoptosis and ER stress-induced apoptosis (*Bax*, *Casp3*, and *CHOP*), and antioxidant defence (*SOD2* and catalase). No gene expression changes were found in the striatum in the




Figure 2. Blood gas pO₂ (a), pCO₂ (b) and cardiovascular parameters (c–f) for lambs in the standard oxygen group (“Standard”, orange), rapid-wean group (“Rapid”, blue), and air group (“21%”, black). Pulmonary artery flow and common carotid artery flow were calculated per unit of organ weight (total lung and total brain weight, respectively). The standard oxygen group tended to have the highest mean carotid artery blood pressure and the highest mean pulmonary artery blood flow in the first 5 minutes after ROSC, but differences between groups were not statistically significant. Lambs in the rapid-wean group had lower mean carotid blood flow both at the fetal timepoint and after ROSC when compared to the standard oxygen group. Error bars represent 95% confidence intervals. Asterisks in red and black represent statistically significant differences between the rapid-wean versus standard oxygen group, and rapid-wean versus air groups, respectively. Comparisons between the standard oxygen and air groups are not shown for clarity ($n = 7-8/\text{group}$). *Adjusted $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ from Tukey test following two-way repeated measures analysis of variance or mixed effects models (when there was missing data).

air cohort relative to standard oxygen (Figure 3(c)), nor in the cerebral cortex (Figure 4(c)).

Discussion

Our findings indicate that tissue hyperoxia occurs with brief oxygen supplementation after ROSC, is driven by

rapid and prolonged increase in cerebral blood flow, and is particularly marked in the striatum. Tissue hyperoxia impairs mitochondrial respiration during the early latent phase in a region-specific manner. Using 100% oxygen during CPR but rapidly weaning oxygen to air following ROSC prevents cerebral overperfusion, hyperoxia and improves mitochondrial respiratory function.




Figure 3. Integrated analysis of the striatum following severe perinatal asphyxia and different re-oxygenation strategies. (a) Mitochondrial respiration rates in the striatum for lambs in the standard oxygen treatment group (“Standard”, orange), rapid-wean group (“Rapid”, blue), and air group (“21%”, black) for leak respiration (CI_L), Complex I-linked respiration (CI_P), CI&CII-linked respiration (CI&CII_P) and CII-linked respiration (CII_E), Complex IV activity, maximal electron transfer (ETS) capacity (Max ETS), Residual Oxygen Consumption, oxidative phosphorylation (OXPHOS) coupling efficiency (OXPHOS Efficiency). (b) Protein abundance of electron transfer system complexes (Complex I [CI], Complex II [CII], Complex III [CIII], Complex IV [CIV], ATP synthase [ATPase]), citrate synthase (CS), 75-kDa glucose-regulated protein (GRP75), and 60-kDa heat shock protein (HSP60). Note: some proteins were probed on the same western blot (CI/CIII/ATPase and GRP75/HSP60). (c) Volcano plot visualizations of key genes involved in mitochondrial biogenesis, apoptosis and ER stress-induced apoptosis, antioxidant defense, and in the generation of free radicals. Continued.

Recent lamb studies found reductions in global cerebral oxygen delivery, as well as lower blood oxidized/reduced glutathione ratio, in lambs that had an abrupt wean in FiO_2 from 100% to air following ROSC compared to a gradual wean.^{10,11} Our results support and extend these findings, providing evidence of marked hyperoxia at the cerebral tissue level in both the striatum and the cerebral cortex, with corresponding impairment of mitochondrial function. These findings provide a biological basis for our recently reported observation that early hyperoxemia was causally related to an increased risk of death or major disability among infants with hypoxic-ischemic encephalopathy enrolled in the Infant Cooling Evaluation trial.²²

The effects of transient hyperoxia may be especially stark in the context of severe asphyxia, where there are currently no clinical studies prospectively evaluating oxygen supplementation strategies.^{6,7} Randomized trials of newborns with mild perinatal hypoxic-ischemia who received respiratory support without chest compressions corroborated the findings of animal models, finding lower mortality when resuscitation commenced in air versus 100% oxygen, informing current recommendations.^{17,23–25} Pre-clinical studies in non-transitioning, severely asphyxiated piglets evaluated a prolonged period (10–30 min) of exposure to 100% oxygen.^{26–28} This duration of hyperoxic exposure is now uncommon in advanced neonatal care settings. Our standard oxygen arm which provided 100% oxygen for 5 min, is more representative of the current spectrum of practice. While some practitioners may wean oxygen more quickly, we estimated that this timeframe mimics the time usually taken to first determine ROSC, obtain a reliable pulse-oximetry signal, and subsequently wean oxygen.

Two previous studies evaluated shorter periods of 100% oxygen use. Perez-de-sa *et al.*²⁹ limited 100% oxygen to 3 minutes from the start of ventilation in less asphyxiated lambs that mostly did not require CPR. Linner *et al.*³⁰ limited 100% oxygen to 3 minutes from commencing CPR in non-transitioning piglets. Consistent with our findings, both studies report that even this brief exposure was insufficient to prevent cerebral hyperoxia. Markers of injury or oxidative stress were not reported. Importantly, we showed that the hyperoxia was primarily driven by cerebral overperfusion, as evident by significantly elevated carotid

blood flow in the standard care group compared to the rapid wean and air groups. The cerebral overperfusion is driven by a rapid increase in pulmonary blood flow, consequent to a rapid oxygen dependent fall in pulmonary vascular resistance, which increases cardiac output. Chandrasekharan and colleagues previously showed higher pulmonary blood flow in the minutes after birth in preterm lambs resuscitated with 100% oxygen compared to those ventilated with 21%.³¹ Our observations suggest that the ability of cerebral vascular tone to autoregulate to a rapid increase in partial pressure of oxygen is severely impaired following acute, severe asphyxia. This is in contrast to the cerebrovascular response during re-oxygenation with supplemental oxygen following partial, prolonged hypoxia, where the brain is protected from an acute rise in cerebral blood flow.³² These findings highlight the critical relationship between oxygen, pulmonary and cerebral perfusion and cerebral hyperoxia immediately after resuscitation.

Our data shows that using 100% oxygen until 5 minutes after ROSC, or using air and increasing oxygen from 5 minutes after ROSC, decreases both Complex I- and Complex II-driven respiration, Complex IV activity, and maximal ETS capacity in the striatum, suggesting that the oxidative machinery is not able to synthesize ATP at rates necessary for maintaining electrochemical gradients to support brain function in the first hour after asphyxia.³³ When respiratory rates were normalized to maximal ETS capacity, they were no longer lower, suggesting that the apparent limitation of mitochondrial OXPHOS capacity observed in these two groups is related to an overall decrease in absolute respiratory rates and ETS capacity. These findings support previous studies in 4-week-old piglets following asphyxia and subsequent exposure to 100% oxygen.^{13–15} Persistent decreases in mitochondrial respiration may ultimately limit cerebral recovery, and play an important role in the development of secondary neurologic injury and contribute to long-term cognitive dysfunction. Indeed, in terms of amino acid neurochemistry 4 days after a partial, prolonged hypoxic insult, re-oxygenation in air following was found to best balance the competing dangers of free-radical formation and ongoing hypoxic cell death.³⁴

Figure 3. Continued.

radicals were quantified in the rapid-wean and air groups compared to standard care. Volcano plots show fold change [$\log_2(\text{fold change})$; i.e., $\log_2(\text{fold change} = 2) = 1$; x-axis] and all gene expression changes [$-\log_2(\text{adjusted P-value})$; y axis]. The grey dashed line indicates the threshold of significance (adjusted $P < 0.05$, using Benjamini-Hochberg correction for multiple comparisons). Genes that were downregulated are presented in red. Asterisks in black represent statistically significant differences between the rapid-wean versus standard oxygen group, and rapid-wean versus air groups, respectively ($n = 5\text{--}7/\text{group}$). * $p < 0.05$, ** $p < 0.01$; t-test, Mann-Whitney U test, or one-way ANOVA, with Tukey post hoc tests as appropriate.




Figure 4. Integrated analysis of the cortex following severe perinatal asphyxia and different re-oxygenation strategies. (a) Mitochondrial respiration rates in the cerebral cortex for lambs in the standard oxygen treatment group (“Standard”, orange), rapid-wean group (“Rapid”, blue), and air group (“21%”, black) for leak respiration (CI_L), Complex I-linked respiration (CI_P), CI&CII-linked respiration ($CI&CII_P$) and CII-linked respiration (CII_E), Complex IV activity, maximal electron transfer (ETS) capacity (Max ETS), Residual Oxygen Consumption, oxidative phosphorylation (OXPHOS) coupling efficiency (OXPHOS Efficiency). (b) Protein abundance of electron transfer system complexes (Complex I [CI], Complex II [CII], Complex III [CIII], Complex IV [CIV], ATP synthase [ATPase]), citrate synthase (CS), 75-kDa glucose-regulated protein (GRP75), and 60-kDa heat shock protein (HSP60). Note: some proteins were probed on the same western blot (CI/CIII/ATPase and GRP75/CS). (c) Volcano plot visualizations of key genes involved in mitochondrial biogenesis, apoptosis and ER stress-induced apoptosis, antioxidant defence, and in the generation of free radicals

Continued.




Figure 5. Mitochondrial respiration rates normalized to maximal electron transfer (ETS) capacity. Relative respiration rates in the striatum (a) and cerebral cortex (b) for lambs in the standard oxygen treatment group (“Standard”, orange), rapid-wean group (“Rapid”, blue), and air group (“21%”, black) for leak respiration (CI_L/ETS), Complex I-linked respiration (CI_P/ETS), CI&CII-linked respiration (CI&CII_P/ETS) and CII-linked respiration (CII_E/ETS). ($n = 5-7/\text{group}$).

In the striatum, genes involved in mitochondrial biogenesis (*PGC-1 α* , *Tfam*), fusion (*Mfn1* and *Opa1*) and fission (*Drp1*) were decreased in the rapid-wean group relative to the standard care group, while mitochondrial Complex II protein abundance was higher. These changes may serve as a mechanism to stabilize existing mitochondrial structures and dampen oxidative stress produced by damaged mitochondria following asphyxia. The downregulation of the NADPH oxidases, major sources of ROS, and key genes involved in apoptosis and ER-stress-induced cell death (*Bax*, *Casp3* and *CHOP*), supports this hypothesis. The increase in Complex II protein abundance points towards an increased tricarboxylic acid (TCA) capacity, and could further serve as a mechanism to help maintain mitochondrial function and sustain ATP production following the asphyxic insult.

Cortical Complex I LEAK to OXPHOS capacity was found to be increased in the rapid-wean group relative to the standard care and air groups. Additionally, there was a decrease in the abundance of GRP75,

which is crucial for mitochondrial homeostasis and stress response, while citrate synthase, a proxy for mitochondrial content, was increased. These changes likely reflect compensatory mechanisms to maintain mitochondrial respiration and reduce mitochondrial stress following asphyxia. These results contrast previous studies in 4-week-old piglets, which reported significant decreases in Complex I and Complex II-driven respiration in the cortex following asphyxia and CPR.^{14,15} These region-specific differences in mitochondrial respiratory capacity and bioenergetics between studies likely reflect differences in cortical maturation, procedural variables such as duration of asphyxia, percent of inspired oxygen provided during CPR, location of tissue extraction, and time post-ROSC of tissue collection.

Our study has some limitations. We used a model of acute-severe asphyxia induced by complete umbilical cord occlusion. The etiology of severe asphyxia in human newborns is varied and may be caused by other acute, or partial, prolonged insults. The cerebrovascular responses

Figure 4. Continued.

were quantified in the rapid-wean and air groups compared to standard care. Volcano plots show fold change [$\log_2(\text{fold change})$; i.e., $\log_2(\text{fold change} = 2) = 1$; x-axis] and all gene expression changes [$-\log_2(\text{adjusted P-value})$; y axis]. No differences in gene expression were observed with the Benjamini-Hochberg correction for multiple comparisons. Asterisks in black represent statistically significant differences between the rapid-wean versus standard oxygen group, and rapid-wean versus air groups, respectively ($n = 5-7/\text{group}$). * $p < 0.05$, ** $p < 0.01$; t-test, Mann-Whitney U test, or one-way ANOVA, with Tukey post hoc tests as appropriate.

following ROSC may be different under these circumstances. Some infants with severe asphyxia may have lung disease such as meconium aspiration syndrome. Supplemental oxygen is likely to assist in the respiratory transition in these instances, but the balance of risks with regard to cerebral hyperoxia is unknown. We acknowledge that the use of an anesthetic may have influenced mitochondrial respiration. However, all experimental groups were administered the same concentration of anaesthetic, therefore, the observed differences in mitochondrial respiration between the groups are likely to remain meaningful. The major strengths include the use of a well-established large-animal model with similar transitional physiology to humans, and instrumentation to measure both regional cerebral tissue oxygenation and corresponding mitochondrial function.^{7,35} Future directions include investigating the potential mechanisms driving mitochondrial dysfunction following hyperoxia-mediated brain injury, such as mitochondrial swelling, other indices of oxidative stress, and the mitochondrial permeability transition pore, a central mediator in the cascade of events leading to mitochondrial dysfunction and tissue damage during oxidative stress.

Our findings indicate that interventions to prevent hyperoxia-mediated brain injury during resuscitation of neonates following acute-profound asphyxia should focus on the period following ROSC. The risk of cerebral hyperoxia occurs not during CPR, but rather during the post-asphyxial overshoot in cerebral blood flow. A “rapid-wean” strategy facilitates the maximizing of blood oxygen content to achieve ROSC,³⁶ while negating the risk of subsequent cerebral tissue hyperoxia and mitochondrial injury. With this strategy, brief exposure to supplemental oxygen in the seconds after ROSC prior to weaning may reduce the need for later supplemental oxygen (as seen in the 21% group), preventing delayed hyperoxia. Further work is needed to evaluate the effect of transient hyperoxia in partial, prolonged asphyxia, its interaction with therapeutic hypothermia, and in preterm asphyxia.^{37,38} Strategies to better target cerebral oxygenation and limit hyperoxia-mediated brain injury are needed.

Acknowledgements

We thank the staff at the Monash Animal Research Platform for their contribution to this research.

Authors' contributions

Shiraz Badurdeen: Conceptualization, Methodology, Investigation, Data Curation, Formal Analysis, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing.

Robert Galinsky: Conceptualization, Methodology, Investigation, Writing – Review & Editing.

Calum Roberts: Conceptualization, Methodology, Investigation, Writing – Review & Editing.

Kelly Crossley: Methodology, Investigation, Project administration, Validation, Writing – Review & Editing.

Valerie Zahra: Methodology, Investigation, Project administration, Validation, Writing – Review & Editing.

Alison Thiel: Methodology, Investigation, Project administration, Validation, Writing – Review & Editing.

Yen Pham: Methodology, Investigation, Validation, Writing – Review & Editing.

Peter Davis: Resources, Supervision, Funding acquisition, Writing – Review & Editing.

Stuart Hooper: Conceptualization, Methodology, Investigation, Resources, Supervision, Funding acquisition, Writing – Review & Editing.

Graeme Polglase: Conceptualization, Methodology, Software, Validation, Investigation, Resources, Supervision, Project administration, Funding acquisition, Writing – Review & Editing.

Emily Camm: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Resources, Supervision, Project administration, Writing – Original Draft Preparation, Writing – Review & Editing.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was supported by the National Health and Medical Research Council (NHMRC) through a Project grant (APP1158494) and Fellowships (CTR: APP1175634, PGD: APP1059111, SBH: APP545921, GRP: APP1105526). SB was supported by an Australian Government Research Training Program Scholarship. The funders had no role in the study design, in the collection, analysis and interpretation of data; in the writing of the manuscript; and in the decision to submit the manuscript for publication.

Declaration of conflicting interests


The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Supplementary material

Supplemental material for this article is available online.

ORCID iDs

Shiraz Badurdeen  <https://orcid.org/0000-0002-3362-6435>

Robert Galinsky  <https://orcid.org/0000-0002-6374-9372>

References

- Wyckoff MH, Greif R, Morley PT, et al. 2022 International consensus on cardiopulmonary

- resuscitation and emergency cardiovascular care science with treatment recommendations: Summary from the basic life support; advanced life support; pediatric life support; neonatal life support; education, implementation, and teams; and first aid task forces. *Resuscitation* 2022; 181: 208–288.
2. neoResus. 2021, <https://resus.org.au/the-arc-guidelines/> (accessed 20 July 2021).
 3. Soar J, Nolan JP, Böttiger BW, et al. European resuscitation council guidelines for resuscitation 2015: section 3. Adult advanced life support. *Resuscitation* 2015; 95: 100–147.
 4. Badurdeen S, Davis PG, Hooper SB, et al. Physiologically based cord clamping for infants $\geq 32+0$ weeks gestation: a randomised clinical trial and reference percentiles for heart rate and oxygen saturation for infants $\geq 35+0$ weeks gestation. *PLoS Med* 2022; 19: e1004029.
 5. Badurdeen S, Gill AW, Kluckow M, et al. Excess cerebral oxygen delivery follows return of spontaneous circulation in near-term asphyxiated lambs. *Sci Rep* 2020; 10: 16443.
 6. Badurdeen S, Roberts C, Blank D, et al. Haemodynamic instability and brain injury in neonates exposed to hypoxia–ischaemia. *Brain Sci* 2019; 9: 49.
 7. Polglase GR, Blank DA, Barton SK, et al. Physiologically based cord clamping stabilises cardiac output and reduces cerebrovascular injury in asphyxiated near-term lambs. *Arch Dis Child Fetal Neonatal Ed* 2018; 103: F530–F538.
 8. Polglase GR, Brian Y, Tantanis D, et al. Endotracheal epinephrine at standard versus high dose for resuscitation of asystolic newborn lambs. *Resuscitation* 2024; 198: 110191.
 9. Saugstad OD. Neonatal chest compressions: time to act. *Pediatr Res* 2021; 90: 510–512.
 10. Sankaran D, Vali P, Chen P, et al. Randomized trial of oxygen weaning strategies following chest compressions during neonatal resuscitation. *Pediatr Res* 2021; 90: 540–548.
 11. Sankaran D, Giusto EM, Lesneski AL, et al. Randomized trial of 21% versus 100% oxygen during chest compressions followed by gradual versus abrupt oxygen titration after return of spontaneous circulation in neonatal lambs. *Child (Basel, Switzerland)* 2023; 10: 575.
 12. De Vries LS and Groenendaal F. Patterns of neonatal hypoxic–ischaemic brain injury. *Neuroradiology* 2010; 52: 555–566.
 13. Lautz AJ, Morgan RW, Karlsson M, et al. Hemodynamic-directed cardiopulmonary resuscitation improves neurologic outcomes and mitochondrial function in the heart and brain. *Crit Care Med* 2019; 47: e241–e249.
 14. Kilbaugh TJ, Sutton RM, Karlsson M, et al. Persistently altered brain mitochondrial bioenergetics after apparently successful resuscitation from cardiac arrest. *J Am Heart Assoc* 2015; 4: e002232.
 15. Marquez AM, Morgan RW, Ko T, et al. Oxygen exposure during cardiopulmonary resuscitation is associated with cerebral oxidative injury in a randomized, blinded, controlled, preclinical trial. *J Am Heart Assoc* 2020; 9: e015032.
 16. Martini S, Austin T, Aceti A, et al. Free radicals and neonatal encephalopathy: mechanisms of injury, biomarkers, and antioxidant treatment perspectives. *Pediatr Res* 2020; 87: 823–833.
 17. Davis PG, Tan A, O'Donnell CPF, et al. Resuscitation of newborn infants with 100% oxygen or air: a systematic review and meta-analysis. *Lancet* 2004; 364: 1329–1333.
 18. Kilkenny C, Browne WJ, Cuthill IC, et al. The ARRIVE guidelines checklist animal research: reporting In vivo experiments. *Br J Pharmacol* 2010; 8: 8–9.
 19. Covert R, Schreiber M, Torgerson L, et al. Prediction of cerebral blood flow in fetal lambs by carotid artery ultrasonic flow transducer. *Reprod Fertil Dev* 1996; 8: 157–162.
 20. Davies KL, Smith DJ, El-Bacha T, et al. Development of cerebral mitochondrial respiratory function is impaired by thyroid hormone deficiency before birth in a region-specific manner. *FASEB J* 2021; 35: e21591.
 21. Davies KL, Smith DJ, El-Bacha T, et al. Cortisol regulates cerebral mitochondrial oxidative phosphorylation and morphology of the brain in a region-specific manner in the ovine fetus. *Biomolecules* 2022; 12: 768.
 22. Badurdeen S, Cheong JLY, Donath S, et al. Early hyperoxemia and 2-year outcomes in infants with hypoxic-ischemic encephalopathy: a secondary analysis of the infant cooling evaluation trial. *J Pediatr* 2024; 267: 113902.
 23. Wyckoff MH, Wyllie J, Aziz K, et al. Neonatal life support: 2020 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Circulation* 2020; 142: S185–S221.
 24. Solberg R, Andresen JH, Escrig R, et al. Resuscitation of hypoxic newborn piglets with oxygen induces a dose-dependent increase in markers of oxidation. *Pediatr Res* 2007; 62: 559–563.
 25. Solberg R, Longini M, Proietti F, et al. Resuscitation with supplementary oxygen induces oxidative injury in the cerebral cortex. *Free Radic Biol Med* 2012; 53: 1061–1067.
 26. Dannevig I, Solevåg A, Sonerud T, et al. Brain inflammation induced by severe asphyxia in newborn pigs and the impact of alternative resuscitation strategies on the newborn central nervous system. *Pediatr Res* 2013; 73: 163–170.
 27. Solevåg A, Dannevig I, Nakstad B, et al. Resuscitation of severely asphyctic newborn pigs with cardiac arrest by using 21% or 100% oxygen. *Neonatology* 2010; 98: 64–72.
 28. Koch JD, Miles DK, Gilley JA, et al. Brief exposure to hyperoxia depletes the glial progenitor pool and impairs functional recovery after hypoxic-ischemic brain injury. *J Cereb Blood Flow Metab* 2008; 28: 1294–1306.

29. Perez-de-Sa V, Cunha-Goncalves D, Nordh A, et al. High brain tissue oxygen tension during ventilation with 100% oxygen after fetal asphyxia in newborn sheep. *Pediatr Res* 2009; 65: 57–61.
30. Linner R, Werner O, Perez-De-Sa V, et al. Circulatory recovery is as fast with air ventilation as with 100% oxygen after asphyxia-induced cardiac arrest in piglets. *Pediatr Res* 2009; 66: 391–394.
31. Chandrasekharan P, Rawat M, Gugino SF, et al. Effect of various inspired oxygen concentrations on pulmonary and systemic hemodynamics and oxygenation during resuscitation in a transitioning preterm model. *Pediatr Res* 2018; 84: 743–750.
32. Richards JG, Todd KG, Emará M, et al. A dose-response study of graded reoxygenation on the carotid haemodynamics, matrix metalloproteinase-2 activities and amino acid concentrations in the brain of asphyxiated newborn piglets. *Resuscitation* 2006; 69: 319–327.
33. Magistretti PJ and Allaman I. A cellular perspective on brain energy metabolism and functional imaging. *Neuron* 2015; 86: 883–901.
34. Jantzie LL, Cheung PY, Obaid L, et al. Persistent neurochemical changes in neonatal piglets after hypoxia-ischemia and resuscitation with 100%, 21% or 18% oxygen. *Resuscitation* 2008; 77: 111–120.
35. Roberts CT, Klink S, Schmölder GM, et al. Comparison of intraosseous and intravenous epinephrine administration during resuscitation of asphyxiated newborn lambs. *Arch Dis Child Fetal Neonatal Ed* 2022; 107: 311–316.
36. Ramachandran S, Bruckner M, Wyckoff MH, et al. Chest compressions in newborn infants: a scoping review. *Arch Dis Child Fetal Neonatal Ed* 2023; 108: 442–450.
37. Dalen ML, Liu X, Elstad M, et al. Resuscitation with 100% oxygen increases injury and counteracts the neuroprotective effect of therapeutic hypothermia in the neonatal rat. *Pediatr Res* 2012; 71: 247–252.
38. Galinsky R, Dhillon SK, Kelly SB, et al. Magnesium sulphate reduces tertiary gliosis but does not improve EEG recovery or white or grey matter cell survival after asphyxia in preterm fetal sheep. *J Physiol* 2023; 601: 1999–2016.