

Received Date : 10-Sep-2016

Accepted Date : 25-Oct-2016

Article type : Research Letter

Corresponding Author Email ID: bircan.erbas@gmail.com

Letter to the Editor

Bircan Erbas¹, Som Kumar Shrestha¹, Shyamali C Dharmage², Connie Katelaris³, Janet Davies⁴ and Michael J. Abramson⁵

¹*La Trobe University, School of Psychology and Public Health, Melbourne, Australia,* ²*The University of Melbourne, Melbourne School of Population and Global Health, Melbourne, Australia,* ³*University of Western Sydney, School of Medicine, Sydney, Australia,* ⁴*Queensland University of Technology, Institute of Health and Biomedical Innovation, South Brisbane, Australia,* ⁵*Monash University, School of Public Health and Preventive Medicine, Melbourne, Australia*

Re: The effects of Air Pollution on asthma Hospital admissions in Adelaide, South Australia, 2003-2013: time series and case-crossover analysis

We read with great interest the recent study by Chen K and colleagues [1] which showed an increase risk of asthma hospital admissions in all age groups with total pollen. However, there are numerous issues that have been overlooked in the analysis of the pollen variable. If these issues were considered, the authors may have observed stronger species-specific effects thereby contributing to the evidence base on which pollen species is the major trigger of asthma exacerbations requiring hospitalization.

Pollens are specific to the local environment and Australia is geographically and climatically diverse; our recent work suggests significant variations in the distribution of airborne pollen across urban regions in Australia [2,3]. The study by Chen and colleagues contribute to the evidence base by showing significant effects of pollen on respiratory disease in Adelaide

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/cea.12847](https://doi.org/10.1111/cea.12847)

This article is protected by copyright. All rights reserved

which is a region with different exposure and climatic distributions compared to Melbourne where most of the Australian pollen studies arise from [4-6] .

It is important to know if grass pollen is the prime trigger in Adelaide too. Our current work analysing data from Sydney using a similar analytical approach confirms our earlier findings on pollen in that instantaneous effects are stronger by different species compared to cumulative effects. For example, an increase from the 75th to 90th percentile in same day grass pollen was associated with childhood asthma hospitalizations in Sydney (2008 to 2013) adjusted odds ratio=1.033 (95% CI 1.0044 to 1.0621) p=0.023 (unpublished data). If we are observing strong grass pollen effects in most regions in Australia we can therefore focus on interventions that minimize exposure to high grass pollen days thereby reducing the likelihood of an asthma exacerbation. It is difficult to assess this in this study from Adelaide when different pollens are collapsed into a total pollen count variable.

The authors observed effects of pollen in cool seasons but not in warm seasons in multi-pollutant models. While they hypothesize that pollen types in Adelaide in the cool season may be more allergenic than those in warm seasons, they do not provide data on pollen species by season in Adelaide. In fact, personal communication with the Adelaide Aerobiology Laboratory [Thomson K 2016, Adelaide Aerobiology Laboratory, oral communication, 19th August] suggest that grass is the prominent pollen during the August to March peak period and other pollens that are less allergenic occur during the cooler season. This chance finding should be interpreted with caution because all pollens are modelled together by season and absence of respiratory viral data during the cooler months would also confound these associations. The authors also do not report the p value from the interaction term in models with the exposure by season interactions. Including readmissions within 28 days is also a potential problem. Our previous work on admissions in children over a 1997 to 2009 suggests that's readmissions within 28 days peak in winter and can be as high as 21% on a weekly basis in children and adolescents. [7] It is difficult to diagnose asthma in the 0 to 2 year age group as the majority of children present with symptoms such as wheezy bronchitis, and recurrent upper respiratory tract infections and should be excluded from the analysis.

Identifying high risk groups is important and by extending the analysis to assess sex differentials may provide further insights. Childhood asthma is more prevalent among boys

and evidence suggests differences in pollen exposure effects between boys and girls. [5,8] Although, this study provides an important contribution a detailed description of the instrument used to collect the pollen, the location, height and counting protocol are important factors that need to be considered. [9] Further studies with uniform measures of pollen and similar analytical methods are necessary to fully understand the impact of different pollen species on allergic respiratory disease in the Australasian region.

References

1. Chen K, Glonek G, Hansen A, Williams S, Tuke J, Salter A, Bi P. The Effects of Air Pollution on Asthma Hospital Admissions in Adelaide, South Australia, 2003–2013: Time- Series and Case- crossover Analyses. *Clinical & Experimental Allergy*. 2016 Aug 1.
2. Beggs PJ, Katelaris CH, Medek D, Johnston FH, Burton PK, Campbell B, Jaggard AK, Vicendese D, Bowman DM, Godwin I, Huete AR. Differences in grass pollen allergen exposure across Australia. *Australian and New Zealand journal of public health*. 2015 Feb 1;39(1):51-5.
3. Haberle SG, Bowman DM, Newnham RM, Johnston FH, Beggs PJ, Buters J, Campbell B, Erbas B, Godwin I, Green BJ, Huete A. The macroecology of airborne pollen in Australian and New Zealand urban areas. *PloS one*. 2014 May 29;9(5):e97925.
4. Erbas B, Chang JH, Dharmage S, Ong EK, Hyndman R, Newbigin E, Abramson M. Do levels of airborne grass pollen influence asthma hospital admissions?. *Clinical & Experimental Allergy*. 2007 Nov 1;37(11):1641-7.
5. Erbas B, Akram M, Dharmage SC, Tham R, Dennekamp M, Newbigin E, Taylor P, Tang ML, Abramson MJ. The role of seasonal grass pollen on childhood asthma emergency department presentations. *Clinical & Experimental Allergy*. 2012 May 1;42(5):799-805.
6. Erbas B, Lowe AJ, Lodge CJ, Matheson MC, Hosking CS, Hill DJ, Vicendese D, Allen KJ, Abramson MJ, Dharmage SC. Persistent pollen exposure during infancy is associated with increased risk of subsequent childhood asthma and hayfever. *Clinical & Experimental Allergy*. 2013 Mar 1;43(3):337-43.
7. Vicendese D, Olenko A, Dharmage S, Tang M, Abramson M, Erbas B. Modelling and predicting low count child asthma hospital readmissions using General Additive Models. *Open Journal of Epidemiology*. 2013 Jul 29;2013.
8. Erbas, B., Dharmage, S. C., Tang, M. L., Akram, M., Allen, K. J., Vicendese, D., ... & Bardin, P. G. (2015). Do human rhinovirus infections and food allergy modify grass

pollen-induced asthma hospital admissions in children?. *Journal of Allergy and Clinical Immunology*, 136(4), 1118.

9. Davies, J. M., Beggs, P. J., Medek, D. E., Newnham, R. M., Erbas, B., Thibaudon, M., ... & Huete, A. R. (2015). Trans-disciplinary research in synthesis of grass pollen aerobiology and its importance for respiratory health in Australasia. *Science of The Total Environment*, 534, 85-96.

Author Manuscript