

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Erbas, B;Malamardi, S;Lambert, KA;Batra, M;Tham, R;Padukudru Anand, M

Title:

A systematic review of the evidence of outdoor air pollution on asthma hospital visits in children and adolescents in South Asia - a call for data

Date:

2021-01-01

Citation:

Erbas, B., Malamardi, S., Lambert, K. A., Batra, M., Tham, R. & Padukudru Anand, M. (2021). A systematic review of the evidence of outdoor air pollution on asthma hospital visits in children and adolescents in South Asia - a call for data. Wellcome Open Research, 6, pp.174-. <https://doi.org/10.12688/wellcomeopenres.16991.1>.

Persistent Link:

<https://hdl.handle.net/11343/290005>

License:

CC BY



SYSTEMATIC REVIEW

A systematic review of the evidence of outdoor air pollution on asthma hospital visits in children and adolescents in South Asia – a call for data [version 1; peer review: 2 approved]

Sowmya Malamardi ^{1,2}, Katrina A. Lambert¹, Mehak Batra¹, Rachel Tham ³, Mahesh Padukudru Anand ^{2*}, Bircan Erbas^{1,4*}

¹School of Psychology & Public Health, College of Science Health and Engineering, La Trobe University, Melbourne, 3086, Australia

²Department of Respiratory Medicine, JSS Academy of Higher Education & Research (JSSAHER), Mysuru, Karnataka, 570023, India

³Mary MacKillop Institute for Health Research, Australian Catholic University, Melbourne, 3086, Australia

⁴Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

* Equal contributors

V1 First published: 06 Jul 2021, 6:174
<https://doi.org/10.12688/wellcomeopenres.16991.1>
 Latest published: 06 Jul 2021, 6:174
<https://doi.org/10.12688/wellcomeopenres.16991.1>

Abstract

Background: Outdoor air pollution and childhood asthma are increasing problems in South Asian countries. However, little is known about the associations between levels of air pollution and severe childhood asthma requiring hospital treatment in these regions.

Methods: We undertook a systematic review to assess the evidence between outdoor air pollution exposure and childhood and adolescent asthma hospitalization in South Asia. MEDLINE, Web of Science, Google Scholar, CINAHL, Embase, Scopus, ProQuest Central databases were searched for peer-reviewed papers, and examination of reference lists was conducted for additional studies. We identified all the literature published in English up to January 2021 for the study population comprised of children aged less than 19 years. The search strategy was designed to identify all the studies and screen them as per the inclusion criteria. The method of qualitative synthesis using the standard tool determined the comprehensiveness of the assessment of bias.

Results: Of the original 367 studies screened three studies were ultimately included from India, Pakistan and Sri Lanka and a narrative synthesis was conducted. Although studies reported adverse effects of outdoor pollution on asthma hospitalizations, limitations in exposure assessments, varying definitions of asthma hospitalizations and limited data analysis were identified.

Conclusions: There is currently limited evidence that can provide meaningful risk estimates of the impact of outdoor air pollution on asthma hospitalizations during childhood and adolescence. Studies with comparable outcome definitions, appropriate exposure

Open Peer Review

Reviewer Status

	Invited Reviewers	
	1	2
version 1		
06 Jul 2021	report	report
1. Sanjeev Nair , Government Medical College, Thiruvananthapuram, India		
2. Balamugesh Thangakunam , Christian Medical College, Vellore, India		

Any reports and responses or comments on the article can be found at the end of the article.

assessments and study designs are needed to inform future public and environmental health policy.

PROSPERO registration: CRD42020156714 (28/04/2020)

Keywords

asthma, air pollution, South Asia, hospitalization



This article is included in the [Wellcome Trust/DBT India Alliance](#) gateway.

Corresponding author: Bircan Erbas (b.eras@latrobe.edu.au)

Author roles: **Malamardi S:** Data Curation, Formal Analysis, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Lambert KA:** Investigation, Methodology, Supervision, Validation; **Batra M:** Data Curation, Formal Analysis, Resources, Validation; **Tham R:** Investigation, Validation, Writing – Review & Editing; **Padukudru Anand M:** Conceptualization, Project Administration, Supervision, Writing – Review & Editing; **Erbas B:** Conceptualization, Methodology, Supervision, Validation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) received no financial support for the research, authorship, and/or publication of this article. Mahesh P A is a fellow of the DBT/Wellcome Trust India Alliance funded by the team science grant IA/TSG/19/1/600026.

Copyright: © 2021 Malamardi S *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Malamardi S, Lambert KA, Batra M *et al.* **A systematic review of the evidence of outdoor air pollution on asthma hospital visits in children and adolescents in South Asia – a call for data [version 1; peer review: 2 approved]** Wellcome Open Research 2021, 6:174 <https://doi.org/10.12688/wellcomeopenres.16991.1>

First published: 06 Jul 2021, 6:174 <https://doi.org/10.12688/wellcomeopenres.16991.1>

Introduction

Asthma is an important global health issue affecting people of all age groups and is one of the most common non-communicable diseases (NCD). Asthma prevalence is increasing and globally, it is the 16th leading cause of years lived with disability (YLD) among all ages and the 4th leading cause among children aged 5–14 years¹. Children bear the highest direct costs associated with asthma (hospital visits, doctor visits, diagnostic tests, and medication) as well as indirect costs such as school and work days lost due to illness and care responsibilities and impaired quality of life, or sometimes, premature death². With this growing trend, it is estimated that by 2025, there will be one hundred million additional cases of asthma³. Although respiratory diseases are expected to be the second highest contributor to economic burden by 2025, developing countries are frequently short of reliable estimates due to limited health system diagnosis and reporting, and impaired quality of care^{4,5}. Afghanistan, Bangladesh, Bhutan, Maldives, Nepal, India, Pakistan and Sri Lanka are developing countries in South Asia where the prevalence of asthma is increasing, but very few have been studied to understand the strongest risk factors leading to this situation^{6,7}.

The overall prevalence of asthma in Asia is 3.5%^{3,4} and has increased⁸ due to growing population, increased prevalence of chronic NCDs, increasing urbanization and rising air pollution levels^{4,8–10}. In South Asia, the overall prevalence of asthma among children aged < 19 years is 1.74%¹¹. However there are wide variations across countries, for example, 1.59% for Bangladesh and 6.14% for Afghanistan¹¹. Studies in Canada and the United Kingdom (UK) have observed an increase in hospitalization among children and young adults of South Asian ethnic groups over a period of 9 years^{12–14}.

Changes in environmental conditions including increasing urbanization, energy use and vehicular traffic are increasing and may contribute to asthma hospitalizations^{15,16}. Lifestyle factors such as smoking and obesity, along with environmental exposures which can include aeroallergens (pollen, fungal spores, dust, animal dander), air pollutants and respiratory viruses, individual characteristics such as being male and having genetic predispositions are associated with asthma hospitalizations in children and adolescents^{17–19}. In terms of major environmental factors, short and long-term exposures to air pollutants such as coarse particulate matter (PM) sized <10µm in aerodynamic diameter (PM₁₀), fine PM sized <2.5µm (PM_{2.5}), ultrafine PM (UFP), ozone (O₃), volatile organic compounds, hydrocarbons, nitrogen dioxide (NO₂) and sulfur dioxide (SO₂) may act as triggers for asthma exacerbation which may lead to hospitalization^{20–22}. South Asian countries are predominantly agrarian countries with extensive crop burning; have increasing motor vehicle use and rapid industrial development contributing to high levels of air pollution. The permissible levels of air pollutants set by public authorities varies across South Asian countries and cities²³ and the evidence around health effects remains inconclusive across multiple geographical locations

and sources of emissions. Children residing in the developing countries of South Asia have a greater risk of severity of disease and mortality due to asthma due to their developing lungs, and high volume of air intake relative to their size and their preponderance to engage in outdoor physical activities^{24–26}. Asthma exacerbations in children are associated with poor lung growth²⁷ which in turn is associated with reduced longevity over their life course^{28,29}. Hence children's respiratory health is critical for their long-term health and wellbeing but little is known about the impact of outdoor air pollution in these geographic locations.

To address this knowledge gap, we conducted a systematic review to assess the current evidence on the relationship between outdoor air pollution and asthma hospitalization of South Asian children and adolescents.

Methodology

This review is registered with the International Prospective Register of Systematic Reviews (PROSPERO Registration no. CRD42020156714, 28th April 2020) and we followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines for conducting reviews³⁰. The inclusion criteria were analytical research articles published in English up to January 2021 with the study population comprising children aged less than 19 years. In studies with adults and children, data specific for children needed to be presented separately in order to be included in the review; studies were conducted in a South Asian country which includes Afghanistan, Bangladesh, Bhutan, Maldives, Nepal, India, Pakistan and Sri Lanka; and studies must have explored at least one outdoor or ambient air pollutant and its relationship to asthma exacerbations requiring attendance to a hospital. There were no restrictions on the study design.

A comprehensive literature search was conducted using the electronic databases: [MEDLINE](#), [Web of Science](#), [Google Scholar](#) (first 10 pages), [CINAHL](#), [Embase](#), [Scopus](#) and [ProQuest Central](#), using combination of key terms and MeSH terms (Extended data, Table 1³⁰). Additional studies were included by examining the references of screened studies. After removing the duplicates, screening titles and abstracts, papers were identified for full text review and their eligibility was independently tested by two authors (SM and MB). Any conflicts between the two reviewers were then discussed with other coauthors before extracting the data.

Details of type of study design, location of study, study settings, year the study was conducted, age range and number of children studied, data on the exposure (primary and secondary considered), outcome, risk estimates with 95% confidence intervals (95% CI), confounders and adjustment covariates, or effect modification or interaction were extracted from the studies.

A quality assessment was conducted on the eligible studies using a validated quality assessment tool by Zaza *et al.*, 2020

with that evidence the quality of study based on study design, study sample, exposure assessment, outcome measures and adjustment for confounders³¹. Two authors (SM and MB) independently assessed each article, scored, and any differences resolved by consensus and inputs by other co-authors. A benchmark score of 75% and above as high-quality studies and below 75% set as low-quality studies.

Results

The electronic search and manual search found 367 peer-reviewed scientific articles after removing the duplicates (Figure 1). Of these, 275 were excluded following review of titles and abstracts. On conducting a full text review of the remaining 92 articles, all but three were excluded as they were found irrelevant and not meeting the inclusion criteria (Extended data, Table 2³⁰). The primary reason for the exclusion of studies was not meeting the outcome definition. For example, Awasthi *et al.*³² reports 157 of the hospitalized 386 cases were for asthma, but they treat this variable as a covariate, not an outcome. Similarly, in Singh *et al.*³³ while the asthmatic cohort after hospital admission enrolled in the study, the prospective six-month study outcome related to air pollution exposure was “asthmatic episode” or “asthma exacerbations” that not necessarily relates to another hospitalization. Also, the authors using the terms interchangeably and do not define them clearly³³. Studies in other countries such as Bangladesh reported asthma prevalence and air pollution associated asthma diagnosed by a physician but did not include asthma hospitalization³⁴. The secondary exclusion was the lack of outdoor air pollution exposure assessment. Some studies were excluded because they did not state the age group of the study participants³⁵ or did not report age stratified analyses specific to children³⁶. No relevant studies met the inclusion criteria from Afghanistan, Bangladesh, Bhutan, Maldives, Myanmar, Nepal or Sri Lanka. Only three papers were ultimately included as they fulfilled all inclusion criteria. Only one study scored fairly well in the quality assessment with a difference in the methodology as a measure of exposure, appropriate statistical methods used, controlling for key confounding variables and interpretation of findings. We were unable to conduct a meta-analysis of the results; instead we conducted a narrative synthesis of the results and present key issues pertaining to the other studies that met the initial search criteria at full text review.

The characteristics of the included studies are presented in Table 1. The three studies of South Asia were conducted in child populations from Kolkata, India³⁷, Lahore, Pakistan³⁸ and Colombo, Sri Lanka³⁹. In the Kolkata study children aged 5 to 18 years and in Lahore study only 2 to 15-year-old were included whereas in Colombo study all children less than 12 years were part of the study. All had a substantially large sample size and used time series analyses to assess the association between outdoor air pollution and the asthma hospitalization. Chakraborty *et al.*³⁷ included both Internal Classification of Diseases 10 (ICD-10) codes J45, J46, and ICD-9-CM code 493.90. The frequency of hospital admissions following an asthma attack or exacerbations were reported in Anwar *et al.*³⁸ Senanayake *et al.*³⁹ reported the total number of visits to an emergency treatment

unit (ETU) made by children presenting with acute wheezing episodes.

The outdoor air pollutants measured were NO₂ (in parts per billion- ppb), SO₂ (ppb), O₃ (ppb), CO (ppb), PM₁₀ (µg/m³), and total suspended particles (TSP) (µg/m³). The three studies reported indirect measure of population exposure to air pollutants measured at fixed monitoring stations as determined by the respective national pollution control board depending on the region. The findings of these studies are shown in Table 2. Chakraborty *et al.*³⁷ studied associations between with daily concentration levels of PM₁₀ and O₃ obtained from the monitoring station of the West Bengal pollution control board over 12 month period and asthma hospitalizations of children aged 5–18 years³⁷. This daily time series study used generalized additive models (GAM) to assess non-linear relationship among pollutants PM₁₀ and O₃. They found that PM₁₀ is associated with increased emergency asthma hospitalization in school aged children [0.0037, 0.007 (estimate, SE), p<0.01]. They also reported associations with O₃ [0.0022, 0.001 (estimate, SE), p<0.01]. They did not report the estimated effects as they modelled the non-linear smooth associations only. The authors adjusted for confounding factors such as humidity, temperature and wind speed. The study did not report any interactions between confounding variables for childhood asthma hospitalization (Table 2).

Anwar and colleagues³⁸ estimated the measure of pollutants: TSP, NO₂, NO, SO₂, and CO using data from the Environmental Protection Department, Lahore, Pakistan. In a study sample of 1135 children aged 2 to 15 years during October 2002 an increase in SO₂ concentration of 80 ppb was associated with increased number of asthma emergency visits. No statistical modelling was used to analyze the associations. They did not provide any age or sex stratification when describing the study participants³⁸.

In Colombo, Senanayake *et al.*³⁹ examined the children attending the ETU for asthma management (N=30,932) and for other than asthma (N=10,100). In the study, estimated concentration levels of SO₂ and oxides of nitrogen (NO_x) coincided with the frequency of the hospital visits and the effect estimates for SO₂ and NO₂ were 0.1- 0.01 parts per million (ppm) and 0.28- 0.04 ppm respectively. The weekly pattern of hospital emergency visits for asthma correlated with days of high pollutant concentration. This study reported correlation results only and did not perform any statistical modelling nor did they adjust for seasonality, age group and gender.

We conducted a quality assessment of the three studies and found significant issues with quality of reporting and risk of bias inherent in the studies. Overall, the outcome variable definition was unclear, and they conducted limited statistical analyses³⁸. Single monitoring station data were used in each study contributing to risk of exposure assessment misclassification^{37–40}. In known cases of asthmatic children, it is not unusual to have repeated asthma attacks leading to hospitalization; these studies did not discuss the issue of repeat admissions. Chakraborty




Figure 1. Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) flow diagram of the included studies in the review, January 2021.

Table 1. Characteristics of the included studies from South Asia evaluating outdoor air pollution concentrations and hospital admission due to asthma in children.
ICD=Internal Classification of Diseases.

Author, study design and analysis methods	Location of study (City, District, and Country)	Study Population	No. of children in the sample	Age group	Exposure (Pollutants) measurement	Exposure measurement	Outcome definition
Chakraborty <i>et al.</i> , (2014) Time series; Generalized Additive models (GAM).	Kolkata, West Bengal, India	Children attending the emergency department of Sir Nirlatan Sarkar Hospital and National Medical College Hospital due to asthma exacerbation, between 2009–2010	2,708	5–18 years	Particulate matter sized <10micrometer (PM10) ($\mu\text{g}/\text{m}^3$) and Ozone (O_3) (ppb)	Daily concentrates (PM_{10}) and Ozone (O_3) collected from West Bengal state pollution control board, India. Location: Victoria Memorial, Kolkata (approx. 10 km from hospitals.)	Asthma hospitalization: ICD-9-CM code 493.90; ICD-10 code J45/4d first discharge diagnosis: Emergency hospital admission due to asthma during 2008–2010
Anwar <i>et al.</i> , (2012); Time series; Frequency distribution and measure	Lahore, Pakistan	Children visiting the physician at Mayo Hospital and Children hospital at Lahore, Pakistan following an acute asthma exacerbation.	1135	2–15 years	Total soluble particles (TSP) $\mu\text{g}/\text{m}^3$, Nitrogen dioxide (NO_2), Nitric Oxide (NO), Sulfur dioxide (SO_2) in ppb and carbon monoxide (CO) ppm.	Monthly data on pollutants TSP, NO_2 , NO , SO_2 , and CO were collected from Emergency Protection Agency department during 2002 to 2003. Location: Urban Lahore, Pakistan	Hospital visits due to asthma and in acute asthma exacerbation cases
Senanayake <i>et al.</i> , (2001); Case control study; Correlational Analysis	Colombo, Sri Lanka	Children presenting with acute wheezing and no wheezing episode attend the emergency treatment unit (ETU) of Lady Ridgeway Hospital (LRH). Visits made between 1 st July 1998 to 30 th June 1999 by the children attending ETU with or without wheezing were studied.	Total 41,032 of children of which Cases=30,932 (with wheezing) Controls=10100 (without wheezing)	< 12 years	Sulfur dioxide (SO_2) (ppm) and oxides of nitrogen (NO_x) in ppm	Daily hourly concentrates of pollutants SO_2 and NO_x collected from air quality monitoring station of Environmental Division of the National Building Research Organization (NBRO), Central Environmental Authority of Sri Lanka. Location: Colombo Fort, Sri Lanka (approx. 3 km from Lady Ridgeway Hospital)	Emergency hospital visits made following an acute episode of wheezing associated asthma for emergency care and management

Table 2. Findings from the included studies from South Asia evaluating outdoor air pollution concentrations and hospital admission due to asthma in children.

Author	Findings/ Risk estimates	Covariates included in the models as Confounders	Interactions
Chakraborty <i>et al.</i> (2014)	Generalized additive models were used to analyze the daily time series data. No direct estimates were provided as they were smoothed. Estimate of particulate matter sized <10micro meter (PM ₁₀): LOESS method-0.0010 (0.0008), p<0.01 and by Spline method is 0.0037 (0.0007), p<0.01 Estimate of Ozone (O ₃): LOESS: 0.0024 (0.0018), p= 0.82 Spline method is 0.0022 (0.0011), p=0.0024. In addition, significant seasonal variation in daily asthma hospitalization was observed. There were two peak seasons, the first one (up to 24/day) was in spring- early summer (last week of March to mid-April) and the second one (up to 23/day) was in autumn (mid-September to October)	Temperature, Humidity, Rainfall and season.	None
Anwar <i>et al.</i> (2012)	No statistical modelling conducted. The recorded cases were proportionately high consistent with the higher concentration of SO ₂ and total soluble particulates (TSP) in the month of July to December and January respectively.	Although the weather data were not considered as confounders asthma attendance were high in the months of July to December corresponding to high humidity and lower temperature.	None
Senanayake <i>et al.</i> , (2001)	Estimated concentration of pollutants varied but followed a consistent pattern, Correlation co-efficient (r>0.6) and the estimates are SO ₂ : 0.1- 0.01 parts per million (ppm) NO ₂ : 0.28- 0.04 ppm Highest visits to the emergency treatment unit (ETU) were made on maximum pollutants (both SO ₂ & NO ₂) concentrations days. (p= 0.05) which co-occurred 16 times in 41 weeks study of SO ₂ and 13 times in 47 weeks study of NO ₂ among cases. The lowest no of visits per day to emergency treatment unit (ETU) associated with least daily pollutant concentrations. Such a presentation manifested 22 events in 48 weeks and 27 events in 49 weeks across SO ₂ and NO ₂ among cases.	None	None

*et al.*³⁷ reported the seasonal variations in the regression models but did not adjust for seasonality, gender or age.

Discussion

Initially, we sought to conduct a systematic review but after implementing the search, screening and conducting the eligibility check only three studies were eligible for inclusion. To our knowledge, this is the first review that has attempted to comprehensively synthesize the evidence, assessed the strengths, limitations and quality of studies reported from South Asia on the effect of outdoor air pollution and asthma related hospitalizations in children and adolescents. Although the three studies indicated some adverse effects, this review identified significant deficiencies in exposure assessments and outcome definitions.

The ecological time series study conducted in Kolkata by Chakraborty *et al.*³⁷ was one of the few studies with clear definitions of outcome definitions and 5 to 18 years age group and assessment of air pollutants from monitoring stations³⁷. They had sufficient sample size (n=2,708) to conduct a rigorous statistical analysis which captured the nonlinear relationships between PM₁₀ exposure and the outcome. Reasonable adjustments for confounding effects of temperature were also considered but no interaction or age, sex strata analyses were presented. The study conducted in Lahore was a cross sectional descriptive study, but we classified it as an ecological study due to the trend of hospitalization related to asthma were studied³⁸. On the other hand, the Colombo study was classified as time series though the author reported it as a case-control study³⁹. These three were focused on children and adolescents

residing in urban areas. Although there were other studies in the initial search with exposure measures from monitoring stations such as Singh *et al.*³³ the outcome definitions were not related to a hospital visit or admission because of asthma.

Seasonality is an important variable to consider when assessing outdoor air pollution exposure and asthma hospitalizations. Studying the effect of periods involving different seasons and weather conditions was not an objective of the included studies^{38,39}. The findings from two of the three included studies show that particulates PM₁₀ and TSP were high in the month of summer^{37,38}. Although most of the asthma-related hospitalization occurred in winter, they also peaked in summer in Chakraborty *et al.*³⁷ and during the transition from one season to another corresponding to October and November as an autumn season in a study by Anwar *et al.*³⁸. Seasonal variation is influenced by other environmental factors such as temperature, humidity, rainfall, wind, pollen grains, fungal spores, viral infections and others. The study at Kolkata controlled for the meteorological factors; average temperature, humidity and wind speed in the model³⁷. Spore concentration per day was adjusted along with covariates PM₁₀ and O₃ in the model and thus spore counts (per cubic meter per day) were associated with increased asthma hospitalizations during spring and autumn³⁷. The effect of temperature was important in both studies but only Chakraborty *et al.*³⁷ adequately modelled it. Neither study adjusted for other variables such as exposure to environmental tobacco smoke (ETS), dust allergens, household ventilation, history and current smoking. Moreover, they did not assess gender and age-specific effects.

Our review included a comprehensive systematic search of eight electronic databases, additional hand searching, and other narrative literature reviews on similar topics in these regions, hence presents unique results but with limited evidence. Although the search terms did include all countries in South Asia, most of the studies that fulfilled the initial search criteria were from India. Studies from Sri Lanka were excluded as their defined study population age varied widely, ranging from infants to adults⁴⁰. Though, respiratory admissions including asthma were described in studies from Nepal but no results or discussion specific to asthma were presented^{41,42}. Studies in our search from Pakistan and Bangladesh, described hospital admissions due to asthma, but studying air pollutants as exposure of interest was not their primary objective^{35,43}. Developing countries facing economic constraints with the growing burden of asthma morbidity may experience greater challenges in the coming years⁴⁴. It is worth understanding that there is a lack of studies in South Asian countries; therefore, there should be urgency among public health authorities in health impact assessment along with appropriate air quality measurement or modelling¹⁶. There is a

need to measure outdoor air pollutants on a real-time monitoring basis and studying trends of hospital admissions to fulfil current gaps; lack of long-term studies is also the highlight of our findings¹³. Our review was limited to studies in English language which might have missed some eligible studies published in other languages. But this is unlikely given that majority of the research work in South Asian countries are published within English language journals. We have presented key limitations in assessing the findings of studies included in our review. Diverse methodologies, including some with no statistical analysis at all and different statistical methods to assess exposure and outcome associations in the studies, made it impossible to conduct a meta-analysis.

In summary, we have shown that the evidence on the impact of air pollution on asthma hospital visits in children and adolescence in South Asia is scarce and a call for data is a priority. Outdoor air pollution is increasingly problematic but because of inconsistencies in exposure assessments, asthma diagnosis and hospitalization coding, controlling for key confounders and statistical analytical methods we were unable to synthesize the current studies and provide clear evidence of such associations. Respiratory health effects from outdoor pollution will continue to be a growing burden and more rigorous studies in South Asia are urgently needed.

Data availability

Underlying data

All data underlying the results are available as part of the article and no additional source data are required.

Extended data

Open Science Framework: Air Pollution and Asthma Hospital visits in Children_South Asia. <https://doi.org/10.17605/OSF.IO/XQRCH>³⁰.

This project contains the following extended data:

- Online Resource 1.docx (Table 1: Search Strategy)
- Online Resource 2.docx (Table 2: Studies from search history excluded from final synthesis with reasons)

Reporting guidelines

Open Science Framework: PRISMA checklist for 'A systematic review of the evidence of outdoor air pollution on asthma hospital visits in children and adolescents in South Asia - a call for data'. <https://doi.org/10.17605/OSF.IO/XQRCH>³⁰.

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

References

1. Global Burden of Disease 2015 Chronic Respiratory Disease Collaborators: **Global, regional, and national deaths, prevalence, disability-adjusted life years, and years lived with disability for chronic obstructive pulmonary disease and asthma, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015.** *Lancet Respir Med.* 2017; 5(9): 691–706. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
2. Bahadori K, Doyle-Waters MM, Marra C, *et al.*: **Economic burden of asthma: a systematic review.** *BMC Pulm Med.* 2009; 9: 24. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. Masoli M, Fabian D, Holt S, *et al.*: **The global burden of asthma: executive summary of the GINA Dissemination Committee report.** *Allergy.* 2004; 59(5): 469–478. [PubMed Abstract](#) | [Publisher Full Text](#)
4. Global Asthma Network: **Global Asthma Report-2018.** Report, Global Asthma Network, Auckland, New Zealand, 2018. [Reference Source](#)
5. World Health Organization: **Air pollution levels rising in many of the world's poorest cities.** 2016. Accessed November 4, 2020. [Reference Source](#)
6. Bishwajit G, Tang S, Yaya S, *et al.*: **Burden of asthma, dyspnea, and chronic cough in South Asia.** *Int J of Chron Obstruct Pulmon Dis.* 2017; 12: 1093–1099. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. Paudel U, Pant KP: **Beyond Smoking: environmental determinants of asthma prevalence in western Nepal.** *J Health Pollut.* 2020; 10(25): 200310. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. Song WJ, Kang MG, Chang YS, *et al.*: **Epidemiology of adult asthma in Asia: toward a better understanding.** *Asia Pac Allergy.* 2014; 4(2): 75–85. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
9. Ramanakumar AV, Aparajita C: **Respiratory disease burden in rural India: a review from multiple data sources.** *Int J Epidemiol.* 2005; 2(2): 2–10.
10. Pawankar R, Wang JY: **Climate Change, air pollution, and biodiversity in Asia Pacific and impact on respiratory allergies.** *Immunol Allergy Clin North Am.* 2020; 41(1): 63–71. [PubMed Abstract](#) | [Publisher Full Text](#)
11. Institute of Health Metric and Evaluation: **Global Burden of Disease, Compare** Seattle, WA: IHME, University of Washington, 2015 [2020, accessed 11 July 2020].
12. Senthilvelan A, Habbick BF: **Increased asthma hospitalizations among registered Indian children and adults in Saskatchewan, 1970–1989.** *J Clin Epidemiol.* 1995; 48(10): 1277–1283. [PubMed Abstract](#) | [Publisher Full Text](#)
13. Hussein S, Partridge M: **Perceptions of asthma in South Asians and their views on educational materials and self-management plans: a qualitative study.** *Patient Educ Couns.* 2002; 48(2): 189–194. [PubMed Abstract](#) | [Publisher Full Text](#)
14. Sheikh A, Steiner MF, Cezard G, *et al.*: **Ethnic variations in asthma hospital admission, readmission and death: a retrospective, national cohort study of 4.62 million people in Scotland.** *BMC Med.* 2016; 14: 3. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Pollock J, Shi L, Gimbel RW: **Outdoor environment and pediatric asthma: an update on the evidence from North America.** *Can Respir J.* 2017; 2017: 8921917. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Dezateux C, Stocks J, Dundas I, *et al.*: **Impaired airway function and wheezing in infancy: the influence of maternal smoking and a genetic predisposition to asthma.** *Am J Respir Crit Care Med.* 1999; 159(2): 403–410. [PubMed Abstract](#) | [Publisher Full Text](#)
17. Shore SA, Fredberg JJ: **Obesity, smooth muscle, and airway hyperresponsiveness.** *J Allergy Clin Immunol.* 2005; 115(5): 925–927. [PubMed Abstract](#) | [Publisher Full Text](#)
18. Beuther DA, Weiss ST, Sutherland ER: **Obesity and asthma.** *Am J Respir Crit Care Med.* 2006; 174(2): 112–119. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Erbas B, Kelly AM, Physick B, *et al.*: **Air pollution and childhood asthma emergency hospital admissions: estimating intra-city regional variations.** *Int J Environ Health Res.* 2005; 15(1): 11–20. [PubMed Abstract](#) | [Publisher Full Text](#)
20. Babin SM, Burkum HS, Holtry RS, *et al.*: **Pediatric patient asthma-related emergency department visits and admissions in Washington, DC from 2001–2004, and associations with air quality, socio-economic status and age group.** *Environ Health.* 2007; 6: 9. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Gleason JA, Bielory L, Fagliano JA: **Associations between ozone, PM_{2.5}, and four pollen types on emergency department pediatric asthma events during the warm season in New Jersey: a case-crossover study.** *Environ Res.* 2014; 132: 421–429. [PubMed Abstract](#) | [Publisher Full Text](#)
22. Friedman MS, Powell KE, Hutwagner L, *et al.*: **Impact of Changes in Transportation and Commuting Behaviors During the 1996 Summer Olympic Games in Atlanta on Air Quality and Childhood Asthma.** *JAMA.* 2001; 285(7): 897–905. [PubMed Abstract](#) | [Publisher Full Text](#)
23. Joss MK, Eeftens M, Gintowt E, *et al.*: **Time to harmonize national ambient air quality standards.** *Int J Public Health.* 2017; 62(4): 453–62. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Lai CKW, Beasley R, Crane J, *et al.*: **Global variation in the prevalence and severity of asthma symptoms: phase three of the International Study of Asthma and Allergies in Childhood (ISAAC).** *Thorax.* 2009; 64(6): 476–483. [PubMed Abstract](#) | [Publisher Full Text](#)
25. Kaiser SV, Johnson MD, Walls TA, *et al.*: **Pathways to Improve Pediatric Asthma Care: A Multisite, National Study of Emergency Department Asthma Pathway Implementation.** *J Pediatr.* 2020; 223: 100–107.e2. [PubMed Abstract](#) | [Publisher Full Text](#)
26. Sheikh SI, Ryan-Wenger NA, Pitts J, *et al.*: **Impact of asthma severity on long-term asthma control.** *J Asthma.* 2021; 58(6): 725–734. [PubMed Abstract](#) | [Publisher Full Text](#)
27. Belgrave DC, Buchan I, Bishop C, *et al.*: **Trajectories of lung function during childhood.** *Am J Respir Crit Care Med.* 2014; 189(9): 1101–1109. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Mannino DM, Diaz-Guzman E, Buist S: **Pre- and post-bronchodilator lung function as predictors of mortality in the Lung Health Study.** *Respir Res.* 2011; 12(1): 136. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Baughman P, Marott JL, Lange P, *et al.*: **Combined effect of lung function level and decline increases morbidity and mortality risks.** *Eur J Epidemiol.* 2012; 27(12): 933–943. [PubMed Abstract](#) | [Publisher Full Text](#)
30. Malamardi S: **Air Pollution and Asthma Hospital visits in Children_South Asia.** 2021. <http://www.doi.org/10.17605/OSF.IO/XQRCH>
31. Zaza S, Wright-De Agüero LK, Briss PA, *et al.*: **Data collection instrument and procedure for systematic reviews in the Guide to Community Preventive Services.** Task Force on Community Preventive Services. *Am J Prev Med.* 2000; 18(1 Suppl): 44–74. [PubMed Abstract](#) | [Publisher Full Text](#)
32. Awasthi S, Gupta S, Maurya N, *et al.*: **Environmental Risk Factors for Persistent Asthma in Lucknow.** *Indian J Pediatr.* 2012; 79(10): 1311–1317. [PubMed Abstract](#) | [Publisher Full Text](#)
33. Singh DK, Kumar S, Singh GV, *et al.*: **Study on impact of air pollution on asthma among school going children residing in urban Agra.** *Indian J Allergy Asthma Immunol.* 2018; 32(2): 65–69. [Publisher Full Text](#)
34. Zaman K, Takeuchi H, Yunus M, *et al.*: **Asthma in rural Bangladeshi children.** *Indian J Pediatr.* 2007; 74(6): 539–543. [PubMed Abstract](#) | [Publisher Full Text](#)
35. Agarwal R, Jayaraman G, Anand S, *et al.*: **Assessing respiratory morbidity through pollution status and meteorological conditions for Delhi.** *Environ Monit Assess.* 2006; 114(1–3): 489–504. [PubMed Abstract](#) | [Publisher Full Text](#)
36. Kumarihamy RMK, Tripathi NK: **Geostatistical predictive modeling for asthma and chronic obstructive pulmonary disease using socioeconomic and environmental determinants.** *Environ Monit Assess.* 2019; 191(Suppl 2): 366. [PubMed Abstract](#) | [Publisher Full Text](#)
37. Chakraborty P, Chakraborty A, Ghosh D, *et al.*: **Effect of airborne *Alternaria* conidia, ozone exposure, PM₁₀ and weather on emergency visits for asthma in school-age children in Kolkata city, India.** *Aerobiologia.* 2014; 30(2): 137–148. [Publisher Full Text](#)
38. Anwar S, Noor-Ul-Ain MAS, Nasi JA: **Seasonal Changes and Children with Bronchial Asthma (2–15 Years Age) Reporting at Asthma Clinics of Mayo Hospital and Children Hospital, Lahore.** *Journal of Fatima Jinnah Medical University.* 2012; 6(3): 3–7. [Reference Source](#)
39. Senanayak MP, Samarakkody RP, Sumanasena SP, *et al.*: **A relational analysis of acute wheezing and air pollution.** *Sri Lanka J Child Health.* 2001; 30(3): 66–68. [Publisher Full Text](#)
40. Thishan Dharshana KG, Coowanitwong N: **Ambient PM₁₀ and respiratory illnesses in Colombo city, Sri Lanka.** *J Environ Sci Health A Tox Hazard Subst Environ Eng.* 2008; 43(9): 1064–1070. [PubMed Abstract](#) | [Publisher Full Text](#)
41. Shrestha SL: **Time series modelling of respiratory hospital admissions and geometrically weighted distributed lag effects from ambient particulate air pollution within Kathmandu Valley, Nepal.** *Environ Model Assess.* 2007; 12(3): 239–251. [Publisher Full Text](#)

42. Gurung A, Son JY, Bell ML: **Particulate matter and risk of hospital admission in the Kathmandu Valley, Nepal: a case-crossover study.** *Am J Epidemiol.* 2017; **186**(5): 573–580.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Ahmed A, Ahmed F, Raza MZ, *et al.*: **A descriptive analysis of asthma exacerbations and it's mortality in Karachi, Pakistan.** *J Aller Ther.* 2013; **11**: 2.
[Publisher Full Text](#)
44. Anenberg SC, Henze DK, Tinney V, *et al.*: **Estimates of the Global Burden of Ambient PM_{2.5}, Ozone, and NO₂ on Asthma Incidence and Emergency Room Visits.** *Environ Health Perspect.* 2018; **126**(10): 107004.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 1

Reviewer Report 07 January 2022

<https://doi.org/10.21956/wellcomeopenres.18761.r47446>

© 2022 Thangakunam B. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Balamugesh Thangakunam 

Department of Pulmonary Medicine, Christian Medical College, Vellore, Tamil Nadu, India

Asthma and respiratory infections are being increasingly recognized as causes of childhood morbidity. One of the proposed reasons is increasing air pollution. The authors have attempted to find out whether there is sufficient medical evidence to correlate air pollution to asthma episodes in children in South Asia. Authors have done a reasonably well strategized method of evidence synthesis and came up with only three studies from the sub-continent. They found that all three studies had significant issues with the quality of reporting and risk of bias and had also used data from a single monitoring station. This review highlights the lacunae of data on the impact of outdoor air pollution on asthma hospitalizations during childhood and adolescence and the urgent need for well conducted studies.

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Yes

Are sufficient details of the methods and analysis provided to allow replication by others?

Yes

Is the statistical analysis and its interpretation appropriate?

Yes

Are the conclusions drawn adequately supported by the results presented in the review?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Tuberculosis, lung cancer

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 25 August 2021

<https://doi.org/10.21956/wellcomeopenres.18761.r44860>

© 2021 Nair S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Sanjeev Nair** 

Department of Respiratory Medicine, Government Medical College, Thiruvananthapuram, Kerala, India

A systematic review in an important area, which is of growing importance due to the increasing morbidity and mortality due to air pollution in South Asia. The search has been done systematically, however the authors have stated the issue of scarcity of studies in this area and lack of standardisation in these studies. Most of the studies sought to measure exposure in terms of environmental measurements from routine measurement of air pollutions from centres close to the institution, which may not be the ideal measure of exposure at an individual level, and the association with outcome in Asthmatic children and adolescents. The association could have been better established if there was direct measurement of household level and individual level exposure to the children to air pollution and then association could be looked for with exacerbations. However, there is lack of data / published studies with this kind of data. With the limitations of available data, the authors have done what was possible in a systematic review. Studies have looked at individual measurements of air pollution in three of countries included in this study and the authors could refer to those studies to point out the opportunity for individual measurements of air pollution and establishing association with outcomes related to Asthma in future studies.¹

References

1. Shupler M, Hystad P, Birch A, Miller-Lionberg D, et al.: Household and personal air pollution exposure measurements from 120 communities in eight countries: results from the PURE-AIR study. *Lancet Planet Health*. 4 (10): e451-e462 [PubMed Abstract](#) | [Publisher Full Text](#)

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Yes

Are sufficient details of the methods and analysis provided to allow replication by others?

Yes

Is the statistical analysis and its interpretation appropriate?

Not applicable

Are the conclusions drawn adequately supported by the results presented in the review?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Tuberculosis, COPD and Asthma, Air pollution

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.
