



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Parameswaran, R;Kalman, JM

Title:

Left atrium and cardiovascular risk: Does functionality matter more than size?

Date:

2018-04-01

Citation:

Parameswaran, R. & Kalman, J. M. (2018). Left atrium and cardiovascular risk: Does functionality matter more than size?. *Journal of the American Heart Association*, 7 (7), <https://doi.org/10.1161/JAHA.117.008930>.

Persistent Link:

<https://hdl.handle.net/11343/254837>

License:

[CC BY-NC-ND](#)

Left Atrium and Cardiovascular Risk: Does Functionality Matter More Than Size?

Ramanathan Parameswaran, MBBS; Jonathan M. Kalman, MBBS, PhD

The left atrium (LA) is often regarded as a biomarker of cardiovascular risk. It serves several important functions: it modulates left ventricular filling and cardiovascular performance by functioning as a reservoir for pulmonary venous return during ventricular systole; it is a conduit for pulmonary venous return during early ventricular diastole; and it functions as a booster pump that augments ventricular filling during late ventricular diastole.¹ It is highly dynamic and demonstrates structural, functional, and electrical remodeling in a time-dependent manner in response to various pathophysiological insults. The hallmark of structural remodeling is atrial dilatation and is frequently accompanied by increased interstitial fibrosis with cardiac structural alterations.² Functional impairment may occur with or without alteration in LA size. Remodeling that occurs in the initial phase is often reversible,³ in contrast to remodeling that occurs with ongoing “stressors” over longer duration.⁴ These maladaptive changes have a strong association with a wide array of conditions, including heart failure,⁴ hypertension,⁵ myocardial ischemia,⁶ obesity,⁷ and obstructive sleep apnea.⁸ In tandem with structural change, electrophysiologic abnormalities develop, including global and regional conduction slowing and changes in refractoriness predisposing to onset of atrial arrhythmias and ultimately atrial fibrillation (AF).

Atrial size and function can be assessed with echocardiography, cardiac computed tomography, and cardiac magnetic

resonance, but because of its safety, relative low cost, and universal availability, echocardiography is the most widely applied first-line modality. Many of the earlier measures of atrial function were only measurable during sinus rhythm, making comparisons in subjects with AF challenging. These measures include the following: (1) peak “A” wave velocity and velocity time integral of transmitral flow in late diastole; (2) estimation of the fractional contribution of atrial contraction to total mitral inflow; and (3) the use of Doppler tissue imaging as a global measure of atrial function. Recent advances in 2- and 3-dimensional echocardiography, including in tissue Doppler imaging, and speckle tracking for strain analysis⁹ have allowed evaluation of atrial function during AF. These varying parameters have distinct strengths and weaknesses and are complementary in differing clinical scenarios. The LA function index (LAFI) uses volumetric measurement of the LA and is a composite rhythm-independent measure of LA function. It is a ratio that incorporates analogues of cardiac output, atrial reservoir function, and LA size and is calculated as follows: $LAFI = (LA \text{ emptying fraction} \times \text{velocity time integral across the left ventricular outflow tract}) / \text{largest LA volume measure in ventricular systole indexed to body surface area}$.¹⁰ Thus, LAFI increases proportionally to LA reservoir function and stroke volume, but it is inversely proportional to LA size.

It has become increasingly recognized that structural and functional remodeling are important factors that determine cardiovascular prognosis and risk stratification. Several studies have demonstrated this association in different population subsets. In a single-center study¹¹ that included ambulatory patients with heart failure and reduced ejection fraction (<40%) who were clinically stable and optimally treated, LAFI was a predictor of long-term survival. Patients in the lower LAFI quartile had more severe heart failure symptoms, renal function, and left ventricular systolic function. Subgroup analysis by heart rhythm (sinus rhythm versus AF) revealed that this remained consistent in both subgroups. LAFI was also noted to be a predictor of heart failure hospitalization in another population-based cohort study with stable coronary artery disease and preserved baseline ejection fraction ($\geq 50\%$).¹² Besides functional remodeling, LA enlargement alone estimated by LA volume indexed to body

The opinions expressed in this article are not necessarily those of the editors or of the American Heart Association.

From the Department of Cardiology, Royal Melbourne Hospital, Melbourne, Australia (R.P., J.M.K.); and Department of Medicine, University of Melbourne, Victoria, Australia (R.P., J.M.K.).

Correspondence to: Ramanathan Parameswaran, MBBS, Department of Cardiology, Royal Melbourne Hospital, Melbourne, Australia 3050. E-mail: ramanathan.parameswaran@mh.org.au

J Am Heart Assoc. 2018;7:e008930. DOI: 10.1161/JAHA.118.008930.

© 2018 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

surface area has also been demonstrated to predict adverse cardiovascular events in patients with stable coronary artery disease,¹³ valvular heart disease,¹⁴ and dilated cardiomyopathy.¹⁵

The association between echocardiographic parameters of atrial remodeling and AF has also been extensively documented. In a case-control cohort, structural and functional LA remodeling assessed by echocardiogram were independently and strongly associated with paroxysmal AF.¹⁶ Assessment of LA size and function has also been useful to predict the success of restoring sinus rhythm in patients with AF after either direct-current cardioversion¹⁷ or catheter ablation.¹⁸ Electrophysiologic mapping studies have demonstrated that these echocardiographic markers of remodeling are associated with both electrical conduction slowing and regional atrial fibrosis; both are also strongly linked to progression of AF.¹⁹ More recently, a retrospective cohort study of patients with AF showed that more severe LA fibrosis, as detected by late gadolinium enhancement on cardiovascular magnetic resonance imaging, was associated with increased major cardiovascular and cerebrovascular event risk, driven primarily by increased risk of stroke or transient ischemic attack.²⁰

The study by Sardana et al,²¹ in this issue of *Journal of the American Heart Association (JAHA)*, provides new insights into the predictive power of LA remodeling in incident AF and cardiovascular disease (CVD). In a prospective analysis of 1786 participants from a longitudinal community-based cohort, the authors investigated how LAFI, as an echocardiographic marker of structural and functional remodeling, may better characterize adverse LA remodeling and predict incident AF and CVD than existing measures. The CHARGE-AF (Cohorts for Heart and Aging Research in Genomic Epidemiology-AF) and the 10-year Framingham Heart Study CVD risk models were used to determine its association with AF and CVD, respectively. There were 3 key observations made over a median follow-up of 8.3 years (range, 7.5–9.1 years). First, LAFI was associated with incident AF (hazard ratio, 3.83) and CVD (hazard ratio, 2.20), an association that persisted after adjustment for both clinical prediction scores and for echocardiographic parameters associated with AF and CVD. More important, the association remained strong, even in patients with normal LA volume. In the current study, the authors also confirmed prior studies demonstrating that the anteroposterior LA diameter was associated with incident AF and CVD events. The LA diameter was measurable in most participants with echocardiographic images suboptimal for the measurement of LA volumes. However, addition of LAFI, LA diameter, or indexed maximum LA volume to prediction models for AF or CVD did not significantly improve model discrimination for either outcome. Thus, although LAFI may represent an important echocardiographic marker of adverse cardiovascular risk, even in the

presence of normal LA size, it does not seem to add further discrimination to already robust clinical prediction models.

The current study reiterates the clinical importance of assessing LA remodeling in cardiovascular prognostication. The study supports the use of LAFI, which provides a composite measure of LA structure and function. In comparison, simpler echocardiographic measures, such as LA volume indexed to body surface area, currently recommended for risk stratification by consensus guidelines, provide a more limited structural evaluation only. Apart from being rhythm independent, LAFI adjusts LA function for LA size and attempts to isolate atrial remodeling from ventricular systolic function. Future large prospective studies will be needed to determine whether LAFI is the ideal tool for evaluation of atrial remodeling, together with incident AF and cardiovascular risk. The current study emphasizes that in left atrial prediction of cardiovascular risk, function may matter more than size alone.

Disclosures

None.

References

- Blume GG, McLeod CJ, Barnes ME, Seward JB, Pellikka PA, Bastiansen PM, Tsang TSM. Left atrial function: physiology, assessment, and clinical implications. *Eur J Echocardiogr*. 2011;12:421–430.
- Kallergis EM, Manios EG, Kanoupakis EM, Mavrakis HE, Arfanakis DA, Maliaraki NE, Lathourakis CE, Chlouverakis GI, Vardas PE. Extracellular matrix alterations in patients with paroxysmal and persistent atrial fibrillation: biochemical assessment of collagen type-I turnover. *J Am Coll Cardiol*. 2008;52:211–215.
- Goette A, Honeycutt C, Langberg JJ. Electrical remodeling in atrial fibrillation: time course and mechanisms. *Circulation*. 1996;94:2968–2974.
- Sanders P, Morton JB, Davidson NC, Spence SJ, Vohra JK, Sparks PB, Kalman JM. Electrical remodeling of the atria in congestive heart failure: electrophysiological and electroanatomic mapping in humans. *Circulation*. 2003;108:1461–1468.
- Medi C, Kalman JM, Spence SJ, Teh AW, Lee G, Bader I, Kaye DM, Kistler PM. Atrial electrical and structural changes associated with longstanding hypertension in humans: implications for the substrate for atrial fibrillation. *J Cardiovasc Electrophysiol*. 2011;22:1317–1324.
- Alasady M, Shipp NJ, Brooks AG, Lim HS, Lau DH, Barlow D, Kuklik P, Worthley MI, Roberts-Thomson KC, Saint DA, Abhayaratna W, Sanders P. Myocardial infarction and atrial fibrillation: importance of atrial ischemia. *Circ Arrhythm Electrophysiol*. 2013;6:738–745.
- Mahajan R, Lau DH, Brooks AG, Shipp NJ, Manavis J, Wood JP, Finnie JW, Samuel CS, Royce SG, Twomey DJ, Thanigaimani S, Kalman JM, Sanders P. Electrophysiological, electroanatomical, and structural remodeling of the atria as consequences of sustained obesity. *J Am Coll Cardiol*. 2015;66:1–11.
- Dimitri H, Ng M, Brooks AG, Kuklik P, Stiles MK, Lau DH, Antic N, Thornton A, Saint DA, McEvoy D, Antic R, Kalman JM, Sanders P. Atrial remodeling in obstructive sleep apnea: implications for atrial fibrillation. *Heart Rhythm*. 2012;9:321–327.
- To ACY, Flamm SD, Marwick TH, Klein AL. Clinical utility of multimodality LA imaging: assessment of size, function, and structure. *JACC Cardiovasc Imaging*. 2011;4:788–798.
- Thomas L, Hoy M, Byth K, Schiller NB. The left atrial function index: a rhythm independent marker of atrial function. *Eur J Echocardiogr*. 2008;9:356–362.
- Sargento L, Vicente Simões A, Longo S, Lousada N, Palma dos Reis R. Left atrial function index predicts long-term survival in stable outpatients with systolic heart failure. *Eur Heart J Cardiovasc Imaging*. 2017;18:119–127.
- Welles CC, Ku IA, Kwan DM, Whooley MA, Schiller NB, Turakhia MP. Left atrial function predicts heart failure hospitalization in subjects with preserved

- ejection fraction and coronary heart disease: longitudinal data from the Heart and Soul Study. *J Am Coll Cardiol*. 2012;59:673–680.
13. Ristow B, Ali S, Whooley MA, Schiller NB. Usefulness of left atrial volume index to predict heart failure hospitalization and mortality in ambulatory patients with coronary heart disease and comparison to left ventricular ejection fraction (from the Heart and Soul Study). *Am J Cardiol*. 2008;102:70–76.
 14. Le Tourneau T, Messika-Zeitoun D, Russo A, Detaint D, Topilsky Y, Mahoney DW, Suri R, Enriquez-Sarano M. Impact of left atrial volume on clinical outcome in organic mitral regurgitation. *J Am Coll Cardiol*. 2010;56:570–578.
 15. Rossi A, Cicoira M, Zanolla L, Sandrini R, Golia G, Zardini P, Enriquez-Sarano M. Determinants and prognostic value of left atrial volume in patients with dilated cardiomyopathy. *J Am Coll Cardiol*. 2002;40:1425–1430.
 16. Schaaf M, Andre P, Altman M, Maucort-Boulch D, Placide J, Chevalier P, Bergerot C, Thibault H. Left atrial remodeling assessed by 2D and 3D echocardiography identifies paroxysmal atrial fibrillation. *Eur Heart J Cardiovasc Imaging*. 2017;18:46–53.
 17. Di Salvo G, Caso P, Lo Piccolo R, Fusco A, Martiniello AR, Russo MG, D'Onofrio A, Severino S, Calabro P, Pacileo G, Mininni N, Calabro R. Atrial myocardial deformation properties predict maintenance of sinus rhythm after external cardioversion of recent-onset lone atrial fibrillation: a color Doppler myocardial imaging and transthoracic and transesophageal echocardiographic study. *Circulation*. 2005;112:387–395.
 18. Hammerstingl C, Schwekendiek M, Momcilovic D, Schueler R, Sinning J-M, Schrickel JW, Mittmann-Braun E, Nickenig G, Lickfett L. Left atrial deformation imaging with ultrasound based two-dimensional speckle-tracking predicts the rate of recurrence of paroxysmal and persistent atrial fibrillation after successful ablation procedures. *J Cardiovasc Electrophysiol*. 2012;23:247–255.
 19. Nattel S, Burstein B, Dobrev D. Atrial remodeling and atrial fibrillation: mechanisms and implications. *Circ Arrhythm Electrophysiol*. 2008;1:62–73.
 20. King JB, Azadani PN, Suksaranjit P, Bress AP, Witt DM, Han FT, Chelu MG, Silver MA, Biskupiak J, Wilson BD, Morris AK, Kholmovski EG, Marrouche N. Left atrial fibrosis and risk of cerebrovascular and cardiovascular events in patients with atrial fibrillation. *J Am Coll Cardiol*. 2017;70:1311–1321.
 21. Sardana M, Lessard D, Tsao CW, Parikh NI, Barton BA, Nah G, Thomas RC, Cheng S, Schiller NB, Aragam JR, Mitchell GF, Vaze A, Benjamin EJ, Vasan RS, McManus DD. Association of left atrial function index with atrial fibrillation and cardiovascular disease: the Framingham Offspring Study. *J Am Heart Assoc*. 2018;7:e008435. DOI: 10.1161/JAHA.117.008435.

Key Words: Editorials • atrial fibrillation • atrium • cardiovascular outcomes • echocardiography • remodeling