

Measurements of tropospheric ice clouds with a ground-based CMB polarization experiment, POLARBEAR

S. TAKAKURA,<sup>1</sup> M. A. O. AGUILAR-FAÚNDEZ,<sup>2,3</sup> Y. AKIBA,<sup>4</sup> K. ARNOLD,<sup>5</sup> C. BACCIGALUPI,<sup>6,7</sup> D. BARRON,<sup>8</sup> D. BECK,<sup>9</sup> F. BIANCHINI,<sup>10</sup> D. BOETTGER,<sup>11</sup> J. BORRILL,<sup>12</sup> K. CHEUNG,<sup>13</sup> Y. CHINONE,<sup>13,1</sup> T. ELLEFLOT,<sup>5</sup> J. ERRARD,<sup>9</sup> G. FABBIAN,<sup>14</sup> C. FENG,<sup>15</sup> N. GOECKNER-WALD,<sup>13</sup> T. HAMADA,<sup>16</sup> M. HASEGAWA,<sup>17</sup> M. HAZUMI,<sup>17,4,1,18</sup> L. HOWE,<sup>5</sup> D. KANEKO,<sup>1</sup> N. KATAYAMA,<sup>1</sup> B. KEATING,<sup>5</sup> R. KESKITALO,<sup>12</sup> T. KISNER,<sup>12,19</sup> N. KRACHMALNICOFF,<sup>6</sup> A. KUSAKA,<sup>20,21</sup> A. T. LEE,<sup>13,20</sup> L. N. LOWRY,<sup>5</sup> F. T. MATSUDA,<sup>1</sup> A. J. MAY,<sup>22</sup> Y. MINAMI,<sup>17</sup> M. NAVAROLI,<sup>5</sup> H. NISHINO,<sup>17</sup> L. PICCIRILLO,<sup>22</sup> D. POLETTI,<sup>6</sup> G. PUGLISI,<sup>23</sup> C. L. REICHARDT,<sup>10</sup> Y. SEGAWA,<sup>4</sup> M. SILVA-FEAVER,<sup>5</sup> P. SIRITANASAK,<sup>5</sup> A. SUZUKI,<sup>20</sup> O. TAJIMA,<sup>24</sup> S. TAKATORI,<sup>17</sup> D. TANABE,<sup>4</sup> G. P. TEPLY,<sup>5</sup> AND C. TSAI<sup>5</sup>

<sup>1</sup>Kavli IPMU (WPI), UTIAS, The University of Tokyo, Kashiwa, Chiba 277-8583, Japan

<sup>2</sup>Departamento de Física, FCFM, Universidad de Chile, Blanco Encalada 2008, Santiago, Chile

<sup>3</sup>Department of Physics and Astronomy, Johns Hopkins University, Baltimore, MD 21218, USA

<sup>4</sup>Department of Particle and Nuclear Physics, SOKENDAI, Hayama, Kanagawa 240-0193, Japan

<sup>5</sup>Department of Physics, University of California, San Diego, CA 92093-0424, USA

<sup>6</sup>International School for Advanced Studies (SISSA), Via Bonomea 265, 34136, Trieste, Italy

<sup>7</sup>The National Institute for Nuclear Physics, INFN, Sezione di Trieste Via Valerio 2, I-34127, Trieste, Italy

<sup>8</sup>Department of Physics and Astronomy, University of New Mexico, Albuquerque, NM 87131, USA

<sup>9</sup>AstroParticule et Cosmologie (APC), Univ Paris Diderot, CNRS/IN2P3, CEA/Irfu, Obs de Paris, Sorbonne Paris Cité, France

<sup>10</sup>School of Physics, University of Melbourne, Parkville, VIC 3010, Australia

<sup>11</sup>Instituto de Astrofísica and Centro de Astro-Ingeniería, Facultad de Física, Pontificia Universidad Católica de Chile, Vicuña Mackenna 4860, 7820436 Macul, Santiago, Chile

<sup>12</sup>Computational Cosmology Center, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

<sup>13</sup>Department of Physics, University of California, Berkeley, CA 94720, USA

<sup>14</sup>Institut d'Astrophysique Spatiale, CNRS (UMR 8617), Univ. Paris-Sud, Université Paris-Saclay, bât. 121, 91405 Orsay, France

<sup>15</sup>Department of Physics, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

<sup>16</sup>Astronomical Institute, Graduate School of Science, Tohoku University, Sendai, Miyagi 980-8578, Japan

<sup>17</sup>Institute of Particle and Nuclear Studies, High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305-0801, Japan

<sup>18</sup>Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency (JAXA), Sagami-hara, Kanagawa 252-0222, Japan

<sup>19</sup>Space Sciences Laboratory, University of California, Berkeley, CA 94720, USA

<sup>20</sup>Physics Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

<sup>21</sup>Department of Physics, The University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan

<sup>22</sup>Jodrell Bank Centre for Astrophysics, University of Manchester, Manchester, M13 9PL, UK

<sup>23</sup>Department of Physics and KIPAC, Stanford University, Stanford, CA 94305, USA

<sup>24</sup>Department of Physics, Kyoto University, Kyoto, Kyoto 606-8502, Japan

(Received XXX; Revised XXX; Accepted XXX)

ABSTRACT

The polarization of the atmosphere has been a long-standing concern for ground-based experiments targeting cosmic microwave background (CMB) polarization. Ice crystals in upper tropospheric clouds scatter thermal radiation from the ground and produce a horizontally-polarized signal. We report the detailed analysis of the cloud signal using a ground-based CMB experiment, POLARBEAR, located at the Atacama desert in Chile and observing at 150 GHz. We observe horizontally-polarized temporal increases of low-frequency fluctuations (“polarized bursts,” hereafter) of  $\lesssim 0.1$  K when clouds appear in a webcam monitoring the telescope and the sky. The hypothesis of no correlation between polarized bursts and clouds is rejected with  $> 24\sigma$  statistical significance using three years of data. We consider many other possibilities including instrumental and environmental effects, and find no other reasons

other than clouds that can explain the data better. We also discuss the impact of the cloud polarization on future ground-based CMB polarization experiments.

*Keywords:* atmospheric effects — scattering — cosmology: observations — cosmic background radiation — polarization

## 1. INTRODUCTION

The atmosphere is an unavoidable foreground in any measurement with a ground-based telescope. Absorption, emission, and scattering by atmospheric molecules define the exploitable wavelength windows for astronomical observations. In addition, turbulence in the troposphere due to convective heat transfer causes variable weather conditions and reduces the observing efficiency.

In particular, cosmic microwave background (CMB) experiments observe the sky for thousands of hours to measure very faint anisotropies from the early universe, such as degree-scale parity-odd ( $B$ -mode) polarization anisotropies generated by primordial gravitational waves (Seljak & Zaldarriaga 1997). Atmospheric fluctuations introduce gradually varying (low-frequency) noise and degrade the CMB anisotropy measurements at large-angular scales (Lay & Halverson 2000). Therefore, the polarization of the atmosphere is a very significant concern for current and future ground-based CMB experiments.

Under the cloudless sky, the atmospheric transmission windows for CMB observation are typically <50, 70–110, 120–180, and 190–320 GHz bands. The atmospheric emission in this frequency range is dominated by oxygen and water vapor (e.g. Westwater et al. 2004). Fortunately, the emission is almost completely unpolarized (Kusaka et al. 2014; Errard et al. 2015), or slightly circularly polarized because of Zeeman splitting due to the Earth’s magnetic field (Rosenkranz & Staelin 1988; Keating et al. 1998; Hanany & Rosenkranz 2003; Spinelli et al. 2011). Although density and temperature fluctuations in the turbulent atmosphere cause significant low-frequency noise for CMB intensity (or temperature) measurements, they do not affect linear polarization measurements if the instrumental polarization leakage is negligible.

However, clouds in the atmosphere could produce linearly polarized microwave radiation. Clouds consist of small ice crystals, water droplets, or both depending on atmospheric conditions, and these small particles scatter the upwelling thermal radiation. The scattered light appears as a horizontally-polarized signal in the line of sight (Troitsky & Osharin 2000; Pietranera et al. 2007). Furthermore, the horizontal alignment of ice crystals having a column or plate shape (Ono 1969; Chepfer 1999) increases the polarization signal (Czekala 1998).

This linearly-polarized signal from anisotropic clouds is a source of low-frequency noise for linear polarization data. It cannot be mitigated even with ideal instruments or by other techniques such as polarization modulation (Brown et al. 2009).

The impact of the polarized signal from clouds for CMB polarization measurements is fully discussed in Pietranera et al. (2007) and partially mentioned in Kuo (2017). Measurements of the signal have been reported in the atmospheric science community using microwave radiometers (Troitsky & Osharin 2000; Troitsky et al. 2003, 2005; Kneifel et al. 2010; Xie et al. 2012; Defer et al. 2014; Xie et al. 2015; Pettersen et al. 2016; Gong & Wu 2017). In the CMB community, the BICEP2 Collaboration (2014) mentions the possibility of low-frequency noise ( $1/f$  noise) from clouds and ABS (Kusaka et al. 2018) reports the existence of noise flare-ups in the polarization signal.

In this paper, we report measurements of the polarization of clouds at POLARBEAR, a ground-based experiment observing CMB polarization at 150 GHz from the Atacama Desert in Chile. To our knowledge, this is the first detailed report of this kind of effect using a CMB instrument. One of the unique features of POLARBEAR is polarization modulation using a continuously-rotating half-wave plate (CRHWP) (Takakura et al. 2017; Hill et al. 2016). This technique mitigates the spurious polarization due to the leakage of unpolarized signals and instrumental temperature variations, and we can clearly measure the polarization from the sky.

In section 2, we briefly explain the basics of the scattering of microwave radiation by ice crystals within the clouds. In section 3, we show an example of polarization measurements during a cloudy day and then look for similar observations in 2.5 years of data. Following the results, we discuss the impact of the clouds on the CMB experiments in section 4 and summarize this study in section 5.

## 2. BASICS OF ICE CLOUDS

The POLARBEAR experiment is located at the James Ax Observatory, at an altitude of 5,200 m on Cerro Toco. This site, in the Atacama Desert in Northern Chile, is in one of the driest regions on the Earth (Suen et al. 2014). However, clouds still do occasionally exist there (e.g. Fig. 7 of Kuo 2017). Clouds form when a moist air

parcel is adiabatically expanded due to a rapid change in elevation and its water vapor content supersaturates. Clouds take various forms, which are typically classified into ten types depending on the atmospheric condition (e.g. Liou & Yang 2016). Since POLARBEAR does not have any dedicated instruments to monitor clouds, we cannot characterize them by ourselves. Here, we follow Erasmus & Van Staden (2001) and assume that the clouds at POLARBEAR site are mostly high clouds (cirrus, cirrocumulus, and cirrostratus), which form at altitudes around 5,000–13,000 m above sea level. The type of clouds will affect the amplitude and polarization fraction of the scattered radiation explained below. In this study, however, we focus on testing just the existence of the polarized signal from clouds, and the evaluation of the cloud properties are out of the scope. The cloud type is not very important for that purpose.

The high clouds consist mainly of ice crystals,<sup>1</sup> which have various properties depending on the atmospheric condition, i.e. temperature and water vapor content, as well as the evolution of the clouds. The mean effective size of an ice crystal is typically  $D_e \simeq 20\text{--}100\ \mu\text{m}$ . The ice water content (IWC), which is the density of ice in the clouds, is about  $10^{-3}\text{--}10^{-1}\ \text{g} \cdot \text{m}^{-3}$  (Rolland et al. 2000). This results in a number density  $n$  of about  $10^4\text{--}10^5\ \text{m}^{-3}$ . The ice water path (IWP), which is the total mass of ice crystals per unit area, is about  $1\text{--}10\ \text{g} \cdot \text{m}^{-2}$  (Kuo 2017). Thus the geometric thickness of the cirrus clouds  $\Delta h$  is about  $10^3\ \text{m}$ . To be precise, the thickness ranges typically from  $0.5 \times 10^3$  to  $7 \times 10^3\ \text{m}$ , but the impact of the variation is very small in our rough estimation of the cloud signal in section 2.1.

Ice crystals in clouds are not spherical. Small, primary crystals take the form of a hexagonal column and evolve to longer columns, larger hexagonal plates, or their aggregates depending on the temperature and humidity (Magono & Lee 1966). The aspherical shapes could cause alignment of the ice crystals due to the drag of the atmosphere (Ono 1969). One can see the signal as characteristic halos such as sundogs, circumzenithal/circumhorizontal arcs, and upper/lower tangent arcs (Cowley & Schroeder 2009).

### 2.1. Rayleigh Scattering

Since the size of ice crystals is sufficiently smaller than the wavelength, the scattering of microwave radiation by ice crystals is described by Rayleigh scattering. The

electric field  $\mathbf{E}_{\text{sc}}$  scattered by an ice crystal located at the origin is expressed as (e.g. Landau & Lifshitz 1960)

$$\mathbf{E}_{\text{sc}}(r\hat{\mathbf{n}}) = -\frac{\omega^2}{c^2 r} \hat{\mathbf{n}} \times (\hat{\mathbf{n}} \times \mathbf{P}), \quad (1)$$

where  $r$  is the distance from the origin,  $\hat{\mathbf{n}}$  is a unit vector toward the propagation direction,  $\omega$  is the angular frequency of the wave, and  $c$  is the speed of light. In general, the electric dipole moment  $\mathbf{P}$  is expressed as

$$\mathbf{P} = V\boldsymbol{\alpha}\mathbf{E}_{\text{in}}, \quad (2)$$

where  $V$  is the volume of the scatterer,  $\mathbf{E}_{\text{in}}$  is the incident electric field, and  $\boldsymbol{\alpha}$  is the polarizability matrix, calculated as

$$\boldsymbol{\alpha} = \frac{\epsilon - 1}{4\pi} \left[ \mathbf{I} + (\epsilon - 1)\boldsymbol{\Delta} \right]^{-1}, \quad (3)$$

where  $\mathbf{I}$  is the identity matrix and  $\boldsymbol{\Delta}$  is the depolarization factor, which is a positive definite symmetric matrix satisfying  $\text{Tr}(\boldsymbol{\Delta}) = 1$  and depends on the shape and orientation of the scatterer. In the case of spheroids, the  $\boldsymbol{\Delta}$  is parameterized as a diagonal matrix  $\text{diag}\{(1 - \Delta_z)/2, (1 - \Delta_z)/2, \Delta_z\}$ , where  $\Delta_z < 1/3$ ,  $\Delta_z = 1/3$ , and  $\Delta_z > 1/3$  correspond to the prolate (column), spherical, and oblate (plate) shapes, respectively. The relative permittivity  $\epsilon$  of ice is about 3.15 for microwave radiation (Warren & Brandt 2008).


In the simple case of spherical particles, we can obtain the total cross-section of the Rayleigh scattering  $\sigma_{\text{R}}$  as

$$\sigma_{\text{R}} = \frac{8\pi}{3} \frac{V^2 \alpha^2 \omega^4}{c^4} \propto D_e^6 \omega^4, \quad (4)$$

where  $\alpha = (3/4\pi)(\epsilon - 1)/(\epsilon + 2)$ . We can see the well-known dependence on the size of the scatterer  $D_e$  and the frequency of the light  $\omega$ . This strong dependence on the particle size and variation of the size distribution in clouds cause huge uncertainty in the prediction of  $\sigma_{\text{R}}$  by orders of magnitude. At the observing frequency of the POLARBEAR,  $\omega/(2\pi) = 150\ \text{GHz}$ , the cross-section  $\sigma_{\text{R}}$  becomes  $\sim 10^{-16}\ \text{m}^2$  for  $D_e = 20\ \mu\text{m}$  and  $\sim 10^{-12}\ \text{m}^2$  for  $D_e = 100\ \mu\text{m}$ . By assuming that all the particles in a cloud have the same size and using the typical number density of ice crystals,  $n \sim 10^4\text{--}10^5\ \text{m}^{-3}$ , and the typical thickness of cirrus clouds,  $\Delta h \sim 10^3\ \text{m}$ , we estimate the optical depth of the clouds as  $\tau \sim 10^{-9}\text{--}10^{-4}$ . Note that the estimate increases for 220 or 280 GHz due to the frequency dependence.

The calculation above has been substantially simplified by ignoring the size and shape distributions but does indicate that larger ice crystals in clouds are the main contributor to the scattering of microwave radiation and that the optical depth would be  $\lesssim 10^{-3}$  at most.

<sup>1</sup> We focus on ice clouds in this study, but the same model with different parameters can be applied to clouds that consist of water droplets. Since the water droplets are spherical and more absorptive than ice, the polarization fraction would be small.




**Figure 1.** Illustration of the microwave signal from clouds. Ice crystals scatter thermal emission from the ground and generate horizontal polarization.

However, scattering by clouds changes the direction of the thermal radiation from the ground and injects it into the line of sight. Since almost half of the solid angle as seen from the cloud is covered by the ground at ambient temperature, scattering of the ground emission at sub-percent levels could cause additional signal at the  $\sim 1$  K level. The clouds are randomly distributed in the sky and are gradually varying and moving due to atmospheric turbulence and wind, which leads to low-frequency variations of the signal from the clouds. Therefore, clouds can become an important source of low-frequency noise (see section 4 for further discussions). The cloud signal is significantly larger than the current detector noise level for CMB observations, which is lower than 1 mK over a few seconds of beam-crossing time, and thus can be detected instantaneously.

## 2.2. Polarization of Ice Clouds

There are mainly two types of effects that polarize the light scattered by the ice crystals. The first is due to the curvature of the ground. The second is due to the horizontal alignment of ice crystals with plate or column shape. We explain the two effects in the following. Our estimate and observation suggest the latter is dominant. Other subsidiary effects are also discussed in section 4.3.

The three-dimensional positional relations among the telescope, the clouds, and the ground produce polarization (Fig. 1). As shown in Eqs. (1) and (2), the polarization of the scattered light is determined by its scattering angle and the polarization of the incident light. By taking spherical coordinates  $(\theta, \phi)$  centered at the clouds and aligning the  $z$ -axis to zenith as shown in Fig. 2, the ground radiation can be expanded in spherical harmon-



**Figure 2.** Sketch showing microwave radiation emitted from the spherical ground, reaching a cloud above the observing site.

ics  $Y_l^m(\theta, \phi)$  with  $m = 0$  because of the axial symmetry. If we assume that the ground is a blackbody with a uniform temperature  $T_g$ , the expansion coefficients  $a_{l,0}$  are obtained as

$$a_{l,0} = 2\pi T_g \int_0^{\frac{\pi}{2} - \delta\theta} Y_l^0(\theta, 0) \sin\theta d\theta. \quad (5)$$

Here,  $\delta\theta$  is the look-down angle of the horizon from the clouds, which is approximately  $\sqrt{2h/R}$ , with the altitude of the clouds' center above sea level  $h$  and the radius of the Earth  $R$ . In particular, the monopole ( $l = 0$ ) and quadrupole ( $l = 2$ ) components are calculated as

$$\frac{a_{0,0}}{\sqrt{4\pi}T_g} = \frac{1 - \sin\delta\theta}{2} \quad \text{and} \quad \frac{a_{2,0}}{\sqrt{4\pi}T_g} = \frac{\sqrt{5}}{4} \sin\delta\theta \cos^2\delta\theta, \quad (6)$$

respectively. Again, if we assume spherical particles to deal with the scattering simply by the optical depth  $\tau$ , the Stokes parameters of the scattered light are calculated as (Hu & White 1997)


$$\begin{pmatrix} I \\ Q \end{pmatrix} \approx \frac{\tau T_g}{\sqrt{4\pi}} \begin{pmatrix} 1 \\ 0 \end{pmatrix} a_{0,0} + \frac{\tau T_g}{10} \begin{pmatrix} Y_2^0(\theta, \phi) \\ -\sqrt{6} Y_2^0(\theta, \phi) \end{pmatrix} a_{2,0}, \quad (7)$$

where  $Y_2^0(\theta, \phi) \equiv \sqrt{5/(16\pi)}(3\cos^2\theta - 1)$ ,  ${}_2Y_2^0(\theta, \phi) \equiv \sqrt{15/(32\pi)}\sin^2\theta$  and the other Stokes parameters,  $U$  and  $V$ , are zero. Here, the polarizations are defined on the usual base vectors  $(\mathbf{e}_\theta, \mathbf{e}_\phi)$ , and the negative Stokes  $Q$  represents horizontal linear polarization. The polarization fraction  $p$  is the ratio of  $Q$  to  $I$ , thus

$$|p| \approx \frac{3}{4\sqrt{5}} \frac{a_{2,0}}{a_{0,0}} \sin^2\theta \approx \frac{3}{4} \sqrt{\frac{h}{2R}} \sin^2\theta. \quad (8)$$

Putting  $h = 10$  km,  $R = 6,400$  km, and  $\theta = 45^\circ$  into Eq. (8) results in  $|p| \sim 1\%$ .

Horizontally aligned ice crystals with column and plate shapes scatter horizontal electric fields more efficiently and can, therefore, produce a larger polarized signal. We approximate the crystal shapes as spheroids and directly calculate Eqs. (1) to (3). In the case of column shape, we set the long axis horizontal but assume



**Figure 3.** Calculated polarization fraction of the light scattered by horizontally aligned ice crystals with a spheroidal shape as a function of the elevation. The dashed black lines show the spherical case. The line color of the other lines represents the shape information parameterizing the  $z$ -component of the depolarization factor  $\Delta_z$ . The column case shows the averaged contribution among randomly oriented particles in azimuth.

that its azimuth is random. The polarization fraction calculated from the model is shown in Fig. 3. Here, all the crystals are assumed to have the same shape and to be horizontally aligned, i.e. the long axis of the spheroid is in the horizontal plane. Thus, these estimates give an upper limit, whereas real polarization fractions are expected to be smaller. However, the amplitude of the polarization fraction is considerably larger than the spherical case shown as the black line. This is because the horizontally aligned crystals have a tendency to scatter horizontally polarized light even without a quadrupolar anisotropy of the incident radiation field.

### 3. MEASUREMENTS

We analyze the data taken by POLARBEAR (The POLARBEAR Collaboration: Ade et al. 2017) and search for signals that appear to be from tropospheric ice clouds. We use transition-edge sensor (TES) bolometers in the POLARBEAR receiver (Arnold et al. 2012) and a webcam monitoring the exterior of POLARBEAR and its surroundings, including the sky. Each TES bolometer is coupled to a dipole-slot antenna and measures a single polarization of the incident light. In addition, POLARBEAR has a CRHWP at the prime focus (Takakura et al. 2017; Hill et al. 2016), which is continuously rotated at 2 Hz to modulate the polarization

signal from the sky. Thus, the modulated timestream of the detector  $d_m(t)$  is expressed as

$$d_m(t) = I(t) + \text{Re}\{[A_0 + Q(t) + iU(t)]e^{-i\omega_m t}\}, \quad (9)$$

where  $I(t)$ ,  $Q(t)$  and  $U(t)$  are variations of the Stokes parameters of the sky,  $A_0$  is the steady polarization from the instruments, and  $\omega_m$  is the modulation frequency. Throughout the following analysis, the Stokes  $Q(t)$  and  $U(t)$  are defined on the instrumental coordinates ( $e_{ZE}, e_{AZ}$ ), where ZE and AZ represent the zenith and azimuth angle, respectively. We demodulate the timestream and extract the polarization signal as a demodulated timestream  $d_d(t)$ :

$$d_d(t) = A_0 + Q(t) + iU(t). \quad (10)$$


We can also obtain the intensity signal by applying a low-pass filter. See Takakura et al. (2017) for more details. For the analysis in section 3.1, no filters are applied anymore because the signal is very significant. For the analysis in section 3.3, we apply the polynomial filter, the scan-synchronous signal filter, and the intensity-to-polarization leakage filter (Takakura et al. 2017) to mitigate spurious contributions such as the responsivity variation of the detectors, polarized ground signals, and instrumental polarization. For the dataset used in this paper, the absolute polarization angle calibration is still preliminary and the calibration uncertainty is about a few degrees. While further work is in progress toward the final calibration for CMB science analysis, this preliminary calibration suffices for the purpose of the study presented in this paper.

The webcam is placed in the control container located 17.2 m north of the telescope. Since it is mainly used to monitor the telescope, its field of view (FOV) covers about  $130^\circ$ – $180^\circ$  in azimuth and  $-10^\circ$ – $20^\circ$  in elevation, and it takes a picture every 5 minutes.

#### 3.1. Example

Fig. 4 shows an example of the bolometer timestreams from a four-hour observation on December 18, 2014. During the observation, the telescope is azimuthally scanning the sky back and forth in an azimuth range of  $133^\circ$ – $156^\circ$  at a constant elevation of  $30^\circ$ . The precipitable water vapor (PWV) increases from  $\sim 0.8$  mm to  $\sim 1.6$  mm during the time of this data set. The PWV is provided by the APEX experiment (Güsten et al. 2006) using a commercial LHATPRO microwave radiometer (Rose et al. 2005). The APEX PWV will be partially correlated with the PWV at the POLARBEAR site 6 km away.

The intensity signal shown in the top panel is continuously fluctuating by  $\pm 3$  K like a random walk, which



**Figure 4.** Example timestreams from a single bolometer during a cloudy observation. The top panel shows the Stokes  $I$  (intensity) signal, and the middle panel shows demodulated Stokes  $Q$  and  $U$  polarization signals, with the offsets due to instrumental polarization. The bottom panel shows the slope of the simple linear regression between  $I$  and  $Q$ , which corresponds to the signed  $Q$  polarization fraction.

is due to atmospheric turbulence. The Stokes  $Q$  in the middle panel, on the other hand, has negatively directed, burst-like structures, down by as much as  $\sim 0.3$  K relative to an offset  $A_0 \sim 0.2$  K. The Stokes  $U$  also shown in the middle panel has much smaller variation than  $Q$ , which means that the burst-like signal is horizontally polarized. This property agrees with the expectation for the cloud signal that we described in section 2.2.


The bottom panel shows the slope of the simple linear regression between the intensity and  $Q$  polarization signals, i.e. the signed  $Q$  polarization fraction. The intensity signal is a combination of the clouds and atmosphere, whose contributions are both a few Kelvin. However, the time scale of the polarized-burst signal is shorter than that of the atmosphere. Thus, in this calculation, we apply a simple high-pass filter by subtracting the baseline for each  $\sim 50$ -second one-way scans and minimize the contribution of the atmosphere. In the absence of the bursts in  $Q$ , the polarization is  $\sim 0.1\%$ , which is consistent with the level expected solely due to instrumental intensity-to-polarization leakage and no atmospheric polarization (Takakura et al. 2017). On the other hand, the polarization fraction significantly increases to 5–10% at the timings of the bursts. The

polarization fraction of 10% at the elevation of  $30^\circ$  is larger than the estimate for the spherical case as Eq. (8) but can be explained by horizontally aligned column or plate crystals as shown in Fig. 3.


Fig. 5 is another illustration of the same data, which is the map of the  $Q$  polarization data as a function of the azimuth and time accompanied with snapshots from the webcam. Each horizontal row of the map corresponds to each leftward or rightward scan that takes 50 seconds. Around the time of 19:45:05, there is a structure within the row, which means that the variation of the polarized-burst signal is more rapid than the scan time. In the next two scans, the structure appears consistently but in different azimuth, which suggests that the source of the signal is moving within the scan area. The time scale of the motion is several minutes. The existence of the polarized-burst signals and their motion from right to left agree well with those of the clouds in the webcam. This result also supports the argument that the origin of the signal is a cloud. Note that the maximum elevation of the webcam FOV is  $20^\circ$  and does not exactly cover the scanning elevation of  $30^\circ$ . However, the size of the clouds in the photos are sufficiently large to cover most of the sky, thus we suppose that the clouds would expand to the line of sight of the telescope.

We have considered other possibilities to create the polarized-burst signal, but none of them can explain the data as good as the clouds. Sudden responsivity variations may couple to the steady instrumental polarization  $A_0$  and cause apparent variations in the  $Q$  timestream. However, it cannot explain the variation of the  $Q(t)$  to negative values in Fig. 4 because the responsivity of the TES detector does not change its sign. Besides, the  $2f$  signal, which is another stable optical signal from the CRHWP, does not exhibit such variations. Temperature variations of the primary mirror could change the instrumental polarization, but the polarization fraction of the effect is expected to be less than 0.1% (Takakura et al. 2017). The far sidelobe of the telescope may have larger polarization leakage and see the ground and another part of the sky. However, the spurious signal from the ground should stay in the same azimuth, while that from the sky should be gradual rather than burst-like fluctuations. Condensation and evaporation of water vapor on the primary mirror may also cause spurious polarization. We do not have concrete evidence to reject it, but the correlation of the polarized burst with the humidity is weaker than that with the cloud detection in section 3.3.

### 3.2. Cloud Detection Using the Webcam



**Figure 5.** Comparison of the bolometer timestream with the webcam. The Stokes  $Q$  polarization timestream is mapped as a function of the azimuth ordered by time and shown as color. There are gaps to tune the instruments every one hour, which are shown as black. The photos of the webcam are shown next to the map at the corresponding times. The azimuth range of the scan roughly corresponds to the left half in each of the photos. The photos show that the white clouds are carried by the wind across the sky from the right to the left. The polarized-burst signals in the map coincide with the photos with clouds and have the same trend from the right to the left.



**Figure 6.** Example of the cloud detection with a webcam image. The shadowed region shows the mask, which is fixed for all the images. The orange and black rectangles show patches with and without cloud detection in this image.

We analyze all of the photos from the webcam and obtain the statistics of the clouds at the POLARBEAR site, in the Atacama Desert in Chile. Note again that the FOV of the webcam covers a small fraction of the sky and that the telescope points to a sky region outside of the FOV in all the observations. The webcam images are not useful during the night. We also removed the pictures taken at dawn or dusk in the following analysis because the gradient of the brightness in the sky increases false detection rate.


The basic idea of the cloud detection algorithm is to find white regions in the picture. First, we mask the telescope, mountain, and ground, and split the sky into 29 patches as shown in Fig. 6. For each patch, we calculate the average Red-Green-Blue color ( $R, G, B$ ) among the pixels. Then, we convert the color into Hue-Saturation-Value color ( $H, S, V$ ), specifically (Smith 1978)

$$S = \frac{\max(R, G, B) - \min(R, G, B)}{\max(R, G, B)}, \quad (11)$$


$$V = \max(R, G, B). \quad (12)$$

We set the thresholds to detect the clouds in each patch as  $S < 0.1$  and  $V < 0.98$ , where the former condition rejects the blue sky and the latter cuts saturated pixels due to the Sun. The performance of the cloud detection is checked by eye for pictures from several days chosen randomly. The algorithm often fails to detect faint clouds as in Fig. 6 but rarely make false detections, which are occasionally caused by ghost images, i.e. images that appear at spurious positions due to multiple reflections.

Fig. 7 shows the rate of cloud detection for each patch, i.e. the number of shots with positive cloud detection in the patch divided by the total number of shots. Although there is a small gradient in elevation, possibly due to the difference in the area of the sky, there is no




**Figure 7.** Cloud detection rate for each patch of the sky shown in Fig. 6.



**Figure 8.** Annual variation of the cloud detection rate. Each line shows the result in the different year. The cloud detection rate increases in January and February every year because of Altiplanic winter.

clear tendency for clouds to appear in any particular region of the FOV. The clouds may be created outside of the webcam FOV, but they are expected to persist sufficiently long enough such that they will pass across the FOV.

Figs. 8 and 9 show annual and daytime variations of the cloud detection rate per shot. Here we flag a shot as “cloudy” when a cloud is detected in at least one patch according to the algorithm specified above. In Fig. 8, we can see the significant cloud detection rate increase around February, which is known as Altiplanic



**Figure 9.** Daytime variation of the cloud detection rate. The horizontal axis represents the time in the daytime, which is scaled with respect to the sunrise and sunset times to adjust their seasonal variations. The blue line shows the result from all the data, and the orange and green lines show the results in the high (January and February) and low (from July to December) seasons, respectively. The shadowed regions show the mask used to cut dawn and dusk.

winter (See Erasmus & Van Staden (2001) for more details about the climate of Northern Chile). During that season, the cloud detection rate seems to increase in the afternoon (Fig. 9). It might be due to the lift of the atmosphere heated up by thermal conduction from the ground, which is also heated by the sunlight. During July through December, on the other hand, the daytime trend is moderate.


Fig. 10 shows the correlation between the cloud detection rate per shot and the APEX PWV. There is a clear correlation between the cloud detection and PWV. This is an expected trend and provides additional supports and validation to our cloud detection method.

The overall cloud detection rate per shot is about 26%. Note that we use only the daytime photos and have no information during the night.

### 3.3. Cloud Detection in the Bolometer Data

By using the polarized-burst signals in the bolometer data, we also detect clouds as shown in Fig. 4. We use the POLARBEAR data from July 2014 to January 2017. The cloud detection results are compared with the webcam study described in section 3.2.

Since the typical size of the clouds should be larger than the FOV of POLARBEAR,  $3^\circ$ , the cloud signal is correlated among all the detectors. Thus, we can improve the sensitivity to detect the clouds by averaging the timestreams among all the detectors. Similar to



**Figure 10.** The dependence of the cloud detection rate on the PWV provided by the APEX experiment. Each blue point shows the rate for data in each PWV bin and the orange line shows the rate for all the data below the PWV. The cumulative fraction of the data volume is also shown as the black dashed line.

Kusaka et al. (2014), we separate the averaged polarization timestreams,  $Q(t)$  and  $U(t)$ , into two components using the principal component analysis (PCA) method as

$$X_1(t) + iX_2(t) = \{Q(t) + iU(t)\} e^{-i\phi}, \quad (13)$$

where the rotation  $\phi$  is determined to maximize the variance of  $X_1(t)$ . Since the secondary component  $X_2(t)$  is dominated by the detector white noise, the signal-to-noise ratio of the cloud is calculated as


$$\text{SNR} = \sqrt{\frac{\sigma(X_1)^2}{\sigma(X_2)^2} - 1}, \quad (14)$$

where the  $\sigma$  denotes the standard deviation. The polarization angle of the signal  $\psi$  is obtained as

$$\psi = \frac{\phi}{2}. \quad (15)$$

Here, we assume that the signal is almost horizontal as shown in Fig. 4, and constrain  $\pi/2 < \phi < 3\pi/2$ , which cannot be determined by PCA due to degeneracy.

Fig. 11 shows the histogram of the polarization angle of the polarized-burst signal  $\psi$  with  $\text{SNR} > 10$  for each scan. If we have any instrumental noise sources other than the clouds, e.g. the intensity leakage due to the detector non-linearity, the polarized ground structure, and the HWP encoder error, they could appear at angles  $\psi \approx 90^\circ$ . Having only the single peak at  $\psi \sim 90^\circ$  supports that such extra noises are not significant and that all the polarized-burst signals are most likely coming



**Figure 11.** The histogram of the polarization angle of the polarized-burst signal  $\psi$ . The mean, median, and root-mean-square (RMS) values are calculated from samples in the range of  $70^\circ < \psi < 110^\circ$ . There are no other bunches outside of the range shown. Note that we have not finalized the absolute angle calibration for this data set yet, and it might have an error of a few degrees.

from clouds. There are still possibilities of the responsivity variation due to electrical noise and the temperature variation of the primary mirror, but they cannot explain the coincidence with the webcam described below. Both the width of the peak by  $\text{RMS} = 2.68^\circ$  and the offset from  $\psi = 90^\circ$  might be due to a systematic error of the cloud signal or the instrument, which is still under investigation (see more discussions in [section 4.3](#)).

Each of the CMB observations typically takes one hour and contains 40–70 left and right scans at a constant elevation. We calculate the SNR and polarization angle  $\psi$  of the cloud signal for each scan and take the values with the highest SNR as representative of the CMB observation. On the other hand, the detection of clouds by the webcam is determined by a detection in at least one of the  $\sim 12$  photos taken during the observation. Note that the CMB observations are performed at all times of the day, but the webcam can be used during the daytime only.

Table 1 shows the coincidence of the cloud detection in the bolometer data with that in the webcam. For the data with the clouds in the webcam, the rate of the polarized-burst detection significantly increases to 46.3% compared to 1.3% for the data without the clouds. Assuming that the appearance of polarized bursts in the bolometer data and the appearance of clouds in the webcam each occur at their observed rates but are also independent, then the significance of observing such a strong

Data	Webcam	Polarized burst
Daytime	All	16.1% (295/1835)
	Cloud	46.3% (279/602)
	No cloud	1.3% (16/1233)
Night	—	9.7% (458/4735)

**Table 1.** Fractions of the one-hour observations containing polarized-burst signals found in the bolometer data for the data sets corresponding either to a concurrent cloud detected in the webcam or to no cloud detected. The fraction of burst-like signals occurring at night is also listed, though clouds cannot be identified by the webcam during the night. The original numbers of one-hour observations are shown in parentheses.

covariance between them is estimated to be  $> 24\sigma$ . Note again that the FOV of the webcam does not cover the sky regions of the CMB observations. That could be the reason for the deviation from the perfect separation. Also, the daytime rate of the polarized burst is 16.1%, which is smaller than the  $\sim 30\%$ , the fraction of data with clouds detected by the webcam ([section 3.2](#)). It is because CMB observations are not performed when  $\text{PWV} > 4$  mm, or the webcam is more sensitive to clouds than bolometers in this analysis using the thresholds above. For the night data, we have no information on the clouds from the webcam, but the polarized-burst signals are detected in 9.7% of the data.

Fig. 12 shows the distribution of the SNR for each data set. Note that it shows the partial data with high SNR (Table 1). Again, it shows the clear difference between the data with and without clouds detected by the webcam. Besides, the SNR distribution for the night data is very similar to that for the daytime data with clouds. As explained in [section 2.2](#), the clouds scatter the thermal emission from the ground, which takes place also at night. Our results support the expectation that the cloud signal also exists during the night.


## 4. DISCUSSION

In the previous section, we have shown that the polarized cloud signals are detected in the CMB observations with POLARBEAR. Here, we discuss the impact of the clouds on CMB polarization measurements.

### 4.1. Degradation of the Statistical Precision

CMB measurements require thousands of hours of observations with high-sensitivity detectors to measure the faint CMB signals. The cloud signal is just noise that lowers the quality of the observations.

One simple approach to reduce the impact of the clouds is to drop noisy data, but that would inevitably



**Figure 12.** The histogram of the maximum SNR of the polarized-burst signal in each observation normalized by the number of the observation. The blue and orange histograms show the results for the daytime data with and without clouds detected by the webcam, and the result for all the daytime data is shown as the dashed line. The result for the night data is also shown as the hatched histogram. The outliers in the blue histogram might be due to clouds outside of the webcam FOV.

reduce the observation efficiency.<sup>2</sup> It is possible to detect cloud signals in the bolometer data to an extent as performed in section 3.3. We find cloud-like polarized-burst signals in 16.1% (9.7%) of one-hour observations (Table 1) and in 5.2% (3.0%) of leftward or rightward scans during the daytime (night). One could use image analysis similar to section 3.2, in which we detect clouds in 26% of all the daytime shots of the webcam. As shown in Fig. 10, the fraction of data without clouds improves if we observe only in good PWV. But, this decreases the total amount of data.

Even with data cuts, we expect residuals from faint clouds below the detection threshold. These residuals will be present at low frequency with a  $1/f$  behavior and degrade the detector performance at large angular scales. Such residuals cannot be mitigated by polarization modulation techniques such as the CRHWP used in POLARBEAR.

If we detect the clouds using the bolometer data as performed in section 3.3, the residual noise level will depend on the SNR threshold of the cloud detection. For example, to achieve an SNR of 10, the power of the cloud signal should be 100 times larger than that of the

<sup>2</sup> Since the cloud signal only affects the Stokes  $Q$  polarization, it may be possible to save the Stokes  $U$  component.

detector noise. In other words, the residual  $1/f$  noise below the threshold could naïvely degrade the sensitivity of the CMB angular power spectra by 100 times in the worst case. Tightening the threshold will mitigate the contamination but also decrease the observation efficiency. Thus, optimization of the threshold is required to maximize performance.

The effect of clouds would depend on the telescope FOV and the detector beam size. Small-aperture telescopes with a large FOV have a high probability of seeing clouds. Large-aperture telescopes with small beams have a high instantaneous SNR of a cloud, because many detectors simultaneously observe the same cloud, which is larger than the beam size.

These studies could inform forecasting and optimization of future CMB experiments, such as CMB-S4 (Abazajian et al. 2016).

#### 4.2. Systematic Errors on CMB Measurements

Systematic errors due to the residual cloud signal are also of concern for CMB measurements.

The cloud signal is horizontally polarized, i.e.  $Q < 0$ , and not symmetric fluctuations between plus and minus. This highly non-Gaussian fluctuation could lead to possible systematics in the map as many map-making algorithms assume Gaussianity for noise fluctuations. This systematics can be mitigated to some extent by parallactic angle rotation.

The cloud signal would affect foreground removal due to its distinct frequency dependence. By using maps at multiple frequencies, we separate the CMB and the other foregrounds that are stationary in the sky, i.e. not associated to any Earth or atmospheric motion, such as the Galactic dust and synchrotron emissions. Since the cloud signal has the Rayleigh scattering spectrum as  $\propto \omega^6$  (see Eq. (4) with additional  $\omega^2$  from the spectrum of the ground emission), it would appear rising in frequency, similar to a dust component (approximately  $\omega^{1.5}$ ), but much steeper. On the other hand, the atmospheric motion would likely de-correlate with astrophysical foregrounds, making the control of the clouds appropriate for a time domain analysis, rather than in maps.

Clouds staying in the same position may also cause systematic errors. However, there is no significant localized cloud feature appearing in Fig. 7. In addition, any localized clouds would be fixed to ground structures, such as mountains, but the rotation of the sky would change the relative positions and suppress systematic errors.

The daytime trend of the cloud detection rate shown in Fig. 9 might cause a systematic difference in the ad-

ditional noise from the residual cloud signals between the morning and afternoon observations. However, the yearly motion of the Earth gradually shifts the observing time of the CMB patch fixed on the sky. For yearlong observations, the difference will be at least partially averaged.

Performing null tests sensitive to the clouds is necessary to validate the data. A rising vs. setting split can test for localized clouds, and a summer vs. winter split can test the impact of the diurnal variation as mentioned above. A low-PWV vs. high-PWV split can test the cloud rate because of their correlation as shown in Fig. 10.

#### 4.3. Polarization Angle Calibration

While ice clouds are a nuisance in CMB observations, the polarized signal from the clouds could be a useful calibrator for the absolute polarization angle. As explained in section 2.2 and demonstrated in Figs. 4 and 11, the signal is horizontally polarized mainly because the column or plate ice crystals are aligned horizontally by gravity. The center of cirrus clouds lie at an altitude of  $\sim 10$  km above sea level, so the distance from the telescope is sufficient to achieve a far-field measurement for POLARBEAR with the 2.5 m diameter aperture observing at 150 GHz. Furthermore, the clouds are diffuse objects, making beam systematics of less concern. These two properties are better than near-field calibrators, such as the sparse wire grid in front of the telescope (Tajima et al. 2012; Kusaka et al. 2018), which need a connection from the near-field measurement to the far-field beam relying on the optics model. Although the spectrum of the cloud signal depends strongly on the observing frequency as  $\propto \omega^6$ , the polarization angle does not depend on the frequency. This addresses an uncertain polarization angle rotation feature of Tau A, a popular polarized celestial source. Compared to calibrators on the ground, cloud calibration would make it possible to operate the detectors with typical sky loading, as opposed to extra loading observing a source near the ground, and it would not have uncertainty in extrapolating the pointing model of the telescope.

In Fig. 11, the precision of the polarization angle calibration for each scan is only  $2.7^\circ$ , but the uncertainty of the mean value can be shrunk by accumulating statistics to  $0.03^\circ$ , provided the errors are independent and Gaussian distributed. That is better than the statistical uncertainty of  $0.16^\circ$  from the polarization angle calibration from nulling the apparent correlation between the CMB  $E$  and  $B$  mode patterns from two years of POLARBEAR data (The POLARBEAR Collaboration: Ade et al. 2017). In addition, the cloud polarization is

absolutely referenced to gravity, and it does not use assumptions about the symmetry properties of the CMB.

On the other hand, the cloud signal may have its own systematic errors. For example, wind and electrification may slightly tilt the ice crystals. The ground emission may not be uniform due to local features, e.g. deserts, mountains, lakes, snowfields, etc. The contribution of the Sun can become non-negligible. We have estimated the systematic error by splitting the data into subsets for various observation conditions: year split, day-night split, scan-direction split, PWV split, outside-temperature split, and wind-speed split. The median value for each subset has about  $0.4^\circ$  variation. However, that value also includes the systematic error of the instrument such as the time constant variation during the observation and imperfection of the pointing model. Further investigation is necessary to separate them, but the possibility of having many measurements with various conditions demonstrates the potential usefulness of the cloud signal as a polarization angle calibrator.

#### 4.4. Prescription for Future Experiments

For future ground-based CMB experiments aimed at more precise measurement of CMB polarization, such as the CMB-S4, steps to mitigate the contamination of the cloud signal will be necessary.

One approach is the remote sensing of the clouds from the ground on site. In section 3.1, we have demonstrated a simple cloud detection technique using the webcam for monitoring the telescope. Even with the limitation of its FOV, the significant coincidence of the cloud detections between the webcam and bolometers is observed as shown in Table 1 and Fig. 12. It can be improved by using a whole-sky camera and a co-mounted infrared camera, which would be useful during the night (e.g. Suganuma et al. 2007). As already mentioned in Pietranera et al. (2007), the most informative but challenging method is polarized lidar (e.g. Lewis et al. 2016), which shoots a laser pulse to the sky, receives the scattered light, and characterizes the atmospheric properties along the line of sight including the shape, size distribution, and orientation of the ice crystals. These tools would enable reliable cloud detection and precise data selection. This would also help with understanding the clouds and reducing the systematic error of polarization angle.

Another approach might be to perform foreground separation in the time domain. The clouds are obviously the frontmost component of the foregrounds for CMB observations. The cloud signal has frequency dependence markedly different from that of the CMB and the other astrophysical foregrounds, i.e., approximately  $\omega^6$  as opposed to  $\omega^{-3}$  for the synchrotron and  $\omega^{1.5}$  for

the dust. Therefore, it would be possible to separate the cloud signal in measurements with multi-frequency bands. Here, it is important to observe the same position at the same time among detectors with different frequency bands. The multi-chroic detector technique used in e.g. the Simons Array experiment (Stebor et al. 2016) would be useful for that purpose.

Of course, satellite missions are the best solution to avoid the clouds. Balloon-borne experiments may see clouds in the mesosphere. But their impact would be small because the particle size of the mesospheric clouds is smaller than the tropospheric clouds.

## 5. SUMMARY

The ice crystals in tropospheric clouds scatter the thermal emission from the ground and produce horizontally-polarized signals. Especially, the column and plate crystals should have the tendency to align horizontally with respect to the ground, which enhances the polarization fraction up to tens of percents.

In this study, we have presented the measurements of the clouds with the POLARBEAR experiment. The horizontal polarization and the significant coincidence between the detectors and webcam strongly support the argument that the polarized-burst signals are actually coming from the clouds. Note that the polarization modulation technique using the CRHWP is essential to perform the clear separation between the intensity and polarization signals.

Dropping data with clouds could decrease the efficiency of CMB observations. In the webcam analysis, clouds are detected in 26% of all the daytime shots. In addition, the residual cloud signal may become a critical source of low-frequency noise and systematic error that cannot be mitigated with polarization modulation techniques. In future experiments, *in situ* measurements of the clouds with extra instruments or a sophisticated analysis combining multi-frequency detectors will help mitigate the contamination.

On the other hand, the cloud signal could potentially be a good calibrator of the absolute polarization angle with  $0.03^\circ$  precision if the systematic errors of  $0.4^\circ$  associated with it can be understood.

S. Takakura was supported by Grant-in-Aid for JSPS Research Fellow JP14J01662 and JP18J02133. The

POLARBEAR project is funded by the National Science Foundation under Grants No. AST-0618398 and No. AST-1212230. The James Ax Observatory operates in the Parque Astronómico Atacama in Northern Chile under the auspices of the Comisión Nacional de Investigación Científica y Tecnológica de Chile (CONICYT). The APEX PWV was obtained from the APEX Weather Query Form of the ESO Science Archive Facility. This research used resources of the Central Computing System, owned and operated by the Computing Research Center at KEK, the HPCI system (Project ID:hp150132), and the National Energy Research Scientific Computing Center, a DOE Office of Science User Facility supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. Kavli IPMU was supported by World Premier International Research Center Initiative (WPI), MEXT, Japan. This work was supported by MEXT KAKENHI grant Nos. 21111002, JP15H05891, and JP18H05539, JSPS KAKENHI grant Nos. JP26220709, JP26800125, and JP16K21744, and the JSPS Core-to-Core Program. MA acknowledges support from CONICYT UC Berkeley-Chile Seed Grant (CLAS fund) Number 77047, Fondecyt project 1130777 and 1171811, DFI postgraduate scholarship program and DFI Postgraduate Competitive Fund for Support in the Attendance to Scientific Events. CB, NK, and DP acknowledge support from the RADIOFOREGROUNDS project (radioforegrounds.eu), funded by the European Commission's H2020 Research Infrastructures under the Grant Agreement 687312, the INDARK INFN Initiative and the COSMOS network of the Italian Space Agency (cosmosnet.it). FB and CR acknowledges support from an Australian Research Council Future Fellowship (FT150100074). D. Boettger gratefully acknowledges support from grant ALMA CONICYT 31140004. GF acknowledges the support of the CNES postdoctoral program. Work at LBNL was supported by the Laboratory Directed Research and Development Program of Lawrence Berkeley National Laboratory under U.S. Department of Energy Contract No. DE-AC02-05CH11231. AK acknowledges the support by JSPS Leading Initiative for Excellent Young Researchers (LEADER). FM acknowledges JSPS International Research Fellowship (Grant number JP17F17025).

## REFERENCES

- Abazajian, K. N., Adshead, P., Ahmed, Z., et al. 2016, ArXiv e-prints, arXiv:1610.02743
- Arnold, K., Ade, P. A. R., Anthony, A. E., et al. 2012, in Proc. SPIE, Vol. 8452, Millimeter, Submillimeter, and Far-Infrared Detectors and Instrumentation for Astronomy VI, 84521D

- BICEP2 Collaboration, Ade, P. A. R., Aikin, R. W., et al. 2014, *ApJ*, 792, 62
- Brown, M. L., Challinor, A., North, C. E., et al. 2009, *MNRAS*, 397, 634
- Chepfer, H. 1999, *JQSRT*, 63, 521
- Cowley, L., & Schroeder, M. 2009, HaloSim3 ray tracing simulator, <https://www.atoptics.co.uk/halosim.htm>, ,
- Czekala, H. 1998, *Geophys. Res. Lett.*, 25, 1669
- Defer, E., Galligani, V. S., Prigent, C., & Jimenez, C. 2014, *Journal of Geophysical Research (Atmospheres)*, 119, 12
- Erasmus, D. A., & Van Staden, C. A. 2001, Cerro Tololo Inter-American Observatory
- Errard, J., Ade, P. A. R., Akiba, Y., et al. 2015, *ApJ*, 809, 63
- Gong, J., & Wu, D. L. 2017, *Atmospheric Chemistry & Physics*, 17, 2741
- Güsten, R., Nyman, L. Å., Schilke, P., et al. 2006, *A&A*, 454, L13
- Hanany, S., & Rosenkranz, P. 2003, *NewAR*, 47, 1159
- Hill, C. A., Beckman, S., Chinone, Y., et al. 2016, in *Proc. SPIE, Vol. 9914, Millimeter, Submillimeter, and Far-Infrared Detectors and Instrumentation for Astronomy VIII*, 99142U
- Hu, W., & White, M. 1997, *PhRvD*, 56, 596
- Keating, B., Timbie, P., Polnarev, A., & Steinberger, J. 1998, *ApJ*, 495, 580
- Kneifel, S., Löhnert, U., Battaglia, A., Crewell, S., & Siebler, D. 2010, *Journal of Geophysical Research (Atmospheres)*, 115, D16214
- Kuo, C.-L. 2017, *ApJ*, 848, 64
- Kusaka, A., Essinger-Hileman, T., Appel, J. W., et al. 2014, *Review of Scientific Instruments*, 85, 024501
- Kusaka, A., Appel, J., Essinger-Hileman, T., et al. 2018, *ArXiv e-prints*, arXiv:1801.01218
- Landau, L. D., & Lifshitz, E. M. 1960, *Course of theoretical physics, Vol. 8, The electrodynamics of continuous media* (Pergamon Press)
- Lay, O. P., & Halverson, N. W. 2000, *ApJ*, 543, 787
- Lewis, J. R., Campbell, J. R., Welton, E. J., Stewart, S. A., & Haftings, P. C. 2016, *Journal of Atmospheric and Oceanic Technology*, 33, 2113
- Liou, K.-N., & Yang, P. 2016, *Light Scattering by Ice Crystals: Fundamentals and Applications* (Cambridge University Press), doi:10.1017/CBO9781139030052
- Magono, C., & Lee, C. W. 1966, *J. Fac. Sci. Hokkaido Univ.*, 2, 321. <https://ci.nii.ac.jp/naid/120000956995/>
- Ono, A. 1969, *Journal of Atmospheric Sciences*, 26, 138
- Pettersen, C., Bennartz, R., Kulie, M. S., et al. 2016, *Atmospheric Chemistry & Physics*, 16, 4743
- Pietranera, L., Buehler, S. A., Calisse, P. G., et al. 2007, *MNRAS*, 376, 645
- Rolland, P., Liou, K. N., King, M. D., Tsay, S. C., & McFarquhar, G. M. 2000, *J. Geophys. Res.*, 105, 11
- Rose, T., Crewell, S., Löhnert, U., & Simmer, C. 2005, *Atmospheric Research*, 75, 183
- Rosenkranz, P. W., & Staelin, D. H. 1988, *Radio Science*, 23, 721
- Seljak, U., & Zaldarriaga, M. 1997, *Physical Review Letters*, 78, 2054
- Smith, A. R. 1978, *SIGGRAPH Comput. Graph.*, 12, 12
- Spinelli, S., Fabbian, G., Tartari, A., Zannoni, M., & Gervasi, M. 2011, *MNRAS*, 414, 3272
- Stebor, N., Ade, P., Akiba, Y., et al. 2016, in *Proc. SPIE, Vol. 9914, Millimeter, Submillimeter, and Far-Infrared Detectors and Instrumentation for Astronomy VIII*, 99141H
- Suen, J. Y., Fang, M. T., & Lubin, P. M. 2014, *IEEE Transactions on Terahertz Science and Technology*, 4, 86
- Suganuma, M., Kobayashi, Y., Okada, N., et al. 2007, *PASP*, 119, 567
- Tajima, O., Nguyen, H., Bischoff, C., et al. 2012, *Journal of Low Temperature Physics*, 167, 936
- Takakura, S., Aguilar, M., Akiba, Y., et al. 2017, *JCAP*, 2017, 008
- The POLARBEAR Collaboration: Ade, P. A. R., Aguilar, M., Akiba, Y., et al. 2017, *ApJ*, 848, 121
- Troitsky, A. V., & Osharin, A. M. 2000, *Radiophysics and Quantum Electronics*, 43, 356
- Troitsky, A. V., Osharin, A. M., Korolev, A. V., & Strapp, J. W. 2003, *Journal of Atmospheric Sciences*, 60, 1608
- Troitsky, A. V., Vostokov, A. V., & Osharin, A. M. 2005, *Radiophysics and Quantum Electronics*, 48, 281
- Warren, S. G., & Brandt, R. E. 2008, *Journal of Geophysical Research (Atmospheres)*, 113, D14220
- Westwater, E. R., Crewell, S., & Matzler, C. 2004, *URSI Radio Science Bulletin*, 2004, 59
- Xie, X., Crewell, S., Löhnert, U., Simmer, C., & Miao, J. 2015, *Journal of Geophysical Research (Atmospheres)*, 120, 6145
- Xie, X., Löhnert, U., Kneifel, S., & Crewell, S. 2012, *Journal of Geophysical Research (Atmospheres)*, 117, D02206