

1 Title: Soil nutrients and microbial biomass in three contrasting Mediterranean forests

2

3 Authors: Cristina Aponte^{1,2*}, Luis Matías^{3,4}, Victoria González-Rodríguez⁵, Jorge Castro³

4 Luis V. García¹, Rafael Villar⁵, Teodoro Marañón¹

5

6 Affiliation:

7 ¹ Instituto de Recursos Naturales y Agrobiología de Sevilla (IRNAS), CSIC. Reina Mercedes, 10. Sevilla, E-
8 41012, Spain.

9 ² Department of Forest and Ecosystem Science; The University of Melbourne
10 500 Yarra Boulevard, Richmond, Victoria 3121, Australia

11 ³ Grupo de Ecología Terrestre, Dpto. de Ecología, Facultad de Ciencias, Universidad de Granada, E-18071
12 Granada, Spain

13 ⁴ Biological and Environmental Sciences, School of Natural Sciences, University of Stirling, FK9 4LA, Stirling,
14 UK

15 ⁵ Área de Ecología, Campus de Rabanales, Universidad de Córdoba, E-14071 Córdoba, Spain

16 * Corresponding author: caponte@unimelb.edu.au

17 Department of Forest and Ecosystem Science; The University of Melbourne
18 500 Yarra Boulevard, Richmond, Victoria 3121, Australia

19 p. +61 3 9035 6862

20

21 ABSTRACT

22 Aims: The extent to which the spatial and temporal patterns of soil microbial and available nutrient pools hold
23 across different Mediterranean forest types is unclear impeding the generalization needed to consolidate our
24 understanding on Mediterranean ecosystems functioning.

25 Methods: We explored the response of soil microbial, total, organic and inorganic extractable nutrient pools (C,
26 N and P) to common sources of variability, namely habitat (tree cover), soil depth and season (summer drought),
27 in three contrasting Mediterranean forest types: a *Quercus ilex* open woodland, a mixed *Q. suber* and *Q.*
28 *canariensis* woodland and a *Pinus sylvestris* forest.

29 Results: Soil microbial and available nutrient pools were larger beneath tree cover than in open areas in both oak
30 woodlands whereas the opposite trend was found in the pine forest. The greatest differences in soil properties
31 between habitat types were found in the open woodland. Season (drought effect) was the main driver of
32 variability in the pine forest and was related to a loss of microbial nutrients (up to 75% loss of N_{mic} and P_{mic}) and
33 an increase in microbial ratios (C_{mic}/N_{mic} , C_{mic}/P_{mic}) from Spring to Summer in all sites. Nutrient pools
34 consistently decreased with soil depth, with microbial C, N and P in the top soil being up to 208%, 215% and
35 274% larger than in the deeper soil respectively.

36 Conclusions: Similar patterns of variation emerged in relation to season and soil depth across the three forest
37 types whereas the direction and magnitude of the habitat (tree cover) effect was site-dependent, possibly related
38 to the differences in tree species composition and forest structure, and thus in the quality and distribution of the
39 litter input.

40

41 Keywords: soil fertility, plant-soil interactions, soil carbon, nitrogen, phosphorus

42

43

44 INTRODUCTION

45 Soil microorganisms, in their role in organic matter decomposition, have the capacity to both mineralize and
46 immobilize nutrients (Singh et al. 1989) thereby influencing soil nutrient availability and plant growth (Lambers
47 et al. 1998). Spatial and temporal changes in soil microbial biomass may determine the patterns of availability
48 of limiting nutrients such nitrogen (N) and phosphorus (P), thus having profound influence on plant
49 communities and ecosystem functioning (Ettema and Wardle 2002; Gallardo and Schlesinger 1994; Sardans et
50 al. 2005; van der Putten et al. 2009).

51 Spatial and temporal variations of soil microbial biomass and activity are related to different biotic and
52 abiotic factors that modulate the temperature, moisture conditions and substrate quality and availability. For
53 instance, vegetation composition and structure control the spatial distribution, quality and quantity of nutrients
54 inputs *via* litter and root exudates (Aponte et al. 2011; Huang et al. 2013; Prescott and Grayston 2013; Ushio et
55 al. 2010). Soil nutrients and microbial activity usually decrease as soil depth increase due to a decline in the
56 quality and quantity of organic matter (Gaudinski et al. 2000; Xiang et al. 2008). Seasonal changes in
57 temperature, water and substrate availability also have a large impact on soil microbial activity and nutrient
58 cycling (Corre et al. 2002; Quilchano and Marañón 2002; Schmidt et al. 1999). In highly seasonal ecosystems,
59 such as Mediterranean forest, the effects imposed by seasonal variations, in particular associated to the summer
60 drought, are especially important for ecosystem functioning (Aponte et al. 2010b; Marañón-Jiménez et al. 2011;
61 Matías et al. 2011).

62 Many studies have described soil nutrient heterogeneity in Mediterranean forests; however, most of them have
63 been conducted at local spatial scales, focused on a single forest type (Barcenas-Moreno et al. 2011; Carreira et
64 al. 1994; Gallardo 2003; García et al. 2006; Maltez-Mouro et al. 2005; Monokrousos et al. 2004). The detection
65 of general patterns across different forest types is necessary to fully understand microbial biomass and nutrients
66 dynamics and their consequences for plant community. At the same time, the emergence of site-dependent
67 effects will be of interest from a modelling and management perspective, to properly determine nutrient pools at
68 wide geographical scales including a mosaic of forest types. Patterns of microbial biomass and nutrient
69 heterogeneity across different forest types have been largely investigated in temperate, boreal and tropical forest
70 (e.g. Hackl et al. 2005; Lindo and Visser 2003; Liu et al. 2012; Zhong and Makeschin 2006), while remain far
71 less studied in Mediterranean forest (but see García et al. 2002; Goberna et al. 2006). This coordinated study
72 addressed this knowledge gap and aimed to evaluate whether the effects of main sources of variability, namely
73 habitat (i.e. tree cover), soil depth and season, in the soil nutrients and microbial C, N and P pools could be

74 generalised across three contrasting Mediterranean forests: a *Quercus ilex* open woodland, a mixed *Q. suber* and
75 *Q. canariensis* woodland and a *Pinus sylvestris* forest. While this study builds upon previous knowledge on soil
76 nutrient heterogeneity at local scales (Aponte et al. 2010b; Matías et al. 2011), it focuses on the comparison
77 among forests with different structure and species composition, thus taking a step forward towards
78 understanding general patterns of soil microbial responses to biotic and abiotic environmental drivers.
79 Explicitly, we aimed to answer the following questions: 1) Is there a common pattern across the three forests in
80 relation to the tree effect, soil depth and seasonal drought? ; 2) Are the interactions between the effects of tree
81 cover, soil depth and season (summer drought) similar across forest types?; 3) What is the quantitative
82 importance of the studied factors (tree effect, soil depth and seasonal drought) on the soil and microbial
83 variables in each forest type? 4) Do the relationships between microbial and soil chemical properties hold when
84 examined across forest types?

85

86 METHODS

87 *Study areas*

88 The study was conducted in three different Mediterranean forest types: a mixed woodland of *Quercus suber* L.
89 (evergreen) and *Q. canariensis* Willd. (deciduous) in Los Alcornocales Natural Park in the extreme south, near
90 the Strait of Gibraltar, an open woodland dominated by the sclerophyllous *Quercus ilex* subsp. *ballota* L. and
91 eventually mixed with other *Quercus* species (*Q. suber*, *Q. pyrenaica* Willd., *Q. faginea* Lam.) in Sierra de
92 Cardeña and Montoro Natural Park (Cardeña), in the south mainland, and a forest mainly comprised of *Pinus*
93 *sylvestris* L. interspersed with *Q. ilex* subsp. *ballota* in Sierra Nevada National Park in the southeast of Spain
94 (Fig. 1). In all three forest types, the main tree species are intermingled with open areas covered by sparse
95 herbaceous vegetation. The study sites vary in altitude, climate and soil conditions (Table 1). The general
96 climate of the three sites is Mediterranean-type, characterized by hot and dry summers, and cold and wet winters
97 with most rainfall occurring from October to May. The sites in Cardeña and Sierra Nevada experience more
98 extreme temperatures due to their continental and altitudinal locations (respectively), while temperatures in
99 Alcornocales site are milder due to the lower elevation and proximity to the Mediterranean Sea and Atlantic
100 Ocean. Mean annual rainfall follows a rising gradient from Cardeña to Alcornocales (Table 1). The sites in
101 Alcornocales and Cardeña stand on a bedrock of sandstone and granite, both producing acidic sandy soils. On
102 the contrary the site in Sierra Nevada stands on limestone, which gives rise to basic loamy soils. Cambisols
103 dominated in Alcornocales and regosols in Cardeña (nomenclature follows WRB 2006), indicating a greater

104 soil development i.e. soil depth, structure, water holding capacity and chemical fertility in the former than the
105 later.

106

107 *Experimental design*

108 At each forest site 10-20 replicates (depending on the site, Table 1) of two main habitat types were identified
109 within a stand: beneath the canopy of the dominant tree species (*Q. suber* and *Q. canariensis* in Alcornocales,
110 *Q. ilex* in Cardeña and *P. sylvestris* in Sierra Nevada), and in open areas with bare soil or sparse herbaceous
111 cover and no tree cover. These habitat types will be referred as ‘Tree’ and ‘Open’ respectively hereafter. At
112 each replicate point, four soil cores (0-16 cm) were extracted using an auger after removing the litter layer,
113 divided between ‘Top soil’ (0-8 cm) and ‘Deeper soil’ (8-16 cm) and homogenized within the same depth to
114 obtain a composite soil sample per habitat type replicate and depth. Soil samples were taken in Spring (May-
115 June) and Summer (August-September) 2007, coinciding with the moment of maximum soil biological activity
116 and maximum water stress in soil, respectively. In total 400 soil samples were taken corresponding to 10-20
117 replicates (Table 1) of 2 habitat types x 2 soil depths x 2 seasons x 3 forest sites. Litter, i.e. dead plant material
118 relatively undecomposed standing on the ground, was collected once in all sampling points using a 10 x 10 cm
119 quadrat (in Sierra Nevada) or a 30 x 30 cm quadrat (in Alcornocales and Cardeña). Litter samples were oven-
120 dried at 60°C for 72 h and weighted.

121

122 *Laboratory analyses*

123 Soil samples were brought to the laboratory in an ice-box, fresh-sieved at 2 mm removing stones, roots and
124 other recognizable plant parts and stored at 4°C for analyses. Water content was determined on a subsample as
125 the difference in weight between fresh and oven dried (105°C) soil.

126 Microbial C, N and P were estimated in fresh soils using a chloroform fumigation-extraction procedure
127 (Brookes et al. 1985; Brookes et al. 1982; Vance et al. 1987). Dissolved organic C (DOC) and N (DON) and
128 inorganic P (P_{inorg}) were determined in non-fumigated and chloroform fumigated soil subsamples (24h).
129 Dissolved C and N were extracted with 0.5M K_2SO_4 , and their concentration was determined using a Shimadzu
130 TOC-V CSH analyzer. Inorganic P was extracted with either 0.025N HCl+0.03N NH_4F (Bray Kurtz 1 method
131 (Bray and Kurtz 1945) for the acidic soils of Alcornocales and Cardeña) or 0.5M $NaHCO_3$ (Olsen method
132 (Olsen et al. 1954) for the basic soils of Sierra Nevada) and its concentration was determined by colorimetry

133 using the ascorbic acid-molybdenum blue method (Sparks 1996). Microbial C (C_{mic}), N (N_{mic}) and P (P_{mic}) were
134 estimated as the difference in DOC, DON and P between fumigated and non-fumigated samples.

135 Inorganic nitrogen (N_{inorg}) was extracted from non-fumigated soils using 2M KCl and the extracts were
136 analyzed for NH_4^+ and NO_3^- by the Kjeldhal method (Bremner and Keeney 1965). Soil total C (C_{tot}) and N (N_{tot})
137 were determined on oven dried soils by combustion at 850°C (Leco TruSpec autoanalyzer) and total inorganic C
138 (C_{inorg}) was measured by acidification with $HClO_4$ in a TIC analyzer (UIC CM-5014). The difference between
139 C_{tot} and C_{inorg} gave the total organic C (C_{org}).

140

141 *Data analysis*

142 Differences among habitat, soil depth and season were analyzed using repeated measurement ANOVAs with
143 season as a within-group effect and habitat and depth as between-group effects. Forest site was also included in
144 the analysis to test for potential interactions with the studied factors. Variables were transformed (log, arcsin)
145 when necessary to meet normality assumptions. To control the type I error inflation resulted from repeated tests,
146 the false discovery rate (FDR), i.e. the expected proportion of tests erroneously declared as significant, was
147 controlled at 5% using a step-up procedure (Benjamini and Hochberg 1995; García 2003). The percentage of
148 the total variance explained by the studied factors (habitat, soil depth and season) was calculated for each
149 variable and site using a repeated measurement ANOVA with no interactions. Patterns in pairwise Pearson's
150 correlations between microbial and soil nutrient fractions were explored using correlation network analysis (R
151 package igraph, Csardi and Nepusz 2006). Multivariate relationships between variables were analysed using
152 Principal Component Analysis (PCA). The 'Broken stick' method (King and Jackson 1999) was used to select
153 significant components. Habitat, soil depth and season were included in the PCA as supplementary variables,
154 i.e. these factors did not participate in the analysis, but were projected on the multivariate space generated by the
155 PCA for the purpose of interpretation.

156

157 RESULTS

158 Overall, the study forests differed in all the analyzed soil and microbial properties (Table 2 and 3). Cardeña was
159 the least fertile site while Alcornocales had the largest fraction of microbial nutrients (from 3 to 6-fold the
160 values of the other sites) and the largest pool of total and dissolved C and N and organic C (Fig. 2 and 3). Sierra
161 Nevada showed the highest inorganic N and P values (~2-fold to 8-fold the values of the other sites), the highest
162 C_{mic}/N_{mic} ratio (2-fold) and the largest litter pool (~6-fold) (Fig. 3). The ratios of nutrients retained in the

163 microbial biomass vs. the pool of available nutrients (N_{mic}/N_{inorg} and P_{mic}/P_{inorg}) as well as the fraction of soil
164 organic carbon and total nitrogen in the microbial biomass (C_{mic}/C_{org} and N_{mic}/N_{tot}) were the highest in
165 Alcornocales and Cardeña and the lowest in Sierra Nevada (Online Resource 1).

166

167 *Effect of habitat*

168 Soil parameters differed significantly between the two habitat types in all forest sites (Table 3, Fig. 2 and 3).
169 However, the magnitude and direction of those differences varied across sites, as the interaction Site \times Habitat
170 was significant for most of the variables (Table 3). In both oak woodlands, Alcornocales and Cardeña, the
171 nutrient pools (microbial, dissolved organic and inorganic) tended to be larger beneath tree canopy than in open
172 areas with the exception of nitrate that showed the opposite trend (Fig. 2 and 3). Greater concentrations of
173 ammonium (156% in both sites), phosphate (120% in Alcornocales and 182% in Cardeña), and microbial
174 nutrients (123 and 166% for C_{mic} ; 126 and 227% for N_{mic} ; 215 and 175% for P_{mic}) were found beneath tree cover
175 than in open areas. Mean soil organic carbon was also greater beneath tree cover than in open areas in Cardeña
176 (1.7% vs. 0.97%; $P < 0.0001$) and Alcornocales (4.1% vs. 3.6%, not significant difference). A different pattern
177 was observed in the pine forest (Sierra Nevada) where most of the soil nutrient pools were similar between the
178 two habitats or even decreased beneath trees as it occurred with N_{mic} and inorganic N and P (Fig. 2 and 3).
179 Organic and inorganic C also decreased significantly from open areas (3.1%, and 2.85% respectively) to
180 beneath pine tree cover (2.9% and 0.93% respectively). Nevertheless, the amount of litter was larger beneath
181 tree canopy than in open areas in all sites, being the values larger in Sierra Nevada than in the other two forests
182 (Table 1). There were no habitat differences in the fractions of microbial values relative to soil pools
183 (N_{mic}/N_{inorg} , P_{mic}/P_{inorg} , C_{mic}/C_{org} and N_{mic}/N_{tot} ; data not shown).

184

185 *Effect of soil depth*

186 In general all variables measured showed a consistent pattern with soil depth in the three forest sites, with values
187 decreasing from Top soil to Deeper soil (Fig. 2 and 3). However, there was a significant Site \times Depth interaction
188 (Table 3) due to the lack of statistical significance of soil depth for many variables in Cardeña (C_{mic} , P_{mic} , DON,
189 NH_4 and P_{inorg}) (Fig. 2 and 3).

190 Microbial C, N and P in Top soil were higher than in Deeper soil with the largest variations found in
191 Alcornocales (208, 215 and 274% respectively) and the smallest changes found in Cardeña (128, 155 and 119%
192) (Fig. 2). On average across sites, the pool of inorganic N and P, DOC, DON and C_{org} was 133% (site mean

193 range 110 – 146%), 155% (129 – 177%), 142% (117-172%), 140% (112-159%) and 118% (114-120%) higher
194 in Top soil than in Deeper soil respectively. As with microbial pools, variation was the least in Cardeña (Fig. 2
195 and 3). Microbial ratios C_{mic}/C_{org} and N_{mic}/N_{tot} showed the largest decrease with soil depth in Alcornocales but
196 remained constant in Cardeña. The ratio of microbial biomass nutrients (C_{mic}/N_{mic} and C_{mic}/P_{mic}) showed no
197 significant variation from Top soil to Deeper soil in any site. The only exception was found for soils in open
198 areas in Cardeña where C_{mic}/P_{mic} increased with soil depth from 55 to 143, as evidenced by a significant Site \times
199 Habitat \times Depth interaction (Table 3).

200

201 *Effect of season*

202 Soil microbial fractions and nutrient pools varied significantly with the season. However, the seasonal patterns
203 of variation were site-dependent as indicated by a significant Site \times Season (Table 3). Seasonal variations were
204 stronger in Sierra Nevada and Alcornocales whereas Cardeña showed the lowest variability between seasons
205 (Fig. 2 and 3). In general, microbial pools were larger in Spring than in Summer, particularly for N_{mic} and P_{mic}
206 which values were on average 237% and 258% higher in Spring (Fig 2). The fraction of microbial C and N
207 relative to soil total pools (C_{mic}/C_{org} and N_{mic}/N_{tot}) decreased from Spring to Summer in Sierra Nevada but not in
208 Cardeña. Microbial ratios (C_{mic}/N_{mic} , C_{mic}/P_{mic}) increased from Spring to Summer in all sites revealing a larger
209 loss of N_{mic} and P_{mic} as compared to C_{mic} .

210 The seasonal variability of N_{mic} , P_{mic} , C_{org} , DOC and DON was larger in soils beneath tree canopy
211 whereas the variation was subdued in the open habitats (Season \times Habitat interaction, Table 3). We also found a
212 strong and significant Site \times Season interaction for N_{inorg} and P_{inorg} (Table 3), which was due to opposite seasonal
213 changes across forest types. For example, the pool of available inorganic nutrients (ammonium and phosphate)
214 as well as DON increased from Spring to Summer in Cardeña and Sierra Nevada, whereas the values decreased
215 in Alcornocales (Fig. 3). Despite the discrepancies in the seasonal dynamics of P_{inorg} , the proportion of P_{mic}
216 relative to P_{inorg} was higher in Spring than in Summer in all sites (data not shown). The observed seasonal
217 patterns were similar at the two soil depths.

218

219 *Variance partitioning among habitat, soil depth and season*

220 As shown in the partition of variance (Fig. 4) and the principal component analysis (Fig. 5) the main drivers
221 of variability differed between sites. Soil depth and season accounted for the largest part of the variability
222 observed in the microbial and soil nutrient pools in Alcornocales and Sierra Nevada. For instance, in

223 Alcornocales soil depth explained 50, 38 and 30% of the variation of microbial C, N and P respectively and
224 season explained 55 and 39% of the variation of N_{inorg} and P_{inorg} . In Sierra Nevada season was the main driver of
225 microbial variability accounting for 26, 58 and 66% of the variation of microbial C, N and P. In contrast, the
226 variability of soil biotic and abiotic properties in Cardeña was mainly driven by habitat type, which explained
227 10, 16 and 4% of microbial C, N and P variation respectively and 9 and 12% of N_{inorg} and P_{inorg} variation.

228

229 *Relations between microbial pools and soil properties*

230 Soil microbial C, N and P were significantly correlated among them in all sites (Fig. 5 and Online Resource 1).
231 Microbial C and N were consistently and strongly coupled ($r > 0.76$ in all sites), whereas P_{mic} was more weakly
232 but still significantly related with C_{mic} (from $r = 0.28$ in Cardeña to $r = 0.69$ in Sierra Nevada) and N_{mic} (from
233 $r = 0.30$ in Cardeña to $r = 0.82$ in Sierra Nevada). Microbial C, N and P were positively related to most of the
234 measured soil properties in each site (Fig 5). The strongest correlations were found with C_{org} , N_{tot} and soil
235 moisture reflecting microbial biomass dependence on substrate and water availability. Microbial C also showed
236 a significant correlation with P_{inorg} in all sites ($r \sim 0.36$). The relationship between litter and C_{org} (r_C) and N_{tot} (r_N)
237 varied across forest sites, being positive in Cardeña ($r_C = 0.43$, $r_N = 0.35$, $P < 0.0001$, seasons and depths
238 pooled), positive in Top soil in Alcornocales ($r_C = 0.28$, $r_N = 0.27$, $P < 0.05$; not significant in Deeper soil) and
239 negative in Sierra Nevada ($r_C = -0.16$, $P < 0.06$; $r_N = -0.36$, $P < 0.0001$). The correlation network was the
240 strongest in Alcornocales, i.e. there was a tight coupling between most variables, and the weakest in Cardeña
241 (Online Resource 1).

242 The multivariate analyses (PCAs) showed similar patterns of covariation among the nutrient pools for
243 all sites (Fig 5). Two main significant gradients (axes) emerged for each PCA from the analysis based on the
244 ‘Broken –stick’ method (King and Jackson 1999). For all sites the first axis was strongly correlated to microbial
245 C, N and P, total N and organic C. In Alcornocales and Sierra Nevada the first axis was also positively related to
246 soil moisture and negatively related to C_{mic}/N_{mic} . The separation of samples along the main axis and the analysis
247 of the supplementary variables indicated that both season and soil depth imposed a similar degree of variability
248 in Alcornocales whereas season was the main driver of variation in Sierra Nevada, which is agreement with our
249 variance partitioning analysis. In Cardeña the first axis was related to litter amount, but not to soil moisture, and
250 separated the samples by habitat type. The second axis in all PCAs was related to the availability of inorganic
251 nutrients (N and/or P), which covariation with other variables was inconsistent across forest types. Higher
252 microbial ratios (C_{mic}/N_{mic}) were consistently associated to lower soil moisture and Summer samples in all sites.

253 The relationship between litter abundance and microbial and total nutrient pools was positive in Alcornocales
254 and Cardeña, but negative in Sierra Nevada.

255 Variables covaried similarly when all three sites were combined in a single PCA (Online Resource 2):
256 the first axis accounted for 34% of the variability and was strongly correlated to most nutrient pools (microbial,
257 dissolved organic and total) and soil moisture. The second axis accounted for 27% of the variability and was
258 mostly related to inorganic N and P. The two axes clearly separated between forest sites, with Cardeña at the
259 poorest end of both axes and Alcornocales and Sierra Nevada at the richest end of the first and second axes,
260 respectively.

261

262 DISCUSSION

263 Overall, the three sources of variability considered (habitat, soil depth and season) had significant effects on the
264 soil microbial pools and nutrient concentrations in the studied forests. However the direction and magnitude of
265 these effects varied across forest types and with the soil parameter examined.

266 The expected positive effect of tree canopy on soil and microbial nutrients was confirmed for the two oak
267 woodlands (Cardeña and Alcornocales) but not for the pine forest (Sierra Nevada), where the soil and microbial
268 nutrients pools were smaller beneath tree canopy than in open areas. The inconsistency of the habitat effect
269 could be attributed to the forests' distinct species composition. Trees generate species-specific effects on soil
270 conditions through multiple pathways, such as changing microclimatic conditions or *via* leaf and root litter input
271 or root exudates (Alameda et al. 2012; Aponte et al. 2013; Aponte et al. 2011; Malchair and Carnol 2009). Tree
272 species changes in soil abiotic properties might in turn affect soil biota (Aponte et al. 2013; Aponte et al. 2010a;
273 Prescott and Grayston 2013). In particular, tree-mediated changes in soil acidity and in the amount and quality
274 of substrate are known to affect microbial communities size and composition (Lucas-Borja et al. 2012; Sagova-
275 Mareckova et al. 2011; Thoms et al. 2010). In Sierra Nevada soil acidity was higher beneath pine cover than in
276 open areas, as evidenced by their distinct pH (7.7 vs. 8.1) and C_{inorg} (0.93% vs. 2.85% respectively), while clay
277 content was lower (18.5 vs. 21.6%). Litter biomass was 15 times greater (8594 vs. 559 g m⁻²) and the amount
278 and quality of the substrate (C_{org} , DOC, N_{tot} , C_{org}/N_{tot}) were significantly lower beneath tree cover (*Pinus*) than
279 in open areas, in agreement with the negative correlation observed between litter and soil C_{org} and N_{tot} .
280 Meanwhile, the opposite was found in the two oak forest sites, i.e substrate quality was higher beneath tree
281 cover (*Quercus*) than in open areas, and it was positively related to litter biomass, thus sustaining the
282 counteracting patterns observed for microbial nutrients. In accordance with our results, previous studies on the

283 effects of tree species on soils have related the lower soil nutrient and microbial values found beneath pine
284 cover, compared to other broadleaves tree species (including *Quercus*), with the poorer quality of the pine litter,
285 and thus to its lower decomposition rate and nutrient release, and its capacity to acidify soils (Augusto et al.
286 2002; Rutigliano et al. 2004; Smolander and Kitunen 2002; Ste-Marie et al. 2007). Nonetheless, the observed
287 differences in soil and microbial nutrients between habitat types should not be solely attributed to vegetation
288 cover. Other soil physicochemical properties, such as soil depth, structure and texture, which may be the
289 underlying reason for the distinct cover type, can also control microbial development (Hassink 1994).

290 Interestingly our results also revealed a difference in the magnitude of the positive tree-effect on soil
291 nutrients between the two oak woodlands, Cardeña and Alcornocales. These two sites significantly differed in
292 their soil type and nutrient content: Cardeña sited over Regosols, i.e. weakly developed soils with a low organic
293 matter content and water holding capacity (WRB 2006) (Table 1). In contrast, soils in Alcornocales were
294 cambisols (also known as Brown forest soils, WRB 2006), they were well structured and presented a thick
295 humic horizon (15-20cm beneath tree canopy; Garcia et al, unpublished data), and a relatively high soil organic
296 matter content (11% in 0-25cm upper soil, Polo 2006). Mean site C_{org} was greater in Alcornocales (3.9%) than
297 in in Cardeña (1.3 %), clay content was 7 times higher in the former (36%) than in the later (5%), and CEC
298 (cation exchange capacity) was two-fold in Alcornocales than in Cardeña (Table 1), all of which supported the
299 distinct soil fertility and microbial nutrient levels observed in both forest sites (Table 2). These two sites also
300 differed in their stand structure, with a lower tree density in Cardeña than in Alcornocales (131 vs. 219 stems ha⁻¹).
301 It is possible that the interaction between their distinct soil types and stand structure could be determining
302 why habitat type was the main driver of variability of soil microbial properties in Cardeña but it was of lesser
303 importance in Alcornocales. The intensity of tree effects on soil properties is modulated by the spatial
304 distribution of tree canopies (Bennett et al. 2009; Ushio et al. 2010). It is well-known that oak trees in
305 Mediterranean savannah-like systems (dehesas) generate islands of fertility beneath their canopies where the
306 leaf litter and root exudates accumulate and build up the soil organic matter that sustain microbial biomass and
307 nutrient cycling (Alameda et al. 2012; Gallardo 2003). In sparse forests, such as Cardeña, trees are scattered in a
308 matrix of open areas and their footprints on soil fertility are expected to be more intense beneath the canopy. In
309 contrast, a more diffuse footprint occurs in dense forests where open areas are intermingled in a matrix of trees.
310 This is consistent with C_{org} and C_{mic} being greater beneath tree cover than in open areas by a factor of 1.7 and 1.7
311 in Cardeña and a factor of 1.1 and 1.2 in Alcornocales respectively. In addition, the small concentrations of

312 substrate (C_{org} , DOC, N_{tot} , DON) in Cardeña could be a limiting factor for microbial biomass and a tree-
313 mediated increase in its availability would render a larger boost of microbial growth than in more fertile sites.

314 Microbial C, N and P showed a common and seasonal pattern, with values decreasing from Spring to
315 Summer in response to summer drought. This response was the weakest in Cardeña, where changes were not
316 significant. Seasonal variation was larger for N_{mic} and P_{mic} than for C_{mic} , rendering a shift in the microbial ratios,
317 as evidenced by the multivariate analyses. The change in C_{mic}/N_{mic} was the largest in Sierra Nevada (from 9 in
318 Spring to 34 in Summer), where a decrease in C_{mic}/C_{org} , a proxy for microbial C assimilation efficiency
319 (Sparling 1992), was also observed. Soil microorganisms in Mediterranean ecosystems have adapted to
320 withstand the seasonal variation in water availability and temperature that define the Mediterranean-type climate
321 (Goberna et al. 2007). Seasonality, in particular the summer drought, may influence microbial biomass directly
322 by inducing microbial metabolic responses to changes in soil moisture and temperature (Chen et al. 2003 ;
323 Jensen et al. 2003), or indirectly by influencing plant productivity, organic matter release and C diffusion in soil,
324 and hence substrate availability (Rey et al. 2002; Xiang et al. 2008). The high microbial values found in Spring
325 may reflect favourable environmental conditions and more labile substrates derived from roots or from materials
326 incorporated into the soil whereas the decrease in Summer might indicate a loss in the total number of
327 organisms. This is consistent with previous work conducted in the same forest stand in Alcornocales, which
328 showed higher soil enzyme activity during the rainy season than in summer (Quilchano and Marañón 2002). On
329 the other hand, summer increases in microbial ratios can be related to an increasing proportion of fungi vs.
330 bacteria (Jensen et al. 2003), since fungi have a higher carbon to nitrogen ratio (C_{mic}/N_{mic}) (related to their
331 lower efficiency, Cleveland and Liptzin 2007); and are more drought-tolerant than bacteria (Wilkinson et al.
332 2002). In addition at low water potentials, fungi are able to increase their cytoplasmic C (thus further increasing
333 C_{mic}/N_{mic} and C_{mic}/P_{mic}) to reduce osmotic pressure and maintain hydration (Schimel et al. 2007). We propose
334 that the loss of N_{mic} and P_{mic} as compared to C_{mic} in all sites could be explained by a net decrease in the size of
335 the microbial biomass, driven by lower substrate (C_{org}) and water availability, together with an increase in the
336 proportional abundance of fungi. However, neither microbial activity nor community composition indicators
337 were measured in our study, thus the underlying mechanisms for the observed seasonal changes remain unclear.

338 Although the microbial pool showed a common trend affected by the summer drought, we observed
339 significant discrepancies on the seasonal dynamics of the available pools. In Alcornocales, nutrient availability
340 was higher in Spring, whereas the opposite was found for the other two sites. Net nutrient pools size is the result
341 of the nutrient release through mineralization, nutrient immobilization and uptake by microorganisms and

342 plants. The rates of N mineralization and nitrification can be more influenced by soil type and soil organic
343 matter quality than by changes in temperature, and the effect of temperature on the rate of P mineralization can
344 vary among soil types (Nadelhoffer et al. 1991). The more severe summer drought in Cardeña and Sierra
345 Nevada might reduce plant uptake capacity (Kozłowski and Pallardi 2002), and increase the proportion of
346 nutrients in the soil when compared to Alcornocales. In a climate change study conducted in the same forest site
347 in Sierra Nevada, Matías et al. (2011) observed that under a dry scenario (30% summer rainfall reduction) soil
348 available nutrients increased and plant and microbial nutrient pools decreased. Thus the contrasting seasonal
349 patterns observed could be the result of different interacting factors such as the activity rates of soil
350 microorganisms, the substrate availability and accessibility, the soil acidity and texture, and the plant nutrient
351 uptake.

352 The effect of soil depth, i.e. decreasing soil and microbial nutrient content from Top soil to Deeper soil, was
353 similar in all forest types. However the magnitude of the change varied among forests, with Cardeña showing
354 the smallest changes. This effect of soil depth has been previously reported for Mediterranean and other forest
355 types (Aponte et al. 2010b; Raubuch and Joergensen 2002; Ross et al. 1996; Wang et al. 2004), the main causes
356 being a decrease in the labile C pools and an increase in the concentration of recalcitrant compounds (Fierer et
357 al. 2003; Goberna et al. 2006). In our study C_{org} decreased with soil depth in all sites, the largest change
358 observed in Alcornocales (from 4.6% to 3.2%) and the smallest one in Cardeña (from 1.5% to 1.2%).The
359 stronger vertical development of cambisol soils in Alcornocales, as evidenced by the deeper soil layer having a
360 significantly lower amount (C_{org}) and quality (C_{org}/N_{tot}) of carbon compounds than the top soil, explains the
361 larger variability of soil properties associated to soil depth observed in this site. This is consistent with the
362 changes observed in microbial C and N values related to soil total pools (C_{mic}/C_{org} and N_{mic}/N_{tot}) in
363 Alcornocales, which are an indicator of a lower efficiency of the microbial biomass to assimilate C and N
364 possibly due to a higher proportion of the soil organic matter being highly recalcitrant (Sparling 1992).
365 Meanwhile, the dominance of shallower and more weakly developed soils in Cardeña (i.e. regosols) underpin
366 the low importance of soil depth as a driver of soil and microbial nutrient content.

367 The size of the microbial pool fell within the ranges observed in other Mediterranean forests (Gallardo et al.
368 2000; Goberna et al. 2006) although it differed significantly between sites. It was not within the scope of this
369 study to investigate the overall differences between forest types, but in general differences among the studied
370 forest types were probably underpinned by the variation in soil types, the amount and quality of soil organic
371 matter, the soil texture and water content, all of them factors constraining the size of the soil microbial biomass

372 (Bohlen et al. 2001; Nielsen et al. 2009). For example, clay content was the highest in Alcornocales (site mean
373 of 35% vs. 8% in Cardeña). Clay content is positively related to microbial biomass and soil organic carbon
374 because it protects microbial biomass from predation by creating refuge microsites. Furthermore, it increases
375 soil organic matter stabilization and soil water retention thus enhancing soil conditions for microbial
376 development (Insam et al. 1989; Sparling 1992).

377

378 *Conclusions*

379 Our findings revealed that across three contrasting Mediterranean forest types with significant differences in
380 soil abiotic conditions, the microbial nutrient pools showed a consistent response in relation to soil depth and
381 seasonal (drought effect) variability, which is indeed mirrored in many other ecosystems at a global scale. In
382 contrast, the direction and magnitude of the variability associated to habitat (tree effect) varied among forest
383 types suggesting a higher complexity in the biotic interactions between the aboveground and belowground
384 components of these ecosystems. Few consistent interactions between factors (tree effect, soil depth and
385 seasonal drought) were observed across forest types.

386 Microbial and soil chemical properties showed similar patterns of covariation in all sites, with microbial
387 biomass responding to variations in the amount and quality of soil organic carbon and soil moisture. Thereby,
388 the quantitative importance of the three studied factors on soil microbial nutrients varied across site, being the
389 most important factor in each case that one which alleviated limitations and imposed the largest variability in
390 substrate and water availability. As such, differences in forest structure and species composition between forest
391 types would underpin the observed inconsistent tree effect on soil microbial properties, since they are related to
392 the amount, quality and spatial and temporal distribution of the resources available to soil microorganisms.

393

394 ACKNOWLEDGEMENTS

395 We thank the Consejería de Medio Ambiente (Andalusian Government) and the Direction of the Los
396 Alcornocales Natural Park, Sierra de Cardeña and Montoro Natural Park and Sierra Nevada National Park for
397 the facilities and support to carry out our field work. We are grateful to Susana Hitos, Eduardo Gutiérrez, Carlos
398 Ros, Raquel Casado, Nacho Villegas, Ramón Ruiz and Eulogio Corral for field and lab assistance. This study
399 was supported by the coordinated Spanish MEC Projects DINAMED (CGL2005-05830-C03), INTERBOS
400 (CGL2008- 4503-C03-01), and DIVERBOS (CGL2011-30285-C02), the Andalusian Projects GESBOME (P06-
401 RNM-1890) and ANASINQUE (PE2010-RNM5782), the Life+ Biodehesa project (11/BIO/ES/000726), three

402 FPI-MEC predoctoral fellowships to CA, LM and VGR, the Subprograma de Técnicos de Apoyo MICINN
403 (PTA2009-1782-I), a MECD postdoctoral grant E-28-2012-0934030 to CA and a EU Marie Curie fellowship
404 (FP7-2011-IEF-300825) to LM.

405

406 REFERENCES

- 407 Alameda D, Villar R, Iriondo JM (2012) Spatial pattern of soil compaction: Trees' footprint on soil physical
408 properties. *Forest Ecol Manag* 283: 128-137. doi: 10.1016/j.foreco.2012.07.018.
- 409 Aponte C, García L, Marañón T (2013) Tree species effects on nutrient cycling and soil biota: a feedback
410 mechanism favouring species coexistence. *Forest Ecol Manag* 309: 36-46. doi:
411 10.1016/j.foreco.2013.05.035.
- 412 Aponte C, García LV, Marañón T, Gardes M (2010a) Indirect host effect on ectomycorrhizal fungi: Leaf fall
413 and litter quality explain changes in fungal communities on the roots of co-occurring Mediterranean oaks.
414 *Soil Biol Biochem* 42: 788-796. doi: 10.1016/j.soilbio.2010.01.014.
- 415 Aponte C, García LV, Pérez-Ramos IM, Gutiérrez E, Marañón T (2011) Oak trees and soil interactions in
416 Mediterranean forests: a positive feedback model. *J Veg Sci* 22: 856-867. doi: 10.1111/j.1654-
417 1103.2011.01298.x.
- 418 Aponte C, Marañón T, García LV (2010b) Microbial C, N and P in soils of Mediterranean oak forests:
419 influence of season, canopy cover and soil depth. *Biogeochemistry* 101: 77-92. doi: 10.1007/s10533-010-
420 9418-5.
- 421 Augusto L, Ranger J, Binkley D, Rothe A (2002) Impact of several common tree species of European
422 temperate forests on soil fertility. *Ann For Sci* 59: 233-253. doi: 10.1051/forest:2002020.
- 423 Barcenas-Moreno G, Garcia-Orenes F, Mataix-Solera J, Mataix-Beneyto J, Baath E (2011) Soil microbial
424 recolonisation after a fire in a Mediterranean forest. *Biol Fert Soils* 47: 261-272. doi: 10.1007/s00374-010-
425 0532-2.
- 426 Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach to
427 multiple testing. *Journal of the Royal Statistical Society* 57: 289-300.
- 428 Bennett LT, Kasel S, Tibbits J (2009) Woodland trees modulate soil resources and conserve fungal diversity in
429 fragmented landscapes. *Soil Biol Biochem* 41: 2162-2169. doi: 10.1016/j.soilbio.2009.07.030.
- 430 Bohlen PJ, Groffman PM, Driscoll CT, Fahey TJ, Siccama TG (2001) Plant-soil-microbial interactions in a
431 northern hardwood forest. *Ecology* 82: 965-978.
- 432 Bray RH, Kurtz LT (1945) Determination of total, organic and available forms of phosphorous in soils. *Soil*
433 *Science* 59: 39-45. doi: 10.1097/00010694-194501000-00006.
- 434 Bremner J, Keeney D (1965) Steam distillation methods for determination of ammonium nitrate and nitrate.
435 *Analytica Chimica Acta*: 485-495. doi: 10.1016/S0003-2670(00)88973-4.
- 436 Brookes PC, Landman A, Pruden G, Jenkinson DS (1985) Chloroform fumigation and the release of soil
437 nitrogen: A rapid direct extraction method to measure microbial biomass nitrogen in soil. *Soil Biol Biochem*
438 17: 837-842. doi: 10.1016/0038-0717(85)90144-0.
- 439 Brookes PC, Powlson DS, Jenkinson DS (1982) Measurement of microbial phosphorus in soil. *Soil Biol*
440 *Biochem* 14: 319-329.

441 Carreira JA, Niell FX, Lajtha K (1994) Soil nitrogen availability and nitrification in Mediterranean shrublands
442 of varying fire history and successional stage. *Biogeochemistry* 26: 189-209. doi: 10.1007/bf00002906.

443 Chen CR, Xu ZH, Blumfield TJ, Hughes JM (2003) Soil microbial biomass during the early establishment of
444 hoop pine plantation: seasonal variation and impacts of site preparation. *Forest Ecol Manag* 186: 213-225.
445 doi: 10.1016/S0378-1127(03)00275-5.

446 Cleveland CC, Liptzin D (2007) C:N:P stoichiometry in soil: Is there a "Redfield ratio" for the microbial
447 biomass? *Biogeochemistry* 85: 235-252. doi: 10.1007/s10533-007-9132-0.

448 Corre MD, Schnabel RR, Stout WL (2002) Spatial and seasonal variation of gross nitrogen transformations
449 and microbial biomass in a Northeastern US grassland. *Soil Biol Biochem* 34: 445-457. doi: 10.1016/S0038-
450 0717(01)00198-5.

451 Csardi G, Nepusz T (2006) The igraph software package for complex network research. *InterJournal Complex*
452 *Systems*: 1695.

453 Ettema CH, Wardle DA (2002) Spatial soil ecology. *Trends Ecol Evol* 17: 177-183. doi: 10.1016/S0169-
454 5347(02)02496-5.

455 Fierer N, Schimel JP, Holden PA (2003) Variations in microbial community composition through two soil
456 depth profiles. *Soil Biol Biochem* 35: 167-176. doi: 10.1016/S0038-0717(02)00251-1.

457 Gallardo A (2003) Effect of tree canopy on the spatial distribution of soil nutrients in a Mediterranean Dehesa.
458 *Pedobiologia* 47: 117-125. doi: 10.1078/0031-4056-00175.

459 Gallardo A, Rodríguez-Saucedo JJ, Covelo F, Fernández-Alés R (2000) Soil nitrogen heterogeneity in a
460 Dehesa ecosystem. *Plant and Soil* 222: 71-82.

461 Gallardo A, Schlesinger WH (1994) Factors limiting microbial biomass in the mineral soil and forest floor of a
462 warm-temperate forest. *Soil Biol Biochem* 26: 1409-1415. doi: 10.1016/0038-0717(94)90225-9.

463 García C, Hernandez T, Roldan A, Martin A (2002) Effect of plant cover decline on chemical and
464 microbiological parameters under Mediterranean climate. *Soil Biol Biochem* 34: 635-642. doi:
465 10.1016/S0038-0717(01)00225-5.

466 García LV (2003) Controlling the false discovery rate in ecological research. *Trends Ecol Evol* 18: 553-554.
467 doi: 10.1016/j.tree.2003.08.011.

468 García LV, Maltez-Mouro S, Pérez-Ramos IM, Freitas H, Marañón T (2006) Counteracting gradients of light
469 and soil nutrients in the understorey of Mediterranean oak forest. *Web Ecology* 6: 67-74. doi: 10.5194/we-6-
470 67-2006.

471 Gaudinski JB, Trumbore SE, Davidson EA, Zheng S (2000) Soil carbon cycling in a temperate forest:
472 radiocarbon-based estimates of residence times, sequestration rates and partitioning of fluxes.
473 *Biogeochemistry* 51: 33-69.

474 Gil Torres J, Rodero Pérez I, Odierna C (2003) Inventario de los suelos de la provincia de Cordoba.
475 Diputacion de Cordoba, Spain.

476 Goberna M, Pascual JA, García C, Sánchez J (2007) Do plant clumps constitute microbial hotspots in semiarid
477 Mediterranean patchy landscapes? *Soil Biol Biochem* 39: 1047-1054. doi: 10.1016/j.soilbio.2006.11.015.

478 Goberna M, Sánchez J, Pascual JA, García C (2006) Surface and subsurface organic carbon, microbial
479 biomass and activity in a forest soil sequence. *Soil Biol Biochem* 38: 2233-2243. doi:
480 10.1016/j.soilbio.2006.02.003.

481 Hackl E, Pfeffer M, Donat C, Bachmann G, Zechmeister-Boltenstern S (2005) Composition of the microbial
482 communities in the mineral soil under different types of natural forest. *Soil Biol Biochem* 37: 661-671. doi:
483 10.1016/j.soilbio.2004.08.023.

484 Hassink J (1994) Effect of soil texture on the size of the microbial biomass and on the amount of c and n
485 mineralized per unit of microbial biomass in dutch grassland soils. *Soil Biol Biochem* 26: 1573-1581. doi:
486 10.1016/0038-0717(94)90100-7.

487 Huang Z, Wan X, He Z, Yu Z, Wang M, Hu Z, Yang Y (2013) Soil microbial biomass, community
488 composition and soil nitrogen cycling in relation to tree species in subtropical China. *Soil Biol Biochem* 62:
489 68-75. doi: 10.1016/j.soilbio.2013.03.008.

490 Insam H, Parkinson D, Domsch KH (1989) Influence of macroclimate on soil microbial biomass. *Soil Biol*
491 *Biochem* 21: 211-221. doi: 10.1016/0038-0717(89)90097-7.

492 Jensen KD, Beier C, Michelsen A, Emmett BA (2003) Effects of experimental drought on microbial processes
493 in two temperate heathlands at contrasting water conditions. *Appl Soil Ecol* 24: 165-176. doi:
494 10.1016/S0929-1393(03)00091-X.

495 King JR, Jackson DA (1999) Variable selection in large environmental data sets using principal components
496 analysis. *Environmetrics* 10: 67-77. doi: 10.1002/(SICI)1099-095X(199901/02)10:1%3C67::AID-
497 ENV336%3E3.0.CO;2-0.

498 Kozłowski TT, Pallardi SG (2002) Acclimation and adaptive responses of woody plants to environmental
499 stresses. *Bot Rev* 68: 270-334. doi: 10.1663/0006-8101(2002)068%5B0270:AAAROW%5D2.0.CO;2.

500 Lambers H, Chapin III F, Pons T (1998) *Plant physiological ecology*. Springer-Verlag New York.

501 Lindo Z, Visser S (2003) Microbial biomass, nitrogen and phosphorus mineralization, and mesofauna in
502 boreal conifer and deciduous forest floors following partial and clear-cut harvesting. *Canadian Journal of*
503 *Forest Research* 33: 1610-1620. doi: 10.1139/x03-080.

504 Liu L, Gundersen P, Zhang T, Mo J (2012) Effects of phosphorus addition on soil microbial biomass and
505 community composition in three forest types in tropical China. *Soil Biol Biochem* 44: 31-38. doi:
506 10.1016/j.soilbio.2011.08.017.

507 Lucas-Borja ME, Candel D, Jindo K, Moreno JL, Andrés M, Bastida F (2012) Soil microbial community
508 structure and activity in monospecific and mixed forest stands, under Mediterranean humid conditions. *Plant*
509 *and Soil* 354: 359-370. doi: 10.1007/s11104-011-1072-8.

510 Malchair S, Carnol M (2009) Microbial biomass and C and N transformations in forest floors under European
511 beech, sessile oak, Norway spruce and Douglas-fir at four temperate forest sites. *Soil Biol Biochem* 41: 831-
512 839. doi: 10.1016/j.soilbio.2009.02.004.

513 Maltez-Mouro S, García L, Marañón T, Freitas H (2005) The combined role of topography and overstorey tree
514 composition in promoting edaphic and floristic variation in a Mediterranean forest. *Ecol Res* 20: 668-677.
515 doi: 10.1007/s11284-005-0081-6.

516 Marañón-Jiménez S, Castro J, Kowalski AS, Serrano-Ortiz P, Reverter BR, Sánchez-Cañete EP, Zamora R
517 (2011) Post-fire soil respiration in relation to burnt wood management in a Mediterranean mountain
518 ecosystem. *Forest Ecol Manag* 261: 1436-1447. doi: 10.1016/j.foreco.2011.01.030.

519 Matías L, Castro J, Zamora R (2011) Soil-nutrient availability under a global-change scenario in a
520 Mediterranean mountain ecosystem. *Glob Change Biol* 17: 1646-1657. doi: 10.1111/j.1365-
521 2486.2010.02338.x.

522 Monokrousos N, Papatheodorou EM, Diamantopoulos JD, Stamou GP (2004) Temporal and spatial variability
523 of soil chemical and biological variables in a Mediterranean shrubland. *Forest Ecol Manag* 202: 83-91. doi:
524 10.1016/j.foreco.2004.07.039.

525 Nadelhoffer KJ, Giblin AE, Shaver GR, Laundre JA (1991) Effects of Temperature and Substrate Quality on
526 Element Mineralization in Six Arctic Soils. *Ecology* 72: 242-253. doi: 10.2307/1938918.

527 Nielsen PL, Andresen LC, Michelsen A, Schmidt IK, Kongstad J (2009) Seasonal variations and effects of
528 nutrient applications on N and P and microbial biomass under two temperate heathland plants. *Appl Soil*
529 *Ecol* 42: 279-287. doi: 10.1016/j.apsoil.2009.05.006.

530 Olsen SR, Cole CV, Watanabe FS, L.A. D (1954) Estimation of available phosphorous in soils by extraction
531 with sodium bicarbonate. In: USDA (ed). USDA, Washington, D.C.

532 Polo A (2006) Heterogeneidad edáfica en una parcela experimental de bosque mixto de *Quercus suber* y
533 *Quercus canariensis* del Parque Natural de los Alcornocales (La Panera). Escuela Universitaria Ingenieros
534 Técnicos Agrícolas. Universidad de Sevilla.

535 Prescott CE, Grayston SJ (2013) Tree species influence on microbial communities in litter and soil: Current
536 knowledge and research needs. *Forest Ecol Manag*. doi: 10.1016/j.foreco.2013.02.034.

537 Quilchano C, Marañón T (2002) Dehydrogenase activity in Mediterranean forest soils. *Biol Fert Soils* 35: 102.
538 doi: 10.1007/s00374-002-0446-8.

539 Raubuch M, Joergensen RG (2002) C and net N mineralisation in a coniferous forest soil: the contribution of
540 the temporal variability of microbial biomass C and N. *Soil Biol Biochem* 34: 841-849. doi: 10.1016/S0038-
541 0717(02)00016-0.

542 Rey A, Pegoraro E, Tedeschi V, De Parri I, Jarvis PG, Valentini R (2002) Annual variation in soil respiration
543 and its components in a coppice oak forest in Central Italy. *Glob Change Biol* 8: 851-866. doi:
544 10.1046/j.1365-2486.2002.00521.x.

545 Ross DJ, Tate KR, Feltham CW (1996) Microbial biomass, and C and N mineralization, in litter and mineral
546 soil of adjacent montane ecosystems in a southern beech (*Nothofagus*) forest and a tussock grassland. *Soil*
547 *Biol Biochem* 28: 1613-1620. doi: 10.1016/S0038-0717(96)00255-6.

548 Rutigliano FA, D'Ascoli R, Virzo De Santo A (2004) Soil microbial metabolism and nutrient status in a
549 Mediterranean area as affected by plant cover. *Soil Biol Biochem* 36: 1719-1729. doi:
550 10.1016/j.soilbio.2004.04.029.

551 Sagova-Mareckova M, Omelka M, Cermak L, Kamenik Z, Olsovska J, Hackl E, Kopecky J, Hadacek F (2011)
552 Microbial communities show parallels at sites with distinct litter and soil characteristics. *Appl Environ*
553 *Microb* 77: 7560-7567. doi: 10.1128/AEM.00527-11.

554 Sardans J, Peñuelas J, Rodà F (2005) Changes in nutrient status, retranslocation and use efficiency in young
555 post-fire regeneration *Pinus halepensis* in response to sudden N and P input, irrigation and removal of
556 competing vegetation. *Trees* 19 19: 233-250. doi: 10.1007/s00468-004-0374-3.

557 Schimel J, Balsler TC, Wallenstein M (2007) Microbial stress-response physiology and its implications for
558 ecosystem function. *Ecology* 88: 1386-1394. doi: 10.1890/06-0219.

559 Schmidt IK, Jonasson S, Michelsen A (1999) Mineralization and microbial immobilization of N and P in
560 arctic soils in relation to season, temperature and nutrient amendment. *Appl Soil Ecol* 11: 147-160. doi:
561 10.1016/S0929-1393(98)00147-4.

562 Singh JS, Raghubanshi AS, Singh RS, Srivastava SC (1989) Microbial biomass acts as a source of plant
563 nutrients in dry tropical forest and savanna. *Nature* 338: 499-500. doi: 10.1038/338499a0.

564 Smolander A, Kitunen V (2002) Soil microbial activities and characteristics of dissolved organic C and N in
565 relation to tree species. *Soil Biol Biochem* 34: 651-660. doi: 10.1016/S0038-0717(01)00227-9.

566 Sparks DL (1996) *Methods of Soil Analysis. Part 3. Chemical Methods* Soil Science Society of America and
567 American Society of Agronomy, Madison, Wisconsin, USA.

568 Sparling GP (1992) Ratio of microbial biomass carbon to soil organic carbon as a sensitive indicator of
569 changes in soil organic matter. *Soil Res* 30: 195-207. doi: /10.1071/SR9920195.

570 Ste-Marie C, Pare D, Gagnon D (2007) The contrasting effects of aspen and jack pine on soil nutritional
571 properties depend on parent material. *Ecosystems* 10: 1299-1310. doi: 10.1007/s10021-007-9098-8.

572 Thoms C, Gattinger A, Jacob M, Thomas FM, Gleixner G (2010) Direct and indirect effects of tree diversity
573 drive soil microbial diversity in temperate deciduous forest. *Soil Biol Biochem* 42: 1558-1565. doi:
574 10.1016/j.soilbio.2010.05.030.

575 Ushio M, Kitayama K, Balser TC (2010) Tree species-mediated spatial patchiness of the composition of
576 microbial community and physicochemical properties in the topsoils of a tropical montane forest. *Soil Biol*
577 *Biochem* 42: 1588-1595. doi: 10.1016/j.soilbio.2010.05.035.

578 van der Putten W, Bardgett R, de Ruiter P, Hol W, Meyer K, Bezemer T, Bradford M, Christensen S, Eppinga
579 M, Fukami T, Hemerik L, Molofsky J, Schädler M, Scherber C, Strauss S, Vos M, Wardle D (2009)
580 Empirical and theoretical challenges in aboveground–belowground ecology. *Oecologia* 161: 1-14. doi:
581 10.1007/s00442-009-1351-8.

582 Vance ED, Brookes PC, Jenkinson DS (1987) An extraction method for measuring soil microbial biomass C.
583 *Soil Biol Biochem* 19: 703-707. doi: 10.1016/0038-0717(87)90052-6.

584 Wang FE, Chen YX, Tian GM, Kumar S, He YF, Fu QL, Lin Q (2004) Microbial biomass carbon, nitrogen
585 and phosphorus in the soil profiles of different vegetation covers established for soil rehabilitation in a red
586 soil region of southeastern China. *Nutr Cycl Agroecosys* 68: 181-189. doi:
587 10.1023/B:FRES.0000017470.14789.2a.

588 Wilkinson SC, Anderson JM, Scardelis SP, Tisiafouli M, Taylor A, Wolters V (2002) PLFA profiles of
589 microbial communities in decomposing conifer litters subject to moisture stress. *Soil Biol Biochem* 34: 189-
590 200. doi: 10.1016/S0038-0717(01)00168-7.

591 WRB IWG (2006) *World reference base for soil resources 2006: A framework for international classification,*
592 *correlation and communication.* In: FAO (ed) *World Soil Resources Reports No 103*, Rome.

593 Xiang SR, Doyle A, Holden PA, Schimel JP (2008) Drying and rewetting effects on C and N mineralization
594 and microbial activity in surface and subsurface California grassland soils. *Soil Biol Biochem* 40: 2281-
595 2289. doi: 10.1016/j.soilbio.2008.05.004.

596 Zhong Z, Makeshin F (2006) Differences of Soil Microbial Biomass and Nitrogen Transformation under Two
597 Forest Types in Central Germany. *Plant and Soil* 283: 287-297. doi: 10.1007/s11104-006-0018-z.

598
599

600 FIGURE CAPTIONS

601

602 **Fig. 1.** Location of the studied forest sites in the Iberian Peninsula.

603 **Fig. 2.** Microbial and soil nutrient fractions in Alcornocales (A), Cardeña (C) and Sierra Nevada (SN).

604 Differences between the levels of each factor are as indicated by repeated measurement ANOVAs with season

605 as a within-group effect and site, habitat and depth as between-group effects followed by Tukey's posthoc

606 comparisons (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$). Abbreviations are: C_{mic} , microbial C; N_{mic} , microbial N;

607 P_{mic} , microbial P; DOC, dissolved organic C; DON, dissolved organic N.

608 **Fig. 3.** Soil inorganic nutrient fractions and organic carbon in Alcornocales (A), Cardeña (C) and Sierra Nevada

609 (SN). Differences between the levels of each factor are as indicated by repeated measurement ANOVAs with

610 season as a within-group effect and site, habitat and depth as between-group effects followed by Tukey's

611 posthoc comparisons (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$). Abbreviations are: P_{inorg} , inorganic available P;

612 C_{org} , organic C.

613 **Fig. 4.** Percentage of the total variance explained by each of the studied factors, habitat (tree effect), soil depth

614 and season, for each variable in each site. C_{mic} , N_{mic} , P_{mic} : microbial C, N and P, respectively; C_{org} : organic C;

615 P_{inorg} : inorganic P; N_{inorg} : inorganic N; DOC, DON: dissolved organic C and N, respectively; C_{mic}/N_{mic} , C_{mic}/P_{mic} :

616 ratios between respective variables; Overall: mean across all variables.

617 **Fig. 5.** PCA ordination plot showing the distribution of Spring and Summer values of each study sites. C_{mic} ,

618 N_{mic} , P_{mic} : microbial C, N and P, respectively; P_{inorg} : inorganic P; DOC, DON: dissolved organic C and N,

619 respectively; N_{tot} : total N; C_{org} : organic C; C_{mic}/P_{mic} , C_{mic}/N_{mic} , N_{mic}/N_{inorg} , P_{mic}/P_{inorg} : ratios between respective

620 variables. Depth, season and habitat (in grey) are supplementary variables included as passive in the analysis.

621

622

623 TABLES

624 Table 1. Characteristics of the studied forest sites. Values are site means (\pm standard deviation, when
 625 provided) and habitat means in square brackets [Open; Tree].

626

	Alcornocales	Cardeña	Sierra Nevada
Coordinates	36°31' N, 5°34' W	38° 15' N, 4° 21' W	37°05' N, 3°28' W
Altitude (m a.s.l.)	545	750	1650
Soil			
Bedrock	sandstone	granite	limestone
pH	acidic [6.34; 6.07] ^a	acidic 5.4 ^b	basic [8.1; 7.7] ^c
Soil type	cambisol	regosol	regosol, cambisol
Texture	sandy	sandy	loamy
Sand (%)	[44; 49] ^a	[80; 79] ^d	[22; 19] ^c
Clay (%)	[39; 33] ^a	[4.8; 4.5] ^d	[29; 37] ^c
CEC (meq 100g ⁻¹)	[23.1; 19.7] ^a	[7.8; 8.6] ^d	[14.7; 18.5] ^c
Litter (g m ⁻²)	[45±38; 936±350]	[282± 259; 1200± 875]	[559±362; 8594±5543]
Climate			
Temperature (°C)			
mean annual	15.5	15.3	12.1
mean min	9.1	7.3	-1.1
mean max	23.6	25.3	29.2
Rainfall (mm)			
annual	1117	752	811
spring	259	151	206
summer	28	39	43
Vegetation			
Tree density (stems ha ⁻¹)	219	131	787
Basal area (m ² ha ⁻¹)	24	13	-
Experimental design (n)	Open (10)	Open (19)	Open (16)
	<i>Q. suber</i> / <i>Q. canariensis</i> (20)	<i>Q. ilex</i> (19)	<i>P. sylvestris</i> (16)

627 ^a Values determined in 0-25 cm deep soil samples (Polo 2006)628 ^b Mean value for regosols in the region (0-15cm Gil Torres et al. 2003)629 ^c Values determined in 0-16 cm deep soil samples (Matías et al, unpublished data)

630 ^d Values determined in 2-14 cm deep soil samples (Alameda et al. 2012/ Alameda et al.,
 631 unpublished results).

632

633 Table 2. Mean (\pm SE) values of the measured soil variables across habitat, season and soil depth by
 634 site. Letters indicate differences between sites ($P < 0.05$).

635


	Alcornocales		Cardeña		Sierra Nevada	
C_{mic} (mg kg ⁻¹)	378 \pm 18	a	58 \pm 4	b	63 \pm 3	b
N_{mic} (mg kg ⁻¹)	46 \pm 3	a	17 \pm 1	b	6.9 \pm 0.5	c
P_{mic} (mg kg ⁻¹)	7.9 \pm 0.6	a	1.0 \pm 0.1	b	4.0 \pm 0.3	c
C_{org} (%)	3.9 \pm 0.1	a	1.3 \pm 0.1	b	3.0 \pm 0.1	c
P_{inorg} (mg kg ⁻¹)	2.7 \pm 0.2	a	1.3 \pm 0.1	b	3.4 \pm 0.2	c
NH_4 (mg kg ⁻¹)	8.1 \pm 0.6	a	2.6 \pm 0.2	b	19 \pm 1	c
NO_3 (mg kg ⁻¹)	2.5 \pm 0.2	a	0.7 \pm 0.1	b	7.9 \pm 0.4	c
DOC (mg kg ⁻¹)	168 \pm 7	a	117 \pm 8	b	41 \pm 3	c
DON (mg kg ⁻¹)	27 \pm 1	a	6.7 \pm 0.3	b	3.9 \pm 0.2	c
N_{tot} (%)	0.28 \pm 0.01	a	0.09 \pm 0.0	b	0.22 \pm 0.01	c
C_{mic}/N_{mic}	9.3 \pm 0.3	a	7.2 \pm 0.7	b	21 \pm 4	c
C_{mic}/P_{mic}	108 \pm 18	a	145 \pm 28	a	23 \pm 2	b
Moisture Spring (%)	22 \pm 1	a	9.3 \pm 0.6	b	13 \pm 1	c
Moisture Summer (%)	11.0 \pm 0.4	a	2.7 \pm 0.2	b	3.4 \pm 0.2	c

Table 3. Repeated measurements ANOVA for the studied soil properties. F and P-values for between effects (forest site, habitat and soil depth), within effect (season) and three way interactions are presented. Significant effects are marked with asterisks (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$). C_{mic} , N_{mic} , P_{mic} : microbial C, N and P, respectively; C_{org} : organic C; N_{inorg} : inorganic N ($NH_4 + NO_3$); P_{inorg} : inorganic P; DOC, DON: dissolved organic C and N, respectively.






Effect	C_{mic}		N_{mic}		P_{mic}		C_{org}		P_{inorg}		N_{inorg}		DOC		DON		C_{mic}/N_{mic}		C_{mic}/P_{mic}		Moisture	
	F	P	F	P	F	P	F	P	F	F	F	P	F	P	F	P	F	P	F	P	F	P
Site	342***		368***		170.4***		11.2***		75.6***		619***		303.3***		590.2***		65.6***		126.9***		265.4***	
Habitat	12.1***		7.79**		14.9***		15.4***		3.83		12.5***		39.2***		16.6***		1.13		5.82*		7.96**	
Depth	41.5***		69.2***		63.5***		6.93**		31.1***		18.3***		45.3***		37.6***		1.73		10.9**		0.93	
Site×Habitat	9.7***		22.3***		13***		7.55**		6.61**		7.36***		3.15		9.12***		6.79***		18.2***		1.71	
Site×Depth	8.1***		3.67*		22.2***		3.96*		3.25		1.87		6.2**		4.76*		0.1		4.86**		8.93***	
Habitat×Depth	0.97		2.14		0.01		0.01		0.02		0.07		2.54		3.52		0.19		8.68**		0.91	
Site×Habitat×Depth	2.54		1.37		0.47		4.05*		2.67		2.58		3.17		1.35		0.22		7.93***		0.9	
Season	8.83**		116.8***		154***		0.14		5.65		0.97		144.9***		11.2***		92.1***		36.4***		1283***	
Season×Site	23.3***		27.2***		31.7***		45.8***		105***		193***		21.2***		22.9***		12.3***		0.40		13.6***	
Season×Habitat	0.05		4.41*		11.4***		6.42*		0.04		5.11*		10.9***		8.13**		6.27*		7.67**		0.06	
Season×Depth	0.04		0.09		0.5		1.54		5.32*		4.27		0.52		0.75		0.89		0.00		13.4***	
Season×Site×Habitat	0.18		0.02		2.45		3.36*		7.48***		3.17		8.67***		5.95**		0.7		0.65		1.45	
Season×Site×Depth	0.28		1.7		0.24		1.46		2.32		3.79		8.61***		5.98**		0.03		0.03		6.61**	
Season×Habitat×Depth	0.08		0.27		3.21		0.84		0.63		0.51		2.97		0.45		0		1.26		0.38	

Site: Alcornocales, Cardeña and Sierra Nevada
Habitat: Tree and Open;
Depth: Top soil (0-8cm) and Deeper soil (8-16cm);
Season: Spring and Summer .

Fig. 1.








Habitat








A C SN

Depth




A C SN

Season




A C SN



Alcornocales

Cardeña



Sierra Nevada



Variance explained (%)

Habitat
 Depth
 Season

Fig 5.



Sierra Nevada

