



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Symonds, MRE;Magrath, MJL;Latty, TM

Title:

Reproductive Consequences of Male Arrival Order in the Bark Beetle, *Ips grandicollis*

Date:

2012-07-01

Citation:

Symonds, M. R. E., Magrath, M. J. L. & Latty, T. M. (2012). Reproductive Consequences of Male Arrival Order in the Bark Beetle, *Ips grandicollis*. *Journal of Insect Behavior*, 25 (4), pp.401-407. <https://doi.org/10.1007/s10905-011-9308-2>.

Persistent Link:

<https://hdl.handle.net/11343/282755>

Reproductive consequences of male arrival order in the bark beetle, *Ips grandicollis*

Matthew R. E. Symonds^{1,2*}, Michael J. L. Magrath³ and Tanya M. Latty^{4, 5}

¹ *Department of Zoology, University of Melbourne, Victoria 3010, Australia*

² *Centre for Integrative Ecology, School of Life and Environmental Sciences, Deakin*

University, 221 Burwood Highway, Burwood, Victoria 3125, Australia (present address)

³ *Department of Wildlife Conservation and Science, Zoos Victoria, Victoria 3052, Australia*

⁴ *Department of Biological Sciences, University of Calgary, 2500 University Dr. NW,
Calgary, Alberta T2N 1N4, Canada*

⁵ *School of Biological Sciences, University of Sydney, New South Wales 2006, Australia
(present address)*

* Corresponding author

Tel: +61 3 9251 7437

Fax: +61 3 9251 7626

Email: symonds@deakin.edu.au

Running head: Arrival order in *Ips grandicollis*

Abstract

For group-living animals the choice of whether to join aggregations or initiate their own is influenced by potential benefits such as group protection and reduced energetic expenditure, as well as costs such as competition for food and mates. The bark beetle *Ips grandicollis* is an invasive pest species that colonises recently felled timber in Australian pine (*Pinus* spp.) plantations. Male beetles initiate colonies by burrowing under the bark of trees and emitting an aggregation pheromone which attracts conspecifics, including a harem of females with whom they mate. We predicted that males that initiated colonies, or who arrived early, would have larger harems than later arrivals (due to decreased competition for females). However, we found the opposite effect with early-arriving males actually associated with *fewer* females than later arriving males, although this may have resulted from some females leaving harems as they get older. We conclude that pioneering does not improve male likelihood of attracting females in *Ips grandicollis*, at least initially, but it may provide advantages for offspring when competing for food during development.

Key words: aggregation, harem size, timing of arrival, reproductive costs, pioneer, Scolytinae

Introduction

Group living individuals are faced with numerous decisions, including the question of which groups to join and when. When aggregations are associated with procuring a resource (habitat, mates, food), the costs and benefits of joining a group may vary depending on the order in which an individual arrives at the group. Initiating an aggregation (pioneering) may allow individuals to secure better quality territories and/or procure an increased proportion of the resource (Bensch and Hasselquist 1991; Candolin and Voigt 2003; Smith and Moore 2005), but may incur costs such as greater energetic expenditure, and mortality or injury from prey defences (Heinsohn and Packer 1995; Pekar et al. 2005). Individuals that join established groups ('joiners') may avoid these costs, but could also experience higher competition and/or a decreased share of the resource (Mangel 1990; Giraldeau and Beauchamp 1999).

Many bark beetle species (subfamily: Scolytinae) live and feed in aggregations beneath the bark of coniferous trees (Wood 1982). In many species aggregation is necessary to overcome tree defences and the fitness of the individuals is contingent on the size of the aggregation (Raffa and Berryman 1983). In the genus *Dendroctonus*, individuals that initiate aggregations or arrive early in the colonisation have higher mortality and lower reproductive success because they have to deal with the toxic effects of tree defences at their strongest (Pureswaran et al. 2006, Latty and Reid 2009). Bark beetles of the genus *Ips*, however, typically attack non-living *Pinus* material (such as freshly windblown trees or cut log billets), which should remove this cost associated with pioneering. Indeed, early arrival may even be beneficial because of reduced competition for mates and food resources (Raffa 2001).

Ips bark beetles have a harem polygynous mating system (Kirkendall 1983). Adult males search for fallen logs, burrowing into the phloem and carving a nuptial chamber where they begin releasing an aggregation pheromone (Wood 1982). This attracts other males to the log, and up to 8 females to the specific boreholes occupied by the males (Kirkendall 1983;

Schlyter and Zhang 1996; Latty et al. 2009). Each female carves her own tunnel radiating outward from the central nuptial chamber (Figure 1). The female then lays eggs in individual niches along the sides of the tunnel. Harem size is related to reproductive success such that males with larger harems produce more offspring than males with small harems (Robertson 1998).

We examined the effect of male arrival order on their subsequent harem size in a South Australian population of the invasive eastern five-spined bark beetle *Ips grandicollis*. Early arriving males may face less competition for females; we therefore tested the hypothesis that these males will have larger harems. We also examined whether time of arrival and harem size were linked to male body size.

Methods and Materials

We examined natural colonisation by *Ips grandicollis* bark beetles of log billets in a 39 year-old *Pinus radiata* plantation at Wirrabara State Forest, South Australia (138° 16' E, 33° 1' S) approximately 250km north of Adelaide. The experiments were carried out in summer 2007, from the 13th to 23rd February. Daily maximum temperatures ranged from 32 – 41°C and weather conditions were fine and sunny during the course of the work.

We cut 50cm-long log billets from freshly felled *Pinus radiata* trees. Log billet diameter ranged from 9 to 23 cm. The billets were put together in 20 piles of three (=60 billets), in order to increase the likelihood of colonisation by beetles. Each pile was spaced 10 metres apart along a transect.

We monitored colonisation by examining the billets every day for fresh *Ips grandicollis* boreholes (easily identified by the presence of orange frass on the outside of the log). New boreholes were marked by placing a date-labelled flat-topped push-pin into the log next to the hole. Some logs are colonised earlier than others, so the push-pin data allowed us to record

not only the arrival order of each male to the log, but also the age of the borehole when its internal characteristics (see below) were measured. We measured arrival order as the day on which the borehole appeared in each log subsequent to the first borehole appearing (i.e. the first boreholes to appear in each log were assigned a day of arrival = 1, the boreholes that appeared the next day had a day of arrival = 2, and so on).

As colonisation continues, and the phloem is progressively consumed and degraded, the patterns of galleries become difficult to distinguish making accurate measurement of male harem size increasingly problematic. Hence, we concentrated on the effects of arrival order on reproductive success in the early stages of colonisation. Whilst this restricts our ability to discuss ultimate effects on survival and success of offspring, it does allow more accurate assessment of the initial costs or benefits of pioneering. Consequently, after 8-10 days, logs were carefully stripped of bark with a chisel, to uncover the nuptial chamber and galleries associated with each borehole. In South Australia, it takes male *Ips grandicollis* 4 days to attract their full complement of females (Morgan 1967). Therefore, we restricted our analysis to males who had been in the logs for at least 5 days. We noted the number of beetles observed under each borehole and the number of galleries. This provided us with two measures of harem size – one based on number of galleries and one based on actual number of females observed. Every borehole was assumed to have one male beetle, with the remaining beetles assumed to be female (Kirkendall 1983).

A subset of males that were physically undamaged during the removal of the bark were collected in labelled plastic vials and stored in the freezer for later body size measurements. Males were generally recognisable in the field as being the one beetle in the gallery system that was not at the far end of a gallery. However, in cases where this was not clear we collected all the beetles from each gallery system, and sexed them under a microscope in the lab using the criteria of Lanier and Cameron (1969). The body length and body width (across

the base of the thorax) of each individual (male or female) was measured in the lab using callipers. Because *Ips* bark beetles are essentially cylindrical in shape we used length and width measures to estimate body volume:

$$Volume = \pi \times length \times \frac{width^2}{4}$$

Relationships between arrival order and both measures of harem size were examined using hierarchical mixed modelling to account for the non-independence and hierarchical structure of the data. Three trees were used as a source of the logs in the experiment. However, source tree did not account for significant variation in response variables so was not included in statistical models.

For analysis, arrival order was calculated in the context of the log pile (days from first borehole in any of the three logs in each pile) because females would most likely detect male aggregation pheromones based on the signal coming from a log pile rather than either individual logs within each pile or the entire population. Nevertheless, we also calculated male arrival order relative to individual log and the whole population to check our results. Analysis was performed using MLwiN 2.02 (Rasbash et al. 2004).

Results

In total we collected data from 95 harems across 43 logs in 19 log-piles. Mean ($\pm$ s.e.) harem size was 3.57 ± 0.16 galleries or 2.96 ± 0.15 females. The analysis of arrival order at all three levels (i.e. relative to log, log pile and population) produced qualitatively the same results. Although trends were stronger (and in the case of the relationship with number of females more highly significant) in the analyses at the log and population levels, we present only the results from the analysis at the log pile level (see justification in the Methods and Materials above).

In 58 harems (61%), the number of galleries equalled the number of females, however in 35 harems (37%), there were fewer females than there were galleries. We found a significant relationship between arrival order and the proportion of females to gallery number ($b = -0.36 \pm 0.15$ s.e.; $P = 0.02$) indicating that females were more likely to be ‘missing’ from galleries the longer the harem had been established. This proportion of females to galleries was not significantly related to the number of galleries in the harem ($b = -0.13 \pm 0.09$ s.e.; $P = 0.16$).

The relationship between arrival order and harem size differed according to the measure of harem size used (Figure 2). We did not find a significant association between arrival order and number of galleries ($b = 0.14 \pm 0.17$ s.e.; $P = 0.43$). However, there were fewer females in the harems of males that arrived earliest ($b = 0.38 \pm 0.16$ s.e.; $P = 0.02$).

There was no significant relationship between male body size (volume) and arrival order ($b = -0.13 \pm 0.13$ s.e.; $P = 0.29$), nor was there any association between male body size and harem size ($b = 0.11 \pm 0.12$ s.e.; $P = 0.39$).

Discussion

Contrary to our prediction, early arriving males did not attract more females than those that arrived later in the aggregation. Indeed, using number of associated females as our measure of harem size, we found that early arriving males were associated with *fewer* females than were late arriving males. However, no relationship between arrival order and harem size was apparent when using number of galleries as the measure. The discrepancy in results using our two measures of harem size stems from the fact that 37% of harems in our study had fewer females than galleries, with the proportion of females to galleries declining in older harems. Therefore the relationship between fewer females and early arrival by males is likely explained by the increase in ‘missing’ females from older boreholes. This loss of females may have one of two explanations. The first is that individual females abandon harems. Re-

emergence is a relatively common phenomenon in bark beetles, and is often related to declines in resource quality or increases in density and competition (Kirkendall 1983; Byers 1989; Anderbrant 1989). However, female densities in logs were low in our study (mean = 0.70 ± 0.09 s.e. females per dm^2), and there were not proportionately more females missing from larger harems which makes it seem unlikely that competition is driving re-emergence here. A second possibility explaining the ‘missing’ females is not that they are missing but that individual females carve out more than one gallery, a phenomenon which is also known to occur in *Ips* species (Reid 1999). If this is the case, then our results suggest that early males may actually be losing out in terms of attracting females. However, because we cannot rule out either possibility it is most appropriate to conclude that there is no benefit to males of arriving early, at least in terms of harem size.

As with previous work on harem size in *Ips* (Schlyter and Zhang 1996; Latty et al. 2009) we found no evidence of an effect of male size on harem size, contrary to what is typically found in vertebrate harem polygynous systems (e.g. Webster 1992; Lindenfors et al. 2002) where larger body size enables males to defend their harems. The lack of a relationship between body size and arrival order likewise suggests that the latter is unrelated to male quality.

Early arrival may, however, have benefits that we could not examine. Most obvious of these is the main theoretical benefit of pioneering: that it reduces the amount of intraspecific larval competition for the offspring of pioneers. Numerous studies of *Ips* have demonstrated negative density effects on larval growth and survival (e.g. Zhang et al. 1992; Lawson et al. 1995; Robins and Reid 1997; Steed and Wagner 2004; Sallé and Raffa 2007). Our experiments only considered the initial stages of the attack, and not the ultimate outcome in terms of offspring survival. It is therefore possible that the offspring of early arriving males experience lower levels of larval competition, and as a result, have higher offspring

191 survivorship. This could result in early arriving males ultimately having a greater number of
192 offspring than late arriving males.

193 In summary, we have shown that pioneering behaviour in *Ips grandicollis* males is not
194 beneficial to them in terms of the number of females they can attract. Further research into the
195 effects of arrival order on larval survivorship would be needed to ascertain whether arrival
196 order is ultimately an important predictor of male reproductive success in *Ips grandicollis*.

197

198

199 **Acknowledgements**

200 Forestry SA allowed us access to the Wirrabara plantations. We thank Charlma Phillips
201 (Forestry SA) for making the arrangements for us to carry out the fieldwork, and the forester
202 Bruno Turrini for logistical assistance and his generous help and enthusiasm. Emile van
203 Lieshout and Kath McNamara provided comments on an earlier draft of this manuscript. This
204 research was funded by a Discovery Project Grant from the Australian Research Council, a
205 Canadian Natural Sciences and Research Council Graduate Scholarship, and travel grants
206 from the Department of Zoology, University of Melbourne and the Western Forest Genetics
207 Association (Canada and USA).

208

209

210 **References**

- 211 Amman GD, Bartos DL (1991) Mountain pine-beetle offspring characteristics associated with
212 females producing 1st and 2nd broods, male presence, and egg gallery length. *Env*
213 *Entomol* 20:1562-1567
- 214 Anderbrant O (1989) Reemergence and second brood in the bark beetle *Ips typographus*.
215 *Holarct Ecol* 12:494-500
- 216 Bensch S, Hasselquist D (1991) Territory infidelity in the polygynous great reed warbler
217 *Acrocephalus arundinaceus*: The effect of variation in territory attractiveness. *J Anim Ecol*
218 60:857-871
- 219 Byers JA (1989) Behavioral mechanisms involved in reducing competition in bark beetles.
220 *Holarct Ecol* 12:466-476
- 221 Candolin U, Voigt HR (2003) Size-dependent selection on arrival times in sticklebacks: Why
222 small males arrive first. *Evolution* 57:862-871
- 223 Clifton KE (1991) Subordinate group members act as food-finders within striped parrotfish
224 territories. *J Mar Biol Ecol* 145:141-148
- 225 Erlandsson A (1988) Food-sharing vs monopolising prey: A form of kleptoparasitism in *Velia*
226 *caprai* (Heteroptera). *Oikos* 53:203-206
- 227 Giraldeau LA, Beauchamp G (1999) Food exploitation: Searching for the optimal joining
228 policy. *Trends Ecol Evol* 14:102-106
- 229 Hansen EM, Bentz BJ (2003) Comparison of reproductive capacity among univoltine,
230 semivoltine, and re-emerged parent spruce beetles (Coleoptera: Scolytidae). *Can Entomol*
231 135:697-712
- 232 Heinsohn R, Packer C (1995) Complex cooperative strategies in group-territorial African
233 lions. *Science* 269:1260-1262

234 Kirkendall LR (1983) The evolution of mating systems in bark and ambrosia beetles
 235 (Coleoptera: Scolytidae and Platypodidae). Zool J Linn Soc 77:293-352

236 Lanier GN, Cameron EA (1969) Secondary sexual characteristics in the North American
 237 species of the genus *Ips* (Coleoptera: Scolytidae). Can Entomol 101:862-870

238 Latty TM, Reid ML (2009) First in line or first in time? Effects of settlement order and arrival
 239 date on reproduction of a group-living beetle, *Dendroctonus ponderosae*. J Anim Ecol
 240 78:549-555

241 Latty TM, Magrath MJL, Symonds MRE (2009) Harem size and oviposition behaviour in a
 242 polygynous bark beetle. Ecol Entomol 34:562-568

243 Lawson SA, Furuta K, Katagiri K (1995) Effect of tree host and beetle density on
 244 reproduction and survival of *Ips typographus japonicus* Nijima (Col., Scolytidae), in
 245 Hokkaido, Japan. J Appl Entomol 119:383-390

246 Lindenfors P, Tullberg BS, Biuw M (2002) Phylogenetic analyses of sexual selection and
 247 sexual size dimorphism in pinnipeds. Behav Ecol Sociobiol 52:188-193

248 Mangel M (1990) Resource divisibility, predation and group formation. Anim Behav
 249 39:1163-1172

250 Morgan FD (1967) *Ips grandicollis* in South Australia. Austral For 31:137-155

251 Pekar S, Hruskova M, Lubin Y (2005) Can solitary spiders (Araneae) cooperate in prey
 252 capture? J Anim Ecol 74:63-70

253 Pureswaran DS, Sullivan BT, Ayres MP (2006) Fitness consequences of pheromone
 254 production and host selection strategies in a tree-killing bark beetle (Coleoptera:
 255 Curculionidae: Scolytinae). Oecologia 148:720-728

256 Raffa KF (2001) Mixed messages across multiple trophic levels: the ecology of bark beetle
 257 chemical communication systems. Chemoecology 11:49-65

258 Raffa KF, Berryman AA (1983) The role of host plant resistance in the colonization behavior
 259 and ecology of bark beetles (Coleoptera: Scolytidae). *Ecol Monogr* 53:27-49
 260 Rasbash J, Steele F, Browne W, Prosser B (2004) A user's guide to MLwiN version 2.0.
 261 Institute of Education, London.
 262 Reid ML (1999) Monogamy in the bark beetle *Ips latidens*: ecological correlates of an
 263 unusual mating system. *Ecol Entomol* 24:89-94
 264 Robertson IC (1998) Paternal care enhances male reproductive success in pine engraver
 265 beetles. *Anim Behav* 56:595-602
 266 Robins GL, Reid ML (1997) Effects of density on the reproductive success of pine engravers
 267 - is aggregation in dead trees beneficial? *Ecol Entomol* 22:329-334
 268 Sallé A, Raffa KF (2007) Interactions among intraspecific competition, emergence patterns,
 269 and host selection behaviour in *Ips pini* (Coleoptera: Scolytinae). *Ecol Entomol* 32:162-
 270 171
 271 Schlyter F, Zhang QH (1996) Testing avian polygyny hypotheses in insects: harem size
 272 distribution and female egg gallery spacing in three *Ips* bark beetles. *Oikos* 76:57-69
 273 Smith RJ, Moore FR (2005) Arrival timing and seasonal reproductive performance in a long-
 274 distance migratory landbird. *Behav Ecol Sociobiol* 57:231-239
 275 Steed BF, Wagner MR (2004) Importance of log size on host selection and reproductive
 276 success from *Ips pini* (Coleoptera: Scolytidae) in ponderosa pine slash of northern Arizona
 277 and western Montana. *J Econ Entomol* 97:436-450
 278 Waterhouse DF, Sands DPA (2001) Classical biological control of arthropods in Australia.
 279 CSIRO Entomology, Canberra.
 280 Webster MS (1992) Sexual dimorphism, mating system and body size in new world
 281 blackbirds (Icterinae). *Evolution* 46:1621-1641

282 Wood SL (1982) The bark and ambrosia beetles of North and Central America (Coleoptera:
283 Scolytidae), a taxonomic monograph. Great Basin Nat Mem 6:1-1359
284 Zhang QH, Byers JA, Schlyter F (1992) Optimal attack density in the larch bark beetle, *Ips*
285 *cembrae* (Coleoptera: Scolytidae). J Appl Ecol 29:672-678
286

Figure Captions

Figure 1. Typical gallery system for *Ips grandicollis*. The male creates the central nuptial chamber, while individual females (here numbering five) bore out the separate galleries along which they lay their eggs. Individual egg notches along the galleries are clearly visible.

Figure 2. Model predictions of harem size in relation to the arrival order of males within log stack. Harem size is shown as both number of galleries and number of females. Male arrival order within each log stack is expressed relative to the arrival day of the first male to colonise the log stack. Predicted effect estimates are shown with 95% confidence interval. The relationship with number of females is statistically significant ($p = 0.02$), and is non-significant with number of galleries ($p = 0.43$) Only males arriving within 5 days of the first male were included ($n = 95$ males colonising 19 log stacks).

301 **Figure 1**



302
303
304

Figure 2

