



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Granger, C;Spittle, AJ;Walsh, J;Pyman, J;Anderson, PJ;Thompson, DK;Lee, KJ;Coleman, L;Dagia, C;Doyle, LW;Cheong, J

Title:

Histologic chorioamnionitis in preterm infants: Correlation with brain magnetic resonance imaging at term equivalent age

Date:

2018-02-15

Citation:

Granger, C., Spittle, A. J., Walsh, J., Pyman, J., Anderson, P. J., Thompson, D. K., Lee, K. J., Coleman, L., Dagia, C., Doyle, L. W. & Cheong, J. (2018). Histologic chorioamnionitis in preterm infants: Correlation with brain magnetic resonance imaging at term equivalent age. *BMC Pediatrics*, 18 (1), <https://doi.org/10.1186/s12887-018-1001-6>.

Persistent Link:

<https://hdl.handle.net/11343/256034>

License:

[CC BY](#)

RESEARCH ARTICLE

Open Access



# Histologic chorioamnionitis in preterm infants: correlation with brain magnetic resonance imaging at term equivalent age

Claire Granger<sup>1,14\*</sup>, Alicia J. Spittle<sup>2,3,4</sup>, Jennifer Walsh<sup>2,4,5,6</sup>, Jan Pyman<sup>7</sup>, Peter J. Anderson<sup>2,8</sup>, Deanne K. Thompson<sup>2,9,10,11</sup>, Katherine J. Lee<sup>9,12</sup>, Lee Coleman<sup>9,13</sup>, Charuta Dagia<sup>9,13</sup>, Lex W. Doyle<sup>2,4,5</sup> and Jeanie Cheong<sup>2,4,5</sup>

## Abstract

**Background:** To explore the associations between histologic chorioamnionitis with brain injury, maturation and size on magnetic resonance imaging (MRI) of preterm infants at term equivalent age.

**Methods:** Preterm infants (23–36 weeks' gestational age) were recruited into two longitudinal cohort studies. Presence or absence of chorioamnionitis was obtained from placental histology and clinical data were recorded. MRI at term-equivalent age was assessed for brain injury (intraventricular haemorrhage, cysts, signal abnormalities), maturation (degree of myelination, gyral maturation) and size of cerebral structures (metrics and brain segmentation). Histologic chorioamnionitis was assessed as a predictor of MRI variables using linear and logistic regression, with adjustment for confounding perinatal variables.

**Results:** Two hundred and twelve infants were included in this study, 47 (22%) of whom had histologic chorioamnionitis. Histologic chorioamnionitis was associated with higher odds of intraventricular haemorrhage (odds ratio [OR] (95% confidence interval [CI]) = 7.4 (2.4, 23.1)), less mature gyral maturation (OR (95% CI) = 2.0 (1.0, 3.8)) and larger brain volume (mean difference in cubic centimeter (95% CI) of 14.1 (1.9, 26.2)); but all relationships disappeared following adjustment for perinatal variables.

**Conclusion:** Histologic chorioamnionitis was not independently associated with IVH, less mature gyral maturation or brain volume at term-equivalent age in preterm infants.

**Keywords:** Chorioamnionitis, Preterm, Magnetic resonance imaging, Intraventricular haemorrhage, Brain volumes

## Background

The developing brain is vulnerable and is subject to a number of antenatal and perinatal stressors. Rapid brain growth and development occurs during the later stages of pregnancy, and any insult to the developing brain during this time may result in adverse neurodevelopmental sequelae [1].

Chorioamnionitis has been implicated as an antecedent of spontaneous preterm labour [2]. Chorioamnionitis refers to inflammation of the placental tissues [3],

and is thought to result from ascending infection into the uterus from the vagina. Pathogens are usually of low virulence and are difficult to culture using traditional methods [3]. Chorioamnionitis is defined as histologic, clinical, or both. The former is considered the “gold standard” [4], whereas the definition for clinical chorioamnionitis is inconsistent and has been shown to correlate poorly with histologic changes in the placenta [4]. Therefore the focus of this study was on histologic chorioamnionitis.

A number of perinatal variables are associated with chorioamnionitis, including earlier preterm delivery, prolonged rupture of membranes, low birthweight and maternal peripartum fever [1, 2, 5]. However, many studies of chorioamnionitis do not include infants born

\* Correspondence: [cmacgilchrist@doctors.org.uk](mailto:cmacgilchrist@doctors.org.uk)

<sup>1</sup>Department of Neonatal Medicine, Royal Children's Hospital, Melbourne, Australia

<sup>14</sup>Neonatal Intensive Care Unit, Princess Royal Maternity Hospital, 16 Alexandra Parade, Glasgow G31 2ER, Scotland

Full list of author information is available at the end of the article



moderate to late preterm (MLPT; 32–36 weeks' gestation). Understanding the effects of chorioamnionitis across the gestational age spectrum is important given the majority of preterm births are moderate or late preterm [6].

Inflammation, including chorioamnionitis, has been implicated in the pathogenesis of brain injury in preterm children. The potential mechanisms include pro-inflammatory cytokine release related to the fetal inflammatory response [7], which, in experimental models, has resulted in brain alterations consistent with impaired myelination, neuronal loss and white matter lesions [8].

Previous studies have attempted to establish a link between chorioamnionitis, and subsequent neurodevelopment and brain abnormalities. The associations between chorioamnionitis, and cystic periventricular leukomalacia and cerebral palsy are well documented. [9] However, the associations between chorioamnionitis and intraventricular haemorrhage, brain volumes and brain maturation are less consistent [10–12].

To address research gaps surrounding chorioamnionitis, we aimed to explore the associations of histologic chorioamnionitis with qualitative/quantitative brain MRI measures at term-equivalent age in preterm infants. We analysed conventional MR images using a modified version of a published reporting system, and we measured brain tissue compartment volumes using an automated method [13, 14].

We hypothesised that preterm infants with histological evidence of chorioamnionitis would have more brain injury at term-equivalent age as evidenced by white matter signal abnormalities, and smaller, less mature brains at term equivalent age compared with those without histologic chorioamnionitis.

## Methods

### Participants

Data were derived from two prospective longitudinal cohort studies of growth and development in preterm infants. Both studies comprised preterm infants born at or transferred to the Royal Women's Hospital neonatal unit, Melbourne, which is a tertiary neonatal unit located within a large perinatal centre. The first study comprised MLPT infants (born between 32<sup>+0</sup> and 36 weeks' gestation [13]), and the second comprised infants born at less than 30 weeks' gestation [15]. These cohorts were selected to answer different research questions; there was no deliberate attempt to exclude infants born at 30 or 31 weeks' gestation. Exclusion criteria did include genetic or congenital problems known to cause long-term developmental problems (Additional file 1).

Extensive perinatal data were collected by chart review. Birthweight was converted to Z-scores using LMS growth reference charts [14].

The Human Research Ethics Committees of the Royal Women's Hospital and the Royal Children's Hospital, Melbourne, approved both studies. Written informed consent was obtained from the parents of all participants. Details of the contact and tracing of the cohorts have previously been described [13, 15].

### Placental pathology

Where available, placentas were sent to the Pathology Department at the Royal Women's Hospital for microscopic/macrosopic analysis. Placental histopathology is performed as part of clinical practice for preterm births. Samples for histopathological examination were fixed in formalin, processed with tissue sections cut to 3–5 mm thickness and stained with haematoxylin and eosin. The placentas were then examined microscopically.

A single investigator (CG) reviewed the histopathology reports and classified the placentas into those with evidence of inflammation (maternal, fetal or both) and those without. Where there was uncertainty in the reports, an anatomical pathologist (JP) reviewed the reports.

### Magnetic resonance imaging (MRI) scan

An MRI brain scan was undertaken at term equivalent age using previously published techniques [13, 15]. MRI scans of the brain were performed using a 3 T Magnetom Trio Tim system (Siemens, Erlangen, Germany) during natural sleep. Conventional T1-weighted and T2-weighted axial images were used for the analysis of this study.

A modified MRI scoring system based on a published system [13, 14] was used to assess MRIs. All images were reviewed by two experienced neuroradiologists (LC and CD), who were unaware of perinatal history. All MRI assessors had undergone similar training. Data were recorded on presence of brain injury (white matter signal abnormalities, periventricular cysts, intraventricular haemorrhage), brain maturation (degree of myelination of the posterior limb of the internal capsule, gyral maturation), brain size (brain biparietal diameter, corpus callosal measures of the genu, midbody and splenium, deep nuclear grey area and transcerebellar diameter) and cerebrospinal fluid measures (lateral ventricular size, inter-hemispheric distance). A global MRI score was also calculated based on an established scoring system using measures of brain injury, maturation and size as listed above [13]. Inter- and intra-rater reliability for items of this scoring system have been published previously [13]. Inter-rater reliability for MRI measures on 10 scans were generally high for all reported MRI metrics (0.82–0.98), with the exception of smaller measures like the genu of the corpus callosum (0.79) and the body of the corpus callosum (0.61). Kappa values of inter-rater reliability for

categorical measures (degree of myelination of the posterior limb of the internal capsule and gyral maturation) were 1.0 [13].

### Infant brain segmentation

An automated Morphologically Adaptive Neonatal Tissue Segmentation (MANTiS) was used to classify the T2 structural MR image of the brain into the following structures: white matter, cortical grey matter, cerebrospinal fluid, subcortical gray matter (including deep nuclear gray matter, hippocampus and amygdala), brainstem, and cerebellum [16]. This technique extends the unified segmentation approach to tissue classification implemented in Statistical Parametric Mapping (SPM) software to infant brain scans. MANTiS combines unified segmentation, template adaptation via morphological segmentation tools and topological filtering. This technique reliably classifies the tissues even in the presence of brain abnormalities common in preterm infants, including enlarged ventricles and white matter signal intensity abnormalities. As the superior aspects of the extracerebral cerebrospinal fluid may not be included in all subjects, the volume calculations potentially underestimate the volumes of cerebrospinal fluid. Thus, we have not included cerebrospinal fluid volumes in the final analysis.

### Data analysis

Data were analysed using Stata version 14 (Stata Corporation, College Station, TX). Participant characteristics were compared between those included in this study (i.e with both placental histology data and MRI) with those not included using means and standard deviations (continuous variables) and proportions (categorical variables).

The relationships between histologic chorioamnionitis and MRI measures were explored using linear and logistic regression fitted to the MRI measures with histologic chorioamnionitis as a predictor. Models were fitted using generalised estimating equations with robust (sandwich) estimation of standard errors to allow for multiple births within a family. Results are presented firstly adjusted for gestation at MRI, and then adjusted for gestation at MRI as well as for gestation at birth, birthweight z-score, prolonged rupture of membranes, vaginal delivery, singleton birth, bronchopulmonary dysplasia (defined as oxygen requirement at 36 weeks' corrected age), necrotizing enterocolitis and sepsis (culture positive or raised inflammatory markers and/or white cell count resulting in at least 5 days' of antibiotics), all of which have been identified as potential confounders of this relationship in previous studies [2, 3, 5]. Results are presented as odds ratios (OR) and 95% confidence intervals (CI) for categorical variables, or mean differences and 95% CI for continuous brain measures.

### Results

Of the 349 preterm infants recruited into the 2 studies, 212 (61%) had both placental histology and MRI performed and are included in this study. Of these, 47 (22%) had histologic chorioamnionitis; 15 had maternal inflammatory response only, 31 had both maternal and fetal inflammatory response, and 1 had mild evidence of fetal inflammatory response only. All MRIs were suitable for qualitative assessment including brain metrics, but only 159 MRIs were suitable for volumetric analysis.

Compared with participants from the original birth cohorts who were not included in the current analysis, participants included in this study had lower mean birthweight z-scores ( $-0.5$  vs  $-0.2$ ) and were less likely to have received postnatal corticosteroids (3.2% vs 8.7%). The other characteristics were similar between those included in this study and those with incomplete data (data not shown). There were 159 participants with MRI suitable for segmentation. Participants with segmentable MRIs were similar in characteristics to those included in the study but who did not have MRI suitable for segmentation (data not shown).

Participant characteristics for the study sample ( $n = 212$ ) are summarised in Table 1. Participants with histologic chorioamnionitis were born at earlier gestational ages and had a lower birthweight z-score than those without histologic chorioamnionitis. Participants with histologic chorioamnionitis were more likely to have had ruptured membranes for  $> 24$  h, be singletons and be delivered by vaginal birth.

Inter-rater reliability for MRI measures on 10 scans were generally high for all reported MRI metrics (0.82–0.98), with the exception of smaller measures

**Table 1** Participant characteristics

	Histologic chorioamnionitis	
	Present ( $n = 47$ )	Absent ( $n = 165$ )
Gestation at birth in weeks, mean (SD)	29.8 (3.6)	32.2 (3.2)
Birthweight in grams, mean (SD)	1505 (728)	1686 (588)
Birthweight z-score, mean (SD)	$-0.1$ (0.9)	$-0.7$ (1.1)
Gestation at MRI in weeks, mean (SD)	42.1 (1.5)	42.0 (1.4)
Antenatal corticosteroids	39 (83.0)	122 (73.9)
Ruptured membranes $> 24$ h	22 (46.8)	13 (8.2)
Vaginal delivery	29 (61.7)	36 (21.8)
Male	24 (51.1)	85 (51.5)
Singleton	37 (78.7)	85 (51.5)
Postnatal corticosteroids	2 (4.9)	4 (2.7)
Bronchopulmonary dysplasia <sup>a</sup>	9 (19.2)	15 (9.1)

Data are  $n$  (%) unless otherwise specified, SD Standard deviation, PVL Periventricular leukomalacia, CI Confidence interval. <sup>a</sup>oxygen requirement at 36 weeks' corrected age

like the genu of the corpus callosum (0.79) and the body of the corpus callosum (0.61). Kappa values of inter-rater reliability for categorical measures (degree of myelination of the posterior limb of the internal capsule and gyral maturation) were 1.0.

#### **Associations of histologic chorioamnionitis with brain MRI measures (Table 2)**

Histologic chorioamnionitis was associated with increased odds of intraventricular haemorrhage and less mature gyral maturation. Both relationships disappeared following adjustment for confounders.

Histologic chorioamnionitis was also associated with larger total brain tissue volumes, white matter, cerebellar and brainstem volumes, although the mean differences in volumes were small (approximately 3–4%). Following adjustment for confounders, there was little evidence for relationships of histologic chorioamnionitis with brain segmented volumes.

There was little evidence for relationships of histologic chorioamnionitis with other parameters of brain injury, brain size or cerebrospinal fluid measures (using metric measurements).

#### **Discussion**

This prospective cohort study of preterm infants identified associations between histologic chorioamnionitis and higher odds of intraventricular haemorrhage, less mature gyral maturation and slightly larger brain size. However, these relationships did not persist following adjustment for perinatal confounding variables suggesting the chorioamnionitis is not an independent predictor of brain injury or size at term equivalent age.

The association of chorioamnionitis with intraventricular haemorrhage has not been consistently demonstrated in previous studies [17]. The current study reported an OR of 1.7 for intraventricular haemorrhage in infants who had histologic chorioamnionitis, which disappeared after adjusting for confounders. This suggests that the association of chorioamnionitis with intraventricular haemorrhage was mediated by other perinatal confounders, such as gestational age at birth and birthweight z-score.

There is, however, more evidence in the literature supporting the associations of histologic chorioamnionitis with cystic periventricular leukomalacia and cerebral palsy, as reported in a recent meta-analysis [18]. A more recent study by Leviton et al [19] reported increased odds for ventriculomegaly and echolucent lesions on cranial ultrasound with histologic chorioamnionitis in infants with gestational ages < 28 weeks. In the current study we were unable to establish associations between histologic chorioamnionitis and cystic periventricular leukomalacia as our prevalence of cystic periventricular

leukomalacia was low. Similarly, our ventricular measures were similar in infants with and without histologic chorioamnionitis. Of note, we measured ventricular size objectively, using MRI methods, rather than cranial ultrasound, as used in the study of Leviton et al. [19]

It is important to note that many of the early studies describing the association between chorioamnionitis (histologic or clinical) and brain injury used cranial ultrasound to report brain injury [20, 21]. Compared with MRI, cranial ultrasound is not sensitive at detecting non-cystic, multifocal white matter injury [1]. More recent studies using MRI to detect brain injury have found little evidence of associations for histologic chorioamnionitis with brain injury, development or growth at term equivalent age. Chau et al [22] studied a cohort of infants born between 24 and 32 weeks' gestation using brain MRI. Infants with histologic chorioamnionitis were not at increased risk of white matter injury, abnormal brain metabolism as measured using magnetic resonance spectroscopy, or abnormal microstructural development. Reiman et al. reported on the relationship between histologic chorioamnionitis and regional brain volumes at term age in a cohort of very low birthweight (< 1500 g) or very preterm (< 32 weeks' gestation) infants [23]. They were also unable to establish any relationships between histologic chorioamnionitis and brain volumes across several brain regions. A recently published study by Anblagan et al [24] looking at histological chorioamnionitis and altered brain microstructure as detected using diffusion MRI in infants < 32 weeks gestation found an association between the two. They concluded that their findings were suggestive of in utero brain inflammation in preterm infants exposed to histologic chorioamnionitis and it would be interesting to validate this finding in future analysis from our cohort. Although our study found a modest increase in brain volumes in those with histologic chorioamnionitis compared with those without, these differences were small and diminished after adjustment for confounding variables.

Data on the potential impact of chorioamnionitis on outcomes in MLPT infants is scarce. Lee et al [25] reported a higher rate of "adverse neonatal outcome" as defined by morbidities in the neonatal period in late preterm infants exposed to histologic chorioamnionitis compared with those who were not. Interestingly, there were no cases of brain injury on cranial ultrasound in either group. As such associations between histologic chorioamnionitis and brain injury, development and size in the MLPT population remain unknown.

Our study has several strengths. We used a prospective cohort of infants that included infants born < 30 weeks as well as those born MLPT among whom brain MRI data are scarce. In addition to advanced image analysis with brain segmentation, we used a simple, validated MRI

**Table 2** Association between histologic chorioamnionitis and MRI measures

Histologic chorioamnionitis	Present (n = 47)	Absent (n = 165)	OR <sup>a</sup> (95% CI) p value	Adjusted OR <sup>b</sup> (95% CI) p value
<b>(I) Brain injury</b>				
Abnormal white matter signal, n (%)	13 (7.9)	4 (8.5)	1.2 (0.4, 4.0) p = 0.78	0.7 (0.2, 2.3) p = 0.59
Cystic periventricular leukomalacia, n (%)	1 (2.1)	0 (0)	–	–
Intraventricular haemorrhage, n (%)	9 (19.2)	5 (3.0)	7.4 (2.4, 23.1) p = 0.001	1.8 (0.5, 7.0) p = 0.41
<b>(II) Brain maturation</b>				
Complete myelination of the PLIC, n (%)	41 (87.2)	139 (84.2)	1.5 (0.3, 3.4) p = 0.58	0.9 (0.2, 4.4) p = 0.99
Gyral maturation < 38 weeks, n (%)	29 (61.7)	71 (43.0)	2.0 (1.0, 3.8) p = 0.05	1.6 (0.7, 3.6) p = 0.24
<b>(III) Brain metrics (a) Brain measures</b>				
			Coefficient <sup>a</sup> (95% CI) p value	Adjusted coefficient <sup>b</sup> (95% CI) p value
Brain biparietal diameter (mm)	83.4 (4.1)	82.5 (4.5)	0.6 (−0.7, 1.9) p = 0.38	0.9 (−0.6, 2.3) p = 0.23
Corpus callosum – genu (mm)	3.9 (1.5)	4.3 (1.4)	−0.3 (−0.8, 0.1) p = 0.14	0.1 (−0.4, 0.5) p = 0.82
Corpus callosum – body (mm)	2.0 (0.6)	2.1 (0.5)	−0.1 (−0.3, 0.1) p = 0.45	0.1 (−0.2, 0.2) p = 0.72
Corpus callosum – splenium (mm)	3.4 (1.2)	3.8 (1.0)	−0.3 (−0.7, 0.1) p = 0.10	−0.1 (−0.4, 0.3) p = 0.82
Deep nuclear gray matter (mm <sup>2</sup> )	1250.4 (104.1)	1235.6 (106.2)	8.9 (−25.0, 42.8) p = 0.61	12.7 (−22.7, 48.1) p = 0.48
Transcerebellar diameter (mm)	55.1 (2.9)	55.5 (3.1)	−0.6 (−1.5, 0.3) p = 0.20	−0.6 (−1.6, 0.3) p = 0.21
<b>(b) Cerebrospinal fluid measures (mm)</b>				
Right lateral ventricle	6.3 (2.5)	5.9 (1.8)	0.4 (−0.3, 1.1) p = 0.30	0.4 (−0.2, 1.1) p = 0.15
Left lateral ventricle	5.9 (2.4)	5.6 (1.8)	0.3 (−0.5, 1.0) p = 0.47	0.2 (−0.4, 0.9) p = 0.44
Interhemispheric distance	3.1 (1.8)	2.8 (1.3)	0.2 (−0.4, 0.8) p = 0.49	−0.4 (−1.0, 0.1) p = 0.14
<b>(IV) Segmented brain volumes (cc)</b>				
	n = 37	n = 122		
Total brain tissue	414.4 (37.9)	399.7 (43.9)	14.1 (1.9, 26.2) p = 0.02	6.4 (−10.1, 22.9) p = 0.45
Cortical grey matter	181.4 (17.4)	176.0 (20.8)	5.0 (−0.7, 10.8) p = 0.09	1.4 (−6.4, 9.2) p = 0.73
White matter	165.3 (15.7)	158.2 (17.4)	7.0 (1.8, 12.1) p = 0.008	4.4 (−2.4, 11.1) p = 0.20
Subcortical grey matter	31.9 (2.8)	31.1 (3.1)	0.7 (−0.3, 1.7) p = 0.15	0.1 (−1.0, 1.3) p = 0.83
Cerebellum	29.4 (3.2)	28.1 (3.6)	0.9 (0.1, 1.7) p = 0.04	0.1 (−1.2, 1.3) p = 0.93
Brainstem	6.5 (0.5)	6.3 (0.6)	0.2 (0.1, 0.4) p = 0.005	0.1 (−0.1, 0.3) p = 0.39
<b>(V) Global MRI score</b>				
	3.6 (2.8)	2.7 (2.3)	0.8 (−0.1, 1.7) p = 0.06	0.1 (−0.7, 0.7) p = 0.98

Results presented as mean (standard deviation) unless specified. CI Confidence interval, OR Odds ratio, PLIC Posterior limb of the internal capsule, MRI Magnetic resonance imaging. <sup>a</sup>Adjusted for gestation at MRI; <sup>b</sup>Adjusted for gestation at MRI, gestation at birth, birthweight z-score, ruptured membranes > 24 h, vaginal birth, singleton, bronchopulmonary dysplasia (oxygen requirement at 36 weeks' corrected age), necrotizing enterocolitis and sepsis (culture positive or raised inflammatory markers and/or white cell count resulting in at least 5 days' of antibiotics)

assessment method [13, 15] suitable for clinicians caring for preterm babies. We used a definition of histologic chorioamnionitis rather than relying on reported clinical chorioamnionitis from obstetric histories.

We acknowledge the limitations as well. We used clinical reports for histology rather than a recognised classification system for case definition. However, these reports were done by a small number of experienced placental histologists at a single tertiary centre, and checked by a senior pathologist (JP) who is a co-author of the paper. In addition, not all participants in the original cohort studies had placental histology and MRI, and thus we may not have ascertained all cases of histologic chorioamnionitis within the cohort.

Further research directions include using advanced MRI techniques to probe white matter microstructure and brain metabolic profiles, which may be more sensitive than brain macrostructural characteristics in establishing associations between chorioamnionitis and brain injury in preterm infants. Correlation of neuroimaging findings with neurodevelopmental outcomes may provide further insight into the pathogenesis of neurodevelopmental deficits described in preterm children exposed to chorioamnionitis.

## Conclusions

In conclusion, this study found evidence that chorioamnionitis is associated with, but not an independent predictor of, IVH, less mature gyral maturation or brain volume. Other perinatal variables appear to contribute to the association between chorioamnionitis and preterm brain injury.

## Additional file

**Additional file 1:** Histologic chorioamnionitis in preterm infants: correlation with brain magnetic resonance imaging at term equivalent age. Supplemental files: Research questions. Copy of research questions used in study. (DOCX 53 kb)

## Abbreviations

CI: Confidence interval; IVH: Intraventricular haemorrhage; MANTIS: Morphologically Adaptive Neonatal Tissue Segmentation; MLPT: Moderate/late preterm; MRI: Magnetic resonance imaging; OR: Odds ratio; SPM: Statistical Parametric Mapping

## Acknowledgements

We thank Jan Pyman for her assistance in the pathology reports. We thank our study coordinators Emma McInnes and Marilyn Bear, our research nurses, and the families for their willingness to participate in this study. We also acknowledge the expertise and efforts of the MRI technologists at the Melbourne Children's MRI Centre, Royal Children's Hospital, Melbourne.

## Funding

Australian National Health and Medical Research Council (Project grant ID 1028822 & 1024516, Centre of Research Excellence ID 1060733, Early Career Fellowship ID 1053787 to JC, Senior Research Fellowship ID 1081288 to PA; Career Development Fellowship ID 1108714 to AS, 1085754 to DT & ID 1053609 to KJL), the Royal Children's Hospital Foundation, and the Victorian Government Operational Infrastructure Support Program.

## Availability of data and materials

Previously published in original publications of both trials [13, 15].

## Authors' contributions

CG, AS, LD and JC were involved in conception and design of the study, data analysis and interpretation, drafting and revising the article; and approval of the final manuscript as submitted. Other authors (JW, JP, PA, DT, KJL, RB, JC, LC and CD) were involved in data analysis and interpretation, revision of the article, and approval of the final manuscript. All authors have read and approved the final version of this manuscript.

## Ethics approval and consent to participate

The Human Research Ethics Committees of the Royal Women's Hospital and the Royal Children's Hospital, Melbourne, approved both studies. Written informed consent was obtained from the parents of all participants. Details of the contact and tracing of the cohorts have previously been described [13, 15].

## Consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Author details

<sup>1</sup>Department of Neonatal Medicine, Royal Children's Hospital, Melbourne, Australia. <sup>2</sup>Victorian Infant Brain Studies, Murdoch Childrens Research Institute, Melbourne, Australia. <sup>3</sup>Department of Physiotherapy, University of Melbourne, Melbourne, Australia. <sup>4</sup>Neonatal Services, Royal Women's Hospital, Melbourne, Australia. <sup>5</sup>Department of Obstetrics & Gynaecology, University of Melbourne, Melbourne, Australia. <sup>6</sup>Paediatric, Infant, Perinatal Emergency Retrieval, Royal Children's Hospital, Melbourne, Australia. <sup>7</sup>Department of Pathology, Royal Women's Hospital, Melbourne, Australia. <sup>8</sup>Monash Institute of Cognitive & Clinical Neurosciences, Monash University, Melbourne, Australia. <sup>9</sup>Department of Paediatrics, University of Melbourne, Melbourne, Australia. <sup>10</sup>Developmental Imaging, Murdoch Childrens Research Institute, Melbourne, Australia. <sup>11</sup>Florey Institute of Neuroscience and Mental Health, Melbourne, Australia. <sup>12</sup>Clinical Epidemiology and Biostatistics, Murdoch Childrens Research Institute, Melbourne, Australia. <sup>13</sup>Department of Radiology, Royal Children's Hospital, Melbourne, Australia. <sup>14</sup>Neonatal Intensive Care Unit, Princess Royal Maternity Hospital, 16 Alexandra Parade, Glasgow G31 2ER, Scotland.

Received: 2 January 2017 Accepted: 22 January 2018

Published online: 15 February 2018

## References

1. Khwaja O, Volpe JJ. Pathogenesis of cerebral white matter injury of prematurity. *Arch Dis Child Fetal Neonatal Ed.* 2008;93:153–61.
2. McElrath TF, Hecht JL, Dammann O, et al. Pregnancy disorders that lead to delivery before the 28th week of gestation: an epidemiologic approach to classification. *Am J Epidemiol.* 2008;168:980–9.
3. Hagberg H, Wennerholm UB, Sävman K. Sequelae of chorioamnionitis. *Curr Opin Infect Dis.* 2002;15(3):301–6.
4. Faye-Petersen OM. The placenta in preterm birth. *J Clin Pathol.* 2008;61(12):1261–75.
5. Grether JK, Nelson KB, Emery ES III, Cummins SK. Prenatal and perinatal factors and cerebral palsy in very low birth weight infants. *J Pediatr.* 1996;128:407–14.
6. Cheong JLY, Doyle LW. Increasing rates of prematurity and epidemiology of late preterm birth. *J Paediatr Child Health.* 2012;48(9):784–8.
7. Kallapur SG, Presicce P, Rueda CM, Jobe AH, Choungnet CA. Fetal immune response to chorioamnionitis. *Semin Reprod Med.* 2014;32(1):56–67.
8. Dammann O, Kuban KC, Leviton A. Perinatal infection, fetal inflammatory response, white matter damage, and cognitive limitations in children born preterm. *Ment Retard Dev Disabil Res Rev.* 2002;8(1):46–50.
9. Shatrow JG, Birch SC, Lam LT, et al. Chorioamnionitis and cerebral palsy: a meta-analysis. *Obstet Gynecol.* 2010;116(2 Pt 1):387–92.

10. Reiman M, Kujara H, Maunu J, Parkkola R, Rikalainen H, Lapinleimu H, Lehtonen L, Haataja L; PIPARI Study Group. Does placental inflammation relate to brain lesions and volume in preterm infants? *J Pediatr* 2008; 152(5): 642–647.e1–2.
11. Bersani I, Thomas W, Speer C. Chorioamnionitis – the good or the evil for neonatal outcome. *J Mat-Fetal and Neonatal Med.* 2012;25:12–6.
12. Chau V, Poskitt KJ, McFadden DE, Bowen-Roberts T, Synnes A, Brant R, Sargent MA, Soulikias W, Miller SP. The effect of chorioamnionitis on brain development and injury in premature newborns. *Ann Neurol.* 2009;66:155–64.
13. Walsh JM, Doyle LW, Anderson PJ, Lee KJ, JLY C. Moderate and late preterm birth: effect on brain size and maturation at term-equivalent age. *Radiology.* 2014;273(1).
14. Kidokoro H, Neil JJ, Inder TE. New MR Imaging assessment tool to define brain abnormalities in very preterm infants at term. *AJNR* 2013; 34:2208–2214.
15. Spittle AJ, Thompson DK, Brown NC, Treyvaud K, Cheong JL, Lee KJ, Pace CC, Olsen J, Allinson LG, Morgan AT, Seal M, Eeles A, Judd F, Doyle LW, Anderson PJ. Neurobehaviour between birth and 40 weeks' gestation in infants born <30 weeks' gestation and parental psychological wellbeing: predictors of brain development and child outcomes. *BMC Pediatr.* 2014;14:111.
16. Beare R, Chen J, Kelly C, Alexopoulos D, Smyser C, Rogers C, Loh WY, Matthews LG, Cheong J, Spittle A, Anderson PJ, Doyle L, Inder T, Seal M, Thompson D. Neonatal brain tissue classification with morphological adaptation and unified segmentation. *Front Neuroinform.* 2016;10:12.
17. Chau V, McFadden DE, Poskitt KJ, Miller SP. Chorioamnionitis in the pathogenesis of brain injury in preterm infants. *Clin Perinatol.* 2014 Mar; 41(1):83–103.
18. Wu YW. Systematic review of chorioamnionitis and cerebral palsy. *Ment Retard Dev Disabil Res Rev.* 2002;8(1):25–9.
19. Leviton A, Allred E, Kubam K, Hecht J, Onderdonk A, O'Shea M, Paneth N. Microbiologic and histologic characteristics of the extremely preterm infant's placenta predict white matter damage and later cerebral palsy. *The ELGAN Study Pediatr Res.* 2010;67(1):95–101.
20. Leviton A, Paneth N, Reuss ML, et al. Maternal infection, fetal inflammatory response, and brain damage in very low birth weight infants. *Developmental Epidemiology Network Investigators Pediatr Res.* 1999;46(5):566–75.
21. Vergani P, Patane L, Doria P, et al. Risk factors for neonatal intraventricular haemorrhage in spontaneous prematurity at 32 weeks gestation or less. *Placenta.* 2000;21(4):402–7.
22. Wu YW. Chorioamnionitis as a risk factor for cerebral palsy. *JAMA.* 2000;284: 1417–24.
23. Reiman M, Kujari H, Maunu J, et al. Does placental inflammation relate to brain lesions and volume in preterm infants? *J Pediatr.* 2008;152(5):642–7.
24. Anblagan D, Pataky R, Evans MJ, Telford EJ, Serag A, Sparrow S, Piyasena C, Semple SI. Association between preterm brain injury and exposure to chorioamnionitis during fetal life. In: Wilkinson AG, Bastin ME, Boardman JP, editors. *Scientific Reports*; 2016.
25. Lee SM, Park JW, Kim BJ, Park CW, Park JS, Jun JK, Yoon BH. Acute histologic chorioamnionitis is a risk factor for adverse neonatal outcome in late preterm birth after preterm premature rupture of membranes. *PLoS One.* 2013;8(12):e79941.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

