

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Deshpande, R; Vinuesa, R

Title:

Streamwise energy-transfer mechanisms in zero- and adverse-pressure-gradient turbulent boundary layers

Date:

2024-10

Citation:


Deshpande, R. & Vinuesa, R. (2024). Streamwise energy-transfer mechanisms in zero- and adverse-pressure-gradient turbulent boundary layers. *Journal of Fluid Mechanics*, 997, <https://doi.org/10.1017/jfm.2024.816>.

Persistent Link:

<https://hdl.handle.net/11343/353547>

License:

[CC BY](#)



Streamwise energy-transfer mechanisms in zero- and adverse-pressure-gradient turbulent boundary layers

Rahul Deshpande^{1,†} and Ricardo Vinuesa²

¹Department of Mechanical Engineering, University of Melbourne, Parkville, VIC 3010, Australia

²FLOW, Department of Engineering Mechanics, KTH Royal Institute of Technology, Stockholm 10044, Sweden

(Received 6 January 2024; revised 8 July 2024; accepted 22 August 2024)

The present study investigates streamwise ($\overline{u^2}$) energy-transfer mechanisms in the inner and outer regions of turbulent boundary layers (TBLs). Particular focus is placed on the $\overline{u^2}$ production, its inter-component and wall-normal transport as well as dissipation, all of which become statistically significant in the outer region with increasing friction Reynolds number (Re_τ). These properties are analysed using published data sets of zero, weak and moderately strong adverse-pressure-gradient (APG) TBLs across a decade of Re_τ , revealing similarity in energy-transfer pathways for all these TBLs. It is found that both the inner and outer peaks of $\overline{u^2}$ are always associated with local maxima in the $\overline{u^2}$ production and its inter-component transport, and the regions below/above each of these peaks are always dominated by wall-ward/away-from-wall transport of $\overline{u^2}$, thereby classifying the $\overline{u^2}$ profiles into four distinct regimes. This classification reveals existence of phenomenologically similar energy-transfer mechanisms in the ‘inner’ and ‘outer’ regions of moderately strong APG TBLs, which meet at an intermediate location coinciding with the minimum in $\overline{u^2}$ profiles. Conditional averaging suggests existence of similar phenomena even in low Re_τ canonical and/or weak APG TBLs, albeit with the outer-region mechanisms weaker than those in the inner region. This explains the absence of their $\overline{u^2}$ outer peak and the dominance of $\overline{u^2}$ wall-normal transport away from the wall, which potentially originates from the inner region. Given that the wall-ward/away-from-wall transport of $\overline{u^2}$ is governed by the Q_4 (sweeps)/ Q_2 (ejections) quadrants of the Reynolds shear stress, it is argued that the emergence of the $\overline{u^2}$ outer peak corresponds with the statistical dominance of Q_4 events in the outer region. Besides unravelling the dynamical significance of Q_2 and Q_4 events in the outer region of

† Email address for correspondence: raadeshpande@gmail.com

TBLs, the present analysis also proposes new phenomenological arguments for testing on canonical wall-turbulence data at very high Re_τ .

Key words: boundary layer structure, turbulent boundary layers

1. Introduction and motivation

The streamwise Reynolds stress ($\overline{u^2}$, where overbar denotes time or ensemble averaging) is one of the most widely analysed statistical quantities in the wall-bounded turbulence literature. This is not only because it is relatively convenient to measure, but also since it is the dominant contributor to the total turbulence fluctuations in any statistically two-dimensional (2-D) wall flow. The magnitude of $\overline{u^2}$, hence, is representative of the fundamental flow physics in different ‘regions’ of a turbulent boundary layer (TBL). For instance, $\overline{u^2}$ in the viscous (sublayer), inner and outer regions of the TBL is representative of the wall-shear stress fluctuations, dominance of the near-wall viscous-scaled streaks, and of the inertial large- and very-large-scale motions, respectively (Schoppa & Hussain 2002; Baars & Marusic 2020). In very low Re_τ zero-pressure-gradient (ZPG) TBLs, the near-wall streaks are the most prominent flow features (Kline *et al.* 1967), leading to the appearance of a local maximum in $\overline{u^2}$ profiles at the location where they are centred (referred to henceforth as the ‘inner peak’). Here, $Re_\tau = \bar{U}_\tau \delta_{99} / \nu$ is the friction Reynolds number defined based on the mean friction velocity (\bar{U}_τ), 99 % TBL thickness (δ_{99}) and kinematic viscosity (ν).

An increase in Re_τ leads to the growth in energy as well as broadening of hierarchy of the large- and very-large-scale motions, which originate in the outer region but have footprints extending down to the wall (Marusic, Mathis & Hutchins 2010; Baars & Marusic 2020). This results in the Re_τ growth of $\overline{u^2}$ magnitude across the TBL, but increasing most strongly in the outer region (Samie *et al.* 2018; Baars & Marusic 2020). A longstanding question (Monkewitz & Nagib 2015; Monkewitz 2022) has been whether this Re_τ growth of $\overline{u^2}$ leads to the emergence of another local maximum farther away from the wall (referred henceforth as the ‘outer peak’), signifying dominance of the inertial large scales. Investigating this, however, requires Re_τ increasing well beyond 10^4 so that the integrated large-scale contributions are distinguishable from those of the small scales (i.e. sufficient scale separation; Alfredsson, Segalini & Örlü 2011). The fact that very few experimental facilities can conduct accurate, well-resolved measurements in such high Re_τ canonical flows (Fiorini *et al.* 2017; Samie *et al.* 2018) has resulted in the discussion around the $\overline{u^2}$ outer peak still remaining an open question, at least for a ZPG TBL.

On the other hand, imposition of an adverse pressure gradient (APG) on a TBL is well known to lead to a discernible outer peak in $\overline{u^2}$, even at low to moderate Re_τ (Pozuelo *et al.* 2022). Previous studies at low Re_τ have found this peak to be a consequence of the enhanced mean shear in an APG TBL, which significantly energizes the large-scale motions in the outer region via production of turbulent kinetic energy (TKE) (Gungor *et al.* 2016; Kitsios *et al.* 2017). While this changes the spatial organization and geometry of the coherent structures in low-to-moderately strong APG TBLs (compared with ZPG TBLs; Lee & Sung 2009), recent investigations have suggested that the dominant coherent structures (Lee 2017) and their inherent energy-transfer mechanisms (Gungor, Maciel & Gungor 2022) remain the same as in ZPG TBLs. Gungor *et al.* (2022) found that imposition of a weak-to-moderately strong pressure gradient predominantly changes only

the intensity of the energy-transfer mechanisms in the inner and outer regions, relative to ZPG TBLs. Notably, even differences in the wall-attached/detached nature of the dynamically significant motions, between ZPG and APG TBLs, were found not to affect their energy-transfer mechanisms (see also the discussion by Dong *et al.* 2017). Inspired by these observations, the present study aims to understand the energy-transfer mechanisms associated with the $\overline{u^2}$ outer peak in weak-to-moderately strong APG TBLs, and assess whether similar mechanisms could be extended to the outer region of high Re_τ ZPG TBLs. This approach can potentially offer new flow-physics-based arguments for future studies supporting/against the emergence of the $\overline{u^2}$ outer peak in ZPG TBLs (after availability of accurate, well-resolved measurements). Looking from a broader perspective, the present approach could potentially provide an alternative pathway to understand the energy dynamics of very high Re_τ TBLs (Vinuesa, Örlü & Schlatter 2017), which are omnipresent in engineering applications but hard to measure.

1.1. Streamwise Reynolds-stress transport

While the standalone analysis of $\overline{u^2}$ has been extensively reported in the literature, its correlation with the individual terms in the transport equation of the streamwise TKE has been relatively limited owing to measurement challenges. For a TBL with an imposed streamwise pressure gradient, the transport equation of streamwise TKE can be expressed as (Gungor *et al.* 2022; Pozuelo *et al.* 2022):

$$\frac{\partial}{\partial t} \overline{u^2} = \mathcal{P}^u + \mathcal{E}^u + \mathcal{D}^u + \mathcal{T}^u - \Pi^u + \Pi_t^u + \mathcal{C}^u, \quad (1.1)$$

where the production term $\mathcal{P}^u = -2\overline{uw}(\partial\overline{U}/\partial z) - 2\overline{u^2}(\partial\overline{U}/\partial x)$, the viscous dissipation $\mathcal{E}^u = -2\nu\sum_{j=1}^3\overline{(\partial u/\partial x_j)^2}$, the viscous diffusion $\mathcal{D}^u = \nu\sum_{j=1}^3(\partial^2\overline{u^2}/\partial x_j^2)$, the pressure strain $\Pi^u = -(2/\rho)\overline{p}(\partial\overline{u}/\partial x)$, the pressure transport $\Pi_t^u = -(2/\rho)(\partial\overline{pu}/\partial x)$, the turbulent transport $\mathcal{T}^u = -(\partial\overline{u^2w}/\partial z)$ and the convection $\mathcal{C}^u = -\overline{U}\partial\overline{u^2}/\partial x - \overline{W}\partial\overline{u^2}/\partial z$. Here, u, v and w represent turbulence fluctuations in the streamwise (x), spanwise (y) and wall-normal (z) directions, respectively. Capital letters (U, V, W) represent instantaneous flow properties (i.e. without subtracting the mean), while overbars (e.g. $\overline{U}, \overline{u^2}$) indicate time or ensemble averaging. Chen & Sreenivasan (2021) have previously considered the energy balance in (1.1) in the very near-wall region of a canonical wall-bounded flow, where $\mathcal{D}^u \approx \mathcal{E}^u$, to motivate the validity of the wall scaling of the $\overline{u^2}$ inner peak magnitude (found at z_{IP}) at $Re_\tau \rightarrow \infty$. In the same region, Tang & Antonia (2023) proposed a dissipation scaling for $\overline{u^2}$ profiles (i.e. based on \mathcal{E}^j and ν), which was found to extend across a larger wall-normal range with increasing Re_τ . Monkewitz & Nagib (2015) estimated the scaling for the $\overline{u^2}$ outer peak (at z_{OP}) in a very high Re_τ canonical flow, based on a simplified balance between the net turbulence production and dissipation, yielding $z_{OP}^+ \approx (0.06\mathcal{E}^j)^{-1}$, where $j = 1-3$. However, it is now well known that the imbalance between \mathcal{P}^j and \mathcal{E}^j in the outer region increases with Re_τ , which is accounted for by the wall-normal transport, \mathcal{T}^j (Cho, Hwang & Choi 2018; Lee & Moser 2019). Considering that our main goal is to understand the high Re_τ dynamics in the outer region, the present study focuses on correlating the wall-normal profiles of relevant Re_τ -dependent terms in (1.1) with $\overline{u^2}$.

In a canonical wall flow, the increase in \mathcal{P}^u (i.e. the TKE source term) is associated with the broadening and energization of the inertial eddy hierarchy with increasing

Re_τ (Marusic *et al.* 2010; Baidya *et al.* 2017), which is accompanied by an increase in energy dissipation (\mathcal{E}^u , i.e. the sink term) that occurs predominantly via the small dissipative scales (Lee & Moser 2019). In terms of variations across the wall-normal direction, the energy pathway from the source to sink comprises the pressure strain term Π^u , representing transfer from $\overline{u^2}$ to $\overline{v^2}$ and $\overline{w^2}$, and the wall-normal transport term $\mathcal{T}^u = \partial(\overline{u^2 w})/\partial z$, representing diffusion of $\overline{u^2}$ from the outer region. Both of these are known to increase with Re_τ alongside \mathcal{P}^u and \mathcal{E}^u (Lee & Moser 2019). Although $\overline{u^2}$ is dependent on the net balance of all the terms in (1.1), the above discussion points towards an increasing influence of \mathcal{P}^u , \mathcal{T}^u , Π^u and \mathcal{E}^u on $\overline{u^2}$ with increasing Re_τ , especially in the outer region. Looking further into the definition of \mathcal{T}^u , the triple product represents a net flux of $\overline{u^2}$ -energy transported in the wall-normal direction, with $\overline{u^2 w} < 0$ and $\overline{u^2 w} > 0$ respectively indicating flux towards or away from the wall (Lee & Moser 2019). In the case of relatively low Re_τ canonical flows, Lee & Moser (2019) noted that the mean energy transport in the outer region is directed away from the wall (i.e. $\overline{u^2 w} > 0$). Based on their Re_τ trends, however, Lee & Moser (2019) hypothesized that a significant increase in \mathcal{P}^u at high Re_τ would be accompanied with a mean energy flux from the intermediate region towards the wall (i.e. $\overline{u^2 w} < 0$). This suggests increasing dominance of the streamwise momentum-carrying motions moving towards the wall (i.e. Q_4 events: $u > 0$, $w < 0$; Deshpande & Marusic 2021; Cremades *et al.* 2024), in very high Re_τ flows where a $\overline{u^2}$ outer peak may emerge.

Interestingly, analogous trends have been noted in the literature for the case of moderately strong APG TBLs, where a $\overline{u^2}$ outer peak is clearly observed (Skaare & Krogstad 1994; Lee 2017). Both these studies found the outer-peak region in \mathcal{P}^u and $\overline{u^2}$ to coincide with dominance of $\overline{u^2}$ energy flux towards the wall (i.e. $\overline{u^2 w} < 0$), with the latter driven by significantly energized Q_4 events. Lee (2017) used conditional analysis to confirm that intensification of these Q_4 events is governed by the large-scale roll modes in the APG TBL outer region. Besides these speculated similarities in \mathcal{T}^u , Gungor *et al.* (2022) recently conducted 2-D spectral analysis of \mathcal{P}^u and Π^u to demonstrate similarities in these transfer mechanisms for canonical flows and moderately strong APG TBLs. Hence, although the dominant mechanisms behind enhancement of the TKE source term (\mathcal{P}^u) are different for APG and ZPG TBLs (at least at low Re_τ), the energy-transfer pathways between \mathcal{P}^u , Π^u and \mathcal{T}^u , on energization of the APG TBL outer region, seem analogous to that hypothesized for very high Re_τ ZPG TBLs. This reaffirms the motivation for the present investigation.

1.2. Present contributions

The present study analyses the energy-transfer mechanisms behind the $\overline{u^2}$ outer peak using published ZPG and APG TBL data sets at various matched Re_τ (described in § 2). The first analysis in § 3 compares all the dominant terms of $\overline{u^2}$ transport with respect to increasing Re_τ , i.e. \mathcal{P}^u , \mathcal{E}^u , \mathcal{T}^u and Π^u . It confirms the similarity in energy-transfer pathways for ZPG and APG TBLs, across a larger Re_τ range than that considered by Gungor *et al.* (2022). This result motivates § 3.1, which reveals the increasing dominance of \overline{uw} in TKE production with increasing Re_τ , for both ZPG and weak-to-moderately strong APG TBLs. Given uw influences both \mathcal{P}^u and \mathcal{T}^u , a comprehensive analysis of the uw contributions from the individual Q_2 (ejections; $u < 0$, $w > 0$) and Q_4 (sweeps; $u > 0$, $w < 0$) events, relative to the streamwise TKE, is discussed in § 4. These two quadrants are the most

dominant out of the four possible quadrants of the Reynolds shear stress in a wall-bounded flow (Q_i , with $i = 1-4$; Wallace 2016). While both Q_2 and Q_4 events contribute positively to the TKE production, the opposite signs in w fluctuations imply that they are individually responsible (Nagano & Tagawa 1990) for $\overline{u^2}$ energy flux towards (Q_4 , $\overline{u^2 w} < 0$) and away from the wall (Q_2 , $\overline{u^2 w} > 0$). Hence, investigating variation in their contributions can potentially explain changes in the direction of $\overline{u^2}$ flux from the outer region, which is noted on increasing APG strength (Skaare & Krogstad 1994) or increasing Re_τ (Lee & Moser 2019; Deshpande & Marusic 2021; Cremades *et al.* 2024). To the authors' knowledge, the present study is the first attempt of using the wall-normal profiles of uw -quadrant contributions to understand the energy dynamics around the $\overline{u^2}$ peaks (Wallace 2016).

2. Experimental and numerical data sets

We consider four previously published experimental and numerical data sets for the present analysis, covering both ZPG and APG TBLs across $290 \lesssim Re_\tau \lesssim 4000$. Three of these data sets are from the high-resolution large-eddy simulations (LES) of a ZPG TBL (Eitel-Amor *et al.* 2014) and APG TBLs (Bobke *et al.* 2017; Pozuelo *et al.* 2022). The fourth is an experimental data set acquired via planar particle image velocimetry (PIV; Sanmiguel Vila *et al.* 2017) in a APG TBL. The high-resolution-LES data considered here has a resolution two times coarser than a typical DNS in the streamwise and spanwise directions. A small body force is added to account for the unresolved TKE dissipation (around 10% of the total), results from which have been shown to agree extremely well with DNS in both canonical (Eitel-Amor *et al.* 2014) and more complex (Negi *et al.* 2018) turbulent flows. The only effect of the lower resolution is a small attenuation of the near-wall $\overline{u^2}$ peak, which however does not influence the present discussion. Our analysis only depends on identification of the $\overline{u^2}$ peak and its z location, which is not affected by use of high-resolution LES (as compared with DNS).

For all the data sets, the boundary-layer thickness (δ_{99}) is estimated via the diagnostic-plot method (see Vinuesa *et al.* 2016). The same definition of δ_{99} has also been used to calculate the Clauser-pressure-gradient parameter, $\beta = (\delta^*/\rho \overline{U_\tau^2})(d\bar{P}/dx)$ (Bobke *et al.* 2017), where $\delta^* = \int_0^{\delta_{99}} (1 - \bar{U}(z)/\overline{U_e}) dz$ is the displacement thickness, $\overline{U_e} = \bar{U}(z = \delta_{99})$, i.e. the edge velocity, ρ is the fluid density and $d\bar{P}/dx$ is the mean streamwise pressure gradient at the x location where β is estimated. All the data sets have been tabulated in table 1, while their β distribution as a function of Re_τ (across the experimental/LES domains) is shown in figure 1 to provide a general overview.

As can be noted from figure 1(a), a unique aspect of the LES data sets of APG TBL is their near-constant β distribution across a substantial portion of the computational domain. In the case of Bobke *et al.* (2017), $\beta \approx 1$ for $300 \lesssim Re_\tau \lesssim 750$, while in the case of Pozuelo *et al.* (2022), $\beta \approx 1.4$ for $700 \lesssim Re_\tau \lesssim 2000$. Subsequently, when combined together, they facilitate investigation of the variation of the flow physics in a near-equilibrium TBL across a broad Re_τ range. Despite the near-constant β , however, the statistics would also be dependent on the unique upstream pressure-gradient history, across $Re_\tau \lesssim 300$ (Bobke *et al.* 2017) and $Re_\tau \lesssim 700$ (Pozuelo *et al.* 2022). Thus, in order to demonstrate that the present conclusions are independent of the upstream history effects, this study analyses turbulence statistics in both nominally constant β as well as varying β sections of the computational domain.

Wall-normal variation of the turbulence statistics is considered at selected x locations of the computational domain, each corresponding to a unique Re_τ , which have been indicated

Data set	TBL	Re_τ	β	No. of cases	Colour shading	Resolution ($\Delta y_m^+ \times \Delta z_m^+$)	Reference
LES	ZPG	290–2000	0	7	Black	10.9×19.6	Eitel-Amor, Örlü & Schlatter (2014)
LES	APG	300–750	0.85–1.0	3	Violet	9.7×8.4	Bobke <i>et al.</i> (2017)
LES	APG	450–2000	0.85–1.6	6	Red	12.5×30.1	Pozuelo <i>et al.</i> (2022)
Exp.	APG	1900–4000	1.3	2	Gold	20×100	Sanmiguel Vila <i>et al.</i> (2017)
		1000–4000	2.4	3	Magenta	20×100	Sanmiguel Vila <i>et al.</i> (2017)

Table 1. Table summarizing the parametric space associated with various published experimental (Exp.) and LES data sets analysed in this study. Definitions/terminologies have been provided in either § 1 or 2. Throughout this paper, light to dark colour shading indicates increasing Re_τ . The pressure-gradient history associated with each case can be visualized in figure 1. In case of LES, Δy_m and Δz_m refer to the numerical grid resolution in the free stream, while it refers to the interrogation window size in case of the ‘Exp.’ data sets.




Figure 1. The β distribution as a function of Re_τ corresponding to the (a) high-resolution LES and (b) experimental data sets analysed in the present study and documented in table 1. Lines represent pressure-gradient histories across the (a) computational domains and (b) wind tunnels associated with these data sets, while symbols represent selected cases for which the turbulence statistics are analysed. Light to dark shading in symbol colours represents increasing Re_τ , with cases at matched Re_τ (across different data sets) represented by the same symbols. Green background shading is used to highlight cases where an outer peak is noted in the $\overline{u^2}$ profiles.

with symbols in figure 1(a). All these selected points are sufficiently far away from the start of the computational domain (refer to Bobke *et al.* 2017; Pozuelo *et al.* 2022). A reference data set for ZPG TBL (Eitel-Amor *et al.* 2014), spanning a Re_τ range similar to that covered by the two APG TBL LES data sets, is also considered. Turbulence statistics are analysed at seven streamwise locations within the ZPG domain, such that their corresponding Re_τ match reasonably well with the selected three and six cases from the APG domains of Bobke *et al.* (2017) and Pozuelo *et al.* (2022), respectively. For convenience in referencing, these cases at matched Re_τ will be represented by the same symbols in the forthcoming analysis (figure 1a). The turbulence statistics presented here are computed by averaging across the time series saved at these selected x positions, as well as along the homogeneous spanwise direction. The computational grid resolution (table 1) has been demonstrated in the original references to be fine enough to ensure that the peaks/maxima in the $\overline{u^2}$ profiles are physical, and not an artefact of spatial-resolution issues (Samie *et al.* 2018). It is worth noting here that the (Δy_m^+ , Δz_m^+) values in table 1 refer to the maximum spanwise and

wall-normal grid spacing. The grid is very fine close to the wall, with seven points below $z^+ < 10$.

The experimental data set for APG TBL was acquired via two-component 2-D PIV in the streamwise–wall-normal (x – z) plane at the Minimum Turbulence Level facility at KTH, Stockholm (Sanmiguel Vila *et al.* 2017). The pressure gradient was imposed by hanging wall inserts in a converging–diverging geometry from the tunnel roof. The converging section initially led to the imposition of a favourable pressure gradient, followed by a region of nominal ZPG (in the straight section) and finally a region of APG (in the diverging section), making its upstream history different than that noted for the high-resolution LES cases. The TBL was measured over a smooth aluminium plate, which was suspended ~ 15 cm from the tunnel floor and had its leading edge (i.e. $x \approx 0$) upstream of the ZPG region. Overall, five different β distributions with unique upstream histories (figure 1*b*) were obtained by using two different roof geometries (indicated by different colours) and three different upstream free-stream speeds of $U_{ref} \sim 6, 12$ and 30 m s^{-1} (indicated by colour shading intensity). Here, U_{ref} is the free-stream speed at $x \approx 0$. Two-dimensional PIV was conducted sufficiently downstream (~ 4.5 m) from the leading edge of the plate, such that a moderately strong APG ($1.3 \lesssim \beta \lesssim 2.4$) was imposed on the TBL. Indeed, the spatial resolution of these PIV experiments is much lower than those of the other simulations considered (table 1). However, the $\overline{u^2}$ and \overline{uw} profiles obtained from these measurements have been demonstrated by Sanmiguel Vila *et al.* (2017) to be sufficiently well resolved in the outer region ($z^+ \gtrsim 100$) of the APG TBL, and we will limit our investigation of this data set only to this region. In their original paper, the authors also employed two post-processing approaches on the same PIV data set to further improve the accuracy of the near-wall-turbulence statistics, which however are not considered here.

3. Similarity in energy-transfer pathways for ZPG and APG TBLs

We begin our analysis by connecting the $\overline{u^2}$ profiles from all the data sets (figure 2*a–d*), with their corresponding terms in the streamwise Reynolds-stress-transport equation: production (\mathcal{P}^{u+} ; figure 2*m–p*), pressure–strain (Π^{u+} ; figure 2*q–s*), viscous dissipation (\mathcal{E}^{u+} ; figure 2*t–v*) and wall-normal turbulent transport (\mathcal{T}^{u+} ; figure 2*w–y*). The intention is to compare the energy-transfer pathways, starting from the TKE source \mathcal{P}^u , for both ZPG and APG TBLs. Here, we only consider the dominant term for streamwise TKE production, $\mathcal{P}^u \approx -2\overline{uw}(\partial\overline{U}/\partial z)$, and neglect the other term included previously in (1.1), $-2\overline{u^2}(\partial\overline{U}/\partial x)$, considering its negligible contribution (Pozuelo *et al.* 2022). While \mathcal{P}^{u+} could be obtained from the PIV data set of the APG TBL (Sanmiguel Vila *et al.* 2017), other properties such as \mathcal{E}^{u+} , \mathcal{T}^{u+} and Π^{u+} were either not possible to estimate or were erroneous owing to their sensitivity to the wall-normal gradient. In figure 2 the locations of the inner (z_{IP}^+) and outer peaks (z_{OP}^+) of $\overline{u^2}$ were identified by estimating the gradients of the profiles, i.e. $d\overline{u^2}/dz = 0$ and $d^2\overline{u^2}/dz^2 < 0$, and the locations of both the peaks are indicated by vertical green lines across all the plots in the figure.

The inner peak of $\overline{u^2}$ for both ZPG and APG TBLs is close to the local maximum of \mathcal{P}^{u+} observed in the near-wall region. This close correspondence is consistent with previous observations (Kline *et al.* 1967; Lee & Moser 2019) and is associated with the intense TKE production by the streaks related to the near-wall cycle (Hamilton, Kim & Waleffe 1995). The significant production of TKE at z_{IP}^+ is connected with a local maximum of $-\Pi^{u+}$ and a local minima in \mathcal{T}^{u+} , for both ZPG and APG TBLs, suggesting similar




Figure 2. Wall-normal profiles for (a–d) u^2+ , (e–h) $\partial U^+/\partial z^+$, (i–l) $-\bar{u}w^+$, (m–p) $\mathcal{P}^{u+} \approx -2\bar{u}w^+\partial U^+/\partial z^+$, (q–s) $-\Pi^{u+}$, (t–v) $-E^{u+}$ and (w–y) T^{u+} for ZPG (a,e,i,m,q,t,w) and APG TBL cases (b,f,j,n,r,u,x,c,g,k,o,s,v,y,d,h,l,p). Symbols/colours correspond to the various cases defined in figure 1, with light to dark shading indicating increasing Re_τ . Note the inconsistent axis ranges.

energy-transfer pathways in their respective inner regions (noted previously for \mathcal{P}^{u+} and $-\Pi^{u+}$ by Gungor *et al.* 2022). Here, the loss in $\overline{u^2}$ via Π^{u+} acts as the gain term for the lateral velocity components $\overline{v^2}$ and $\overline{w^2}$ via Π^{v+} and Π^{w+} , respectively. Similarly, a negative value at the minima of \mathcal{T}^{u+} represents wall-normal transport of $\overline{u^2}$ from z_{IP}^+ to other regions of the TBL (Lee & Moser 2019).

While the \mathcal{P}^{u+} profiles for ZPG TBL seem to collapse reasonably well in inner scaling for a major portion of the TBL, that is not the case in the far outer region (figure 2*m*). In fact, the bulk TKE production (i.e. $z^+\mathcal{P}^{u+}$; figure 3*a*) in the outer region increases with Re_τ , thereby driving the Re_τ growth of $\overline{u^2}$ evident in figure 2(*a*). This increase in \mathcal{P}^{u+} leads to an increase in magnitude of \mathcal{T}^{u+} , as well as enhanced losses in $\overline{u^2}$ via \mathcal{E}^{u+} and Π^{u+} with Re_τ (Lee & Moser 2019), which can also be noted from figure 3(*d,g,j*) depicting transport terms premultiplied by z^+ . While such a qualitative comparison of the premultiplied transport terms is reasonable, due precaution is required when interpreting them quantitatively. Premultiplication with z^+ artificially dampens the energy in the inner region while exaggerating that in the outer region, an example of which is evident from the ‘outer peak’ in $z^+\mathcal{P}^{u+}$ (figure 3*a*) that however does not exist in figure 2(*m*).

The outer region of the APG TBLs exhibit significantly enhanced \mathcal{P}^{u+} even at low Re_τ (figure 2*n-o*), albeit due to a different mechanism than for high Re_τ ZPG TBLs. This enhanced \mathcal{P}^{u+} is again accompanied with significantly high magnitudes of \mathcal{T}^{u+} as well as greater losses in $\overline{u^2}$ via \mathcal{E}^{u+} and Π^{u+} , when compared with ZPG TBLs at matched Re_τ . Qualitatively, the energization of the outer region in the APG TBLs is consistent with that exhibited by ZPG TBLs on increasing Re_τ , except the fact that statistics for moderately strong APG TBLs exhibit a distinct maximum (owing to greater energization of the outer region). Notably, this similarity in the energy-transfer mechanisms of the outer region has also been discussed previously by Gungor *et al.* (2022), but is confirmed for all major Reynolds-stress-transport terms in figures 2 and 3. Harun *et al.* (2013), Pozuelo *et al.* (2022) and Gungor *et al.* (2022) have also confirmed that this energy enhancement is not an artefact of the use of inner scaling (i.e. with \bar{U}_τ), but is physical and associated with energization of the outer region on imposition of the APG.

Interestingly, analogous to the energy-transfer mechanisms noted near the inner peak, there is a local maximum in \mathcal{P}^{u+} as well as $-\Pi^{u+}$ at the $\overline{u^2}$ outer peak location, z_{OP}^+ (see vertical green lines in figure 2*b-d,n-p,r,s*). The outer peak is also close to the minima in \mathcal{T}^{u+} that is again negative (figure 2*x,y*), while no maxima is noted in $-\mathcal{E}^{u+}$ (figure 2*u,v*). On the other hand, there is no local maximum in \mathcal{P}^{u+} and $-\Pi^{u+}$ for scenarios without an outer peak of $\overline{u^2}$ (see dash-dotted profiles), reaffirming the one-to-one connection between these statistics. Hence, similar to the scenario noted at z_{IP}^+ , the significant TKE production at z_{OP}^+ is also accompanied by enhanced energy transfer from $\overline{u^2}$ to $\overline{v^2}$ and $\overline{w^2}$ (via the pressure-strain mechanism), and the wall-normal transport of $\overline{u^2}$ to other regions of the APG TBL. Future work would focus on investigating these statistics for very high Re_τ canonical flow data (where a $\overline{u^2}$ outer peak may emerge), to check for quantitative similarity with moderately strong APG TBLs (having significantly enhanced outer-region energy). However, considering the paucity of such data at the present time, we employ conditional analysis on relatively low Re_τ ZPG TBL data in § 4.2, to unravel the qualitative similarity between ZPG and APG TBLs. Furthermore, we present elaborate analyses on the similarity in the wall-normal energy transport, between the TBL inner and outer regions, in §§ 4.1 and 4.2. However, first we focus on understanding the Re_τ variations




Figure 3. Wall-normal profiles for $\overline{u^2}$ -transport terms premultiplied by z^+ : (a–c) $z^+ \mathcal{P}^{u+} \approx -2z^+ \overline{uw}^+ \partial U^+ / \partial z^+$, (d–f) $-z^+ \Pi^{u+}$, (g–i) $-z^+ \mathcal{E}^{u+}$ and (j–l) $z^+ \mathcal{T}^{u+}$ for ZPG (a,d,g,j) and APG TBL cases. Symbols/colours correspond to the various cases defined in figure 1, with light to dark shading indicating increasing Re_τ . Note the inconsistent range of ordinate values.

in \mathcal{P}^{u+} for ZPG and APG TBLs in § 3.1, given it is the source of the streamwise TKE that leads to both the $\overline{u^2}$ peaks (Harun *et al.* 2013; Gungor *et al.* 2016; Kitsios *et al.* 2017).

3.1. Assessing the enhancement of TKE production with Reynolds number

Considering its mathematical definition, variations in \mathcal{P}^{u+} can be caused by either changes in the mean shear (figure 2e–h), magnitude of the Reynolds shear stress (\overline{uw} ; figure 2i–l) or a combination of both. The mean shear is highest in the near-wall region of both ZPG and weak-to-moderate APG TBLs, and hence, is predominantly responsible for their inner peak in \mathcal{P}^{u+} and $\overline{u^2}$ (Kline *et al.* 1967). However, the mean shear is significantly lower in the outer region of the ZPG TBLs and reduces gradually with increasing Re_τ (figure 2e). The increase in bulk production with Re_τ , hence, can be exclusively associated with the gradual increase and broadening of the quasi-constant \overline{uw} plateau (figure 2i) predominantly around the overlap region of the ZPG TBL (i.e. a subset of the outer region). This has been extensively demonstrated via both experimental and numerical data in the literature (Marusic *et al.* 2010; Baidya *et al.* 2017), and is associated with the energization and broadening of the inertia-dominated eddy hierarchy with increasing Re_τ . Here, the increase in \overline{uw} can be connected with the increase in magnitude of both u fluctuations as well as w fluctuations in the overlap region, and is depicted for the latter by $\overline{w^2}^+$ plotted for ZPG TBLs in figure 4(a). This growth can be traced back to the source term for $\overline{w^2}$, i.e. Π^{w+} (figure 4d), which also clearly increases in the outer region with Re_τ owing to the increased losses from $\overline{u^2}$, via Π^{u+} (also refer to $z^+\Pi^{u+}$, $z^+\Pi^{w+}$ in figures 3(d) and 4(g)).

In the case of low Re_τ APG TBLs, interestingly, the enhanced \mathcal{P}^{u+} in the outer region (relative to ZPG TBLs) can be associated with relatively high mean shear (figure 2e–h) as well as \overline{uw} (figure 2i–l). However, an increase in Re_τ results in a drop in the mean shear in their outer region, and hence, the sustained high magnitudes and peaks of \mathcal{P}^{u+} at high Re_τ can be associated with the increasing \overline{uw} magnitudes noted for low-to-moderately strong APG TBLs. This scenario is consistent with the Re_τ trends noted in the case of ZPG TBLs, which is also confirmed from the high magnitudes of w fluctuations (figure 4b,c) sourced from the significantly enhanced Π^{w+} and $-\Pi^{u+}$ in the outer region (figures 2–4). Indeed, one can note a clear one-to-one connection between \mathcal{P}^{u+} , $-\Pi^{u+}$, Π^{w+} , $\overline{w^2}^+$ and $-\overline{uw}^+$ profiles for APG TBLs in figures 2 and 4, which is reflected by the close alignment of their respective outer peaks with z_{OP}^+ , as also noted previously in high Re_τ APG TBLs (Romero *et al.* 2022). Consistent Re_τ trends are also exhibited by the bulk energy-transfer terms in the outer region (figures 3 and 4), corresponding to the transport of both $\overline{u^2}$ and $\overline{w^2}$.

The aforementioned discussion on the energy-transfer pathway, from TKE production of $\overline{u^2}$ to inter-component transfer to $\overline{w^2}$, explains the consistent variation exhibited by the outer region of $\overline{u^2}$ (figure 2b–d) and \overline{uw} (figure 2i–l) with increasing Re_τ . This consistency is noted for both ZPG and APG TBLs, suggesting that a comprehensive analysis of their \overline{uw} characteristics could further unravel similarities between their energy-transfer pathways. This would be particularly significant considering that the correlation between u and w influences the magnitude and direction of the $\overline{u^2}$ wall-normal transport ($\mathcal{T}^u = -\partial \overline{u^2 w} / \partial z$; see figures 2(w–y) and 3(j–l)). To this end, the next section compares the wall-normal variation of uw contributions from the dominant quadrants Q_2 and Q_4 , with $\overline{u^2}$ and $\overline{w^2}$.




Figure 4. Wall-normal profiles for (a–c) \overline{w}^{2+} , (d–f) Π^{w+} and (g–i) $z^+ \Pi^{w+}$ for ZPG (a,d,g) and APG TBL cases. Symbols/colours correspond to the various cases defined in figure 1, with light to dark shading indicating increasing Re_τ . Note the inconsistent range of ordinate values. Vertical green lines indicate u^2 outer peak locations.

Notably, while both these quadrants contribute positively and significantly to \mathcal{P}^u , their contributions are significant but opposing for $\overline{u^2 w}$ (Nagano & Tagawa 1990).

4. Classification of the streamwise Reynolds-stress profiles based on energy-transfer mechanisms

4.1. Empirical observations based on mean statistics

Figure 5(a–d) compares the wall-normal profiles of $\overline{u^2}$ and the wall-normal flux of streamwise TKE ($\overline{u^2 w}$) for selected ZPG (in black) and APG TBL (in red) cases at matched Re_τ . Here, z_{IP}^+ and z_{OP}^+ are estimated following the same methodology as used previously for figure 2. These profiles are compared with the fractional contributions from Q_2 and Q_4 quadrants to \overline{uw} across the respective TBLs (figure 5e–h). Their contributions are defined by following Wallace (2016) and computing $\sum_i (uw)_{Q_i} / \sum uw$ at each z , where the

Energy-transfer mechanisms in turbulent boundary layers




Figure 5. (a–d) Streamwise Reynolds stress ($\overline{u^2}^+$) and wall-normal flux of streamwise TKE ($\overline{u^2w}^+$), (e–h) fractional contribution to Reynolds shear stress (\overline{uw}^+) from quadrants Q_2 and Q_4 , and (i–l) fraction of the total samples associated with Q_2 and Q_4 , across various ZPG (black) and APG (red) TBL cases at matched Re_τ (see top of each column). The background shading represents different regimes (A–D, B^* , D^*) defined in table 2, based on the relative predominance of Q_2 or Q_4 contributions to \overline{uw} for APG TBLs. Note that z_{AB}^+ , $z_{AB^*}^+$, z_{BC}^+ , z_{CD}^+ and $z_{B^*D^*}^+$ represent locations associated with regime transitions, while z_{IP}^+ , z_{OP}^+ and z_M^+ represent the location of the inner peak, outer peak and local minimum of the $\overline{u^2}^+$ profiles, respectively.

numerator represents summation across samples associated exclusively with Q_2 ($u < 0, w > 0$) or Q_4 ($u > 0, w < 0$) events. Also plotted in figure 5(i–l) are the wall-normal variations for the fraction of the total samples associated with Q_2 and Q_4 events, in both the TBLs. Combined together, all these figures depict a consistent connection between the behaviour of $\overline{u^2}$ and $\overline{u^2w}$ profiles, with statistical dominance of Q_2 or Q_4 , which is discussed here.

Figure 5(a,e,i) compares statistics of a weak APG TBL with ZPG TBL at very low Re_τ , and their respective wall-normal profiles look qualitatively similar. For $z^+ < z_{IP}^+$, $d\overline{u^2}^+/dz^+ > 0$, and this is associated with significantly higher contributions from Q_4 events to \overline{uw} than Q_2 (figure 5e). Interestingly, these statistically significant Q_4 events correspond to a smaller fraction of the total uw signal than Q_2 (figure 5i), suggesting occurrence of intense ‘splating’ events in this near-wall region (Kim, Moin & Moser 1987). This also results in the mean wall-normal flux of $\overline{u^2}$ directed towards the wall (i.e. $\overline{u^2w} < 0$). The relative dominance of Q_4 over Q_2 reverses for $z^+ \gtrsim z_{IP}^+$, which sees Q_2 events contributing significantly more to \overline{uw} and occurring across a smaller fraction of the total uw signal (than Q_4). As a consequence, the mean wall-normal flux of $\overline{u^2}$ is directed away from the wall (i.e. $\overline{u^2w} > 0$), and interestingly, this region is associated with




Figure 6. Scatter plot demonstrating the one-to-one correlation between (a) z_{IP}^+ and z_{AB}^+ (or $z_{AB^*}^+$), (b) z_M^+ and z_{BC}^+ , and (c) z_{OP}^+ and z_{CD}^+ , associated with $\overline{u^2}^+$ and \overline{uw}^+ from various ZPG and APG TBL data sets. Definitions are given in § 4.1, table 2 and figure 5. Symbols and colour shading correspond to the various cases and data sets defined in figure 1 and table 1.

$\frac{d\overline{u^2}^+}{dz^+} < 0$. Based on the variation in predominance of Q_2 and Q_4 , we can classify the near-wall region of ZPG TBLs and weak APG TBLs into regimes A ($0 \lesssim z^+ \lesssim z_{AB^*}^+$) and B^* ($z_{AB^*}^+ \lesssim z^+ \lesssim z_{B^*D^*}^+$). Here, $z_{AB^*}^+$ is the location for the ‘switch’ in relative predominance between Q_2 and Q_4 , and this is well known to coincide with z_{IP}^+ (Kim *et al.* 1987; Wallace 2016). Figure 6(a) reaffirms this one-to-one correspondence by comparing $z_{AB^*}^+$ and z_{IP}^+ for the various ZPG and weak APG TBL cases considered. Furthermore, as can be confirmed from the various Re_τ cases in figure 5, the above discussion on the behaviour of the flow is valid for all ZPG TBL cases ($Re_\tau \lesssim 2000$). A noticeable difference between ZPG and weak APG TBLs, however, is that Q_4 contributions to \overline{uw} are significantly greater in the outer region of APG TBLs (figure 5e). This is expected considering the previous findings of Skaare & Krogstad (1994) and Lee (2017). Next, we investigate how the dominance of Q_4 varies with further energization of $\overline{u^2}$ in the APG TBL outer region. To this end, we define $z_{B^*D^*}^+$ as the location corresponding to a local maximum of Q_4 contributions in the outer region of ZPG and weak APG TBLs, and the regime D^* defined across $z_{B^*D^*}^+ \lesssim z^+ \lesssim \delta_{99}^+$.

On considering an APG TBL at a moderately strong β (≈ 1.35) and slightly higher Re_τ in figure 5(b, f, j), an outer peak can be noted in the u^2 profile. Although the appearance of this outer peak is subtle, it is accompanied by distinct changes in the u^2w profile as well as enhancement of Q_4 contributions to \overline{uw} . Notably, there is a small intermediate regime at $z^+ \gg z_{IP}^+$, wherein Q_4 contributions to \overline{uw} become greater than those from Q_2 (figure 5f). This enhanced Q_4 strength is accompanied by a drop in the fraction of the uw signal associated with Q_4 events (figure 5j), suggesting occurrence of intense Q_4 activity in the outer region of the APG TBLs, analogous to the behaviour observed for $z^+ \lesssim z_{IP}^+$. This explains the mean wall-normal flux of $\overline{u^2}$ directed towards the wall in this intermediate regime (figure 5b), as also observed previously by Skaare & Krogstad (1994). On the other hand, the energy-transfer mechanisms in the inner region of this moderately strong APG TBL are similar to those noted in weak APG and ZPG TBLs.

The dominance of Q_4 over Q_2 in an intermediate regime, in addition to the very near-wall region, means that the u^2 profiles of moderately strong APG TBLs can be classified into four regimes. The regions associated with predominance of Q_4 are defined as regimes A ($0 \lesssim z^+ \lesssim z_{AB}^+$) and C ($z_{BC}^+ \lesssim z^+ \lesssim z_{CD}^+$), while those corresponding to

Regimes	TBL	Q_i dominance	z^+ range	Lower bound	Upper bound
A	ZPG, weak APG	$\overline{uw} _{Q4} > \overline{uw} _{Q2}$	$0 \leq z^+ \leq z_{AB}^+$	Wall	$\overline{uw} _{Q4} = \overline{uw} _{Q2}$
	Moderate APG	$\overline{uw} _{Q4} > \overline{uw} _{Q2}$	$0 \leq z^+ \leq z_{AB}^+$	Wall	$\overline{uw} _{Q4} = \overline{uw} _{Q2}$
B*	ZPG, weak APG	$\overline{uw} _{Q2} > \overline{uw} _{Q4}$	$z_{AB}^+ \leq z^+ \leq z_{B^*D^*}^+$	$\overline{uw} _{Q4} = \overline{uw} _{Q2}$	Maximum $\overline{uw} _{Q4}$
B	Moderate APG	$\overline{uw} _{Q2} > \overline{uw} _{Q4}$	$z_{AB}^+ \leq z^+ \leq z_{BC}^+$	$\overline{uw} _{Q4} = \overline{uw} _{Q2}$	$\overline{uw} _{Q4} = \overline{uw} _{Q2}$
C	Moderate APG	$\overline{uw} _{Q4} > \overline{uw} _{Q2}$	$z_{BC}^+ \leq z^+ \leq z_{CD}^+$	$\overline{uw} _{Q4} = \overline{uw} _{Q2}$	$\overline{uw} _{Q4} = \overline{uw} _{Q2}$
D	Moderate APG	$\overline{uw} _{Q2} > \overline{uw} _{Q4}$	$z_{CD}^+ \leq z^+ \leq \delta^+$	$\overline{uw} _{Q4} = \overline{uw} _{Q2}$	TBL edge
D*	ZPG, weak APG	$\overline{uw} _{Q2} > \overline{uw} _{Q4}$	$z_{B^*D^*}^+ \leq z^+ \leq \delta^+$	Maximum $\overline{uw} _{Q4}$	TBL edge

Table 2. Table summarizing the definitions of various regimes classified in § 4.1, based on the relative dominance of Q_2 or Q_4 Reynolds shear stress events in both ZPG and APG TBLs.

predominance of Q_2 are defined as regimes B ($z_{AB}^+ \lesssim z^+ \lesssim z_{BC}^+$) and D ($z_{CD}^+ \lesssim z^+ \lesssim \delta_{99}^+$). Table 2 summarizes the definitions of these limits, which are used to classify the various regimes based on the relative dominance of Q_2 or Q_4 . Here, z_{BC}^+ and z_{CD}^+ correspond to the two locations in the outer region at which there is a switch in the relative predominance of Q_4 and Q_2 contributions, while z_{AB}^+ corresponds to the same in the inner region (figure 5f). Analogous to the correlations noted in the inner region, where z_{IP}^+ coincides with z_{AB}^+ , we find z_{CD}^+ to coincide with z_{OP}^+ in the outer region. In between the two peaks, there is a minimum in the $\overline{u^2}$ profile at z_M^+ ($= d^2\overline{u^2}^+/dz^{+2} > 0$ and $d\overline{u^2}^+/dz^+ = 0$), which is found to coincide with z_{BC}^+ . Remarkably, both regimes A and C are associated with $\overline{u^2w} < 0$ and $d\overline{u^2}^+/dz^+ > 0$, while regimes B and D are associated with $\overline{u^2w} > 0$ and $d\overline{u^2}^+/dz^+ < 0$. Hence, it can be argued that the wall-normal energy transport across the ‘outer’ region ($z_{BC}^+ \lesssim z^+ \lesssim \delta_{99}^+$) is phenomenologically similar to that across the ‘inner’ region ($0 \lesssim z^+ \lesssim z_{BC}^+$), with Q_4 and Q_2 respectively dominating below and above each of the $\overline{u^2}$ peaks. This is consistent with previous discussions in § 3, based on analysis of the Reynolds-stress-transport terms, wherein the energy-transfer pathways were found to be similar near both the inner and outer peaks of $\overline{u^2}$. Here, note that the use of ‘inner’ and ‘outer’ is not per the classical definitions in the literature. Interestingly, this connection between the sign of $d\overline{u^2}^+/dz^+$, and the relative dominance of Q_4 or Q_2 , has also been noted previously in rough wall flows (Katul *et al.* 2006), hinting at universality in the influence of these quadrants on $\overline{u^2}$.

The aforementioned hypothesis can be tested by analysing the same statistics, at similar moderately strong APGs ($\beta \approx 1.59, 1.13$), in TBLs at relatively high Re_τ values of 700 (figure 5c,g,k) and 2000 (figure 5d,h,l). A higher Re_τ leads to increased separation between scales associated with the inner and outer peaks, which should result in widening of the intermediate regime C ($z_{BC}^+ \lesssim z^+ \lesssim z_{CD}^+$). As a consequence, z_{IP}^+ , z_{OP}^+ as well as z_M^+ can be distinctly noted in each of these relatively high Re_τ cases, and indeed these maxima and minima locations respectively coincide with (z_{AB}^+ , z_{CD}^+) and z_{BC}^+ defined based on relative predominance of Q_2 or Q_4 (table 2). This is reaffirmed in figure 6(a–c), which demonstrates consistency in $\overline{u^2}$ profile classification across all moderately strong APG TBL cases, despite their different upstream histories. Figures 5(c,g,k) and 5(d,h,l), thus, confirm that the underlying wall-normal energy-transport mechanisms discussed for figure 5(b,f,j) remain valid for all moderately strong APG scenarios. That is, regimes A and C correspond to dominant contributions of Q_4 to \overline{uw} , which are associated with

smaller fractions of the total uw signal (compared with Q_2). The mean wall-normal flux of $\overline{u^2}$, consequently, is directed towards the wall in both these regimes, and these again coincide with $d\overline{u^2}^+/dz^+ > 0$. On the other hand, the flow phenomena is entirely opposite in regimes B and D , where the Q_2 events predominate over Q_4 events leading to $\overline{u^2}w > 0$. Hence, phenomenologically, z_{BC}^+ appears to be the intermediate location where wall-ward $\overline{u^2}$ flux from the outer, and away-from-wall flux from the inner region ‘equalize’, plausibly yielding a minimum in $\overline{u^2}$.

The quadrant analysis in this section reveals that the mean wall-normal flux of $\overline{u^2}$ is characteristically different in the outer region of moderately strong APG TBLs, compared with that in low Re_τ ZPG and weak APG TBLs. However, this contrasts with the similar energy-transfer pathways noted for all these TBLs in § 3. We hypothesize this inconsistency is owing to the statistically weak TKE production in the outer region of low Re_τ ZPG and weak APG TBLs, owing to which the associated energy-transfer mechanisms get masked by much stronger transfer mechanisms originating in their inner region (owing to stronger TKE production). We investigate this hypothesis next via conditional analysis, to check if the dominance of Q_4 events noted in moderately strong APG TBLs, is also unravelled on conditionally averaging the outer regions of ZPG and weak APG TBLs.

4.2. Conditional analysis of the outer region

Here, we report conditional averaging based on statistically significant TKE production events detected in the outer region, to bring out the energy-transfer processes associated with the outer region. This is achieved by deploying the same time-series thresholding criteria as that used by Deshpande & Marusic (2021) and Narasimha *et al.* (2007), i.e. $|uw(z; t)| > k(uw(z; t))_{sd}$, which identifies time-series segments associated with strong Reynolds shear stress events (i.e. contributing to strong TKE production). Here, $(uw(z; t))_{sd}$ represents the standard deviation of the uw signal at z , t represents time and k represents the constant threshold, following the original proposal of Narasimha *et al.* (2007). Since we are interested in analysing the wall-normal profiles of conditional-averaged statistics, the entire flow field will be conditioned based on the uw signal at a single z location. For the present analysis, $k = 0.50$ is chosen, however, we have checked that the conclusions do not depend on the threshold when $k \gtrsim 0.25$. The threshold, $k = 0.25$, was found by Deshpande & Marusic (2021) to recover 99 % of the mean Reynolds shear stress despite neglecting ~ 43 % of the total uw signal (meaning this 43 % essentially contributed insignificantly to \overline{uw}). Here, since we want to extract only the most statistically dominant uw signal, we have opted for $k = 0.50$ that leads to disconsideration of ~ 62 % of the uw signal.

We begin by first demonstrating the utility of this conditional-averaging analysis by applying it to a moderately strong APG TBL case at $Re_\tau \approx 2000$, for which similarity in $\overline{u^2}$ wall-normal transport between the inner and outer regions has already been demonstrated (illustrated by the relative dominance of Q_2 and Q_4 in figure 5*d*). Figure 7(*a,e*) reports the conditionally averaged statistics (indicated by angular brackets; plotted using green lines) that are compared with the time-averaged Reynolds shear stresses as well as fractional contributions from Q_4 and Q_2 to \overline{uw} (same as those plotted in figure 5*d,h*). For figures 7(*a,e*), the z location for conditioning is chosen the same as the location of the outer peak, i.e. $z_{CD}^+ = z_{OP}^+$, and we can note the conditionally averaged Q_4 and Q_2 contributions to the Reynolds shear stress to be qualitatively similar to that noted in the time-averaged statistics. To demonstrate that the choice of flow conditioning location does not influence

Energy-transfer mechanisms in turbulent boundary layers




Figure 7. Time-averaged (in red or black) and conditionally averaged (in green) (a–d) Reynolds shear stresses and (e–h) fractional contributions to Reynolds shear stresses from Q_2 and Q_4 , plotted for various ZPG and APG TBL cases considered previously in figure 5. The flow conditioning criteria has been adopted from Deshpande & Marusic (2021) and defined in § 4.2, with the z location for conditioning indicated by the vertical blue dashed lines in (a–d). Dashed green lines with arrows in (e–h) indicate the z^+ range in the outer region where $Q_4 > Q_2$ contributions to $\langle uw \rangle$.

this analysis, it is repeated on the same data set but conditioned at the local maxima of Q_4 contributions (figure 7b,f), which is close to z_{BC}^+ for moderately strong APG TBLs. While the conditionally averaged statistics change quantitatively on changing the conditioning location from z_{CD}^+ to z_{BC}^+ , the relative dominance of the Q_4 and Q_2 contributions remains the same qualitatively, and also in the same wall-normal range as reported based on the time-averaged statistics. For example, the dominance of Q_4 over Q_2 in the outer region is found to exist in a similar z^+ range in both figure 7(e,f), which is indicated by the dashed green lines with arrows. Importantly, independence from the specific choice of flow conditioning location permits us to deploy this analysis technique next on to ZPG and weak APG TBL data, i.e. without discernible $\overline{u^2}$ outer peaks or dominance of Q_4 in the outer region.

Figure 7(c,g) depicts the conditionally averaged statistics computed for a weak APG TBL case shown previously in figure 5(a,e,i), for which the flow is conditioned at the local maxima of Q_4 contributions (i.e. $z_{B^*D^*}^+$). Conditioning the flow based on statistically significant uw signals, in the outer region, indeed brings out the dominance of Q_4 over Q_2 contributions to $\langle uw \rangle$ (marked by green arrows), which is otherwise not observed in the time-averaged statistics. The same analysis is then extended onto ZPG TBL data at $Re_\tau \approx 2000$ (figure 7d,h). Note that the time-averaged statistics for this ZPG case were previously plotted in figure 5(d,h). Remarkably, conditionally averaging the flow at $z_{B^*D^*}^+$ again reveals the dominance of Q_4 over Q_2 contributions to $\langle uw \rangle$, providing evidence in support of this hypothesis from Lee & Moser (2019), Deshpande & Marusic (2021) and Cremades *et al.* (2024). This unravels the existence of similar wall-normal energy-transfer mechanisms in the outer region, as those noted in the inner region, for both ZPG and APG TBLs. It also confirms that energy-transfer mechanisms in the outer region of low Re_τ ZPG and weak APG TBLs are statistically weak in a time-averaged sense, and are consequently masked by the much stronger mechanisms associated with the inner region.

5. Conceptual picture of the streamwise energy-transfer mechanisms

Having gained significant insights based on the empirical trends reported in figures 2–7, here we attempt to represent the relationship between key energy-transfer mechanisms and $\overline{u^2}$ profiles via a generalized conceptual sketch. This will be discussed first for APG TBLs, and then for ZPG TBLs. Developing this general understanding will help to propose a hypothesis for energy-transfer mechanisms in very high Re_τ ZPG TBLs, for future investigations. In figure 8(b) the energy-transfer mechanisms in moderately strong APG TBLs are depicted to be governed by two dominant ‘epicentres’, at z_{AB}^+ ($= z_{IP}^+$) and z_{CD}^+ ($= z_{OP}^+$). These epicentres correspond to the coexisting local maxima noted for \mathcal{P}^u and $-\Pi^u$, as well as equal contributions from Q_4 and Q_2 to \overline{uw} , in both the inner and outer regions (§§ 3 and 4). This justifies referencing them as epicentres, considering these locations correspond to intense $\overline{u^2}$ production as well as its transfer to $\overline{w^2}$ (via Π^u) and wall-normal transport ($\overline{u^2 w}$). However, by no means should these be interpreted as the only z^+ locations for TKE production and inter-component energy transfer, which is well known to occur across a range of scales and wall-normal locations (Cossu & Hwang 2017; Lee & Moser 2019). In figure 8(b), z_{AB}^+ is the epicentre in the near-wall region, associated with the energy of the inner-scaled streaks (which is part of the near-wall cycle proposed by Hamilton *et al.* 1995). While, z_{CD}^+ is the epicentre in the outer region and is associated with either the mean shear or shear stresses based on inertial motions (depending on the Re_τ). The concentration of intense streamwise TKE production at both these locations explains their coincidence with $\overline{u^2}$ peaks, and this results in the positive and negative gradients of $\overline{u^2}$ below and above either of these peaks, respectively. The TKE produced near these epicentres is respectively transported towards and away from the wall by Q_4 ($w < 0$) and Q_2 ($w > 0$) events. This explains the relative predominance of Q_4 and Q_2 below and above these TKE production peaks, respectively. The wall-ward flux of $\overline{u^2}$ from the outer epicentre (carried by Q_4), merges with the flux directed away from the wall (Q_2), from the inner epicentre, at an intermediate location z_{BC}^+ ($= z_M^+$). This explains the coincidence of the $\overline{u^2}$ minima with the balance between Q_2 and Q_4 contributions at z_{BC}^+ , distinguishing it from the epicentres at z_{AB}^+ and z_{CD}^+ . In this way, figure 8(b) summarizes the classification of the $\overline{u^2}$ profiles (into regimes A–D) based on the energy-transfer mechanisms noted for moderately strong APG TBLs, found to be independent of the history effects.

On the other hand, a weak APG implies that the TKE production in the outer region is not ‘intense enough’ to generate an outer peak in the $\overline{u^2}$ profile (figure 2), a hypothesis that was validated in § 4.2. We found that although there is a weak epicentre in the outer region (figure 8a) associated with significant TKE production (figure 2f), the wall-ward flux of $\overline{u^2}$ from this epicentre (via Q_4 events) is not strong enough to balance against the $\overline{u^2}$ flux away from the wall, carried by Q_2 events originating near z_{AB}^+ . As a consequence, $Q_2 > Q_4$ for all $z^+ > z_{AB}^+$ in weak APG TBLs. Hence, based on the evidence from time-averaged (figures 2–6) and conditionally averaged statistics (figure 7) of weak and moderately strong APG TBLs, we argue that the presence of a $\overline{u^2}$ outer peak directly corresponds with the predominance of Q_4 contributions to \overline{uw} in an intermediate regime C. Indeed, the emergence of the $\overline{u^2}$ outer peak with an increase in APG strength can be tracked by investigating the growth of Q_4 contributions at $z_{B^*D^*}^+$ (i.e. maxima of Q_4), which increases beyond the contributions from Q_2 as the outer peak emerges (figure 5a–h).

Next, we extend the same argument to ZPG TBLs, to discuss the extent of similarity in energy-transfer mechanisms with APG TBLs. While the mean shear in the outer region




Figure 8. Conceptual sketches describing the energy-transfer mechanisms inferred based on figure 2–7 for (a) weak APG and low-to-high Re_τ ZPG TBLs, and (b) moderately strong APG TBLs. Solid golden circles indicate z^+ locations where Q_2 and Q_4 events are equally strong and occupy the same fraction of the total signal. These solid circles are referred to as ‘epicentres’ owing to the maxima in \mathcal{P}^u and $-\Pi^u$. Here T/NTI refers to the turbulent/non-turbulent interface.

of ZPG TBLs is not as strong as in the case of APGs, an increase in Re_τ leads to the energization and broadening of the inertial eddy hierarchy, which increases the TKE production in the outer region via growth in the Reynolds shear stresses (§ 3). Based on figures 2–5, it is thus reasonable to conclude that the energy-transfer mechanisms in a low-to-high Re_τ ZPG TBL (where no distinct $\overline{u^2}$ outer peak is observed) are analogous to those in a weak APG TBL, which has been conceptualized in figure 8(a). As demonstrated via figure 7(d,h) for a moderate $Re_\tau \sim 2000$ of a ZPG TBL, the streamwise TKE production in the outer region is statistically insignificant at this Re_τ , owing to which the wall-ward flux of $\overline{u^2}$ energy (from the outer epicentre) is not strong enough to yield predominance of Q_4 contributions in the intermediate region (over Q_2 originating from the inner region). However, it is plausible that a significant increase in Re_τ could enhance the outer TKE production (Marusic *et al.* 2010) such that the energy-transfer mechanisms for very high Re_τ ZPG TBLs become analogous to those conceptualized in figure 8(b), for moderately strong APG TBLs. This hypothesis, however, can only be confirmed in the future after availability of canonical data sets at very high Re_τ .

6. Summary and concluding remarks

This study investigates energy-transfer mechanisms associated with the Re_τ increment of the streamwise Reynolds-stress profiles ($\overline{u^2}$) in the outer region of TBLs. Published data

sets of weak and moderately strong APG TBLs are analysed alongside those of ZPG TBLs, to offer phenomenological arguments based on the energy-transfer mechanisms prominent in the outer region. Connections between the wall-normal profiles of $\overline{u^2}$, its production (\mathcal{P}^u), inter-component (Π^u) and wall-normal transport (\mathcal{T}^u), as well as viscous dissipation (\mathcal{E}^u) are investigated, all of which increase in statistical significance with increasing Re_τ (Marusic *et al.* 2010; Cho *et al.* 2018; Lee & Moser 2019). The present analysis reveals similarity in the energy-transfer pathways between the inner and outer regions of ZPG and APG TBLs. The energy pathway originates from the production of $\overline{u^2}$ via \mathcal{P}^u , which is then either (i) transferred to $\overline{v^2}$ or $\overline{w^2}$ by Π^u , (ii) transported in the wall-normal direction by \mathcal{T}^u , or (iii) lost to heat through \mathcal{E}^u . The enhancement of these energy-transfer processes (with Re_τ) for both ZPG and APG TBLs is predominantly associated with the energization and broadening of the inertial eddy hierarchy, which is indicated by the growth of the Reynolds shear stresses. This similarity noted between ZPG and low-to-moderately strong APG TBLs is consistent with previous findings of Gungor *et al.* (2022). They obtained similar conclusions on comparing transfer mechanisms between canonical channel flows and APG TBLs, thereby supporting extrapolation of the present findings from APG TBLs on to all canonical flows.

For moderately strong APGs, which are characterized by both inner and outer $\overline{u^2}$ peaks, similar energy-transfer mechanisms are found to be centred around either of these peaks (in a phenomenological sense). Both inner (z_{AB}^+) and outer peaks (z_{CD}^+) correspond with a local maximum in \mathcal{P}^u and $-\Pi^u$ profiles, and are accompanied by the predominance of $\overline{u^2}$ energy flux ($\overline{u^2 w}$) towards/away from the wall in the region immediately below/above each of them. This inspires the classification of the $\overline{u^2}$ profile into four distinct regimes, the definition of which is independent of the upstream history effects. The ‘inner’ region has an epicentre of \mathcal{P}^u and $-\Pi^u$ at z_{AB}^+ that regulates $\overline{u^2}$ energy flux towards (in regime A; $0 \lesssim z^+ \lesssim z_{AB}^+$) and away from the wall (in regime B; $z_{AB}^+ \lesssim z^+ \lesssim z_{BC}^+$). This wall-normal transport is governed respectively by the Q_4 and Q_2 quadrants of the Reynolds shear stress. A similar conceptual picture is observed in the ‘outer’ region, with the energy-transfer mechanisms maximum around the outer epicentre z_{CD}^+ , which defines the other two regimes: C ($z_{BC}^+ \lesssim z^+ \lesssim z_{CD}^+$) and D ($z_{CD}^+ \lesssim z^+ \lesssim \delta_{99}^+$). These two ‘inner’ and ‘outer’ regions merge at an intermediate location corresponding to the minimum in the $\overline{u^2}$ profiles, which is located where the $\overline{u^2}$ energy flux from the two epicentres ‘equalize’.

In scenarios of low-to-high Re_τ ZPG TBLs and weak APG TBLs, however, no distinct outer peak is noted in $\overline{u^2}$, \mathcal{P}^u or Π^u , with the wall-normal flux of $\overline{u^2}$ directed away from the wall (i.e. Q_2 dominance) across the outer region. Considering, however, the similarity exhibited by the energy-transfer pathways between moderately strong APG TBLs and ZPG TBLs, it is hypothesized that the outer-energy mechanisms in the latter are statistically weaker than those in their inner region. Conditional averaging is implemented to unravel the flow phenomena associated exclusively with the outer-region energy mechanisms, in weak APG and ZPG TBLs; it reveals relative dominance of Q_4 and Q_2 in their outer region to be consistent with that observed for moderately strong APG TBLs. These trends suggest that a u^2 outer peak can emerge for ZPG TBLs only at very high Re_τ , when the outer-energy mechanisms become sufficiently significant to overcome the influence of the inner-energy mechanisms. This hypothesis, however, can only be confirmed in the future after availability of well-resolved measurements at very high Re_τ , potentially unravelling new flow physics associated with these energy-transfer mechanisms.

To conclude, the present study improves our fundamental understanding of both ZPG and APG TBLs on several fronts. Namely, it brings to light a direct connection between

the z variations in their $\overline{u^2}$ profiles and the Reynolds shear stress-carrying (Q_4 , Q_2) events, and the influential role played by the latter in the Re_τ variation of the former. Furthermore, this study also suggests that moderately strong APG TBLs can be potentially used to qualitatively understand the trends in energy-transfer mechanisms in very high Re_τ ZPG TBLs. Since the fundamental mechanisms causing the energization of the outer region are typically centred at distinctly different locations in these two TBL types, a combination of both parameters (i.e. high Re_τ APG TBLs) will be expected to have interesting flow physics (Vinuesa *et al.* 2017; Romero *et al.* 2022; Deshpande *et al.* 2023). The present analysis and arguments, hence, provide a plausible approach towards understanding this unexplored regime of weak-to-moderately strong APG TBLs at high Re_τ .

Acknowledgements. The authors are grateful to Drs R. Pozuelo and C. Sanmiguel Vila for sharing their data sets, and thank Professors H. Nagib, I. Marusic, A. Smits and M. Hultmark for their encouraging comments and helpful discussions.

Funding. R. Deshpande is grateful to the University of Melbourne's Postdoctoral Fellowship that funded his visit to KTH Stockholm. R. Deshpande also acknowledges the Office of Naval Research (ONR) and ONR Global grant no. N62909-23-1-2068. R. Vinuesa acknowledges the financial support from ERC grant no. '2021-CoG-101043998, DEEPCONTROL'. Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

Declaration of interests. The authors report no conflict of interest.

Author ORCIDs.

 Rahul Deshpande <https://orcid.org/0000-0003-2777-2919>;

 Ricardo Vinuesa <https://orcid.org/0000-0001-6570-5499>.

REFERENCES

- ALFREDSSON, P.H., SEGALINI, A. & ÖRLÜ, R. 2011 A new scaling for the streamwise turbulence intensity in wall-bounded turbulent flows and what it tells us about the 'outer' peak. *Phys. Fluids* **23** (4), 041702.
- BAARS, W.J. & MARUSIC, I. 2020 Data-driven decomposition of the streamwise turbulence kinetic energy in boundary layers. Part 2. Integrated energy and A_1 . *J. Fluid Mech.* **882**, A26.
- BAIDYA, R., PHILIP, J., HUTCHINS, N., MONTY, J.P. & MARUSIC, I. 2017 Distance-from-the-wall scaling of turbulent motions in wall-bounded flows. *Phys. Fluids* **29** (2), 020712.
- BOBKE, A., VINUESA, R., ÖRLÜ, R. & SCHLATTER, P. 2017 History effects and near equilibrium in adverse-pressure-gradient turbulent boundary layers. *J. Fluid Mech.* **820**, 667–692.
- CHEN, X. & SREENIVASAN, K.R. 2021 Reynolds number scaling of the peak turbulence intensity in wall flows. *J. Fluid Mech.* **908**, R3.
- CHO, M., HWANG, Y. & CHOI, H. 2018 Scale interactions and spectral energy transfer in turbulent channel flow. *J. Fluid Mech.* **854**, 474–504.
- COSSU, C. & HWANG, Y. 2017 Self-sustaining processes at all scales in wall-bounded turbulent shear flows. *Phil. Trans. R. Soc. A* **375** (2089), 20160088.
- CREMADES, A., HOYAS, S., DESHPANDE, R., QUINTERO, P., LELLEP, M., LEE, W.J., MONTY, J.P., HUTCHINS, N., LINKMANN, M., MARUSIC, I. & VINUESA, R. 2024 Identifying regions of importance in wall-bounded turbulence through explainable deep learning. *Nat. Commun.* **15** (1), 3864.
- DESHPANDE, R., BOGAARD, A., VINUESA, R., LINDIĆ, L. & MARUSIC, I. 2023 Reynolds-number effects on the outer region of adverse-pressure-gradient turbulent boundary layers. *Phys. Rev. Fluids* **8** (12), 124604.
- DESHPANDE, R. & MARUSIC, I. 2021 Characterising momentum flux events in high Reynolds number turbulent boundary layers. *Fluids* **6** (4), 168.
- DONG, S., LOZANO-DURÁN, A., SEKIMOTO, A. & JIMÉNEZ, J. 2017 Coherent structures in statistically stationary homogeneous shear turbulence. *J. Fluid Mech.* **816**, 167–208.
- EITEL-AMOR, G., ÖRLÜ, R. & SCHLATTER, P. 2014 Simulation and validation of a spatially evolving turbulent boundary layer up to $Re_\theta = 8300$. *Intl J. Heat Fluid Flow* **47**, 57–69.

- FIORINI, T., BELLANI, G., ÖRLÜ, R., SEGALINI, A., ALFREDSSON, P.H. & TALAMELLI, A. 2017 Turbulent pipe flow near-wall statistics. In *Progress in Turbulence VII: Proceedings of the iTi Conference in Turbulence 2016* (ed. R. Örlü, A. Talamelli, M. Oberlack & J. Peinke), pp. 89–94. Springer.
- GUNGOR, A.G., MACIEL, Y., SIMENS, M.P. & SORIA, J. 2016 Scaling and statistics of large-defect adverse pressure gradient turbulent boundary layers. *Intl J. Heat Fluid Flow* **59**, 109–124.
- GUNGOR, T.R., MACIEL, Y. & GUNGOR, A.G. 2022 Energy transfer mechanisms in adverse pressure gradient turbulent boundary layers: production and inter-component redistribution. *J. Fluid Mech.* **948**, A5.
- HAMILTON, J.M., KIM, J. & WALEFFE, F. 1995 Regeneration mechanisms of near-wall turbulence structures. *J. Fluid Mech.* **287**, 317–348.
- HARUN, Z., MONTY, J.P., MATHIS, R. & MARUSIC, I. 2013 Pressure gradient effects on the large-scale structure of turbulent boundary layers. *J. Fluid Mech.* **715**, 477–498.
- KATUL, G., POGGI, D., CAVA, D. & FINNIGAN, J. 2006 The relative importance of ejections and sweeps to momentum transfer in the atmospheric boundary layer. *Boundary-Layer Meteorol.* **120**, 367–375.
- KIM, J., MOIN, P. & MOSER, R. 1987 Turbulence statistics in fully developed channel flow at low Reynolds number. *J. Fluid Mech.* **177**, 133–166.
- KITSIOS, V., SEKIMOTO, A., ATKINSON, C., SILLERO, J.A., BORRELL, G., GUNGOR, A.G., JIMÉNEZ, J. & SORIA, J. 2017 Direct numerical simulation of a self-similar adverse pressure gradient turbulent boundary layer at the verge of separation. *J. Fluid Mech.* **829**, 392–419.
- KLINE, S.J., REYNOLDS, W.C., SCHRAUB, F.A. & RUNSTADLER, P.W. 1967 The structure of turbulent boundary layers. *J. Fluid Mech.* **30** (4), 741–773.
- LEE, J.H. 2017 Large-scale motions in turbulent boundary layers subjected to adverse pressure gradients. *J. Fluid Mech.* **810**, 323–361.
- LEE, J.-H. & SUNG, H.J. 2009 Structures in turbulent boundary layers subjected to adverse pressure gradients. *J. Fluid Mech.* **639**, 101–131.
- LEE, M. & MOSER, R.D. 2019 Spectral analysis of the budget equation in turbulent channel flows at high Reynolds number. *J. Fluid Mech.* **860**, 886–938.
- MARUSIC, I., MATHIS, R. & HUTCHINS, N. 2010 High Reynolds number effects in wall turbulence. *Intl J. Heat Fluid Flow* **31** (3), 418–428.
- MONKEWITZ, P.A. 2022 Asymptotics of streamwise Reynolds stress in wall turbulence. *J. Fluid Mech.* **931**, A18.
- MONKEWITZ, P.A. & NAGIB, H.M. 2015 Large-Reynolds-number asymptotics of the streamwise normal stress in zero-pressure-gradient turbulent boundary layers. *J. Fluid Mech.* **783**, 474–503.
- NAGANO, Y. & TAGAWA, M. 1990 A structural turbulence model for triple products of velocity and scalar. *J. Fluid Mech.* **215**, 639–657.
- NARASIMHA, R., KUMAR, S.R., PRABHU, A. & KAILAS, S.V. 2007 Turbulent flux events in a nearly neutral atmospheric boundary layer. *Phil. Trans. R. Soc. A* **365** (1852), 841–858.
- NEGI, P.S., VINUESA, R., HANIFI, A., SCHLATTER, P. & HENNINGSON, D.S. 2018 Unsteady aerodynamic effects in small-amplitude pitch oscillations of an airfoil. *Intl J. Heat Fluid Flow* **71**, 378–391.
- POZUELO, R., LI, Q., SCHLATTER, P. & VINUESA, R. 2022 An adverse-pressure-gradient turbulent boundary layer with nearly constant $\beta \simeq 1.4$ up to $Re_\theta \simeq 8700$. *J. Fluid Mech.* **939**, A34.
- ROMERO, S., ZIMMERMAN, S., PHILIP, J., WHITE, C. & KLEWICKI, J. 2022 Properties of the inertial sublayer in adverse pressure-gradient turbulent boundary layers. *J. Fluid Mech.* **937**, A30.
- SAMIE, M., MARUSIC, I., HUTCHINS, N., FU, M.K., FAN, Y., HULTMARK, M. & SMITS, A.J. 2018 Fully resolved measurements of turbulent boundary layer flows up to $Re_\tau = 20\,000$. *J. Fluid Mech.* **851**, 391–415.
- SANMIGUEL VILA, C., ÖRLÜ, R., VINUESA, R., SCHLATTER, P., IANIRO, A. & DISCETTI, S. 2017 Adverse-pressure-gradient effects on turbulent boundary layers: statistics and flow-field organization. *Flow Turbul. Combust.* **99**, 589–612.
- SCHOPPA, W. & HUSSAIN, F. 2002 Coherent structure generation in near-wall turbulence. *J. Fluid Mech.* **453**, 57–108.
- SKAARE, P.E. & KROGSTAD, P. 1994 A turbulent equilibrium boundary layer near separation. *J. Fluid Mech.* **272**, 319–348.
- TANG, S.L. & ANTONIA, R.A. 2023 Similarity for dissipation-scaled wall turbulence. *J. Fluid Mech.* **960**, A18.
- VINUESA, R., BOBKE, A., ÖRLÜ, R. & SCHLATTER, P. 2016 On determining characteristic length scales in pressure-gradient turbulent boundary layers. *Phys. Fluids* **28** (5), 055101.
- VINUESA, R., ÖRLÜ, R. & SCHLATTER, P. 2017 Characterisation of backflow events over a wing section. *J. Turbul.* **18** (2), 170–185.
- WALLACE, J.M. 2016 Quadrant analysis in turbulence research: history and evolution. *Annu. Rev. Fluid Mech.* **48**, 131–158.