

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Aloe, CA;Leong, TLT;Wimalaswaran, H;Papagianis, PC;McQualter, JL;McDonald, CF;Khor, YH;Hoy, RF;Ingle, A;Bansal, V;Goh, NSL;Bozinovski, S

Title:

Excess iron promotes emergence of foamy macrophages that overexpress ferritin in the lungs of silicosis patients

Date:

2022-06-01

Citation:

Aloe, C. A., Leong, T. L. T., Wimalaswaran, H., Papagianis, P. C., McQualter, J. L., McDonald, C. F., Khor, Y. H., Hoy, R. F., Ingle, A., Bansal, V., Goh, N. S. L. & Bozinovski, S. (2022). Excess iron promotes emergence of foamy macrophages that overexpress ferritin in the lungs of silicosis patients. *Respirology*, 27 (6), pp.427-436. <https://doi.org/10.1111/resp.14230>.












Persistent Link:

<https://hdl.handle.net/11343/301803>

License:

[CC BY-NC-ND](#)

Excess iron promotes emergence of foamy macrophages that overexpress ferritin in the lungs of silicosis patients

Christian Anthony Aloe¹  | Tracy Li-Tsein Leong^{2,3,4}  | Hari Wimalleswaran^{2,3,5}  |
 Paris Clarice Papagianis¹  | Jonathan Luke McQualter¹  | Christine Faye McDonald^{2,3}  |
 Yet Hong Khor^{2,3,6}  | Ryan Francis Hoy^{7,8}  | Aviraj Ingle⁹ | Vipul Bansal⁹  |
 Nicole Soo Leng Goh^{2,3}  | Steven Bozinovski¹ 

¹School of Health & Biomedical Sciences, RMIT University, Bundoora, Victoria, Australia

²Department of Respiratory Medicine, Austin Health, Heidelberg, Victoria, Australia

³Institute for Breathing and Sleep, Melbourne, Victoria, Australia

⁴The Walter and Eliza Hall Institute of Medical Research, Parkville, Victoria, Australia

⁵Faculty of Medicine, Dentistry and Health Sciences, University of Melbourne, Parkville, Victoria, Australia

⁶Respiratory Research@Alfred, Central Clinical School, Monash University, Melbourne, Victoria, Australia

⁷Department of Epidemiology and Preventive Medicine, School of Public Health and Preventive Medicine, Monash University, Melbourne, Victoria, Australia

⁸Department of Respiratory Medicine, Alfred Hospital, Melbourne, Victoria, Australia

⁹Sir Ian Potter NanoBioSensing Facility, NanoBiotechnology Research Laboratory, School of Science, RMIT University, Melbourne, Victoria, Australia

Correspondence

Steven Bozinovski
 Email: steven.bozinovski@rmit.edu.au

Nicole Soo Leng Goh
 Email: nicole.goh@austin.org.au

Funding information

Open access publishing facilitated by RMIT University, as part of the Wiley - RMIT University agreement via the Council of Australian University Librarians.

Associate Editor: Alexander Neil Larcombe;

Handling Editor: Philip Bardin

Abstract

Background and objective: Inhalation of high concentrations of respirable crystalline silica (RCS) can lead to silicosis. RCS contains varying levels of iron, which can cause oxidative stress and stimulate ferritin production. This study evaluated iron-related and inflammatory markers in control and silicosis patients.

Methods: A cohort of stone benchtop industry workers ($n = 18$) were radiologically classified by disease severity into simple or complicated silicosis. Peripheral blood and bronchoalveolar lavage (BAL) were collected to measure iron, ferritin, C-reactive protein, serum amyloid A and serum silicon levels. Ferritin subunit expression in BAL and transbronchial biopsies was analysed by reverse transcription quantitative PCR. Lipid accumulation in BAL macrophages was assessed by Oil Red O staining.

Results: Serum iron levels were significantly elevated in patients with silicosis, with a strong positive association with serum ferritin levels. In contrast, markers of systemic inflammation were not increased in silicosis patients. Serum silicon levels were significantly elevated in complicated disease. BAL macrophages from silicosis patients were morphologically consistent with lipid-laden foamy macrophages. Ferritin light chain (FTL) mRNA expression in BAL macrophages was also significantly elevated in simple silicosis patients and correlated with systemic ferritin.

Conclusion: Our findings suggest that elevated iron levels during the early phases of silicosis increase FTL expression in BAL macrophages, which drives elevated BAL and serum ferritin levels. Excess iron and ferritin were also associated with the emergence of a foamy BAL macrophage phenotype. Ferritin may represent an early disease

Christian Anthony Aloe and Tracy Li-Tsein Leong have contributed equally to this study.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Respirology* published by John Wiley & Sons Australia, Ltd on behalf of Asian Pacific Society of Respirology.

marker for silicosis, where increased levels are independent of inflammation and may contribute to fibrotic lung remodelling.

KEYWORDS

ferritin, foam cells, interstitial lung disease, iron, silicosis

INTRODUCTION

Silicosis is an irreversible and potentially fatal interstitial lung disease caused by repeated inhalation of respirable crystalline silica (RCS). Silicosis can be divided into acute, accelerated and chronic phenotypes depending on the intensity and duration of exposure.¹ The demand for domestic kitchen benchtops fabricated from engineered (also known as artificial or reconstituted) stone—typically >90% of silica content—over the past decades has seen a wave of silicosis cases in stone benchtop industry workers. These workers typically develop disease after exposure to large amounts of silica over a short period of time (5–15 years).^{2,3} Radiographic changes of silicosis consist of diffuse nodules and thoracic lymphadenopathy. While these nodules are typically small (<1 cm) and distributed throughout both lung fields in early disease, these changes can transition from a ‘simple’ to a ‘complicated’ form as nodules coalesce into large fibrotic masses termed progressive massive fibrosis.⁴ With no effective treatments, and an alarming spike in cases, a greater understanding of the cellular and molecular drivers of this disease is urgently needed.

Alveolar macrophages are frontline cellular responders to inhaled irritants including RCS through engagement of pattern recognition and scavenger receptors. Phagocytosis of RCS activates the NALP3 inflammasome pathway, which promotes secretion of the pro-inflammatory cytokines IL-1 β and IL-18.^{1,5} IL-1 β and IL-18 stimulate the production of a broad suite of acute-phase proteins. Ferritin is well recognized as an acute-phase reactant that is elevated in a wide range of inflammatory conditions.^{6–9} Of significance, high serum ferritin levels are related to worse prognosis in patients with acute exacerbations of idiopathic pulmonary fibrosis, where immunohistochemical staining identified alveolar macrophages as an important cellular source of ferritin.¹⁰ NALP3 inflammasome activation by silica also stimulates the generation of reactive oxygen species (ROS) in a dose-dependent manner.¹¹ The extent of ROS production by RCS is dependent on iron complexed with silica, as high amounts of iron promote greater ROS production and lipid peroxidation in pre-clinical models of RCS exposure.¹² Iron can also independently stimulate the production of ferritin, which is a multimeric protein composed of 24 heavy (FTH) and light chains (FTL). The FTL subunit is primarily responsible for removal of excess free iron from the body, where iron–ferritin complexes are sequestered by macrophages to prevent iron from catalysing the production of free radicals via Fenton chemistry.^{13,14} Elevated serum ferritin levels have been detected

SUMMARY AT A GLANCE

Silicosis is an aggressive and incurable lung disease. In this study, serum iron levels were increased in silicosis patients, and these levels were strongly associated with serum ferritin levels. Lipid-laden bronchoalveolar lavage macrophages were identified as a major source of ferritin, whereas markers of inflammation were not increased.

in simple and complicated silicosis compared to individuals exposed to silica without disease.¹⁵ However, little is known about the driver of excess ferritin production in silicosis.

In this study, we hypothesized that excess iron in silicosis patients is primarily responsible for elevated serum ferritin levels, and that this acute-phase response is stimulated early in the disease process. To address this, we have analysed a panel of serum inflammatory/acute-phase reactants and iron markers in a cohort of silicosis patients stratified by disease severity (simple vs. complicated). In addition, we investigated the cellular source of ferritin whereby matching bronchoalveolar lavage (BAL) macrophages and transbronchial biopsies (TBBx) were analysed for FTH and FTL expression by reverse transcription quantitative PCR (RT-qPCR).

METHODS

Study participants

This study was conducted at the Austin Hospital, Heidelberg, Victoria, Australia, and at RMIT University, Bundoora, Victoria, Australia. Recruitment of cases and controls occurred between January 2020 and October 2021. Stone benchtop industry workers who had worked with engineered stone were eligible for inclusion. The study groups consisted of patients with (1) radiographic changes characteristic of silicosis and (2) no radiological disease. Patients with silicosis were further stratified into ‘simple’ and ‘complicated’ study groups based on radiological findings, where all imaging was read by International Labour Office (ILO)-accredited expert chest radiologists using The International Classification of High-resolution Computed Tomography for Occupational and Environmental Respiratory Diseases (ICOERD) grading scale. The scans were then

TABLE 1 Patient characteristics

	No disease (<i>n</i> = 5)	Simple silicosis (<i>n</i> = 8)	Complicated silicosis (<i>n</i> = 10)
Females/males (<i>n</i>)	4/1	0/8	0/10
Age (years)	55 (46–64)	40 (34–52)	44 (27–55)
BMI (kg/m ²)	34.8 (32–39)	26.7 (21–36)	24.3 (21–38)
Smoking status: (never/ex/current)	4/0/0	1/1/6	2/4/4
FEV ₁ (% of predicted)	84	85 (69–106)	71 (60–94)
FVC (% of predicted)	87 (80–94)	79 (68–100)	93 (62–99)
DLCO (% of predicted)	80 (77–83)	72 (40–97)	79 (65–103)
Artificial stone exposure (years)	NA	13 (5–17)	13 (4–29)

Note: All numerical data are presented as median (range), if not indicated otherwise.

Abbreviations: DLCO, diffusion capacity of the lungs for carbon monoxide; FEV₁, forced expiratory volume in 1 s; FVC, forced vital capacity.

categorized according to the described changes. Patients referred for bronchoscopy to assess chronic cough and who had normal computed tomography (CT) chest and lung function tests were recruited as healthy controls, forming the ‘no disease’ group. Clinical information, demographics, detailed occupational history, spirometry, diffusion capacity of the lungs for carbon monoxide (DLCO) and HRCT chest imaging were obtained. Peripheral blood sampling, BAL and TBBx (in the case of the silicosis group) were also obtained.

Biomarker analysis

Peripheral blood sampling was performed to obtain white blood cell counts and erythrocyte sedimentation rate (ESR). Serum was collected to measure C-reactive protein (CRP), serum amyloid A (SAA), procalcitonin (PCT), ferritin, iron, total iron-binding capacity, silicon, total bilirubin, alanine aminotransferase (ALT), alkaline phosphatase, gamma-glutamyl transferase, creatinine and urea (as outlined in Appendices S2–S5 in the Supporting Information). BAL (as collected in Appendix S1 in the Supporting Information) was utilized to perform a differential BAL count and evaluate macrophage morphology (as outlined in Appendices S6 and S7 in the Supporting Information). To investigate the cellular sources of ferritin in the lung, FTH and FTL subunit mRNA expression was evaluated by RT-qPCR in BAL cells and TBBx (as outlined in Appendix S8 in the Supporting Information).

Statistical analysis

Data were presented as mean values ± SEM. One-way non-parametric analysis of variance was used for the comparison among three groups and the Kruskal–Wallis *H*-test was used for post analysis. Following chest HRCT, patients were divided into their study groups according to their disease severity. GraphPad Prism 9.0 software (GraphPad Software, San Diego, USA) was utilized for graph production and data analysis, where a *p*-value of less than 0.05 was considered as statistical significance.

RESULTS

Table 1 summarizes the clinical characteristics of the study cohort. Twenty-three participants were recruited; 18 (78%) with silica exposure and radiological features of silicosis; and five without radiological features of silicosis that consisted of five (22%) healthy controls. Of the 18 patients with radiological manifestations, eight (44%) presented with early-stage ‘simple’ disease, with the remaining 10 (56%) exhibiting late-stage ‘complicated’ disease. The median time of exposure to dust from engineered stone was 13 ± 6 years, which were not significantly different between patients with simple and complicated disease.

Systemic ferritin is differentially elevated in patients with early-stage silicosis

Serum ferritin levels were significantly elevated in patients with simple silicosis, and there was no significant difference between the simple and complicated silicosis groups (Figure 1A). A similar pattern was observed for serum iron levels and total iron-binding capacity, with levels significantly increased in the simple group and levels remaining elevated in those with advanced disease (Figure 1B,C). There was no evidence for iron overload as transferrin saturation levels fell within normal ranges (Figure 1D). Serum silicon levels were significantly higher in silicosis patients, which increased with disease stage (Figure 1E). Correlation analysis revealed that serum ferritin levels did not associate with serum silicon levels, whereas there was a strong association between serum ferritin and serum iron levels (Figure 1F,G).

Peripheral blood analysis (Figure 2A–D) revealed the absence of a persistent systemic inflammatory response in all groups as total white blood cell, lymphocyte, neutrophil and eosinophil counts were within normal ranges. The classic inflammatory acute-phase reactant, CRP, was not significantly increased in silicosis patients, nor was SAA (Figure 2E,F). Similarly, ESR levels were not significantly altered across the groups and fell within the normal ranges (Figure 2H). Serum PCT levels were also observed to fall




FIGURE 1 Serum ferritin is significantly elevated in patients with simple silicosis (SS) compared to patients with complicated silicosis (CS) and those with no disease (ND) (A). Similarly, serum iron and total iron-binding capacity (TIBC) levels were elevated in patients with SS, with levels remaining elevated in patients with complicated disease (B, C). However, iron overload is not evident as the transferrin saturation % (TS%) levels are within the normal range of 15%–50% (D). Quantification of silicon levels in serum shows significantly higher silicon levels in patients with CS (E). Correlation analysis reveals that ferritin does not associate with serum silicon levels (F), whereas there was a strong association between serum ferritin and serum iron levels (G). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, one-way non-parametric analysis of variance (Kruskal–Wallis test), data represent mean \pm SEM. Horizontal dashed lines represent the limit/s of the normal range

within the normal range in control and silicosis patients (Figure 2G). Ferritin levels can also be altered because of liver or kidney dysfunction; however, assessment of liver and renal functions fell within the normal range in both patients with simple and complicated silicosis (Figure 3), with the exception of serum ALT that was increased in complicated silicosis patients.

Foamy macrophages produce FTL, which increases circulating ferritin levels

BAL cell differential analysis revealed similar profiles in patients with no disease versus patients with simple and complicated silicosis (Figure 4A–D), where BAL




FIGURE 2 Haematological analysis of white blood cell counts and an acute-phase reactant panel demonstrated the absence of a persistent systemic inflammatory response in non-diseased (ND), and in simple (SS) and complicated silicosis (CS) patients. This is evident as the total WCC, blood lymphocyte (Lymphs), neutrophil (Neuts), and eosinophil (Eos) numbers were within normal ranges (A–D). Markers of systemic inflammation, C-reactive protein (CRP), serum-amyloid A (SAA), procalcitonin (PCT) and erythrocyte sedimentation rate (ESR) are not significantly altered across the groups (E–H). * $p < 0.05$, one-way non-parametric analysis of variance (Kruskal–Wallis test), data represent mean \pm SEM. Horizontal dashed lines represent the limit/s of the normal range

macrophages were the dominant immune cell. However, BAL macrophages were morphologically different across the study groups (Figure 4E–G). Macrophages in the silicotic lung appeared both enlarged and vacuolated, where a foam cell-like appearance was attributed to lipid loading, confirmed by Oil Red O Staining (Figure 4H–J). Quantitative assessment demonstrated that there was a $32 \pm 7\%$ and $26 \pm 9\%$ increase in lipid-loaded macrophages in patients with simple and complicated silicosis, respectively, relative to those with no disease (Figure 4K).

Local lung ferritin levels in the BAL fluid were significantly increased in both simple and complicated groups, and positively correlated with circulating serum ferritin levels (Figure 5A,B). FTH mRNA expression in BAL cells from patients with simple silicosis was not increased (Figure 5C), whereas FTL subunit expression was significantly increased (Figure 5D). Median FTL levels were lower in the complicated silicosis group; however, they were not significantly different to the simple silicosis group (Figure 5D). FTH expression in BAL cells did not correlate with serum ferritin levels, whereas there was a strong positive association between FTL mRNA expression in BAL cells and serum ferritin levels (Figure 5E,F). FTH and FTL expression in TBBx were not significantly different in simple versus complicated silicosis (Figure 5G). Paired analysis of FTH and FTL mRNA expression in matching BAL and

TBBx revealed that BAL cells were the main ferritin source in the lung (Figure 5H,I).

DISCUSSION

Our study demonstrated that serum ferritin levels were increased in patients with silicosis, with the highest levels observed in simple silicosis, suggesting that levels rise early during the disease process. Although seven (78%) of nine patients with complicated silicosis displayed serum ferritin levels above those in the no disease control group, a larger sample size may be required to attain significance. In contrast, markers of inflammation and hepatic/renal dysfunction were not increased in patients with simple silicosis, which can also drive increased ferritin production. Our findings are in contrast with a recent study by Blanco-Pérez, where increased serum ferritin levels were detected in silicosis patients in conjunction with elevated levels of inflammatory markers including CRP.¹⁵ However, this study was characterized by high incidence of active tuberculosis infection, which is an independent cause of systemic inflammation and confound findings of elevated ferritin.¹⁶ In our study, there were no cases of active tuberculosis and a single case of latent tuberculosis as evidenced by a positive QuantiFERON-TB Gold assay.




FIGURE 3 Markers of liver and renal dysfunction are not altered in the serum of simple (SS) and complicated silicosis (CS) patients. With the exception of alanine aminotransferase (ALT) in CS, total bilirubin, ALT, alkaline phosphatase (ALP), gamma-glutamyl transferase (GGT), creatinine and urea levels fell within the normal range in patients with no disease (ND), SS and CS (A-F). * $p < 0.05$, one-way non-parametric analysis of variance (Kruskal-Wallis test), data represent mean \pm SEM. Horizontal dashed lines represent the limit/s of the normal range

We also found that serum ferritin levels strongly associated with serum iron levels. Whilst iron is vital in a wide range of metabolic processes, including oxygen transport and DNA synthesis, iron dysregulation can produce free radicals causing oxidative damage in a broad spectrum of diseases.^{13,17} Silicates complex iron onto their surfaces, including inorganic iron and iron derived from biological systems.^{18,19} Ghio et al. demonstrated that the ability of silicates to catalyse oxidant generation in vitro is proportional to the amount of ferric ion complexed onto their surfaces. As such, surface complexation of iron onto silicates may be an important determinant of their ability to cause silicosis.^{12,18} In contrast to iron, serum ferritin levels did not correlate with serum silicon levels. Intriguingly, serum silicon levels were increased in complicated silicosis patients and there was a subset of patients with high serum silicon levels in simple disease. It would therefore be important to establish whether simple silicosis patients with high serum silicon levels progress more rapidly in future studies.

A secondary consequence of excess iron is the generation of oxidative stress including the accumulation of

oxidized phospholipids. Intriguingly, we observed a change in macrophage morphology where alveolar macrophages were enlarged and vacuolated in patients with silicosis. This morphological change was attributed to lipid loading, confirmed by Oil Red O staining. Mechanisms mediating foam cell formation in silicosis are not well defined. Hou et al. reported that foam cells were associated with a dramatic increase in levels of oxidized low-density lipoprotein (ox-LDL).²⁰ Interestingly, both iron and ferritin have been implicated in promoting apoptosis of foam cells in human atheroma,²¹ where iron-laden macrophages associate with LDL oxidation, angiogenesis and nitric oxide production.²² Yuan et al. also demonstrated that iron-laden macrophages exocytosed both iron and ferritin into culture medium, where exposure to ox-LDL, but not high-density lipoprotein, resulted in apoptosis of iron-laden human macrophages.²² It is plausible that iron loading may be driving lipid accumulation and apoptosis in lipid-laden BAL macrophages in a progressive manner, although further work is needed to confirm this.




FIGURE 4 Bronchoalveolar lavage (BAL) cell profiles are not altered in patients with simple (SS) and complicated silicosis (CS) compared to patients with no disease (ND). This is evident as no statistical differences in BAL macrophages (Macs), lymphocytes (Lymphs), neutrophils (Neuts) and eosinophils (Eos) are observed across the study groups when BAL cytosps are stained with Kwik-Diff™ (ThermoFisher Scientific) and quantified (A–D). A change in macrophage morphology is observed across the study groups (E–G), where alveolar macrophages in silicosis-diseased patients are found to be lipid-laden when stained with Oil Red O (H–K). * $p < 0.05$, one-way non-parametric analysis of variance (Kruskal–Wallis test), data represent mean \pm SEM

Our study is the first to show that BAL macrophages are an important source of ferritin in the silicotic lung where FTL, but not FTH, expression correlates with BAL and serum ferritin levels. Iron is oxidized to the ferric state by FTH, while FTL stabilizes the complex, favouring

mineralization during iron storage in macrophages. Hence, ferritin is post-transcriptionally increased during exposure to excess iron through an iron response element (IRE) contained within an untranslated region of the ferritin gene. IRE is regulated by binding of a cytoplasmic protein, iron




FIGURE 5 Ferritin light chain (FTL) is overexpressed in bronchoalveolar lavage (BAL) macrophages and positively associated with circulated ferritin levels. Ferritin in the BAL fluid (BALF) is elevated in patients with both simple (SS) and complicated silicosis (CS) compared to patients with no disease (ND) (A), where local ferritin levels positively correlated with circulating serum ferritin (B). Evaluation of ferritin heavy chain (FTH) and FTL subunit mRNA expression by RT-qPCR revealed the cellular source of ferritin to be BAL macrophages, where FTL is significantly elevated in patients with SS (C, D). FTH expression in BAL cells did not correlate with serum ferritin levels, whereas FTL mRNA expression in BAL cells revealed a strong positive association with serum ferritin levels (E, F). Ferritin subunit expression was also evaluated in transbronchial biopsies (TBBx), where FTH and FTL levels were not significantly different in SS versus CS (G). Paired analysis of FTH and FTL mRNA expression in matching BAL and TBBx revealed that BAL cells are the main ferritin source in the lung (H, I). * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$, one-way non-parametric analysis of variance (Kruskal–Wallis test) or paired t -test was conducted where appropriate, data represent mean \pm SEM

regulatory protein (IRP-1), where their binding is stimulated by multiple cellular signals including cellular iron levels and oxidative stress.²³ Consistent with our clinical observations, silica complexed with iron generates greater production of FTL in human alveolar macrophages compared to silica alone under in vitro conditions.²⁴ Hence, we propose that dysfunctional lung macrophages produce excessive levels of ferritin in response to iron-loaded silica exposure, where ferritin spills over into the lung vasculature to increase serum ferritin levels.

Excess iron and ferritin may also directly contribute to fibrotic remodelling through modulation of pathogenic macrophages. Macrophages exhibit distinct phenotypic and functional states in response to environmental cues, which give rise to a range of effector functions including tissue repair, regeneration and fibrosis.^{25,26} There may be distinct macrophage phenotypes in silicosis, which differentially respond to silica nano- and micro-particle uptake and toxicity.^{27,28} In a murine model of silicosis, Zhao et al. demonstrated that polarization of pulmonary macrophages was dysregulated during silicosis development, where pro-inflammatory macrophages were induced in early disease and anti-inflammatory macrophages were found to promote dysfunctional repair and fibrosis in late disease.²⁹ Emerging evidence demonstrates that iron has the potential to directly influence macrophage polarization.^{19,30}

Iron, in the form of ferric ammonium citrate (FAC), can drive the differentiation of THP-1 macrophages towards a wound-healing phenotype. Additionally, FAC induced a significant increase in ferritin expression.³⁰ Of significance, coculture of FAC pre-treated macrophages with human dermal fibroblasts (HDFs) led to the upregulation of type-1 and -3 collagen mRNA expression in HDFs.³⁰ FTL may also directly promote fibrosis, where FTL knockdown can dramatically repress endothelial-to-mesenchymal transition and migration by regulating AKT/GSK3 β / β -catenin signalling.³¹ Further studies are needed to establish whether FTL can exhibit similar effects in the context of chronic lung diseases including silicosis.

Our study has a number of limitations. Whilst we did observe a differential increase in serum ferritin levels in patients with simple silicosis, the sample size is small. Additionally, the present study represents data from a single time-point. Future studies should ideally focus on longitudinally tracking ferritin levels in the serum and BAL to determine whether ferritin levels are predictive of patients who progress to complicated silicosis as determined by HRCT. Such studies may delineate whether ferritin is valuable as a prognostic biomarker. Moreover, BAL and serum longitudinal analysis of silica-exposed workers who are yet to develop radiological disease may be of great insight into the development and progression of silicosis. Whilst the NALP3 inflammasome pathway can acutely stimulate the production of inflammatory and acute-phase mediators following RCS exposure, we found that systemic inflammation did not persist in patients with simple or complicated silicosis. Potentially of greater significance to the development of silicosis is the iron and ferritin status, where iron-loaded RCS particles stimulate a

phenotypic shift in BAL macrophages, which develop a foam cell morphology and chronically overexpress FTL. Excess iron and FTL overexpressing macrophages have the potential to contribute to lung fibrosis via promoting the accumulation of lipid-loaded foam macrophages that stimulate collagen deposition and promoting endothelial-to-mesenchymal transition through β -catenin signalling. Future work is needed to investigate the significance of excess iron and FTL in the development of silica-induced lung fibrosis.

ACKNOWLEDGEMENTS

Aviraj Ingle acknowledges RMIT University for a Research Stipend Scholarship towards his PhD. Vipul Bansal acknowledges the support of Ian Potter Foundation towards establishing Sir Ian Potter NanoBioSensing Facility at RMIT University that allowed silicon quantification studies. Tracy Li-Tsein Leong is supported by the David Bickart Clinician Research Fellowship from the University of Melbourne.

CONFLICT OF INTEREST

This study was supported by a research grant from Multi-plex Australia.

AUTHOR CONTRIBUTION

Christian Anthony Aloe: Data curation (equal); formal analysis (equal); methodology (equal); project administration (equal); writing – original draft (supporting). **Tracy Li-Tsein Leong:** Data curation (equal); formal analysis (equal); investigation (equal); writing – review and editing (equal). **Hari Wimalaswaran:** Formal analysis (supporting); investigation (supporting); writing – review and editing (supporting). **Paris Clarice Papagianis:** Formal analysis (equal); investigation (equal); writing – review and editing (equal). **Jonathan Luke McQualter:** Formal analysis (equal); investigation (equal); project administration (equal); writing – review and editing (equal). **Christine Faye McDonald:** Formal analysis (equal); supervision (equal); writing – review and editing (equal). **Yet Hong Khor:** Formal analysis (equal); validation (equal); writing – review and editing (equal). **Ryan Francis Hoy:** Writing – review and editing (equal). **Aviraj Ingle:** Formal analysis (equal); investigation (equal); methodology (equal); writing – review and editing (equal). **Vipul Bansal:** Formal analysis (equal); methodology (equal); resources (equal); supervision (equal); writing – review and editing (equal). **Nicole Soo Leng Goh:** Formal analysis (equal); investigation (equal); writing – original draft (equal). **Steven Bozinovski:** Conceptualization (lead); formal analysis (equal); investigation (equal); project administration (lead); writing – original draft (lead).

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

HUMAN ETHICS APPROVAL DECLARATION

This study was performed in accordance with the Declaration of Helsinki and was approved by the Austin Health

Human Research Ethics Committee – approval: 54116/2019. All adult participants provided written informed consent to participate in this study.

ORCID

Christian Anthony Aloe  <https://orcid.org/0000-0002-7646-0535>

Tracy Li-Tsein Leong  <https://orcid.org/0000-0002-1950-1505>

Hari Wimalaswaran  <https://orcid.org/0000-0003-3727-5000>

Paris Clarice Papagianis  <https://orcid.org/0000-0002-4709-6017>


Jonathan Luke McQualter  <https://orcid.org/0000-0002-3844-8009>

Christine Faye McDonald  <https://orcid.org/0000-0001-6481-3391>

Yet Hong Khor  <https://orcid.org/0000-0002-5434-9342>

Ryan Francis Hoy  <https://orcid.org/0000-0001-9150-9440>

Vipul Bansal  <https://orcid.org/0000-0002-3354-4317>

Nicole Soo Leng Goh  <https://orcid.org/0000-0003-2065-4346>

Steven Bozinovski  <https://orcid.org/0000-0001-6533-8641>

REFERENCES

- Leung CC, Yu ITS, Chen W. Silicosis. *Lancet*. 2012;379(9830):2008–18.
- Hoy RF, Baird T, Hammerschlag G, Hart D, Johnson AR, King P, et al. Artificial stone-associated silicosis: a rapidly emerging occupational lung disease. *Occup Environ Med*. 2018;75(1):3–5.
- Hoy RF. Artificial stone silicosis. *Curr Opin Allergy Clin Immunol*. 2021;21(2):114–20.
- Barnes H, Goh NSL, Leong TL, Hoy R. Silica-associated lung disease: an old-world exposure in modern industries. *Respirology*. 2019;24(12):1165–75.
- Tsugita M, Morimoto N, Tashiro M, Kinoshita K, Nakayama M. SR-B1 is a silica receptor that mediates canonical inflammasome activation. *Cell Rep*. 2017;18(5):1298–311.
- Gabay C, Kushner I. Acute-phase proteins and other systemic responses to inflammation. *N Engl J Med*. 1999;340(6):448–54.
- Feelders RA, Vreugdenhil G, Eggermont AM, Kuiper-Kramer PA, van Eijk HG, Swaak AJ. Regulation of iron metabolism in the acute-phase response: interferon gamma and tumour necrosis factor alpha induce hypoferraemia, ferritin production and a decrease in circulating transferrin receptors in cancer patients. *Eur J Clin Invest*. 1998;28(7):520–7.
- Nishiya K, Kawabata F, Ota Z. Elevated urinary ferritin in lupus nephritis. *J Rheumatol*. 1989;16(11):1513–4.
- Blake DR, Bacon PA, Eastham EJ, Brigham K. Synovial fluid ferritin in rheumatoid arthritis. *BMJ*. 1980;281(6242):715–6.
- Enomoto N, Oyama Y, Enomoto Y, Mikamo M, Karayama M, Hozumi H, et al. Prognostic evaluation of serum ferritin in acute exacerbation of idiopathic pulmonary fibrosis. *Clin Respir J*. 2018;12(8):2378–89.
- Cassel SL, Eisenbarth SC, Iyer SS, Sadler JJ, Colegio OR, Tephly LA, et al. The Nalp3 inflammasome is essential for the development of silicosis. *Proc Natl Acad Sci U S A*. 2008;105(26):9035.
- Castranova V, Vallyathan V, Ramsey DM, McLaurin JL, Pack D, Leonard S, et al. Augmentation of pulmonary reactions to quartz inhalation by trace amounts of iron-containing particles. *Environ Health Perspect*. 1997;105(5):1319–24.
- Harrison PM, Arosio P. The ferritins: molecular properties, iron storage function and cellular regulation. *BBA-Bioenergetics*. 1996;1275(3):161–203.
- Gozzelino R, Soares MP. Coupling heme and iron metabolism via ferritin H chain. *Antioxid Redox Signal*. 2014;20(11):1754–69.
- Blanco-Pérez JJ, Blanco-Dorado S, Rodríguez-García J, Gonzalez-Bello ME, Salgado-Barreira Á, Caldera-Díaz AC, et al. Serum levels of inflammatory mediators as prognostic biomarker in silica exposed workers. *Sci Rep*. 2021;11(1):13348.
- Miranda P, Gil-Santana L, Oliveira MG, Mesquita ED, Silva E, Rauwerdink A, et al. Sustained elevated levels of C-reactive protein and ferritin in pulmonary tuberculosis patients remaining culture positive upon treatment initiation. *PLoS One*. 2017;12(4):e0175278.
- Hintze KJ, Theil EC. Cellular regulation and molecular interactions of the ferritins. *Cell Mol Life Sci*. 2006;63(5):591.
- Ghio AJ, Kennedy TP, Whorton AR, Crumbliss AL, Hatch GE, Hoidal JR. Role of surface complexed iron in oxidant generation and lung inflammation induced by silicates. *Am J Physiol*. 1992;263(5 Pt 1):L511–8.
- Agoro R, Taleb M, Quesniaux VFJ, Mura C. Cell iron status influences macrophage polarization. *PLoS One*. 2018;13(5):e0196921.
- Hou X, Summer R, Chen Z, Tian Y, Ma J, Cui J, et al. Lipid uptake by alveolar macrophages drives fibrotic responses to silica dust. *Sci Rep*. 2019;9(1):399.
- Yuan XM. Apoptotic macrophage-derived foam cells of human atheromas are rich in iron and ferritin, suggesting iron-catalysed reactions to be involved in apoptosis. *Free Radic Res*. 1999;30(3):221–31.
- Yuan XM, Li W, Baird SK, Carlsson M, Meleforts O. Secretion of ferritin by iron-laden macrophages and influence of lipoproteins. *Free Radic Res*. 2004;38(10):1133–42.
- Torti FM, Torti SV. Regulation of ferritin genes and protein. *Blood*. 2002;99(10):3505–16.
- Ghio AJ, Carter JD, Samet JM, Quay J, Wortman IA, Richards JH, et al. Ferritin expression after in vitro exposures of human alveolar macrophages to silica is iron-dependent. *Am J Respir Cell Mol Biol*. 1997;17(5):533–40.
- Lawrence T, Natoli G. Transcriptional regulation of macrophage polarization: enabling diversity with identity. *Nat Rev Immunol*. 2011;11(11):750–61.
- Wynn TA, Vannella KM. Macrophages in tissue repair, regeneration, and fibrosis. *Immunity*. 2016;44(3):450–62.
- Hoppstädter J, Seif M, Dembek A, Cavalius C, Huwer H, Kraegeloh A, et al. M2 polarization enhances silica nanoparticle uptake by macrophages. *Front Pharmacol*. 2015;6:55.
- Herd HL, Bartlett KT, Gustafson JA, McGill LD, Ghandehari H. Macrophage silica nanoparticle response is phenotypically dependent. *Biomaterials*. 2015;53:574–82.
- Zhao Y, Hao C, Bao L, Wang D, Li Y, Qu Y, et al. Silica particles disorganize the polarization of pulmonary macrophages in mice. *Ecotoxicol Environ Saf*. 2020;193:110364.
- Wilkinson HN, Roberts ER, Stafford AR, Banyard KL, Matteucci P, Mace KA, et al. Tissue iron promotes wound repair via M2 macrophage polarization and the chemokine (C-C motif) ligands 17 and 22. *Am J Pathol*. 2019;189(11):2196–208.
- Liu J, Gao L, Zhan N, Xu P, Yang J, Yuan F, et al. Hypoxia induced ferritin light chain (FTL) promoted epithelia mesenchymal transition and chemoresistance of glioma. *J Exp Clin Cancer Res*. 2020;39(1):137.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Aloe CA, Leong TL-T, Wimalaswaran H, Papagianis PC, McQualter JL, McDonald CF, et al. Excess iron promotes emergence of foamy macrophages that overexpress ferritin in the lungs of silicosis patients. *Respirology*. 2022;1–10. <https://doi.org/10.1111/resp.14230>