

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Hamza, SA;Paolini, R;Moore, C;Unemo, M;Yap, T;McCullough, M;Mayfosh, A;Rau, T;Kleid, S;Hocking, JS;Kong, FYS;Celentano, A

Title:

Fatty acids-based compounds as novel prophylaxis or treatment for oropharyngeal *Neisseria gonorrhoeae*: an in vitro study

Date:

2026

Citation:

Hamza, S. A., Paolini, R., Moore, C., Unemo, M., Yap, T., McCullough, M., Mayfosh, A., Rau, T., Kleid, S., Hocking, J. S., Kong, F. Y. S. & Celentano, A. (2026). Fatty acids-based compounds as novel prophylaxis or treatment for oropharyngeal *Neisseria gonorrhoeae*: an in vitro study. *Emerging Microbes & Infections*, 15 (1), pp.1-23. <https://doi.org/10.1080/22221751.2026.2678644>.

Persistent Link:

<https://hdl.handle.net/11343/368972>

License:

CC BY-NC-ND



Fatty acids-based compounds as novel prophylaxis or treatment for oropharyngeal *Neisseria gonorrhoeae*: an *in vitro* study

Syed Ameer Hamza , Rita Paolini , Caroline Moore , Magnus Unemo , Tami Yap , Michael McCullough , Alyce Mayfosh , Thomas Rau , Stephen Kleid , Jane S Hocking , Fabian YS Kong & Antonio Celentano

To cite this article: Syed Ameer Hamza , Rita Paolini , Caroline Moore , Magnus Unemo , Tami Yap , Michael McCullough , Alyce Mayfosh , Thomas Rau , Stephen Kleid , Jane S Hocking , Fabian YS Kong & Antonio Celentano (22 May 2026): Fatty acids-based compounds as novel prophylaxis or treatment for oropharyngeal *Neisseria gonorrhoeae*: an *in vitro* study, Emerging Microbes & Infections, DOI: [10.1080/22221751.2026.2678644](https://doi.org/10.1080/22221751.2026.2678644)

To link to this article: <https://doi.org/10.1080/22221751.2026.2678644>



© 2026 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group, on behalf of Shanghai Shangyixun Cultural Communication Co., Ltd



[View supplementary material](#)



Accepted author version posted online: 22 May 2026.



[Submit your article to this journal](#)



Article views: 64



[View related articles](#)



[View Crossmark data](#)

Publisher: Taylor & Francis & The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group, on behalf of Shanghai Shangyixun Cultural Communication Co., Ltd

Journal: *Emerging Microbes & Infections*

DOI: 10.1080/22221751.2026.2678644



Fatty acids-based compounds as novel prophylaxis or treatment for oropharyngeal *Neisseria gonorrhoeae*: an *in vitro* study

Syed Ameer Hamza^a, Rita Paolini^a, Caroline Moore^a, Magnus Unemo^{b,c}, Tami Yap^a, Michael McCullough^a, Alyce Mayfosh^d, Thomas Rau^d, Stephen Kleid^e, Jane S Hocking^f, Fabian YS Kong^{f*}, Antonio Celentano^{a*}

^aMelbourne Dental School, The University of Melbourne, Carlton, Victoria, Australia

^bWHO Collaborating Centre for Gonorrhoea and other STIs, Örebro University, Örebro, Sweden

^cInstitute for Global Health, University College London, London, UK

^dWintermute Biomedical, Victoria, Australia

^eMasada Private Hospital, Ramsay Health, Victoria, Australia

^fCentre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne, Carlton, Victoria, Australia

*Joint senior author

Corresponding author: A/Prof Antonio Celentano, Address: Melbourne Dental School, The University of Melbourne, 720 Swanston Street, Carlton, Victoria, Australia, 3053. E-mail: antonio.celentano@unimelb.edu.au Telephone: +61 3 9341 1495

Abstract

Oropharyngeal gonorrhoea remains a global challenge due to rising antimicrobial resistance and limited therapeutic options, necessitating urgent calls for new treatments and prophylaxis. This study evaluates the *in vitro* activity of arginine-based fatty acids against *Neisseria gonorrhoeae* (NG) strains in a validated oropharyngeal model. Five oropharyngeal cell lines (tonsillar/buccal/gingival/floor of mouth/uvular) were infected with two NG strains. Intracellular infection was assessed at 15/30/60 minutes. MIC₉₀ and cytotoxicity of five arginine-based fatty acids were determined. Promising candidates were evaluated for NG clearance at 30/60/120 minutes (post-infection) and prevention of infection (pre-infection) after 30/120 minutes drug exposure. Arginine-undecanoate (AU) and arginine-laurate (AL) were the most effective fatty-acids, showing potent bactericidal activity, with no cytotoxicity in any cell lines up to 300 µg/mL. In post-exposure experiments, AL at 150 µg/mL and at MIC_{x3} achieved rapid, dose-dependent clearance of ≥94% for FA1090 and ≥90% for WHO R respectively within 30 minutes in all cell lines except in uvular cells. AU showed time-dependent killing with clearance of ≥95% for

FA1090 and $\geq 88\%$ for WHO-R at 120 minutes at 150 $\mu\text{g/mL}$ and 108 $\mu\text{g/mL}$ (MIC $\times 3$) respectively. In pre-exposure experiments, AL at maximum non-cytotoxic concentration (300 $\mu\text{g/mL}$) was more effective than AU, achieving 83.3-97.4% clearance in all cell lines within 30 minutes. AL and AU can be promising, non-toxic candidates for the prevention or treatment of oropharyngeal NG. Use, pre- and post-sex via topical formulations, could potentially reduce oropharyngeal NG transmission.

Keywords: Oropharyngeal gonorrhoea, *Neisseria gonorrhoeae*, antimicrobial resistance, fatty acid antimicrobials

Introduction

Gonorrhoea, a sexually transmitted infection caused by *Neisseria gonorrhoeae* (NG), represents a significant public health concern, with at least 82 million new cases among adolescents and adults annually.[1] Rapid emergence of antimicrobial resistance (AMR) in NG to all antibiotic classes, including to last-line treatment ceftriaxone further limits infection management.[2] While emerging therapies such as zoliflodacin and gepotidacin, show promise for urogenital gonorrhoea, cure rates in oropharyngeal infections remain below the World Health Organization (WHO)'s 95% efficacy threshold.[3, 4]

The oropharynx plays a central role in both treatment failure and the development of AMR. Oropharyngeal infections are often asymptomatic, complicating detection and timely treatment, and systemic antibiotics may achieve suboptimal concentrations at the oropharyngeal site.[5] Moreover, commensal bacterial species in the oropharynx facilitate AMR via horizontal gene transfer.[6]

Current preventive strategies have proven ineffective: condoms are limited by low compliance,[7] commercial mouthwashes have shown no benefit in clinical trials[8] and interventions such as doxycycline post-exposure prophylaxis (doxyPEP)[9] and vaccination using meningococcal (4CMenB) vaccines[10] have also failed to control oropharyngeal infections. These challenges highlight the urgent need for novel approaches to control oropharyngeal infections. One novel and innovative strategy is the use of topical oral products administered before or after oral sex to prevent incident oropharyngeal infections. NG primarily infects superficial epithelial tissue,[11] making topical interventions – such as chewing gums or sprays – potentially more effective than systemic drugs, which face pharmacokinetic barriers in achieving adequate concentrations at the oropharyngeal epithelium.[5] Moreover, oropharyngeal NG infections have lower bacterial loads than urogenital or anorectal infections,[12] suggesting that lower drug concentrations may suffice for prevention. Lastly, many oropharyngeal NG will have been naturally cleared within three months without treatment[13] and missed during routine 3-monthly STI screening. Therefore, intervening at the time of oral sex is one possible intervention to prevent ongoing transmission.

Fatty acids have potent antimicrobial properties that are gaining attention for their therapeutic potential.[14] Medium- to long-chain fatty acids disrupt bacterial membranes

by embedding in lipid bilayers, increasing permeability and causing lysis, or impairing energy metabolism and enzyme function, diminishing bacterial viability.[15] Fatty acids have additional advantages over conventional antibiotics with low resistance development potential due to their non-specific and dynamic, structural modes of action.[15, 16] Lauric acid exhibits strong bactericidal and antiviral properties,[17] while undecanoic acid also exhibits antifungal properties.[18] Fatty acids suffer from low aqueous solubility but combining them with L-arginine can improve solubility without compromising antimicrobial potency.[19] Fatty acids are well-suited for mucosal applications, as they are naturally present in saliva, and are classified as Generally Recognized as Safe (GRAS) by the U.S. Food and Drug Administration.[19] Lastly, as fatty acids are found naturally in human diets, marketing authorization of a final therapeutic product as a food or vitamin rather than a medicine, can accelerate market access. These properties make them attractive candidates for incorporation into portable mucosal delivery systems such as chewing gums and sprays, allowing for direct exposure at the site of infection.

In this study, we investigated the *in vitro* activity and cellular toxicity of five arginine-based fatty acid compounds and the two most promising compounds (arginine-undecanoate (AU) and arginine-laurate (AL)) were tested against clinically relevant NG strains (FA1090 and WHO R/FC428) using our validated *in vitro* 2D human oropharyngeal infection model of NG.[20] This was performed to identify promising candidates for the prevention or treatment of oropharyngeal gonorrhoea, and to our best knowledge, this is the first *in vitro* study evaluating fatty acid derivatives for prevention or treatment of oropharyngeal gonorrhoea, addressing a critical unmet need in gonorrhoea control.

Materials and Methods

Briefly, we started with initial screening of five compounds using the three cell lines we used in our validated *in vitro* 2D human oropharyngeal infection model of NG.[20] After that we expanded our *in vitro* oropharyngeal infection model[20] with addition of two epithelial cell lines (buccal and uvular). So, in total, five human oropharyngeal epithelial cell lines (tonsillar, buccal, gingival, floor of mouth (FOM), uvular) were infected / co-cultured (co-culture supplementary methods) with two strains of NG (antibiotic-susceptible FA1090 and multidrug-resistant WHO R/FC428). Infected cells were treated with AU and AL to assess intracellular clearance (post-exposure experiments). In separate experiments, uninfected cells were pre-exposed to AU or AL and then challenged with NG (pre-exposure experiments). Extracellular bacteria were eliminated with gentamicin prior to cell lysis, so colony-forming units (CFUs) reflect intracellular bacteria only. After lysis, intracellular bacteria were quantified by plating cell lysates and after incubation counting CFUs with an automated colony counter (Scan1200™, Interscience, France).

Fatty acid-based compounds

Five fatty acid-based compounds (AU, AL, undecylenate (undecylenic acid), linoleate (linoleic acid), and monodecanoate (decanoic acid)) were screened for efficacy (MIC₉₀) and cytotoxicity. All compounds were supplied by Wintermute Biomedical (<https://www.wintermutebiomedical.com>).

Oral epithelial cell lines and culture conditions

Five epithelial cell lines (tonsillar, buccal, gingival, FOM, and uvular) representing the anatomical variation across the human oral and oropharyngeal axis (supplementary figure 1) were used. Their characteristics and culturing conditions are summarized in supplementary table 1. Uvular cells are not commercially available, so we isolated and immortalized primary human uvular cells in our lab. Isolation and immortalization protocols are available in supplementary methods.

Ethics approval

This study was conducted in accordance with the ethical principles outlined in the Declaration of Helsinki. This project was approved by University of Melbourne Human Research Ethics Committee (approval 2023-27876-47484-3), the Masada Ramsay National Research Unit (2023/PID/0213; 2023/RGO/0240). Written informed consent was obtained from the donors of primary human uvular cells before their participation in the study.

Minimum inhibitory concentration (MIC₉₀)

MIC₉₀ values (mg/L) for the fatty acids were determined by microbroth dilution (MBD) as previously described. [20] Briefly, bacterial suspensions (OD₆₀₀ = 1.0 in PBS) were diluted 1:25 to obtain working stocks. Fatty acids were serially diluted using modified Fastidious Broth (mFB) in 96-well plates (Corning, 3370), and 100 µL of inoculum was added to each well. Plates were sealed with gas-permeable film (Merck, BR701364) and incubated for 24 h using a NIVO plate reader (Revity Group) with continuous shaking (300 rpm), and optical density (600 nm) was recorded hourly. MIC₉₀ was defined as the lowest concentration inhibiting ≥90% bacterial growth as shown in (table 1). Two reference strains (WHO F and ATCC 49226) were also used for quality control. MIC values were further validated using agar dilution according to WHO methodology.

Cytotoxicity assay

Cytotoxicity was assessed by lactate dehydrogenase (LDH) release. Oropharyngeal epithelial cells were seeded in 48-well plates (CLS-3338, Corning®, Sigma-Aldrich, Australia) in triplicate (uvular, gingival, and FOM: 1×10⁴ cells/well; tonsillar and buccal: 2×10⁴ cells/well). Once 40–60% confluent (approximately 6×10⁴ cells/well), cells were treated with fatty acids (50–500 µg/mL to determine maximal non-cytotoxic concentrations) or at 1×, 2×, and 3× MIC (to determine therapeutic concentrations) for 0.5, 1, 2, or 4 hours. Forty-five minutes before each endpoint, a lysis buffer was added to one replicate to serve as the maximum lysis control. After incubation, 50 µL of supernatant was transferred to a 96-well plate, followed by 50 µL LDH reagent, 30 minutes incubation at 37°C, and addition of 50 µL stop solution. Absorbance was read at 490 nm using a Biotek 800 TS microplate reader (BioTek, Currumbin, QLD, Australia).

Microscopy visualization of NG in oropharyngeal cells

Oropharyngeal epithelial cells (1×10^5 cells/well) were cultured in chamber slides (#54526, Thermo Fisher Scientific, Australia) and infected as previously described. [20] Briefly, following infection, cells were treated with fatty acid-based compounds (AU, AL), then fixed with 4% paraformaldehyde. Cells were blocked with 1% BSA and subjected to fluorescence staining using WGA–Alexa Fluor 594, FITC-conjugated *N. gonorrhoeae* antibody, and DAPI. Slides were mounted and imaged using an Invitrogen™ FLoid™ Cell Imaging Station at 80× magnification. Representative images of buccal cells under control and AU/AL treatments are shown in Figure 1.

Statistical analysis

Statistical analyses were performed using GraphPad Prism version 10.6.0 (GraphPad Software, Boston, USA). Each experiment was conducted in technical triplicate, and data are presented as mean $\pm$ standard deviation. For intracellular CFU assays, results were normalized to untreated controls and analyzed using two-way ANOVA to assess the effects of pre-exposure (30 and 120 min at 300 μ g) and post-exposure (30, 60, and 120 min) treatments at different concentrations. Cytotoxicity data were analyzed by one-way ANOVA, normalized to maximum lysis, and compared to untreated controls.

Role of the funding source

The funders had no role in study design, data collection, analysis, interpretation, or writing of the report of this study.

Results

Initial cytotoxicity screening and MICs of fatty acids

Five fatty acid-based compounds were initially screened across three oropharyngeal epithelial cell lines (gingival, tonsillar, and floor-of-mouth) to assess cytotoxicity (Supplementary Table 3), and their minimum inhibitory concentrations (MICs) were determined (Table 1). While all five compounds exhibited antimicrobial activity, three demonstrated unacceptable cytotoxicity toward host cells and were therefore excluded. In contrast, arginine laurate (AL, C12) and arginine undecanoate (AU, C11), both medium-chain fatty acid derivatives, showed no detectable cytotoxicity across all tested cell lines while retaining antimicrobial activity. These compounds were therefore selected for further evaluation against *Neisseria gonorrhoeae* strains in the 2D human oropharyngeal infection model, as they provided the most favourable balance between efficacy and host cell compatibility.

Fatty acids (mg/L)	Agar (or microbroth) dilution MIC (mg/L)			
	FA1090	WHO R	WHO F (control)	ATCC 49226 (control)
Arginine undecanoate * (AU, undecanoic acid)	16 (21)	64 (36)	32	64

Arginine laurate * (AL, lauric acid)	32 (23)	32 (49)	32	32
Arginine-linoleate (linoleic acid)	Not tested (>50)	Not tested (33)	Not tested	Not tested
Arginine undecylenate (undecylenic acid)	32 (46)	128 (31)	64	128
Arginine monodecanoate (decanoic acid)	64 (48)	128 (31)	64	64

Table 1. MIC₉₀ values (mg/L) using microbroth and agar dilution assay. *Final compounds tested in the 2D human oropharyngeal infection model[20]

Cytotoxic assessment

The cytotoxicity of AU and AL at 1–3× MIC was assessed across five oropharyngeal cell lines. As shown in figure 2A, neither compound induced substantial cytotoxicity in any cell line. Based on these results, an additional concentration (150 µg/mL) was tested for FA1090 to align with the 147 µg/mL concentration used for WHO R. To define the upper safe limit for pre-exposure, a broader concentration range (50–500 µg/mL) was evaluated. AU and AL were non-cytotoxic up to 300 µg/mL at all time points; however, higher concentrations (400–500 µg/mL) induced significant cytotoxicity exclusively in gingival cells after 1 hour (figure 2B). Accordingly, 300 µg/mL was selected for subsequent pre-exposure experiments.

pH of culture media

The pH of the culture media containing fatty acid compounds remained within physiological ranges, indicating that these compounds did not disrupt the oropharyngeal epithelial cell environment (supplementary table 2).

Assessment of bacterial clearance by applying compounds before and after infecting oropharyngeal cells with *Neisseria gonorrhoeae*:

Neisseria invasion into oropharyngeal epithelial cells

The five oropharyngeal epithelial cell lines (tonsillar, buccal, gingival, FOM, uvular) were infected with NG strains (FA1090 or WHO R) and commensal *N. oralis* as a negative control.

At 15 minutes, FA1090 infected all cell lines (figure 3A). Highest invasion was seen in buccal (28 CFU), tonsillar (11 CFU), and gingival (10 CFU) cells, and lowest in FOM (1 CFU) and uvular (1 CFU) cells. Similar results were seen for WHO R at 15 minutes with highest invasion in tonsillar (27 CFU), gingival (20 CFU), and buccal (15 CFU) cells, but no invasion in FOM and uvular cells.

At 30 minutes (figure 3A), invasion increased significantly in a time-dependent manner ($p < 0.05$) for FA1090 in buccal (141 CFU), tonsillar (122 CFU), and gingival (121 CFU) cells, while for WHO R only in buccal cells (128 CFU; $p < 0.05$). Invasion in FOM and

uvular cells remained low for both NG strains (FA1090: 10 and 23 CFU; WHO R: 5 CFU and 30 CFU) respectively.

At 60 minutes (figure 3B), FA1090 invasion peaked in tonsillar (457 CFU) and buccal (430 CFU) cells, followed by gingival cells (345 CFU, $p < 0.05$). Invasion in FOM (240 CFU, $p < 0.05$) and uvular cells (184 CFU, $p < 0.05$) was significantly lower. Similarly, WHO R invaded tonsillar (453 CFU) and buccal (620 CFU) cells significantly higher, while gingival (47 CFU, $p < 0.05$), FOM (67 CFU, $p < 0.05$), and uvular (37 CFU, $p < 0.05$) cells showed significantly lower invasion.

For *N. oralis*, at 60 minutes invasion was minimal compared to NG strains. Tonsillar cells exhibited the highest invasion (103 CFU), followed by gingival (38 CFU, $p < 0.05$) and uvular (18 CFU, $p < 0.05$) cells. FOM (8 CFU, $p < 0.05$) and buccal (1 CFU, $p < 0.05$) cells showed negligible invasion.

Post-exposure clearance of *Neisseria gonorrhoeae* strains (FA1090 and WHO R) in oropharyngeal cells by fatty acids:

Clearance of FA1090 infected cells (post-exposure)

Overall, among FA1090 infected cells, AL exhibited the most effective antimicrobial activity, demonstrating both time- and dose-dependent killing, whereas AU showed time-dependent activity only (figure 4A; supplementary table 4A).

For AL, at the highest concentration (150 $\mu\text{g/mL}$), early bacterial clearance at 30 minutes of $>94\%$ was seen in all cell lines except the uvular - tonsillar (98.3% clearance), gingival (97.0%), FOM (95.6%) and buccal (94.1%) cells and lowest in uvular cells (61.8%) as compared to control. Similar results were seen at 60 minutes, however at 120 minutes, AL cleared $>99\%$ of FA1090 from all cell lines.

In contrast, for AU at 150 $\mu\text{g/mL}$, early clearance at 30 minutes of $\geq 93\%$ was only seen in tonsillar and FOM cells (95.8% and 93.0% respectively). Similar results were seen at 60 minutes but with the addition of high clearance in buccal cells (94.6%). At 120 minutes, AU cleared $>95\%$ of FA1090 from all cell lines.

Clearance of WHO R infected cells (post-exposure)

Overall, and similar to FA1090 infected cells, AL displayed the most effective activity against WHO R, showing both time- and dose-dependent killing, while AU was time-dependent only (figure 4B; supplementary table 4B).

For AL at the highest concentration (147 $\mu\text{g/mL}$), early clearance at 30 minutes of $>97\%$ was observed in gingival (100%), tonsillar (97.8%), and uvular (97.7%) cells, and lowest in buccal (90%) and FOM (79.3%) cells, compared with the control. At 60 minutes, clearance of $\geq 94\%$ was observed in all cells (100% for tonsillar and gingival), while at 120 minutes, AL cleared $>99\%$ of WHO R from all cell lines except buccal (96.7%).

For AU at 108 µg/mL (MIC x 3) and 30 minutes, higher clearance was observed only in FOM cells (94.1%), while all other cell lines showed <88% clearance. At 60 minutes, the results remained similar across all cell types, except for the addition of higher clearance in buccal cells (92.0%). At 120 minutes, AU cleared >94% of WHO R only from FOM (97.1%), uvular (96.0%) and buccal (94.2%) cells, while clearance was lower for tonsillar (89.3%) and gingival (88.8%) cells.

Clearance of *Neisseria gonorrhoeae* (FA1090) in oropharyngeal cells pre-treated with fatty acids (pre-exposure)

Similar to the above experiments, AL exhibited the most effective antimicrobial activity. In experiments using the highest non-cytotoxic concentration of 300 µg/mL of AL, early clearance at 30 minutes of >94% was seen in all cell lines except for the buccal cells, i.e. gingival (99.2% clearance), tonsillar (97.4%), uvular (97.4%), FOM (94.4%) and for buccal (83.3%) cells compared to control (figure 4C; supplementary table 4C). At 120 minutes, AL cleared >97% of FA1090 from all cell lines.

In contrast, AU at 300 µg/mL, early clearance at 30 minutes was only seen in uvular cells (93.3%) and lower in gingival (88.3%), tonsillar (73.3%), buccal (55.2%) and FOM (52.5%) cells. At 120 minutes, AU cleared >94% in gingival (99.7%), uvular (98.3%) and tonsillar (94.3%) cells and <88% for buccal (84.1%) and FOM (87.6%) cells.

Discussion

The results of this study demonstrate the promising antimicrobial potential of AL and AU against oropharyngeal NG. Gonorrhoea represents a major global public health challenge due to increasing AMR, including resistance to the last-line agent ceftriaxone,[2] and the suboptimal efficacy in the oropharynx also of newer drugs such as zoliflodacin and gepotidacin (91.3% and 87.5% respectively).[3, 4] Other proposed interventions, e.g. antimicrobial mouthwashes,[8] doxyPEP,[9] 4CMenB vaccines,[10] have shown to be ineffective, underscoring the urgent need for alternative therapeutic strategies. Because many oropharyngeal NG infections are asymptomatic and many clear spontaneously within three months,[13] three-monthly screening recommendations would often fail to detect and treat infections and transmission will have already occurred. A topical preventative product around the time of oral sex would therefore represents a novel intervention.

However, for a topical intervention to be effective, a clear understanding of NG colonization within the oropharynx is essential to ensure adequate exposure to the proposed fatty acids. Significant gaps remain regarding NG colonization and growth sites in the oropharynx.[21]

Early studies suggested tonsillar involvement but reported inconsistent findings for other oropharyngeal regions[11] and provided limited data on intracellular invasion. Building on our prior observations in three oropharyngeal cell lines,[20] the present study assessed five distinct oropharyngeal cell types and identified strain-dependent differences in invasion. FA1090 showed rapid intracellular invasion in tonsillar, buccal, and gingival cells within 15 minutes, with lower invasion in FOM and uvular cells. These findings challenge

assumptions that the tonsils are the primary site of oropharyngeal infection.[13] They highlight the need for pharmacokinetic studies to assess drug distribution across infected sites and ensure effective exposure to maximize cure and minimize AMR emergence.[5] A proposed controlled human infection model may further clarify colonization patterns and support evaluation of new interventions.[22]

Previous studies have shown that free fatty acids and monoglycerides possess antimicrobial activity against *Neisseria gonorrhoeae*, primarily demonstrated in simplified in vitro systems lacking host-cell components and without assessment of cytotoxicity toward human tissues. [23] In contrast, the present study evaluates arginine-conjugated fatty acid derivatives within a 2D human oropharyngeal epithelial infection model, enabling simultaneous assessment of antimicrobial efficacy and host cell compatibility. This approach provides a more physiologically relevant framework and highlights the potential of these compounds for topical application in oropharyngeal infection. Fatty acids have also been explored as potential prophylactics, including for prevention of ophthalmia neonatorum, highlighting relevance beyond genital infection.[24] Building on this rationale, we evaluated two novel arginine-conjugated fatty acid derivatives, AL and AU, designed for topical application.[19] In post-exposure experiments, both compounds showed time- and dose-dependent antimicrobial activity against both NG strains across all oropharyngeal cell types, with AL demonstrating superior efficacy against FA1090 compared with AU.

In pre-exposure experiments, mimicking topical use before oral sex, AL at the maximum non-cytotoxic concentration (300 $\mu\text{g/mL}$) cleared >94% of FA1090 across all cell lines after 30 minutes, with slightly lower clearance in buccal cells (83.3%). In post-exposure experiments, AL at 150 $\mu\text{g/mL}$ cleared >94% of FA1090 in all cell lines except uvular cells (61.8%). Together, these data suggest that a topical regimen using AL at 300 $\mu\text{g/mL}$ before and after oral sex may be optimal, assuming salivary dilution does not reduce concentrations below 150 $\mu\text{g/mL}$. Pre-exposure efficacy is further supported by the rapid intracellular invasion of FA1090 observed within 15 minutes in our 2D human oropharyngeal infection model.

Although pre-exposure experiments focused on the susceptible reference strain FA1090, post-exposure assays demonstrated substantial clearance of the multidrug-resistant WHO R strain ($\geq 90\%$ at 30 minutes in most cell lines, except FOM cells [79.3%]), suggesting activity against circulating resistant gonococci of growing global concern.

Regarding delivery, mucoadhesive formulations such as medicated chewing gums may offer prolonged mucosal contact and sustained local drug concentrations; however, challenges remain,[25] and additional delivery approaches such as sprays may be required to ensure coverage of the posterior oropharynx.[26]

Safety is another essential consideration for any oral prophylactic product to be addressed. In our experimental setting, AL and AU showed no cell cytotoxicity at concentrations up to 300 $\mu\text{g/mL}$, with only moderate toxicity observed above 400 $\mu\text{g/mL}$, but only in gingival cells. This favorable safety profile aligns with the GRAS classification of many fatty acids

and addresses a gap in previous research, which often lacked cytotoxicity assessments in mucosal epithelial cells.[14, 23, 27] Importantly, the antimicrobial effects observed in our study were not attributable to acidification, as the culture medium pH remained within physiological range. This finding is consistent with prior reports showing that fatty acids retain antimicrobial activity under buffered conditions, supporting a membrane-disruptive rather than pH-dependent mechanism.[23] Future studies are needed to examine the impacts on healthy oral microbiota. For example, the Belgium PreGo study examining the impact of Listerine Cool Mint® mouthwash on NG infection, also found a negative effect on healthy oral microbiota.[28] It is also worth highlighting that many lifestyle behaviors can also have a negative impact on oral health[29] and the use of a topical fatty acid should be placed in this context and a topical fatty acid is likely to have substantial benefits over a systemic antibiotic. On the other hand, one of the most significant advantages of fatty acid-based antimicrobials over conventional antibiotics is their low likelihood of resistance development, given their broad and multi-target membrane-disruptive activity.[30] Lastly, our pre-exposure experiments also provided some insights into the pharmacokinetics of tested fatty acids. As cells were washed before they were exposed to NG, any subsequent reduction in intracellular CFUs is likely from fatty acids within the cells, suggesting these fatty acids were able to cross the oral cell membranes. This is encouraging as the fatty acids were conjugated to arginine to increase their water solubility, which can reduce passive diffusion of these fatty acids across lipid-based cell membranes in human tissues. More studies are needed to understand if intracellular uptake was by active or passive drug transport systems or if the fatty acids block cellular uptake of NG.

The present study has limitations. Although our 2D human oropharyngeal infection model uses a normal epithelial cell line and improves upon cancer-derived models, it does not fully recapitulate *in vivo* physiology, including salivary flow, immune components, and epithelial stratification. Advanced 3D mucosal models will be important to validate these findings and inform product development.

In conclusion, AL and AU demonstrate rapid, potent, and site-specific antimicrobial activity against oropharyngeal NG, including AMR strains, with favorable cytotoxicity profiles and a low predicted risk of AMR development. Together, these findings position AL and AU as strong candidates for future topical interventions aimed at preventing or treating oropharyngeal gonorrhoea. Further translational studies, formulation development, and clinical validation are warranted.

Acknowledgments

We would like to acknowledge The Peter Doherty Institute for Infection and Immunity for providing the NG FA1090 strain and Marshall Centre for Infectious Disease for providing the *N. oralis* used in these experiments. We would also like to acknowledge the Prof. John Silke and Ms. Natasha Silke from Walter and Eliza Hall Institute (WEHI), for providing the SV40 virus, EGF expression proteins, and DNA verification protocols.

Funding

The project is supported by the Australian Research Council Industrial Transformation Research Hub to combat Antimicrobial Resistance (Project ID IH190100021; 2022-2206). F.Y.S.K and J.S.H are funded by a National Health and Medical Research Council Grant (GNT2033078, 2025960). Syed Ameer Hamza is supported by a Melbourne Research Scholarship, Rowden White Scholarship, MDS internal grant, and Brian and Ann Dooley Bequest from the University of Melbourne.

Declaration of interests

A.C. and F.Y.S.K have received funding from Wintermute Biomedical to screen their compounds. A.M. and T.R. are employees of Wintermute Biomedical. Other authors declare no competing interest.


References


1. Unemo, M., et al., *Gonorrhoea*. Nat Rev Dis Primers, 2019. **5**(1): p. 79.
2. Unemo, M., et al., *WHO global gonococcal antimicrobial surveillance programmes, 2019-22: a retrospective observational study*. Lancet Microbe, 2025. **6**(10): p. 101181.
3. Ross, J.D.C., et al., *Oral gepotidacin for the treatment of uncomplicated urogenital gonorrhoea (EAGLE-1): a phase 3 randomised, open-label, non-inferiority, multicentre study*. Lancet, 2025. **405**(10489): p. 1608-1620.
4. Luckey, A., et al., *Zoliflodacin versus ceftriaxone plus azithromycin for treatment of uncomplicated urogenital gonorrhoea: an international, randomised, controlled, open-label, phase 3, non-inferiority clinical trial*. The Lancet, 2025.
5. Kong, F.Y.S., et al., *Optimisation of treatments for oral Neisseria gonorrhoea infection: Pharmacokinetics Study (STI-PK project) – study protocol for non-randomised clinical trial*. BMJ Open, 2022. **12**(11).
6. Laumen, J.G.E., et al., *Antimicrobial susceptibility of commensal Neisseria in a general population and men who have sex with men in Belgium*. Sci Rep, 2022. **12**(1): p. 9.
7. Cornelisse, V.J., et al., *Risk factors for oropharyngeal gonorrhoea in men who have sex with men: an age-matched case-control study*. Sex Transm Infect, 2018. **94**(5): p. 359-364.
8. de Vries, H.J.C. and M.F.S. van der Loeff, *Antiseptic mouthwashes against sexually transmitted infections*. The Lancet Infectious Diseases, 2021. **21**(5): p. 583-584.
9. Traeger, M.W., et al., *Doxycycline Postexposure Prophylaxis and Bacterial Sexually Transmitted Infections Among Individuals Using HIV Preexposure Prophylaxis*. JAMA Intern Med, 2025. **185**(3): p. 273-281.
10. Molina, J.M., et al., *Doxycycline prophylaxis and meningococcal group B vaccine to prevent bacterial sexually transmitted infections in France (ANRS 174 DOXYVAC): a multicentre, open-label, randomised trial with a 2 x 2 factorial design*. Lancet Infect Dis, 2024. **May 23:S1473-3099(24)00236-6**.
11. Veien, N.K., E. From, and S.A. Kvorning, *Microscopy of tonsillar smears and sections in tonsillar gonorrhoea*. Acta Otolaryngol, 1976. **82**(5-6): p. 451-4.
12. van der Veer, B.M.J.W., et al., *Men and Women Have Similar Neisseria gonorrhoeae Bacterial Loads: a Comparison of Three Anatomical Sites*. Journal of Clinical Microbiology, 2020. **58**(11): p. 10.1128/jcm.01171-20.

13. Chow, E.P.F., C.K. Fairley, and F.Y.S. Kong, *STI pathogens in the oropharynx: update on screening and treatment*. Current Opinion in Infectious Diseases, 2024. **37**(1): p. 35-45.
14. Casillas-Vargas, G., et al., *Antibacterial fatty acids: An update of possible mechanisms of action and implications in the development of the next-generation of antibacterial agents*. Prog Lipid Res, 2021. **82**: p. 101093.
15. Desbois, A.P. and V.J. Smith, *Antibacterial free fatty acids: activities, mechanisms of action and biotechnological potential*. Applied Microbiology and Biotechnology, 2010. **85**(6): p. 1629-1642.
16. Davies, J. and D. Davies, *Origins and evolution of antibiotic resistance*. Microbiol Mol Biol Rev, 2010. **74**(3): p. 417-33.
17. Matsue, M., et al., *Measuring the Antimicrobial Activity of Lauric Acid against Various Bacteria in Human Gut Microbiota Using a New Method*. Cell Transplant, 2019. **28**(12): p. 1528-1541.
18. Rossi, A., et al., *Reassessing the Use of Undecanoic Acid as a Therapeutic Strategy for Treating Fungal Infections*. Mycopathologia, 2021. **186**(3): p. 327-340.
19. Mayfosh, A.J., et al., *GS-2: A Novel Broad-Spectrum Agent for Environmental Microbial Control*. Biomolecules, 2022. **12**(9).
20. Hamza, S.A., et al., *Development and validation of a human 2D in vitro model of oral Neisseria gonorrhoeae infection*. Sexually Transmitted Infections, 2025.
21. Liljemark, W. and R. Gibbons, *Ability of Veillonella and Neisseria species to attach to oral surfaces and their proportions present indigenously*. Infection and immunity, 1971. **4**(3): p. 264-268.
22. Williams, E., et al., *Rationale and Ethical Assessment of an Oropharyngeal Gonorrhea Controlled Human Infection Model*. The Journal of Infectious Diseases, 2025. **231**(4): p. 841-848.
23. Bergsson, G., O. Steingrímsson, and H. Thormar, *In vitro susceptibilities of Neisseria gonorrhoeae to fatty acids and monoglycerides*. Antimicrob Agents Chemother, 1999. **43**(11): p. 2790-2.
24. Churchward, C.P., et al., *Prevention of Ophthalmia Neonatorum Caused by Neisseria gonorrhoeae Using a Fatty Acid-Based Formulation*. mBio, 2017. **8**(4).
25. Chaudhary, S.A. and A.F. Shahiwala, *Medicated chewing gum – a potential drug delivery system*. Expert Opinion on Drug Delivery, 2010. **7**(7): p. 871-885.
26. Maddaford, K., et al., *Sites in the oropharynx reached by different methods of using mouthwash: clinical implication for oropharyngeal gonorrhoea prevention*. Sexually Transmitted Infections, 2020. **96**(5): p. 358.
27. Palace, S.G., et al., *Identification of bile acid and fatty acid species as candidate rapidly bactericidal agents for topical treatment of gonorrhoea*. J Antimicrob Chemother, 2021. **76**(10): p. 2569-2577.
28. Laumen, J.G.E., et al., *The effect of daily usage of Listerine Cool Mint mouthwash on the oropharyngeal microbiome: a substudy of the PReGo trial*. J Med Microbiol, 2024. **73**(6).
29. Papavergos, M., *Lifestyle matters: holistic oral health for patients and practitioners*. BDJ Team, 2025. **12**(8): p. 380-381.
30. Zang, M., et al., *The Impact of Omega-3 Fatty Acids on the Evolution of Acinetobacter baumannii Drug Resistance*. Microbiology Spectrum, 2021. **9**(3): p. e01455-21.

Figure Legends:

Figure 1: Fluorescence microscopy confirming *Neisseria gonorrhoeae* invasion of buccal mucosa cells. Epithelial cells were infected with *N. gonorrhoeae* (FA1090 and WHO-R strains) and treated with arginine-undecanoate (AU) and arginine-laurate (AL) at 1×, 2×, and 3× MIC. Cells were stained for membranes (WGA–Alexa Fluor 594, red), nuclei (DAPI, blue), and gonococci (NGab–FITC, green). Representative 3× images (AAA, BBB, etc.) were acquired using the Invitrogen™ FLoid™ Cell Imaging Station. Bright-field images were captured at 80× magnification with 21% brightness, and corresponding fluorescence images were captured at 80× magnification with 40% brightness.









Figure 4: A: Clearance of FA1090 in oropharyngeal cells using arginine-undecanoate (AU) and arginine-laurate (AL) at MIC 1, 2, 3 and 150 μ g/mL; B: Clearance of WHO-R in oropharyngeal cells using AU and AL at MIC 1, 2, and 3; C: Pre-exposure clearance of FA1090 in oropharyngeal cells using AU and AL at MIC 300 μ g/mL.

Data are normalized and presented as mean $\pm$ SD. Statistical significance ($p < 0.05$) is denoted as follows: *compared to control, # indicates dose dependence, and ^ indicates time dependence.



Supplementary Material

Table of contents

Supplementary Methods	17
BACTERIA AND ORAL EPITHELIAL CELLS CO-CULTURE	17
Neisseria gonorrhoeae bacterial strains	17
Uvular cells isolation, Ethics and immortalization	Error! Bookmark not defined.

Ethics	4
Isolation and immortalization of primary human oropharyngeal cells (uvular).....	18
Supplementary Table 1	18
Supplementary Table 2	19
Supplementary Table 3	20
Supplementary Table 4A	20
Supplementary Table 4B.....	21
Supplementary Table 4C	21
Supplementary Figure 1.....	Error! Bookmark not defined.
Supplementary Figure 2.....	Error! Bookmark not defined.
Supplementary Figure 3.....	Error! Bookmark not defined.

Supplementary Methods

BACTERIA AND ORAL EPITHELIAL CELLS CO-CULTURE

Neisseria gonorrhoeae bacterial strains

Three well-characterised *Neisseria* strains were used; NG FA1090 and WHO R/FC428,^{21,22} and *N. oralis* (commensal species as negative control). Bacteria were maintained in mFB and harvested during exponential growth for infection assays. Inocula were prepared in tissue culture media (supplementary Table 1), and bacterial counts were verified by plating on chocolate agar supplemented with 1% IsoVitalex. Clearance of FA1090 was the focus of our experiments.

Co-culture / infection assay

Oropharyngeal cells were seeded in 48-well plates (CLS-3338, Corning®, Sigma-Aldrich, Australia) in triplicate: uvular, gingival, and FOM cells at 3×10^4 cells/well, and tonsillar and buccal cells at 6×10^4 cells/well. Upon reaching 60–80% confluence ($\sim 1 \times 10^5$ cells/well), cells were infected with NG (FA1090 or WHO R) or *N. oralis* at a multiplicity of infection (MOI) of 1:15 for 1 hour. For pre-exposure treatment, oropharyngeal cells were incubated with AU and AL (300 $\mu\text{g/mL}$ – the maximum non-cytotoxic concentration) for 30 or 120 minutes before infection. For post-infection treatment, cells were infected and then treated with AU or AL at 1 $\times$, 2 $\times$, or 3 $\times$ their MIC₉₀ for 30, 60, or 120 minutes. For FA1090, an additional concentration (150 $\mu\text{g/mL}$) for AL was tested post-infection to match the concentration used for WHO R experiments. To remove extracellular bacteria, cells were washed with PBS and treated with gentamicin (500 $\mu\text{g/mL}$) for 1 hour, followed by lysis with 1% saponin for 15 minutes. Lysates were plated on agar, and intracellular CFUs were quantified.

The pH of cell culture media containing AU and AL (50–500 $\mu\text{g/mL}$) was measured using a calibrated benchtop pH meter (EUTECH Instruments, Thermo Fisher Scientific, USA), calibrated with standard buffers (pH 4.0, 7.0, 10.0) (supplementary table 2) to determine if they changed the media pH.

Isolation and immortalization of primary human oropharyngeal cells (uvular)

Fresh tissue was collected from a patient undergoing standard care pharyngeal palatoplasty at Masada Private Hospital, Ramsay Health, St Kilda East, Victoria and placed in DMEM with 10% fetal bovine serum (FBS) for transportation to lab. Samples were then washed in PBS containing 2% penicillin-streptomycin and 500 µg/mL gentamicin for 10 min, minced into ~2–3 mm pieces, and incubated overnight at 4°C in 2.4 U/mL Dispase II (7105-041, Gibco; Life Technologies™). Epidermal sheets were digested with 0.25% trypsin at 37°C for 10 min, neutralized with growth medium, filtered through a 40 µm strainer, centrifuged at 200xg, and seeded in keratinocyte serum-free medium (K-SFM) containing 2% penicillin-streptomycin. At ~60% confluence, primary cells were spin-infected with SV40 virus (4 µg/mL polybrene, 2500 rpm, 90 min, 30°C) and expanded. Successful immortalization was confirmed by doxycycline-inducible EGFP expression (GFP fluorescence, supplementary figure 2) and by PCR detection of SV40 sequences in genomic DNA (DNeasy Kit, Qiagen; GoTaq® Green MasterMix, Promega). PCR products were resolved on a 1% agarose gel in 1X TAE at 90 V. Representative SV40-specific bands were observed at passages 10 and 25 (supplementary figure 3)

Supplementary Table 1

Characteristics and subculturing procedures of the human oral cells used in experiments					
Cell line name	Origin/ Company	Product code	Anatomical site	Type	Media characteristics and subculturing procedures
OKF6	Oral Health Cooperative Research Centre (OHCRC), The University of Melbourne, Australia	OKF6	Floor of the mouth	Immortalised normal epithelial keratinocytes	Grown to 60–80% confluence before being further subcultured. OKF6 cells were cultured using keratinocyte serum-free medium (K-SFM) (#17005-042, Thermo Fisher Scientific, Scoresby, VIC, Australia) containing 25 µg/mL bovine pituitary extract and 0.2 ng/mL human recombinant epidermal growth factor (as per manufacturer's instructions), 0.4 mM CaCl ₂ , and 1% penicillin streptomycin mixture (P4333, Sigma-Aldrich, Castle Hill, NSW, Australia). OKF6 cells were incubated at 37°C, 5% CO ₂ for 5 to 7 days to reach 80% confluency. Epithelial cells grown to 80% confluency were subsequently detached via a pre-treatment of 10 mM EDTA for 10 min, followed subsequently with incubation with a 0.25% trypsin in 1mM EDTA solution (T4049, Sigma-Aldrich, Castle Hill, NSW, Australia) for 3-5 min.
Phk (Uvular)	Melbourne Dental school, The University of Melbourne, Australia	Phk-05	Uvula	Immortalised normal epithelial keratinocytes	Grown to 60–80% confluence before being further subcultured. PhK cells were cultured using keratinocyte serum-free medium (K-SFM) (#17005-042, Thermo Fisher Scientific, Scoresby, VIC, Australia) containing 25 µg/mL bovine pituitary extract and 0.2 ng/mL human recombinant epidermal growth factor (as per manufacturer's instructions), 0.4 mM CaCl ₂ , and 1% penicillin streptomycin mixture (P4333, Sigma-Aldrich, Castle Hill, NSW, Australia). OKF6 cells were incubated at 37°C, 5% CO ₂ for 5 to 7 days to reach 80% confluency. Epithelial cells grown to 80% confluency were subsequently detached via a pre-treatment of 10 mM EDTA for 10 min, followed subsequently with incubation with a 0.25% trypsin in 1mM EDTA solution (T4049, Sigma-Aldrich, Castle Hill, NSW, Australia) for 3-5 min.
hTERT TIGs	ATCC, Australia	CRL-3397™	Gingiva/	hTERT-immortalised	Grown to 60–80% confluence before being further subcultured. Gingival cells were cultured using

			alveolar process	normal epithelial keratinocytes	ATCC gingival media (ATCPCS200030, InVitro technologies, Australia) and respective growth supplements as per manufacturer's instructions (ATCPCS200040, InVitro technologies, Australia) containing 1% penicillin streptomycin mixture (P4333, Sigma-Aldrich, Castle Hill, NSW, Australia). Gingival cells were incubated at 37°C, 5% CO ₂ for about 3-5 days to reach 80% confluency. Epithelial cells grown to 80% confluency were subsequently detached via a pretreatment of 10 mM EDTA for 5 min, followed subsequently with incubation with a 0.25% trypsin in 1mM EDTA solution (T4049, Sigma-Aldrich, Castle Hill, NSW, Australia) for 1-3 min.
HTepiC	ScienCell Research Laboratories, Wangara, Western Australia, Australia	#2560	Tonsil	Primary normal epithelial keratinocytes	Grown to 60–80% confluence before being further subcultured. Tonsil cells were cultured using TEpiCM (2561, ScienCell) basal media and respective growth supplements as per manufacturer's instructions (#2301, ScienCell, Wangara, Western Australia, Australia) containing 1% penicillin streptomycin mixture (P4333, Sigma-Aldrich, Castle Hill, NSW, Australia). Tonsil cells were incubated at 37°C, 5% CO ₂ for about 7 days to reach 80% confluency. Epithelial cells grown to 80% confluency were subsequently detached via a pretreatment of 10 mM EDTA for 3-5 min, followed subsequently with incubation with a 0.25% trypsin in 1mM EDTA solution (T4049, Sigma-Aldrich, Castle Hill, NSW, Australia) for 1 min.
Buccal	ScienCell Research Laboratories, Wangara, Western Australia, Australia	#2610	Buccal	Primary normal epithelial keratinocytes	Grown to 60–80% confluence before being further subcultured. Buccal cells were cultured using Oral Keratinocytes medium (2611, ScienCell) basal media and respective growth supplements as per manufacturer's instructions (#2652, ScienCell, Wangara, Western Australia, Australia) containing 1% penicillin streptomycin mixture (P4333, Sigma-Aldrich, Castle Hill, NSW, Australia). Buccal cells were incubated at 37°C, 5% CO ₂ for about 7 days to reach 80% confluency. Epithelial cells grown to 80% confluency were subsequently detached via a pretreatment of 10 mM EDTA for 3-5 min, followed subsequently with incubation with a 0.25% trypsin in 1mM EDTA solution (T4049, Sigma-Aldrich, Castle Hill, NSW, Australia) for 1 min.

Supplementary Table 2

pH measurements of culture media containing fatty acid-based compounds (AU and AL)				
	Tonsillar	Buccal	Floor of mouth / Uvular	Gingival
Arginine Laurate / Arginine undecanoate (µg/mL)				
500	7.3/7.26	7.42/7.39	7.3/7.16	7.18/7.16

400	7.31/7.26	7.42/7.39	7.21/7.16	7.19/7.16
300	7.35/7.27	7.42/7.39	7.25/7.17	7.15/7.14
200	7.33/7.28	7.35/7.36	7.27/7.28	7.18/7.20
150	7.29/7.25	7.27/7.25	7.29/7.25	7.21/7.17
50	7.31/7.27	7.13/7.15	7.09/7.17	7.11/7.16

Supplementary Table 3

Screening of five fatty acid for cytotoxic effects in representative oropharyngeal epithelial cell lines			
Compounds $\mu\text{g/mL}$	Gingival	Floor of mouth	Buccal
Arginine undecanoate (undecanoic Acid)	-	-	-
Arginine laurate (lauric acid)	-	-	-
Arginine-linoleate (linoleic acid)	+	+	+
Arginine undecylenate (undecylenic acid)	-	-	+
Arginine monodecanoate (decanoic acid)	-	+	+

“+” stands for cytotoxic to oropharyngeal epithelial cell lines and “-” stands for non-cytotoxic to oropharyngeal epithelial cell lines.

Supplementary Table 4A

Post exposure clearance of FA1090 in oropharyngeal cells using AU and AL at MIC 1,2,3 and 150µg/mL								
Cell line	AU				AL			
Time (30min)								
	MIC 1 (21 µg/mL)	MIC 2 (42 µg/mL)	MIC 3 (63 µg/mL)	MIC 7 (150 µg/mL)	MIC 1 (23 µg/mL)	MIC 2 (46 µg/mL)	MIC 3 (69 µg/mL)	MIC 7 (150 µg/mL)
Tonsillar	84.0	85.1	89.4	95.8	86.3	90.6	95.9	98.3
Buccal	41.5	64.3	48.5	61.9	63.2	79.3	74.2	94.1
FOM	52.3	88.7	88.7	93.0	75.6	84.1	90.8	95.6
Uvular	48.7	41.8	66.7	64.0	54.5	50.9	65.7	61.8
Gingival	52.5	71.9	78.1	76.6	82.9	91.7	93.3	97.0
Time (60min)								
Tonsillar	89.2	88.0	94.1	96.5	87.6	92.2	97.5	99.5
Buccal	60.8	65.9	83.6	94.6	72.1	81.5	80.6	93.5
FOM	82.2	90.6	93.6	95.1	83.0	92.5	96.4	96.3

Uvular	49.7	70.1	70.0	80.8	50.4	68.4	50.7	68.0
Gingival	60.2	75.7	72.6	74.5	92.8	92.8	91.3	100
Time (120min)								
Tonsillar	96.6	95.9	95.3	97.5	96.3	96.8	98.6	99.6
Buccal	74.5	75.8	85.7	95.4	93.0	95.1	95.9	99.1
FOM	90.2	98.7	98.9	99.8	91.9	91.4	94.7	99.5
Uvular	54.6	84.9	95.3	96.8	92.9	94.1	90.4	99.1
Gingival	92.7	91.9	97.8	97.9	93.6	94.8	93.2	100

Green presenting $\geq 95\%$ clearance and orange representing 90-94.9% clearance

Supplementary Table 4B

Post exposure clearance of WHO-R in oropharyngeal cells using AU and AL at MIC 1,2, and 3						
Cell line	AU			AL		
Time (30min)						
	MIC 1 (36 µg/mL)	MIC 2 (72 µg/mL)	MIC 3 (108 µg/mL)	MIC 1 (49 µg/mL)	MIC 2 (98 µg/mL)	MIC 3 (147 µg/mL)
Tonsillar	82.3	89.5	82.6	87.3	91.9	97.8
Buccal	84.9	91.7	82.1	91.0	92.6	90.0
FOM	91.6	93.6	94.1	90.3	90.0	79.3
Uvular	77.5	77.5	87.6	87.0	75.8	97.7
Gingival	84.4	82.9	80.9	78.6	96.5	100
Time (60min)						
Tonsillar	84.8	90.9	84.5	92.3	88.6	100
Buccal	87.9	86.2	92.0	85.3	88.2	95.6
FOM	92.4	93.1	92.1	81.7	86.1	94.0
Uvular	71.3	84.2	81.4	69.6	73.5	96.0
Gingival	82.9	84.1	75.6	86.8	100	100
Time (120min)						
Tonsillar	87.2	89.6	89.3	94.8	100	100
Buccal	93.3	96.5	94.2	68.7	79.0	96.7
FOM	90.1	95.7	97.1	93.4	95.5	99.7
Uvular	91.0	95.5	96.0	81.4	91.0	100
Gingival	91.8	93.1	88.8	96.7	100	100

Green presenting $\geq 95\%$ clearance and orange representing 90-94.9% clearance


Supplementary Table 4C

Pre exposure clearance of FA1090 in oropharyngeal cells using AU and AL at 300 $\mu\text{g/mL}$		
Cell line	AU	AL
Time (30min)		
	MIC (300 $\mu\text{g/mL}$)	MIC (300 $\mu\text{g/mL}$)
Tonsillar	73.3	97.4
Buccal	55.2	83.3
FOM	52.5	94.4
Uvular	93.3	97.4
Gingival	88.3	99.2
Time (120min)		
Cell line	MIC (300 $\mu\text{g/mL}$)	MIC (300 $\mu\text{g/mL}$)
Tonsillar	94.3	98.7
Buccal	84.1	99.1
FOM	87.6	97.5


Uvular	98.3	97.4
Gingival	99.7	99.9


Green presenting $\geq 95\%$ clearance and orange representing 90-94.9% clearance

Supplementary Figure 1: Representative micrographs of oropharyngeal cells. Images (FLoid cell imaging system, Life Technologies) with magnification at 80x. Scale bar of 125 μ m applies to all panels.



Supplementary Figure 2: Isolation and immortalization of Primary Human Oropharyngeal Cells (Uvular Tissue): Panels A–E show representative images confirming successful immortalization of primary oropharyngeal keratinocytes using a doxycycline-inducible EGFP lentiviral system. Cells were immortalized with SV40 virus and subsequently transduced with an EGFP-expressing virus. EGFP expression was induced with 0.1% doxycycline, while the control group received no doxycycline. GFP fluorescence confirmed successful gene expression. Images were captured using the Invitrogen™ FLoid™ Cell Imaging Station. Bright-field images were taken at 80 $\times$ magnification with 21% brightness; corresponding green fluorescence images were acquired at 80 $\times$ magnification with 40% brightness.





Supplementary Figure 3: PCR detection of SV40 expression in immortalized uvula cells. PCR amplification of the SV40 large T antigen was performed using genomic DNA from uvula-derived epithelial cells at different passages. PCR products were separated on a 1% agarose gel in 1X TAE buffer and visualized under UV illumination. Lane M: DNA ladder; Lane P1: passage 10 cells; Lane P2: passage 25 cells; Lane C: positive control (SV40 viral DNA).