



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Cameron, KL;Cheong, JLY;Spittle, AJ

Title:

Predicting poor physical activity outcomes in children born extremely preterm or extremely low birthweight

Date:

2021-01-01

Citation:

Cameron, K. L., Cheong, J. L. Y. & Spittle, A. J. (2021). Predicting poor physical activity outcomes in children born extremely preterm or extremely low birthweight. *Acta Paediatrica International Journal of Paediatrics*, 110 (1), pp.357-358. <https://doi.org/10.1111/apa.15524>.

Persistent Link:

<https://hdl.handle.net/11343/276195>

MS. KATE L CAMERON (Orcid ID : 0000-0001-5447-594X)

Article type : EBNEO Review

TITLE OF WRITE-UP

Predicting poor physical activity outcomes in children born extremely preterm or extremely low birthweight

REVIEWED BY

Kate L Cameron, DPT

Department of Physiotherapy, The University of Melbourne
Clinical Sciences, Murdoch Children's Research Institute

Prof Jeanie LY Cheong, MD

Clinical Sciences, Murdoch Children's Research Institute
Newborn Research, Royal Women's Hospital
Obstetrics and Gynaecology, The University of Melbourne

Prof Alicia J Spittle, PhD

Department of Physiotherapy, The University of Melbourne
Clinical Sciences, Murdoch Children's Research Institute
Newborn Research, Royal Women's Hospital

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/APA.15524](https://doi.org/10.1111/APA.15524)

This article is protected by copyright. All rights reserved

CORRESPONDING AUTHOR

Name: Kate L Cameron

Email: kate.cameron@mcri.edu.au

Review template

TITLE OF WRITE-UP

Predicting poor physical activity outcomes in children born extremely preterm or extremely low birthweight

MANUSCRIPT CITATION

Engan M, Engeseth MS, Fevang S, Vollsæter M, Eide GE, Røksund OD, Halvorsen T, Clemm H. Predicting physical activity in a national cohort of children born extremely preterm. *Early Hum Dev.* 2020; 145: 105037. PMID 32438296

REVIEWED BY

Kate L Cameron, DPT

Department of Physiotherapy, The University of Melbourne
Clinical Sciences, Murdoch Children's Research Institute

Prof Jeanie LY Cheong, MD

Clinical Sciences, Murdoch Children's Research Institute
Newborn Research, Royal Women's Hospital
Obstetrics and Gynaecology, The University of Melbourne

Prof Alicia J Spittle, PhD

Department of Physiotherapy, The University of Melbourne
Clinical Sciences, Murdoch Children's Research Institute
Newborn Research, Royal Women's Hospital

This article is protected by copyright. All rights reserved

COMMENTARY

With increased survival of children born EP/ELBW it is important to understand long-term health and development,(1) with many international cohort studies reporting poorer neurodevelopmental outcomes across multiple domains for children born EP/ELBW compared with term.(2, 3) Less is known, however, about functional outcomes, including PA participation, for children born EP/ELBW.

Using a purpose-built questionnaire at 11 years of age, Engan and colleagues found children born at term had better PA outcomes compared with children born EP/ELBW. Within the EP/ELBW group, those without NDD had better PA outcomes. Motor, cognitive and behavioural difficulties at 5 years were considered as possible predictors of PA for children born EP/ELBW, with all three predictors associated with poorer PA outcomes at 11 years.(4)

The authors specifically considered children born EP/ELBW without NDD, choosing to focus on 'more subtle problems', that are often overlooked but can have significant implications.

Previous cohort studies have measured PA related outcomes, including exercise capacity and PA minutes for children born preterm compared with term; with variable findings including; modest differences in PA time (5) and no difference in PA time but reduced exercise capacity.(6) As acknowledged by Engan and colleagues, differences in methodology and cohort characteristics make it difficult to make comparisons.(4)

The authors identified follow-up rates as a study limitation; the children without PA data from the EP/ELBW group were more likely to have NDD than children with data, which may have decreased the true magnitude of group differences.(4) A strength of this study was the consideration of PA as an outcome measure, focusing on what children are doing in a real-world context rather than how they perform under test conditions. Children born EP/ELBW without NDD have greater rates of developmental difficulties compared with children born at term, and this has implications for PA participation.

During childhood, PA participation promotes motor skill development and provides opportunity for social and emotional development. PA has physical (cardiovascular fitness, muscle strength etc.) and mental health benefits,(7) and may improve a range of health and developmental difficulties for which individuals born EP/ELBW are at greater risk than those born at term. (2, 8) This research is valuable in that it looks beyond disability or impairment in children born EP/ELBW, and towards PA participation, a construct more closely aligned with meaningful quality of life. This study contributes towards emerging research considering PA outcomes for children born EP/ELBW, as well as predictors of poor PA participation, essential for providing timely intervention. This is important in the EP/ELBW population with known risk factors for poor PA participation, such as motor impairment.(2) Clinicians working with children born EP/ELBW should consider assessing PA, and work with families to promote meaningful participation in activities preferred by the child. Further research on PA, using large prospective cohort studies with EP/ELBW and term children, are needed.(1) Where possible, objective PA measures (such as accelerometry) and

questionnaires with established psychometric properties should be used; though it is acknowledged that there is no consensus on the best way to measure PA participation.

FUNDING

Kate Cameron's PhD candidature is supported by an Australian Government Research Training Program Scholarship and the National Health and Medical Research Council of Australia Centre of Research Excellence in Newborn Medicine (#1153176); JC and AS are supported by Career Development Fellowships from the Medical Research Future Fund of Australia (#1141354) and the National Health and Medical Research Council of Australia (#1108714) respectively.

CONFLICTS OF INTEREST

None

REFERENCES,

1. Doyle LW, Anderson PJ, Battin M, Bowen JR, Brown N, Callanan C, et al. Long term follow up of high risk children: who, why and how? *BMC Pediatr.* 2014;14(1):1.
2. Spittle AJ, Cameron K, Doyle LW, Cheong JL. Motor Impairment Trends in Extremely Preterm Children: 1991-2005. *Pediatrics.* 2018.
3. Moore T, Hennessy EM, Myles J, Johnson, SJ, Draper ES, Costeloe, KL et al. Neurological and developmental outcome in extremely preterm children born in England in 1995 and 2006: the EPICure studies. *BMJ* 2012; 345: e7961: e10.1136/bmj.e7961.
4. Engan M, Engeseth MS, Fevang S, Vollsæter M, Eide GE, Røksund OD, et al. Predicting physical activity in a national cohort of children born extremely preterm. *Early Hum Dev.* 2020;145:105037.
5. Lowe J, Watkins WJ, Kotecha SJ, Kotecha S. Physical Activity and Sedentary Behavior in Preterm-Born 7-Year Old Children. *PLoS One.* 2016;11(5):e0155229.

6. Welsh L, Kirkby J, Lum S, Odendaal D, Marlow N, Derrick G, et al. The EPICure study: maximal exercise and physical activity in school children born extremely preterm. *Thorax*. 2010;65(2):165-72.
7. Timmons BW, Naylor PJ, Pfeiffer KA. Physical activity for preschool children — how much and how? *Canadian Journal of Public Health / Revue Canadienne de Sante'e Publique* 2007; 98: S122.
8. Doyle LW, Irving L, Haikerwal A, Lee K, Ranganathan S, Cheong J. Airway obstruction in young adults born extremely preterm or extremely low birth weight in the postsurfactant era. *Thorax*. 2019;74(12):1147-53.

Author Manuscript