



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Zahir, SF;Griffin, A;Veerman, JL;Magliano, DJ;Shaw, JE;Cao, KAL;Mehdi, AM

Title:

Exploring the association between BMI and mortality in Australian women and men with and without diabetes: the AusDiab study

Date:

2019-05-01

Citation:

Zahir, S. F., Griffin, A., Veerman, J. L., Magliano, D. J., Shaw, J. E., Cao, K. A. L. & Mehdi, A. M. (2019). Exploring the association between BMI and mortality in Australian women and men with and without diabetes: the AusDiab study. *Diabetologia*, 62 (5), pp.754-758. <https://doi.org/10.1007/s00125-019-4830-4>.

Persistent Link:

<https://hdl.handle.net/11343/251124>

License:

[CC BY](#)



Exploring the association between BMI and mortality in Australian women and men with and without diabetes: the AusDiab study

Syeda F. Zahir¹ · Alison Griffin² · J. Lennert Veerman³ · Dianna J. Magliano⁴ · Jonathan E. Shaw⁴ · Kim-Anh Lê Cao⁵ · Ahmed M. Mehdi¹

Received: 17 July 2018 / Accepted: 22 January 2019 / Published online: 26 February 2019
© The Author(s) 2019

Abstract

Aims/hypothesis There is conflicting evidence about the obesity paradox—the counterintuitive survival advantage of obesity among certain subpopulations of individuals with chronic conditions. It is believed that results supporting the obesity paradox are due to methodological flaws, such as collider bias. The aim of this study was to examine the association between obesity and mortality in Australian men and women. In addition, we explored whether obesity would appear to be protective if the analysis was restricted to a subpopulation with disease, and to discuss the potential role of collider bias in producing such a result.

Methods The examined cohort included 10,575 Australian adults (4844 men and 5731 women) aged 25–91 years who were recruited for the AusDiab baseline survey in 1999 and followed-up through 2014. The main predictor variable was BMI categorised as normal weight (18.5 to <25 kg/m²), overweight (25 to <30 kg/m²) and obese (≥30 kg/m²), and the outcome of interest was all-cause mortality. Hazard ratios were estimated from Cox proportional hazards regression models in the entire cohort and then in subpopulations with and without diabetes.

Results A total of 1477 deaths occurred during 145,384 person-years (median 14.6 years) of follow-up. Mortality was higher in obese than in normal-weight individuals for the full population (HR 1.18; 95% CI 1.05, 1.32). When an interaction between diabetes status and BMI category was added to the model, there was no evidence of an interaction between BMI and diabetes status ($p = 0.92$). When participants with and without diabetes were analysed separately, there was no evidence of an association between obesity and mortality in those with diabetes (HR 0.91; 95% CI 0.62, 1.33).

Conclusions/interpretation In the entire AusDiab cohort, we found a significantly higher mortality among obese participants as compared with their normal-weight counterparts. We found no difference in the obesity–mortality association between individuals with and without diabetes.

Keywords Collider bias · Diabetes · Men · Mortality · Obesity paradox · Women

Kim-Anh Lê Cao and Ahmed M. Mehdi are joint senior authors.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00125-019-4830-4>) contains peer-reviewed but unedited supplementary material, which is available to authorised users.

✉ Syeda F. Zahir
s.zahir1@uq.edu.au

✉ Ahmed M. Mehdi
a.mehdi@uq.edu.au

¹ The University of Queensland Diamantina Institute, Faculty of Medicine, The University of Queensland, Level 6, Translational Research Institute, 37 Kent Street, Woolloongabba, QLD 4102, Australia

² Independent Researcher, Brisbane, QLD, Australia

³ School of Medicine, Griffith University, Gold Coast, QLD, Australia

⁴ Baker Heart and Diabetes Institute, Melbourne, VIC, Australia

⁵ Melbourne Integrative Genomics, School of Mathematics and Statistics, University of Melbourne, Melbourne, VIC, Australia

Research in context

What is already known about this subject?

- Over the past two decades, several observational studies have reported an obesity paradox in individuals with chronic conditions, which led some researchers to question current weight management guidelines
- Few studies, and none using an Australian cohort, have explored collider bias as an explanation for the obesity paradox

What is the key question?

- What effect does restricting the analysis to individuals with diabetes have on the association between obesity and mortality?

What are the new findings?

- We found no evidence of the obesity paradox in the AusDiab cohort
- No significant difference in the obesity–mortality association was found between those with and without diabetes

How might this impact on clinical practice in the foreseeable future?

- Based on the results of our study in the AusDiab cohort, achievement of a healthy weight should remain a cornerstone of diabetes management

Abbreviations

AusDiab	Australian Diabetes, Obesity and Lifestyle Study
IFG	Impaired fasting glucose
IGT	Impaired glucose tolerance

Introduction

There is conflicting evidence on the ‘obesity paradox’, the putative survival advantage of obesity among individuals with chronic conditions [1, 2]. Current weight management guidelines recommend that a BMI <25 kg/m² should be maintained [3]. However, in light of these paradoxical findings, some researchers have advocated the need for revision of these guidelines for individuals with chronic diseases [4]. It is therefore crucial to understand whether these findings reflect true causal effects or result from methodological flaws in the studies reporting the paradox.

Although many hypotheses have been put forward to explain the counterintuitive survival advantage of obesity, the evidence is still inconclusive [5–7]. Collider bias has emerged as the most recent explanation of the obesity paradox [2, 8]. Collider bias is defined as bias due to conditioning on a variable affected by exposure and sharing common causes with the outcome. While previous studies have shown that bias caused by stratification by diabetes status might be responsible for the obesity paradox in people with chronic conditions [2, 8], no such study has been conducted in an Australian cohort. Our primary research objective was to examine the association

between obesity and mortality in Australian men and women and to explore the effect of restricting the analysis to individuals with diabetes. In addition, we explored the BMI–mortality association in individuals with and without diabetes relative to normal-weight individuals without diabetes.

Methods

Study participants The cohort included 10,575 adult participants (4844 men and 5731 women, aged 25 to 91 years) enrolled in the Australian Diabetes, Obesity and Lifestyle (AusDiab) study. A flow chart of study population derivation is given in ESM Fig. 1. Participants were followed from the date of their baseline examination until 31 December 2014, or until death if sooner. All participants provided written informed consent and the study protocol was approved by the ethics committee of the Baker Heart and Diabetes Institute. Additional details are presented in the ESM Methods.

Variables BMI at baseline was the primary exposure variable for this study. During biomedical examination, standard anthropometric measures were obtained by trained staff. Participants were classified as per WHO guidelines as normal weight (18.5 to <25 kg/m²), overweight (25 to <30 kg/m²) and obese (≥30 kg/m²). Participants were classified as having diabetes based on venous plasma glucose levels as recommended by WHO (ESM Table 1) [9], or if they were currently being treated with insulin or oral glucose-lowering drugs (see ESM variables in ESM Methods for details). The outcome of this

study was all-cause mortality, which was defined as death from any cause until 31 December 2014. Mortality status was identified by linking the AusDiab data to the Australian National Death Index.

Statistical analysis Cox proportional hazards regression was used to model the association between BMI category and all-cause mortality. Models were adjusted for sex, level of education, weekly income, smoking status, physical activity and cluster of census collection district area. The baseline hazard function in the model was stratified by age and marital status, as the proportional hazards assumption was satisfied after their inclusion.

We estimated hazard ratios for (1) the full population; (2) for participants with and without diabetes separately; (3) for men and women separately; and (4) to compare mortality for each BMI/diabetes status relative to ‘normal-weight participants without diabetes’, the addition of diabetes status to the model with an interaction term with BMI category. Evidence for effect modification of the association between BMI category and mortality by diabetes status was examined by testing the significance of this interaction term (please see ESM statistical analysis for additional details).

Sensitivity analyses included consideration of diabetes status, BMI, smoking and physical activity as time-dependent variables. Variables used in time-varying analyses were measured at baseline, and then at the two follow-up surveys (2004–2005 and 2011–2012). Additional sensitivity analyses of our final models (1) and (2) were conducted after exclusion of ever smokers; exclusion of deaths in the first 3 and 5 years of follow-up; exclusion or reclassification of individuals with impaired fasting glucose (IFG) and impaired glucose tolerance (IGT) as having diabetes. All data analyses were conducted using Stata (version 13.1, StataCorp, College Station, TX, USA) or R (version 3.4.0, <https://cran.r-project.org/src/base/R-3/R-3.4.0.tar.gz>).

Results

In total, 10,575 participants (54.2% women) were included in the analysis, of whom 860 (8.1%) had diabetes at baseline (ESM Tables 2, 3).

Table 1 Hazard ratios for all-cause mortality by BMI category for all participants and participants without and with diabetes

Population	Deaths/ <i>n</i>	BMI category: HR (95% CI)			<i>p</i> value
		Normal weight	Overweight	Obese	
Total	1438/10394	1.00	0.97 (0.87,1.09)	1.18 (1.05,1.32)	0.001
Without diabetes	1131/9557	1.00	1.00 (0.88,1.13)	1.16 (1.01,1.34)	0.06
With diabetes	307/837	1.00	0.86 (0.60,1.21)	0.91 (0.62,1.33)	0.65

Adjusted for sex, educational attainment, weekly income, smoking status, physical activity, cluster, and strata of age group and marital status

Association between BMI and all-cause mortality Participants were observed for a total of 145,384 person-years (median 14.6 years). A total of 1477 deaths occurred during follow-up, giving a mortality rate of 102 per 10,000 person-years. The number of deaths by diabetes status and BMI category are reported in ESM Table 4.

For the full population, mortality was higher in obese than in normal-weight individuals (HR 1.18; 95% CI 1.05, 1.32) (Table 1). When diabetes status and an interaction term between diabetes and BMI category were included in the model, there was no evidence of an interaction between BMI and diabetes status ($p = 0.92$) (ESM Table 5). When participants with and without diabetes were analysed separately, obesity was found to be associated with higher mortality in those without diabetes (HR 1.16; 95% CI 1.01, 1.34). However, there was no evidence of an obesity–mortality association in those with diabetes (HR 0.91; 95% CI 0.62, 1.33) (Table 1).

Similar results were obtained in sensitivity analyses when (1) time-varying covariates were included in the model; (2) analysis was restricted to never smokers; (3) after excluding deaths within the first 3 and then 5 years; (4) after excluding or reclassifying individuals with IFG and IGT as having diabetes; and when BMI was used as a continuous variable with relaxation of linearity (ESM Tables 6–10, ESM Fig. 2).

Sex-specific analysis For the full population (participants with and without diabetes), obesity appeared to be associated with higher mortality in women (HR 1.31; 95% CI 1.07, 1.61), but not in men (HR 1.10; 95% CI 0.91, 1.34) (Table 2). However, the interaction term was non-significant ($p = 0.38$), indicating no statistical evidence of a difference between men and women with regard to the association between obesity and mortality.

Discussion

Our findings illustrate that obesity was associated with higher mortality in the entire cohort, and there was no evidence of a protective effect of obesity on mortality when the analysis was restricted to individuals with diabetes. There was no statistical

Table 2 Sex-specific hazard ratios for all-cause mortality by BMI category in all participants and participants without and with diabetes

Population	Women			<i>p</i> value	Men			<i>p</i> value
	BMI category: HR (95% CI)				BMI category: HR (95% CI)			
	Normal weight	Overweight	Obese		Normal weight	Overweight	Obese	
Total	1.00	0.96 (0.81,1.15)	1.31 (1.07,1.61)	0.002	1.00	1.01 (0.85,1.20)	1.10 (0.91,1.34)	0.57
Without diabetes	1.00	0.98 (0.80,1.21)	1.32 (1.05,1.64)	0.01	1.00	1.05 (0.86,1.28)	1.08 (0.84,1.38)	0.82
With diabetes	1.00	0.79 (0.47,1.34)	0.93 (0.54,1.62)	0.56	1.00	0.80 (0.48,1.35)	0.74 (0.43,1.28)	0.55

Adjusted for sex, educational attainment, weekly income, smoking status, physical activity, cluster, and strata of age group and marital status

evidence that the obesity–mortality association was different between those with and without diabetes ($p = 0.92$ for interaction between BMI and diabetes status). However, if the study had only included participants with diabetes, these findings could have been interpreted as evidence of a lack of association between obesity and mortality in individuals with diabetes.

Studies reporting the obesity paradox have been criticised for restricting analysis to individuals with disease [2, 8], which could result in a form of selection bias termed collider bias. When analyses are conducted on a selected group of individuals (including people with diabetes and ignoring those without diabetes in the same population), conditioning on the collider (diabetes in this case) occurs, which affects the exposure–outcome association in an unpredictable way. Our study is consistent with previous studies that confirm that restricting analysis in this way could lead to an apparent protective association or loss of association between obesity and mortality. However, due to the small sample size and non-significant interaction term, we are unable to conclude that the lack of association is due to collider bias or that obesity is behaving differently in those with or without diabetes. There is some scepticism in the field about collider bias, as the factor biasing the result is unknown and hence cannot be accounted for [10]. Previous studies have demonstrated that the obesity paradox can be only partially explained by collider bias [6, 11].

While in this study it appeared that there was no association between obesity and mortality among men, the non-significant interaction term for sex indicated that the significant association between obesity and mortality in the whole population likely applied equally to men and women. The differences in the obesity–mortality association between the sexes in this study might have been due to chance. A meta-analysis previously confirmed an association between obesity and higher mortality in both men and women [12].

Strengths of our study include measured weight and height, long-term follow-up and detailed assessment of clinical and socioeconomic factors. Limitations of the study include the relatively small number of deaths, the use of BMI as a surrogate measure of adiposity, self-reported smoking status and

the inclusion of smokers in analyses. In addition, lack of data at baseline precluded adjustment for current medical history or exclusion of participants with chronic conditions, which could result in some bias due to reverse causation.

To summarise, studies reporting the obesity paradox present a confusing message for clinicians and policy makers, leading to a risk of misinforming obese individuals about healthy lifestyle management plans. In this study we found no evidence of the obesity paradox in individuals with diabetes in the AusDiab cohort, and found no difference between participants with and without diabetes with regard to the association between obesity and mortality.

Acknowledgements The AusDiab study co-ordinated by the Baker Heart and Diabetes Institute, Melbourne, VIC, Australia gratefully acknowledges the support and assistance given by:

K. Anstey (School of Psychology, University of New South Wales, Sydney, NSW, Australia); B. Atkins (Kidney Disease Prevention Unit, Monash University, Melbourne, VIC, Australia); B. Balkau (Centre for Research in Epidemiology and Population Health, Renal and Cardiovascular Epidemiology, French Institute of Health and Medical Research, France); E. Barr (Department of Clinical Diabetes and Epidemiology, Baker Heart and Diabetes Institute, Melbourne, VIC, Australia); A. Cameron (School of Health & Social Development, Deakin University, Melbourne, VIC, Australia); S. Chadban (Kidney Node Laboratory, Charles Perkins Centre, The University of Sydney, Sydney, NSW, Australia); M. de Courten (Centre for Chronic Disease, Victoria University, Melbourne, VIC, Australia); D. Dunstan (Department of Physical Activity Research, Baker Heart and Diabetes Institute, Melbourne, VIC, Australia); A. Kavanagh (Melbourne School of Population and Global Health, University of Melbourne, VIC, Australia); S. Murray (Department of Clinical Diabetes and Epidemiology, Baker Heart and Diabetes Institute, Melbourne, VIC, Australia); N. Owen (Department of Behavioural Epidemiology, Baker Heart and Diabetes Institute, Melbourne, VIC, Australia); K. Polkinghorne (Department of Nephrology, Monash Medical Centre, Melbourne, VIC, Australia); T. Welborn (Department of Endocrinology and Diabetes, Sir Charles Gairdner Hospital, Perth, WA, Australia); P. Zimmet (Department of Medicine, Monash University, Melbourne, VIC, Australia); and all the study participants.

For funding or logistical support of the AusDiab study, we are grateful to: National Health and Medical Research Council (NHMRC grants 233200 and 1007544); Australian Government Department of Health and Ageing; Abbott Australasia Pty Ltd.; Alphapharm Pty Ltd.; Amgen Australia; AstraZeneca; Bristol-Myers Squibb; City Health Centre, Diabetes Service, Canberra, ACT; Department of Health and

Community Services, NT; Department of Health and Human Services, TAS; Department of Health, NSW; Department of Health, WA; Department of Health, SA; Department of Human Services, VIC; Diabetes Australia; Diabetes Australia, NT; Eli Lilly Australia; Estate of the Late Edward Wilson; GlaxoSmithKline; Jack Brockhoff Foundation; Janssen-Cilag; Kidney Health Australia; Marian & EH Flack Trust; Menzies Research Institute, University of Tasmania; Merck Sharp & Dohme, Novartis Pharmaceuticals; Novo Nordisk Pharmaceuticals; Pfizer Pty Ltd.; the Pratt Foundation; Queensland Health; Roche Diagnostics Australia; Royal Prince Alfred Hospital, Sydney, NSW; Sanofi Aventis; Sanofi-Synthélabo; and the Victorian Government's OIS Program.

Data availability The data that support the findings of this study are available from AusDiab but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available.

Funding SFZ is supported by an Endeavour Postgraduate Scholarship (4712). KALC is supported by a National Health and Medical Research Council (NHMRC) Career Development fellowship (GNT1087415). AMM is supported by a Postdoctoral Fellowship from JDRF (3-PDF-2016-198-A-N). JES is supported by a National Health and Medical Research Council (NHMRC) Senior Research Fellowship (APP1079438). DJM is supported by a National Health and Medical Research Council (NHMRC) Senior Fellowship (APP1118161). The funding sources were not involved in the design of the study, collection, analysis, and interpretation of data, writing the manuscript or the decision to submit it for publication.

Duality of interest The authors declare that there is no duality of interest associated with this manuscript.

Contribution statement SFZ contributed to study conception and design, data acquisition, analysis and interpretation of data, and was the major contributor to writing this manuscript. AG analysed and interpreted data, wrote statistical analysis section of this manuscript and provided feedback on the entire manuscript. DJM, JES and JLV contributed to the interpretation of results and critically reviewed the manuscript. KALC and AMM were involved in study supervision, study conception, interpretation of results and drafting the manuscript. All authors read and approved the final manuscript. SFZ is responsible for the integrity of work as a whole.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

1. Carnethon MR, De Chavez PJ, Biggs ML et al (2012) Association of weight status with mortality in adults with incident diabetes. *JAMA* 308:581–590
2. Lajous M, Bijon A, Fagherazzi G et al (2014) Body mass index, diabetes, and mortality in French women: explaining away a “paradox”. *Epidemiology* 25(1):10–14. <https://doi.org/10.1097/EDE.0000000000000031>
3. Jensen MD, Ryan DH, Apovian CM et al (2014) 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults. *Circulation* 129(25 suppl 2):S102–S138. <https://doi.org/10.1161/01.cir.0000437739.71477.ee>
4. Brown RE, Kuk JL (2015) Consequences of obesity and weight loss: a devil's advocate position. *Obes Rev* 16(1):77–87. <https://doi.org/10.1111/obr.12232>
5. Lavie CJ, Milani RV (2003) Obesity and cardiovascular disease: the Hippocrates paradox? *J Am Coll Cardiol* 42(4):677–679. [https://doi.org/10.1016/S0735-1097\(03\)00784-8](https://doi.org/10.1016/S0735-1097(03)00784-8)
6. Sperrin M, Candlish J, Badrick E, Renehan A, Buchan I (2016) Collider bias is only a partial explanation for the obesity paradox. *Epidemiology* 27(4):525–530. <https://doi.org/10.1097/EDE.0000000000000493>
7. Preston SH, Stokes A (2014) Obesity paradox: conditioning on disease enhances biases in estimating the mortality risks of obesity. *Epidemiology* 25(3):454–461. <https://doi.org/10.1097/EDE.0000000000000075>
8. Banack HR, Kaufman JS (2014) The obesity paradox: understanding the effect of obesity on mortality among individuals with cardiovascular disease. *Prev Med* 62:96–102. <https://doi.org/10.1016/j.ypmed.2014.02.003>
9. World Health Organization (1999) Definition, diagnosis and classification of diabetes mellitus and its complications: report of a WHO consultation. Part 1: diagnosis and classification of diabetes mellitus. WHO, Geneva
10. Peeters A (2018) Journals should no longer accept ‘obesity paradox’ articles. *Int J Obes* 42(3):584–585. <https://doi.org/10.1038/ijo.2017.259>
11. Badrick E, Sperrin M, Buchan IE, Renehan AG (2017) Obesity paradox and mortality in adults with and without incident type 2 diabetes: a matched population-level cohort study. *BMJ Open Diabetes Res Care* 5(1):e000369. <https://doi.org/10.1136/bmjdr-2016-000369>
12. Berrington de Gonzalez A, Hartge P, Cerhan JR et al (2010) Body-mass index and mortality among 1.46 million white adults. *N Engl J Med* 363(23):2211–2219. <https://doi.org/10.1056/NEJMoa1000367>