



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Selman, C;Mainzer, R;Lee, K;Anderson, P;Burnett, A;Garland, SM;Patton, GC;Pigdon, L;Roberts, G;Wark, J;Doyle, LW;Cheong, JLY

Title:

Health-related quality of life in adults born extremely preterm or with extremely low birth weight in the postsurfactant era: A longitudinal cohort study

Date:

2023-03-30

Citation:

Selman, C., Mainzer, R., Lee, K., Anderson, P., Burnett, A., Garland, S. M., Patton, G. C., Pigdon, L., Roberts, G., Wark, J., Doyle, L. W. & Cheong, J. L. Y. (2023). Health-related quality of life in adults born extremely preterm or with extremely low birth weight in the postsurfactant era: A longitudinal cohort study. *Archives of Disease in Childhood Fetal and Neonatal Edition*, 108 (6), pp.581-587. <https://doi.org/10.1136/archdischild-2022-325230>.

Persistent Link:

<https://hdl.handle.net/11343/337223>

# Health-Related Quality of Life in Adults Born Extremely Preterm or Extremely Low Birthweight in the Post-Surfactant Era: A Longitudinal Cohort Study

## Authors

Chris Selman, MBIostats<sup>1</sup>

Rheanna M. Mainzer, PhD<sup>1</sup>

Katherine J. Lee, PhD<sup>1,2</sup>

Peter J Anderson, PhD<sup>3,4</sup>

Alice C Burnett, PhD<sup>2,3,5</sup>

Suzanne Garland, MD<sup>6,7,8</sup>

George C Patton, MD<sup>2,9</sup>

Lauren Pigdon, DPsych<sup>3,10</sup>

Gehan Roberts, PhD<sup>2,11,12</sup>

John D Wark, PhD<sup>13,14</sup>

Lex W. Doyle, MD<sup>2,3,5,6</sup>

Jeanie L.Y. Cheong, MD<sup>3,6,10</sup>, for the Victorian Infant Collaborative Study Group\*.

## Affiliations:

<sup>1</sup> Clinical Epidemiology and Biostatistics Unit, Murdoch Children's Research Institute, Melbourne, Australia

<sup>2</sup> Dept of Paediatrics, University of Melbourne, Melbourne, Australia

<sup>3</sup> Clinical Sciences, Murdoch Children's Research Institute, Melbourne, Australia

<sup>4</sup> School for Psychological Sciences, Turner Institute for Brain and Mental Health, Monash University, Melbourne, Australia

<sup>5</sup> Premature Infant Follow-Up Program, Royal Women's Hospital, Melbourne, Australia

<sup>6</sup> Dept of Obstetrics and Gynaecology, University of Melbourne, Melbourne, Australia

<sup>7</sup> Infection and Immunity, Murdoch Children's Research Institute

<sup>8</sup> Centre for Women's Infectious Diseases, Royal Women's Hospital, Melbourne, Australia

<sup>9</sup> Centre for Adolescent Health, Murdoch Children's Research Institute, Melbourne, Australia

<sup>10</sup> Newborn Research, Royal Women's Hospital, Melbourne, Australia

<sup>11</sup> Centre for Community Child Health, The Royal Children's Hospital, Melbourne, Australia

<sup>12</sup> Population Health, Murdoch Children's Research Institute, Melbourne, Australia.

<sup>13</sup> Dept of Medicine, University of Melbourne (Royal Melbourne Hospital), Melbourne, Australia.

<sup>14</sup> Bone and Mineral Medicine, Dept of Diabetes & Endocrinology, Royal Melbourne hospital, Melbourne, Australia.

\*Members listed in acknowledgements.

**Corresponding author:** Chris Selman, Level 4, CEBU, Murdoch Children's Research Institute, Parkville, VIC 3052, Australia. [chris.selman@mcri.edu.au](mailto:chris.selman@mcri.edu.au)

**Short title:** Health-Related Quality of Life in Extremely Preterm Survivors

**Conflicts of Interest:** The authors have no conflicts of interest relevant to this article to disclose.

**Funding source:** Supported by grants from the National Health and Medical Research Council of Australia (Centre of Clinical Research Excellence #546519; Centre of Research Excellence #1060733 & #1153176; Project Grant #108702; Career Development Fellowship #1127984 to KL; Investigator Grant #1176077 to PJA), the Medical Research Future Fund of Australia (Career Development Fellowship #1141354 to JC), and the Victorian Government's Operational Infrastructure Support Program. The Murdoch Children's Research Institute is supported by the Victorian Government's Operational Infrastructure Support Program.

**Role of the funder:** The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

**List of key words not in the title:** intellectual quotient, preterm birth, multiple imputation

**Ethics:** The Human Research Ethics Committees at the Royal Women's Hospital, Monash Medical Centre and Mercy Hospital for Women. HREC number is: Project 08/06 (ext). Project approved by the RHW HREC.

**Contributions:** JC, RM, LW, PA, AB, SG, GP, LP, GR, JW, KL conceived the study and CS wrote the first draft of the manuscript. JC, SG, JW acquired funding. JC was also involved with methodology, data curation, investigation, supervision/oversight and resources. RM, KL, and LW were also involved with methodology, data curation, investigation, formal analysis and supervision/oversight. GP, PA, JW, AB, SG, LP, GR provided supervision/oversight. All authors contributed to the revision of the manuscript, and take public responsibility for its content.

**Abbreviations:**

HRQoL - health-related quality of life.

HUI3 - Health Utilities Index Mark 3.

EP/ELBW - extremely preterm/extremely low birthweight.

NBW - normal birthweight.

IQ - intelligence quotient.

MD - median difference.

## **Abstract**

**Objectives:** To compare health-related quality of life (HRQoL) at 25 and 18 years in individuals born extremely preterm (EP; <28 weeks' gestation) or extremely low birthweight (ELBW; birthweight <1000 g) with term-born ( $\geq 37$  weeks) controls. Within the EP/ELBW cohort, to determine whether HRQoL differed between those with lower and higher intelligent quotients (IQ).

**Methods:** HRQoL was self-reported using the Health Utilities Index-3 (HUI3) at 18 and 25 years by 297 EP/ELBW and 251 controls born in 1991-92 in Victoria, Australia. Median differences (MD) between groups were estimated using multiple imputation to handle missing data.

**Results:** Adults born EP/ELBW had lower HRQoL (median utility 0.89) at 25 years than controls (median utility 0.93; MD -0.040), but with substantial uncertainty in the estimate (95% confidence interval [CI] -0.088, 0.008), and a smaller reduction at 18 years (MD -0.016; 95% CI -0.061, 0.029). On individual HUI3 items, there was suboptimal performance on Speech (odds ratio (OR) 9.28; 95% CI 3.09, 27.9) and Dexterity (OR 5.44; 95% CI 1.04, 28.4) in the EP/ELBW cohort. Within the EP/ELBW cohort, individuals with lower IQ had lower HRQoL compared with those with higher IQ at 25 (MD -0.031; 95% CI -0.126, 0.064) and 18 years (MD -0.034; 95% CI -0.107, 0.040), but again with substantial uncertainty in the estimates.

**Conclusions:** Compared with term-born controls, young adults born EP/ELBW reported poorer HRQoL, as did those with lower IQ compared with those with higher IQ among the EP/ELBW cohort. Given the uncertainties, our findings need corroboration.

**What's already known on this subject**

The evidence for group differences in health-related quality of life (HRQoL) in adulthood between survivors born EP/ELBW and term-born controls, and how differences in HRQoL might alter with increasing age of survivors are unclear.

**What this study adds**

In our cohort of 25-year-olds born EP/ELBW in the surfactant era, HRQoL was lower than in term-born controls, and was a larger difference than when the cohorts were 18 years of age.

**How this study might affect research, practice, or policy**

This study suggests that there may be a reduction in HRQoL in survivors born EP/ELBW compared with term-born controls in the era since surfactant has been introduced into clinical practice. However, given the uncertainties in the estimates, the findings should be interpreted with caution until further research on this vulnerable population is completed. Whether the current findings persist into later adulthood remains to be determined.

## **Introduction**

Improvements in neonatal intensive care, especially the introduction of surfactant replacement therapy, have contributed to dramatic increases in the survival of infants born extremely preterm (EP, <28 weeks' gestation) or extremely low birthweight (ELBW, <1000 g) since the early 1990s.<sup>1-3</sup> However, survivors born EP/ELBW are more likely to develop motor, cognitive, educational and socio-emotional problems than their term-born ( $\geq 37$  weeks) and normal birthweight (NBW; >2499 g) peers,<sup>1-6</sup> which might affect their health-related quality of life (HRQoL). The evidence for differences in HRQoL between individuals born EP/ELBW and at term in adulthood is less clear,<sup>7-8</sup> with most studies derived from cohorts born prior to the introduction of surfactant in the 1990s which did not include the most immature EP/ELBW infants due to lower survival rates in earlier eras. In addition, it is unclear whether HRQoL changes from adolescence to adulthood, when additional challenges may arise. Furthermore, individuals born EP/ELBW have lower intellectual quotients (IQs) than those born at term, which may affect their HRQoL.<sup>9-11</sup>

To address these gaps in the literature, this study aimed to (i) compare HRQoL in a geographic cohort of EP/ELBW individuals born in the era after surfactant was introduced into clinical care with term-born controls at 25 years, (ii) explore whether the differences in HRQoL between EP/ELBW and controls changed between late adolescence (18 years) and young adulthood (25 years), and (iii) determine whether individuals with a lower IQ within the EP/ELBW group reported different HRQoL at 18 and 25 years compared with those with a higher IQ. We hypothesised that adults born EP/ELBW would report worse HRQoL than controls, and, based on the literature,<sup>12-14</sup> the disparities in HRQoL between the groups would widen between 18 and 25 years of age. We also hypothesised that within the EP/ELBW group, those with lower IQ would report worse HRQoL than those with higher IQ.

## Methods

Participants comprised all survivors born EP/ELBW in the state of Victoria, Australia during 1991 and 1992 ( $n=297$ ), and matched NBW and term-born controls ( $n=251$ ) who survived to 25 years of age. Controls were randomly selected from births in the three tertiary maternity hospitals in Victoria and were matched for the expected date of birth of a survivor born EP/ELBW, the sex of the participant, whether English was the primary language spoken in the mother's country of birth, and the mother's health insurance status as a proxy for social class. Perinatal data were collected prospectively in the newborn period, as were outcome data from follow-up assessments of participants at 2, 5, 8, 18 and 25 years of age, corrected for prematurity.<sup>5</sup>

The Human Research Ethics Committees at the Royal Women's Hospital, Monash Medical Centre and Mercy Hospital for Women approved the initial and each of the follow-up studies (ethics approval for 25-year follow-up study is Project 08/06 (*ext*)). Participants gave written informed consent at ages 18 and 25 years.

Participants completed the Health Utilities Index Mark 3 (HUI3), a self-reported measure of HRQoL, at age 18 and 25 years. The HUI3 is a validated and reliable tool that can detect changes in quality of life over time,<sup>15</sup> and covers eight attributes: pain, cognition, emotion, dexterity, ambulation, speech, hearing, and vision. Each attribute is rated on a 5 or 6-point scale, with 1 representing normal function and 5 (or 6) indicative of severe impairment.<sup>16</sup> Each attribute is defined as sub-optimal if the level of function is reported as 2 or above.<sup>17-18</sup> The attribute scores are combined to an overall utility score that ranges from 0 to 1 (where 0 represents death and 1 represents perfect health), although values below 0 are possible, reflecting that some health states are identified as being worse than death. The HUI3 is an established tool that accurately measures HRQoL and is commonly used in population

health surveys.<sup>19</sup> Differences in utilities larger than 0.03 are deemed to be clinically important.<sup>20-21</sup>

Lower IQ was defined as having an IQ <-1 SD relative to the mean IQ of the term-born controls measured at age 18 years using the two-subtest version of the Wechsler Abbreviated Scale of Intelligence.<sup>22</sup>

Data were analysed using Stata Release 16.1.<sup>25</sup> The estimand of interest was the difference in median utilities between the birth groups, since the distribution of utilities was negatively skewed. Directed acyclic graphs (DAGs) were used to identify variables to control for confounding (see Supplementary Material Figure 1). Differences in median utilities at age 18 and 25 were estimated using quantile regression adjusted for the confounders of lower social class and lower maternal education. A similar approach was used to estimate the median differences in utilities between those with lower and higher IQ in the EP/ELBW group at age 18 and 25 years. Sub-optimal attributes at age 18 and 25 years were also compared between birth groups and between those with lower and higher IQ among the EP/ELBW participants using odds ratios [ORs] estimated using logistic regression, adjusted for lower social class and lower maternal education. A possible lack of independence of data from individuals arising from multiple births within the same family was accounted for by reporting all estimates with robust standard errors. In recognition of the increased potential for false positive findings that arise through the analysis of multiple outcomes, findings should be interpreted cautiously and in context with one another rather than in isolation.

Missingness DAGs were used to depict missingness assumptions and guide the treatment of missing data (see Supplementary Material Figure 2).<sup>26</sup> Based on the missingness DAG, multiple imputation was used to handle missing data in the primary analysis, although we also report the results from a complete case analysis for comparison. Imputations were generated using chained equations with 50 imputations and 20 iterations between each

imputation. Imputation was performed separately for each aim (and for the first aim, separately within each birth group) and for each attribute comparison. Imputation models included HRQoL at age 18 and 25 years, IQ, social class, maternal education, and auxiliary variables that were predictors of missingness and associated with HRQoL (further details of the multiple imputation approach are provided in the Supplementary Material)<sup>27</sup>. Estimates of the median difference [MD] and OR were obtained using Rubin's rules,<sup>28</sup> and are reported with 95% confidence intervals (CIs) and *p*-values.

As missingness in HRQoL may be associated with HRQoL itself (see Supplementary Figure 2), a sensitivity analysis was conducted where we assumed individuals with missing data had a mean utility 0.01, 0.03, 0.05, 0.10 and 0.20 lower than the individuals with observed data based on potential differences informed by clinicians. This analysis was performed using a pattern-mixture approach<sup>29</sup> (see Supplementary Material for details).

## **Results**

HRQoL data at age 25 years were available for 56% (165/297) of individuals born EP/ELBW and 52% (131/251) of controls, and at age 18 years for 65% (194/297) of individuals born EP/ELBW and 59% (148/251) of controls. Compared with adults born EP/ELBW who did not report HRQoL data at 25 years, those with data had a higher mean IQ at 8 and 18 years, and less disability at 8 years, but had similar neonatal and demographic characteristics (Table 1). Adults from the control group who provided data at 25 years were less likely to have a mother with less education or be from a lower social class. HRQoL at 8 and 18 years was similar for both EP/ELBW and control groups between those with and without HRQoL data at 25 years.

Based on the primary analysis using multiple imputation, adults born EP/ELBW reported a median utility of 0.89 at 25 years, compared with a median utility of 0.93 reported

by controls. Overall, HRQoL was lower for individuals born EP/ELBW than controls at age 25 years (MD -0.040; 95% CI -0.088, 0.008;  $p=0.10$ ). Although a clinically important difference, the width of the CI indicates large uncertainty in this estimate as it includes values in which both birth groups have similar HRQoL ( $\Delta$ ). The difference was smaller and the evidence weaker in the complete-case (Table 2) and sensitivity analyses (Supplementary Table 1). There was a smaller difference in median utilities at 18 years between EP/ELBW (0.92) and control groups (0.94; MD at 18 years -0.016; 95% CI -0.061, 0.029;  $p=0.47$ ) (Figure 1A). The EP/ELBW group were more likely to report sub-optimal function for speech (OR 5.44, 95% CI 1.04, 28.4) and dexterity (OR 9.28; 95% CI 3.09, 27.9) than the control group at 25 years of age. The disparities in these attributes between groups were less obvious at age 18 years (Table 3 and Supplementary Table 3).

Among the EP/ELBW cohort, individuals with lower IQ rated their HRQoL lower than those with higher IQ at age 25 years (median utilities 0.86 vs 0.90, respectively), but with considerable uncertainty around the estimate (MD -0.031; 95% CI -0.126, 0.064;) (Figure 1B). Similar results were obtained from the complete case (Table 2) and sensitivity analyses (Supplementary Table 1). Utilities were also lower in those with lower IQ groups at 18 years (median utilities 0.88 vs 0.93, respectively; MD -0.034; 95% CI -0.107, 0.040). In the EP/ELBW cohort, those with a lower IQ had similar odds of sub-optimal performance in all areas of the HUI3 than those with higher IQ, including in the complete-case analysis (Supplementary Table 3).

For both birth and IQ groups, there were minimal differences in attributes such as vision, hearing, pain, and emotion (Table 3).

## **Discussion**

In the era after the introduction of surfactant to clinical care, individuals born EP/ELBW in this geographical cohort study had lower overall HRQoL compared with term-born controls

at age 25 years. The median utilities indicate that there is a tendency for at least one attribute to be at a sub-optimal level in both groups, but that these may be either readily corrected or do not prevent any activities.<sup>16</sup> The median difference in utilities between the two groups at age 25 years represents a clinically important difference, and the disparity between EP/ELBW and controls increased from late adolescence, implying that a change in the management of HRQoL for adults born EP/ELBW may be necessary. However, there was substantial uncertainty in the estimates reflected in the width of the 95% CIs. Further, within the EP/ELBW group, those who had lower IQ reported lower HRQoL compared with those who had a higher IQ, again with uncertainty in the estimate, and the gap between these groups remained stable since late adolescence. Within individual areas of the HUI3, those born EP/ELBW reported more problems than controls in Speech and Dexterity at 25 years of age only, although there were minimal differences in attributes such as vision, hearing, pain and emotion.

Poorer performance between birth groups at age 25 years for speech and dexterity, but not at 18 years, might relate to the increasing demands associated with age, including needing more self-reliance for independent living. Technology and the internet have enabled individuals to become increasingly connected, allowing adults born EP/ELBW to compare their QoL with an ever-growing reference group of individuals who are similar to themselves rather than the whole population, which parents may use. Of note, several studies have shown that reports by parents and self-reports differ,<sup>13,17-18</sup> emphasising the importance of informant ratings when evaluating HRQoL in cohorts born preterm.<sup>30</sup>

There are conflicting reports on HRQoL of cohorts born preterm compared with term-born controls. In a systematic review of 18 studies reporting HRQoL outcomes in late adolescence to early adulthood of very preterm (<32 weeks' gestation) or very low birthweight (<1500 g birthweight) cohorts compared with term/NBW controls 11 studies

concluded there was little to no difference between groups, three were inconclusive, and four reported lower HRQoL in the preterm group compared with controls.<sup>7</sup> That study was not able to perform a meta-analysis due to the heterogeneity within the cohorts, and to the different measures used to assess HRQoL. All but two of the cohorts were born prior to the introduction of surfactant into clinical care; as noted above, these results may not be generalisable to cohorts born after the introduction of surfactant. Four studies in the systematic review used a version of the HUI to assess HRQoL; in 3 studies there was little evidence for group differences (one of which is the 18-year outcomes of the cohort of the present study),<sup>12 14 23</sup> while the other study reported poorer HRQoL in the preterm group than controls.<sup>13</sup> Since then, the EPICure study in the UK has reported poorer HRQoL at age 19 years in those born <26 weeks' gestation compared with term-born controls as rated by parents and participants themselves.<sup>30</sup>

Other studies have also explored differences in HRQoL in subgroups within the preterm population, especially those with disability.<sup>7</sup> The findings are also conflicting,<sup>7</sup> which may in part be explained by the different definitions of “disability”, comprising varying combinations of cerebral palsy, developmental disorders, sensorineural problems involving vision and hearing, and intellectual impairment, although no study has explored mild cognitive differences using the HUI3. In the present study, we found on average that EP/ELBW individuals with lower IQ had lower HRQoL at 18 and 25 years than EP/ELBW individuals with a higher IQ, a difference that remained stable between adolescence and young adulthood, but at both ages there was uncertainty around the estimated differences.

There are few reports of trajectories of HRQoL. Saigal et al reported trajectories of HRQoL across 3 time points, 12-16 years, 22-26 years and 29-36 years, in a Canadian cohort of ELBW participants born in the late 1970s.<sup>14</sup> ELBW participants with neurosensory impairment had poorer HRQoL than ELBW participants without neurosensory impairment,

and both of these subgroups had lower HRQoL than NBW controls at all 3 time points. There was a similar rate of decline with increasing age for all three groups. The EPICure cohort also provided evidence of a decline in parent-reported HRQoL between age 11 and 19 years.<sup>30</sup>

Strengths of our study include the geographic cohort of all infants born EP/ELBW and contemporaneously recruited term-born controls in Victoria, with follow-up to 25 years of age. Limitations relate to the attrition rate with increasing age of follow-up. Given the length of follow-up, study complexity, and considering that many families had left the state of Victoria, these follow-up rates were comparable to similar studies spanning decades.<sup>31-33</sup> The missing data in HRQoL at age 18 and 25 years were accounted for by using multiple imputation and we performed a comprehensive sensitivity analysis a range of assumptions about the missing data in these variables. Although our study is as large or larger than most others, there is still uncertainty in whether there is a clinically important difference in HRQoL between groups, reflected in the width of the 95% CI.

In an era with high survival outcomes, individuals born EP/ELBW are rating their HRQoL lower than term-born controls at age 25 years, which is a larger difference than at age 18 years. However, given the uncertainties in the estimated differences, our findings need corroboration. Longer-term surveillance is imperative, as life's challenges increase through adulthood, so that we can gain a holistic understanding of how HRQoL in survivors born EP/ELBW might change later in life.

## Acknowledgement Section

We thank the study participants and families for their involvement in this study.

Members of the Victorian Infant Collaborative Study Group

Convenor: Jeanie Cheong.<sup>1,2,3,4</sup> Collaborators (in alphabetical order): Niranjan Abraham,<sup>5</sup> Peter Anderson,<sup>2,4,6</sup> Merilyn Bear,<sup>2,4</sup> Rosemarie Boland,<sup>2,3</sup> Alice Burnett,<sup>2,4,7,8</sup> Margaret Charlton,<sup>9</sup> Marissa Clark,<sup>9</sup> Noni Davis,<sup>4</sup> Lex Doyle,<sup>1,2,3,4,7</sup> Julianne Duff,<sup>4</sup> Leah Hickey,<sup>2,7,8</sup> Emily Johnston,<sup>9</sup> Elisha Josev,<sup>2,7,10</sup> Katherine Lee,<sup>7,11</sup> Marion McDonald,<sup>4</sup> Rheanna Mainzer,<sup>11</sup> Bronwyn Novella,<sup>10</sup> Joy Olsen,<sup>2,4</sup> Gillian Opie,<sup>3,10</sup> Lauren Pigdon,<sup>2,4</sup> Gehan Roberts,<sup>2,4,7,12,13</sup> Katherine Scott,<sup>10</sup> Alicia Spittle,<sup>1,2,14</sup> Penelope Stevens,<sup>9</sup> Alice Stewart,<sup>9</sup> Anne-Marie Turner.<sup>10</sup>

<sup>1</sup> Neonatal Services, Royal Women's Hospital, Melbourne, Australia

<sup>2</sup> Clinical Sciences, Murdoch Children's Research Institute, Melbourne, Australia

<sup>3</sup> Department of Obstetrics & Gynaecology, University of Melbourne, Melbourne, Australia

<sup>4</sup> Premature Infant Follow Up Program, Royal Women's Hospital, Melbourne, Australia

<sup>5</sup> Newborn Services, Joan Kirner Women's and Children's Hospital, Melbourne, Australia

<sup>6</sup> Turner Institute for Brain and Mental Health, Monash University, Melbourne, Australia

<sup>7</sup> Department of Paediatrics, University of Melbourne, Melbourne, Australia

<sup>8</sup> Department of Neonatal Medicine, Royal Children's Hospital, Melbourne, Australia

<sup>9</sup> Department of Neonatology, Monash Medical Centre, Melbourne, Australia

<sup>10</sup> Neonatal Services, Mercy Hospital for Women, Melbourne, Australia

<sup>11</sup> Clinical Epidemiology and Biostatistics, Murdoch Children's Research Institute, Melbourne, Australia

<sup>12</sup> Centre for Community and Child Health, Royal Children's Hospital, Melbourne, Australia

<sup>13</sup> Population Health, Murdoch Children's Research Institute, Melbourne, Australia

<sup>14</sup> Department of Physiotherapy, University of Melbourne, Melbourne, Australia

## References

- [1] Saigal S, Doyle LW. An overview of mortality and sequelae of preterm birth from infancy to adulthood. *Lancet*. 2008;371(9608):261-269. doi:10.1016/S0140-6736(08)60136-1
- [2] Stoll BJ, Hansen NI, Bell EF, Walsh MC, Carlo WA, Shankaran S, et al. Trends in Care Practices, Morbidity, and Mortality of Extremely Preterm Neonates, 1993-2012. *JAMA*. 2015;314(10):1039-1051. doi:10.1001/jama.2015.10244
- [3] Joseph KS, Kramer MS, Allen AC, Cyr M, Fair M, Ohlsson A, et al. Gestational age- and birthweight-specific declines in infant mortality in Canada, 1985-94. Fetal and Infant Health Study Group of the Canadian Perinatal Surveillance System. *Paediatr Perinat Epidemiol*. 2000;14(4):332-339. doi:10.1046/j.1365-3016.2000.00298.x
- [4] Hack M, Fanaroff AA. Outcomes of children of extremely low birthweight and gestational age in the 1990's. *Early Hum Dev*. 1999;53(3):193-218. doi:10.1016/s0378-3782(98)00052-8
- [5] Cheong JLY, Wark JD, Cheung MM, Irving L, Burnett AC, Lee KJ, et al. Impact of extreme prematurity or extreme low birth weight on young adult health and well-being: the Victorian Infant Collaborative Study (VICS) 1991-1992 Longitudinal Cohort study protocol. *BMJ Open*. 2019;9(5):e030345. Published 2019 May 9. doi:10.1136/bmjopen-2019-030345
- [6] Marlow N, Wolke D, Bracewell MA, Samara M; EPICure Study Group. Neurologic and developmental disability at six years of age after extremely preterm birth. *N Engl J Med*. 2005;352(1):9-19. doi:10.1056/NEJMoa041367

[7] van der Pal S, Steinhof M, Grevinga M, Wolke D, Verrips GE. Quality of life of adults born very preterm or very low birth weight: A systematic review. *Acta Paediatr.* 2020;109(10):1974-1988. doi:10.1111/apa.15249

[8] Vederhus BJ, Markestad T, Eide GE, Graue M, Halvorsen T. Health related quality of life after extremely preterm birth: a matched controlled cohort study. *Health Qual Life Outcomes.* 2010;8:53. Published 2010 May 23. doi:10.1186/1477-7525-8-53

[9] Roberts G. Quality of life after extremely preterm birth. *Arch Dis Child.* 2019;104(4):311-312. doi:10.1136/archdischild-2018-315708

[10] Doyle LW, Spittle A, Anderson PJ, Cheong JLY. School-aged neurodevelopmental outcomes for children born extremely preterm. *Arch Dis Child.* 2021;106(9):834-838. doi:10.1136/archdischild-2021-321668

[11] Simões C, Santos S. Comparing the quality of life of adults with and without intellectual disability. *J Intellect Disabil Res.* 2016;60(4):378-388. doi:10.1111/jir.12256

[12] Saigal S, Stoskopf B, Pinelli J, Streiner D, Hoult L, Paneth N, Goddeeris J. Self-perceived health-related quality of life of former extremely low birth weight infants at young adulthood. *Pediatrics.* 2006 Sep;118(3):1140-8. 1977-82 assessed at 22-25 yrs.

[13] Baumann N, Bartmann P, Wolke D. Health-Related Quality of Life Into Adulthood After Very Preterm Birth. *Pediatrics.* 2016 Apr;137(4):e20153148.

[14] Saigal S, Ferro MA, Van Lieshout RJ, Schmidt LA, Morrison KM, Boyle MH. Health-Related Quality of Life Trajectories of Extremely Low Birth Weight Survivors into Adulthood. *J Pediatr*. 2016;179:68-73.e1. doi:10.1016/j.jpeds.2016.08.018

[15] Horsman J, Furlong W, Feeny D, Torrance G. The Health Utilities Index (HUI): concepts, measurement properties and applications. *Health Qual Life Outcomes*. 2003;1:54. Published 2003 Oct 16. doi:10.1186/1477-7525-1-54

[16] Feng Y, Bernier J, McIntosh C, Orpana H. Validation of disability categories derived from Health Utilities Index Mark 3 scores. *Health Rep*. 2009;20(2):43-50.

[17] Wolke D, Chernova J, Eryigit-Madzwamuse S, Samara M, Zwierzynska K, Petrou S. Self and parent perspectives on health-related quality of life of adolescents born very preterm. *J Pediatr*. 2013;163(4):1020-6.e2. doi:10.1016/j.jpeds.2013.04.030

[18] Ni Y, Johnson S, Marlow N, Wolke D. Reduced health-related quality of life in children born extremely preterm in 2006 compared with 1995: the EPICure Studies. *Arch Dis Child Fetal Neonatal Ed*. 2022;107(4):408-413. doi:10.1136/archdischild-2021-322888

[19] Feeny D, Furlong W, Torrance GW, Goldsmith CH, Zhu Z, DePauw S, et al. Multiattribute and single-attribute utility functions for the health utilities index mark 3 system. *Med Care*. 2002;40(2):113-128. doi:10.1097/00005650-200202000-00006

[20] Drummond M. Introducing economic and quality of life measurements into clinical studies. *Ann Med*. 2001;33(5):344-349. doi:10.3109/07853890109002088

[21] Samsa G, Edelman D, Rothman ML, Williams GR, Lipscomb J, Matchar D. Determining clinically important differences in health status measures: a general approach with illustration to the

Health Utilities Index Mark II. *Pharmacoeconomics*. 1999;15(2):141-155. doi:10.2165/00019053-199915020-00003

[22] Wechsler, D. 1999. Wechsler Abbreviated Scale of Intelligence. San Antonio, TX: The Psychological Corporation.

[23] Roberts G, Burnett AC, Lee KJ, Cheong J, Wood SJ, Anderson PJ, et al. Quality of life at age 18 years after extremely preterm birth in the post-surfactant era. *J Pediatr*. 2013;163(4):1008-13.e1. doi:10.1016/j.jpeds.2013.05.048

[24] Peart S, Cheong JLY, Roberts G, Davis N, Anderson PJ, Doyle LW, et al. Changes over time in quality of life of school-aged children born extremely preterm: 1991-2005. *Arch Dis Child Fetal Neonatal Ed*. 2021;106(4):425-429. doi:10.1136/archdischild-2020-320582

[25] StataCorp 2019. Stata Statistical Software: Release 16. College Station, TX: StataCorp LLC.

[26] Moreno-Betancur M, Lee KJ, Leacy FP, White IR, Simpson JA, Carlin JB. Canonical Causal Diagrams to Guide the Treatment of Missing Data in Epidemiologic Studies. *Am J Epidemiol*. 2018;187(12):2705-2715. doi:10.1093/aje/kwy173

[27] Morris, T.P., White, I.R. & Royston, P. Tuning multiple imputation by predictive mean matching and local residual draws. *BMC Med Res Methodol*. 2014; 14, 75.

<https://doi.org/10.1186/1471-2288-14-75>

[28] Sterne JA, White IR, Carlin JB, Spratt M, Royston P, Kenward MG, et al. Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. *BMJ*. 2009;338:b2393. Published 2009 Jun 29. doi:10.1136/bmj.b2393

[29] Tompsett DM, Leacy F, Moreno-Betancur M, Heron J, White IR. On the use of the not-at-random fully conditional specification (NARFCS) procedure in practice. *Stat Med*. 2018;37(15):2338-2353. doi:10.1002/sim.7643

[30] Ni Y, O'Reilly H, Johnson S, Marlow N, Wolke D. Health-Related Quality of Life from Adolescence to Adulthood Following Extremely Preterm Birth. *J Pediatr*. 2021 Oct;237:227-236.e5.

[31] Verrips G, Brouwer L, Vogels T, Taal E, Drossaert C, Feeny D, et al. Long term follow-up of health-related quality of life in young adults born very preterm or with a very low birth weight. *Health Qual Life Outcomes*. 2012;10:49. Published 2012 May 15. doi:10.1186/1477-7525-10-49

[32] Gray R, Petrou S, Hockley C, Gardner F. Self-reported health status and health-related quality of life of teenagers who were born before 29 weeks' gestational age. *Pediatrics*. 2007;120(1):e86-e93. doi:10.1542/peds.2006-2034

[33] van Lunenburg A, van der Pal SM, van Dommelen P, van der Pal-de Bruin KM, Bennebroek Gravenhorst J, Verrips GH. Changes in quality of life into adulthood after very preterm birth and/or very low birth weight in the Netherlands. *Health Qual Life Outcomes*. 2013;11:51. Published 2013 Mar 26. doi:10.1186/1477-7525-11-51

**Table 1:** Comparison of participants with and without HRQoL data at age 25 years

	<b>EP/ELBW</b>		<b>Term-Born Controls</b>	
	<b>HRQoL Data at age 25 <i>n</i> = 165</b>	<b>No HRQoL data at age 25 <i>n</i> = 132</b>	<b>HRQoL Data at age 25 <i>n</i> = 131</b>	<b>No HRQoL data at age 25 <i>n</i> = 120</b>
<b><i>Baseline - neonatal</i></b>				
Male - <i>n</i> (%)	74 (45%)	63 (48%)	53 (41%)	68 (57%)
Gestational age (weeks) - mean (SD)	26.6 (2.0)	26.8 (1.8)	39.3 (1.3)	39.3 (1.3)
Birthweight (g) - mean (SD)	884 (159)	894 (163)	3394 (462)	3411 (409)
Cystic PVL - <i>n</i> (%)*	7 (4%)	11 (8%)	-	-
Intraventricular hemorrhage - <i>n</i> (%)*	7 (4%)	11 (8%)	-	-
Surgery in the newborn period - <i>n</i> (%)	39 (24%)	38 (29%)	0 (0%)	0 (0%)
Postnatal corticosteroids - <i>n</i> (%)*	56 (34%)	41 (31%)	-	-
Supplemental oxygen at 36 weeks - <i>n</i> (%)*	65 (39%)	56 (42%)	-	-
<b><i>8-year outcomes</i></b>				
IQ - mean (SD)	97.8 (16.1) – <i>N</i> = 158	91.8 (15.3) – <i>N</i> = 100	108.0 (11.4) – <i>N</i> = 127	99.7 (16.3) – <i>N</i> = 84
HUI-2 - median (IQR)	0.95 (0.87 – 1.00) – <i>N</i> = 153	0.93 (0.84 – 1.00) – <i>N</i> = 104	1.00 (0.95 – 1.00) – <i>N</i> = 127	0.97 (0.88 – 1.00) – <i>N</i> = 85
Lower maternal education – <i>n</i> (%)	81 (51%) – <i>N</i> = 160	59 (55%) – <i>N</i> = 108	34 (27%) – <i>N</i> = 127	44 (52%) – <i>N</i> = 85
Lower social class - <i>n</i> (%)	44 (28%) – <i>N</i> = 158	39 (35%) – <i>N</i> = 110	18 (14%) – <i>N</i> = 128	24 (28%) – <i>N</i> = 86
No disability - <i>n</i> (%)	106 (66%) – <i>N</i> = 161	51 (42%) – <i>N</i> = 122	119 (93%) – <i>N</i> = 128	67 (75%) – <i>N</i> = 89
<b><i>18-year outcomes</i></b>				
HUI3 - median (IQR)	0.92 (0.82 - 1.00) – <i>N</i> = 134	0.96 (0.87 - 1.00) – <i>N</i> = 52	0.95 (0.88 - 1.00) – <i>N</i> = 101	0.94 (0.82 - 1.00) – <i>N</i> = 32
<b><i>Sub-optimal attributes – n (%)</i></b> :				
Vision	35 (26%) – <i>N</i> = 136	6 (12%) – <i>N</i> = 52	23 (22%) – <i>N</i> = 106	5 (15%) – <i>N</i> = 33
Hearing	4 (3%) – <i>N</i> = 137	2 (4%) – <i>N</i> = 52	1 (1%) – <i>N</i> = 106	1 (3%) – <i>N</i> = 33
Speech	27 (20%) – <i>N</i> = 137	7 (13%) – <i>N</i> = 52	13 (12%) – <i>N</i> = 105	5 (15%) – <i>N</i> = 33
Ambulation	4 (3%) – <i>N</i> = 137	4 (8%) – <i>N</i> = 52	0 (0%) – <i>N</i> = 105	3 (9%) – <i>N</i> = 33
Dexterity	8 (6%) – <i>N</i> = 135	3 (6%) – <i>N</i> = 52	1 (1%) – <i>N</i> = 105	0 (0%) – <i>N</i> = 33

Emotion	41 (30%) – N = 135	13 (25%) – N = 52	28 (27%) – N = 103	9 (27%) – N = 33
Cognition	46 (34%) – N = 134	15 (29%) – N = 52	24 (23%) – N = 105	11 (35%) – N = 31
Pain	29 (21%) – N = 135	9 (17%) – N = 52	31 (30%) – N = 105	10 (31%) – N = 32
IQ - mean (SD)	98.0 (16.2) – N = 149	89.9 (15.3) – N = 75	108.6 (12.5) – N = 112	100.2 (14.6) – N = 43
IQ <-1 SD – n (%)	57 (38%) – N = 149	45 (60%) – N = 75	12 (11%) – N = 112	14 (33%) – N = 43
<b>25-year outcomes</b>				
HUI3 – median (IQR)	0.91 (0.74 – 1.00)	-	0.95 (0.85 – 1.00)	-
<i>Sub-optimal attributes – n (%)</i> :				
Vision	55 (33%) – N = 165	-	42 (32%) – N = 131	-
Hearing	6 (4%) – N = 165	-	2 (2%) – N = 131	-
Speech	28 (17%) – N = 165	-	3 (2%) – N = 131	-
Ambulation	8 (5%) – N = 165	-	2 (2%) – N = 131	-
Dexterity	11 (7%) – N = 165	-	0 (0%) – N = 131	-
Emotion	60 (36%) – N = 165	-	44 (34%) – N = 131	-
Cognition	59 (36%) – N = 165	-	29 (22%) – N = 131	-
Pain	57 (35%) – N = 165	-	40 (31%) – N = 131	-

Notes: \*Perinatal characteristics specific to EP/ELBW born infants (not measured for controls); IQR = interquartile range; SD = standard deviation. CHQ = child health questionnaire, WRAT-4 = Wide Range Achievement Test, Edition 4; PVL = periventricular leukomalacia; M-ABC = Movement Assessment Battery for Children; HUI = Health Utilities Index; SD = standard deviation. No Disability = no cerebral palsy, blindness, or deafness, and an IQ >=-1SD relative to controls. Sample sizes for outcomes at age 8 and 18 years listed (N) as not all individuals were assessed at those ages.

**Table 2:** Median differences (95% confidence intervals [CI]) in Health-Related Quality of Life compared between birth groups at age 18 and 25 years, and between those with lower and higher IQ among participants born EP/ELBW

Method of Analysis	Age 18 Years		Age 25 Years	
	Median Difference (95% CI)	<i>p</i> - value	Median Difference (95% CI)	<i>p</i> - value
<b>A. ELBW/EP vs controls</b>				
Primary analysis: multiple imputation	-0.016 (-0.061, 0.029)	0.47	-0.040 (-0.088, 0.008)	0.10
Complete case analysis	-0.014 (-0.050, 0.023)	0.46	-0.026 (-0.065, 0.013)	0.20
<b>B. Lower intellectual quotient vs higher intellectual quotient measured at 18 years among EP/ELBW cohort</b>				
Primary analysis: multiple imputation	-0.034 (-0.107, 0.040)	0.36	-0.031 (-0.126, 0.064)	0.51
Complete case analysis	-0.055 (-0.120, 0.011)	0.10	-0.063 (-0.170, 0.044)	0.25

EP/ELBW = extremely preterm/extremely low birthweight.

**Table 3:** Odds ratios (95% confidence intervals [CI]) of sub-optimal function for each sub-scale attribute compared between birth groups at age 18 and 25 years, and between those with lower and higher IQ among participants born EP/ELBW, using multiple imputation to handle missing data, and adjusting for lower social class and lower maternal education

	<i>Age 18 Years</i>	<i>Age 25 Years</i>
<b>Method of Analysis</b>	<b>Odds Ratio (95% CI)</b>	<b>Odds Ratio (95% CI)</b>
<b>A. ELBW/EP vs controls</b>		
Vision	1.14 (0.68, 1.89)	1.08 (0.67, 1.76)
Hearing*	1.48 (0.31, 7.09)	2.71 (0.62, 11.8)
Speech*	1.46 (0.83, 2.55)	9.28 (3.09, 27.9)
Ambulation	1.50 (0.41, 5.54)	4.17 (0.89, 19.7)
Dexterity	3.42 (0.82, 14.2)	5.44 (1.04, 28.4)
Emotion	0.79 (0.44, 1.43)	1.01 (0.62, 1.63)
Cognition	1.24 (0.67, 2.31)	1.55 (0.95, 2.51)
Pain	0.66 (0.37, 1.17)	1.23 (0.76, 2.01)
<b>B. Lower intellectual quotient vs higher intellectual quotient measured at 18 years among EP/ELBW cohort</b>		
Vision	1.02 (0.50, 2.06)	1.08 (0.55, 2.12)
Hearing*	2.30 (0.44, 12.0)	2.10 (0.42, 10.6)
Speech	1.85 (0.75, 4.52)	1.48 (0.64, 3.45)
Ambulation*	1.26 (0.27, 5.82)	3.41 (0.78, 14.9)
Dexterity	1.04 (0.20, 5.38)	1.94 (0.56, 6.71)
Emotion	1.00 (0.46, 2.20)	0.77 (0.39, 1.52)
Cognition	1.37 (0.63, 2.94)	1.56 (0.82, 3.00)
Pain	1.26 (0.64, 2.50)	1.69 (0.90, 3.17)

EP/ELBW = extremely preterm/extremely low birthweight. Results from a logistic regression models adjusted for lower social class and maternal education.

\*Lower social class omitted as *mi estimate* would not converge using logistic regression (likely due to social class being a perfect predictor).

