



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Idrose, NS;Tham, RCA;Lodge, CJ;Lowe, AJ;Bui, D;Perret, JL;Vicendese, D;Newbigin, EJ;Tang, MLK;Aldakheel, FM;Waidyatillake, NT;Douglass, JA;Abramson, MJ;Walters, EH;Erbas, B;Dharmage, SC

Title:

Is short-term exposure to grass pollen adversely associated with lung function and airway inflammation in the community?

Date:

2021-04-01

Citation:

Idrose, N. S., Tham, R. C. A., Lodge, C. J., Lowe, A. J., Bui, D., Perret, J. L., Vicendese, D., Newbigin, E. J., Tang, M. L. K., Aldakheel, F. M., Waidyatillake, N. T., Douglass, J. A., Abramson, M. J., Walters, E. H., Erbas, B. & Dharmage, S. C. (2021). Is short-term exposure to grass pollen adversely associated with lung function and airway inflammation in the community?. *Allergy European Journal of Allergy and Clinical Immunology*, 76 (4), pp.1136-1146. <https://doi.org/10.1111/all.14566>.

Persistent Link:

<https://hdl.handle.net/11343/276301>

1

2 DR. CAROLINE LODGE (Orcid ID : 0000-0002-2342-3888)

3 DR. ADRIAN LOWE (Orcid ID : 0000-0002-4691-8162)

4 DR. FAHAD MOHAMMED ALDAKHEEL (Orcid ID : 0000-0001-8047-8650)

5 DR. BIRCAN ERBAS (Orcid ID : 0000-0001-9597-418X)

6 DR. SHYAMALI C DHARMAGE (Orcid ID : 0000-0001-6063-1937)

7

8

9 Article type : Original Article: Asthma and Lower Airway Disease

10

11

12 **Title: Is short-term exposure to grass pollen adversely associated with lung function and**  
13 **airway inflammation in the community?**

14 **Short title:** Outdoor grass pollen and respiratory health

15 N. Sabrina Idrose, MSc<sup>a</sup>, Rachel C.A. Tham, PhD<sup>b</sup>, Caroline J Lodge, PhD<sup>a</sup>, Adrian J Lowe,  
16 PhD<sup>a</sup>, Dinh Bui, PhD<sup>a</sup>, Jennifer L Perret, PhD<sup>a</sup>, Don Vicendese, PhD<sup>a,c</sup>, Edward J Newbigin,  
17 PhD<sup>d</sup>, Mimi L.K. Tang, MD, PhD<sup>e,f</sup>, Fahad M Aldakheel, PhD<sup>g</sup>, Nilakshi T Waidyatillake,  
18 MD, PhD<sup>a</sup>, Jo A Douglass, MD<sup>h,i</sup>, Michael J Abramson, MD, PhD<sup>j</sup>, E. Haydn Walters, DM<sup>a,k</sup>,  
19 \*Bircan Erbas, PhD<sup>l</sup>, \*Shyamali C Dharmage, MD, PhD<sup>a</sup>

20 \*Equally contributed

21 a. Allergy and Lung Health Unit, Melbourne School of Population and Global Health,  
22 The University of Melbourne, 207 Bouverie St, Carlton, Victoria, 3053, Australia.

23 b. Mary MacKillop Institute for Health Research, Australian Catholic University, 215  
24 Spring St, Melbourne, Victoria, 3000, Australia.

25 c. Department of Mathematics and Statistics, La Trobe University, Bundoora, 3086,  
26 Victoria, Australia.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/ALL.14566](https://doi.org/10.1111/ALL.14566)

This article is protected by copyright. All rights reserved

- 27 d. School of BioSciences, The University of Melbourne, Parkville, Victoria, 3052,  
28 Australia.
- 29 e. Allergy and Immunology, Murdoch Children's Research Institute, Royal Children's  
30 Hospital, Melbourne, Victoria, 3052, Australia.
- 31 f. Department of Pediatrics, University of Melbourne, Melbourne, VIC, Australia.
- 32 g. Department of Clinical Laboratory Sciences, College of Applied Medical Sciences,  
33 King Saud University, Riyadh, Saudi Arabia.
- 34 h. Department of Clinical Immunology and Allergy, Royal Melbourne Hospital,  
35 Parkville, 3050, Victoria, Australia.
- 36 i. Department of Medicine, University of Melbourne, Victoria, 3052, Australia.
- 37 j. School of Public Health & Preventive Medicine, Monash University, Melbourne,  
38 3004, Australia.
- 39 k. School of Medicine, University of Tasmania, Hobart, Tasmania, 7000, Australia.
- 40 l. School of Psychology and Public Health, La Trobe University, Bundoora, 3086,  
41 Victoria, Australia.

42  
43  
44 **Corresponding Author:** Shyamali Dharmage  
45 Centre of Epidemiology and Biostatistics,  
46 Melbourne School of Population and Global Health,  
47 The University of Melbourne,  
48 207 Bouverie St, Carlton, VIC 3052  
49 Tel: +61 3 83440737  
50 Fax : +61 3 93495815  
51 Email: [s.dharmage@unimelb.edu.au](mailto:s.dharmage@unimelb.edu.au)

52  
53 **Authors' contributions:** **Sabrina Idrose:** Investigation, Writing- Original Draft. **Rachel**  
54 **Tham:** Investigation, Writing- Review & Editing, Supervision. **Caroline Lodge:** Writing-  
55 Review & Editing. **Adrian Lowe:** Writing- Review & Editing. **Dinh Bui:** Writing- Review  
56 & Editing. **Jennifer Perret:** Writing- Review & Editing. **Don Vicendese:** Writing- Review  
57 & Editing. **Edward Newbigin:** Writing- Review & Editing. **Mimi Tang:** Writing- Review &  
58 Editing. **Fahad Aldakheel:** Investigation, Writing- Review & Editing. **Nilakshi**  
59 **Waidyatillake:** Writing- Review & Editing. **Jo Douglass:** Writing- Review & Editing.  
60 **Michael Abramson:** Writing- Review & Editing. **Haydn Walters:** Writing- Review &

61 Editing. **Bircan Erbas:** Conceptualization, Investigation, Writing- Review & Editing,  
62 Supervision. **Shyamali Dharmage:** Conceptualization, Investigation, Writing- Review &  
63 Editing, Supervision.

64 **Acknowledgements:** We acknowledge the early contributions of the original MACS  
65 investigators: Dr. David Hill, Dr. Cliff Hosking and Christine Axelrad. We also thank the  
66 participants for their ongoing support. Fahad M Aldakheel extends his appreciation to the  
67 College of Applied Medical Sciences Research Centre and the Deanship of Scientific  
68 Research at King Saud University.

69 **Potential conflicts of interest:** Michael Abramson holds investigator-initiated grants from  
70 Pfizer and Boehringer-Ingelheim. He has undertaken an unrelated consultancy for and  
71 received assistance with conference attendance from Sanofi. He has also received a speaker's  
72 fee from GSK. Jennifer Perret has received a travel grant from Boehringer Ingelheim in 2018.  
73 Jo Douglass reports grants from CSL-Behring LTD, personal fees from Astra-Zeneca,  
74 Novartis and GSK as well as contracted research from Sanofi-Aventis, Grifols P/L, BioCryst  
75 P/L and Equilium P/L outside the submitted work.

76 **Funding sources:** Australian National Health and Medical Research Council (NHMRC)

77 **Word count (Abstract):** 241

78 **Word count (Text):** 3660

79

80

81

82

83

84

85

86

87

88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99

100 Abstract

101 **Background:** The association between grass pollen exposure and early markers of asthma  
102 exacerbations such as lung function changes and increase in airway inflammation is limited.  
103 We investigated the associations between short-term grass pollen exposure and lung function  
104 and airway inflammation in a community-based sample, and whether any such associations  
105 were modified by current asthma, current hay fever, pollen sensitization, age and other  
106 environmental factors.

107 **Methods:** Cross-sectional and short-term analyses of data from the Melbourne Atopy Cohort  
108 Study (MACS) participants (n=936). Lung function was assessed using spirometry. Airway  
109 inflammation was assessed by fractional exhaled nitric oxide (FeNO), and exhaled breath  
110 condensate pH and nitrogen oxides (NO<sub>x</sub>). Daily pollen counts were collected using a  
111 volumetric spore trap. The associations were examined by linear regression.

112 **Results:** Higher ambient levels of grass pollen 2 days before (lag 2) were associated with  
113 lower mid-forced expiratory flow (FEF<sub>25-75%</sub>) and FEV<sub>1</sub>/FVC ratio (Coef. [95% CI] = -119 [-  
114 226, -11] mL/s and -1.0 [-3.0, -0.03] %, respectively) and also 3 days before (lag 3).

115 Increased levels of grass pollen a day before (lag 1) was associated with increased FeNO  
116 (4.35 [-0.1, 8.7] ppb) and also at lag 2. Adverse associations between pollen and multiple  
117 outcomes were greater in adults with current asthma, hay fever and pollen sensitization.

118 **Conclusion:** Grass pollen exposure was associated with eosinophilic airway inflammation 1-  
119 2 days after exposure and airway obstruction 2-3 days after exposure. Adults and individuals  
120 with asthma, hay fever and pollen sensitization may be at higher risk.

121 **Keywords:** Airway inflammation, environmental health, grass, lung function, pollen

122 **Abbreviations:** ATS/ERS: American Thoracic Society/European Respiratory Society; BDR:  
123 bronchodilator response; CI: confidence interval; EBC: exhaled breath condensate; ED:  
124 emergency department; FEF<sub>25-75%</sub>: mid-forced expiratory flow; FeNO: fractional exhaled  
125 nitric oxide; FEV<sub>1</sub>: forced expiratory volume in the first second; FVC: forced vital capacity;  
126 IgE: Immunoglobulin E; ISAAC: International Study of Asthma and Allergies in Childhood;  
127 MACS: Melbourne Atopy Cohort Study; MAPCAH: Melbourne Air Pollen Children and  
128 Adolescent Health; NO<sub>x</sub>: nitrogen oxide; RV: residual volume; SD: standard deviation; SPT:  
129 skin-prick test; Th2: T helper 2 lymphocytes

## 130 Introduction

131 Asthma is currently estimated to affect 339 million people worldwide and has become a  
132 global public health problem over the last few decades<sup>1</sup>. It affects people of all ages and one  
133 of the top contributors to preventable hospitalizations due to acute exacerbations<sup>2</sup>.

134 Identification of acute asthma triggers is a priority to inform preventive strategies to  
135 minimize individual episodes as well as societal healthcare costs.

136 Exposure to temperate grass pollen is being recognized as an increasingly important trigger  
137 for acute asthma including thunderstorm asthma<sup>3-5</sup>. Temperate grass pollen contains several  
138 allergens, particularly Group 1 and 5, which have considerable homology across grass  
139 species<sup>6,7</sup>. While whole grass pollen grains (10-100 $\mu$ m) usually get trapped in the upper  
140 airways, rupture of grass pollen grains during thunderstorms produces hundreds of tiny  
141 allergenic starch granules (0.1-5 $\mu$ m) that can penetrate deeper into the lower airways<sup>8</sup>.  
142 Furthermore, with changing climatic conditions, the intensity and duration of the grass  
143 pollenating period is increasing in many parts of the world<sup>9</sup>, increasing the potential impact  
144 of grass pollen exposure on health.

145 While exposure to grass pollen is increasingly recognized as a trigger of acute asthma attacks  
146 leading to hospital attendance, this is only the “tip of the iceberg” as there is likely to be a  
147 much higher burden of more moderate asthma exacerbations that do not lead to  
148 hospitalization within the general community<sup>10</sup>. Despite this, community-based studies  
149 investigating the role of pollen on early markers of asthma exacerbations such as lung  
150 function changes and increases in airway inflammation have been inconsistent and limited in  
151 terms of methodology.

152

153 Most studies<sup>11-13</sup> have included selected samples with small number of participants, limiting  
154 the study power and generalizability. Of the very few population-based studies with adequate  
155 sample sizes, key confounding variables such as temperature and humidity were not  
156 considered<sup>14,15</sup>. More importantly, lack of investigation into factors that may modify these  
157 associations such as current asthma, pollen sensitization, hay fever and age has made it  
158 difficult to establish high-risk groups who need to be targeted for monitoring or interventions  
159 during grass pollen seasons. Furthermore, it has been suggested that the impact of grass  
160 pollen exposure is higher in the presence of fungi<sup>16,17</sup>, rainfall<sup>18</sup>, air pollutants<sup>19</sup>, higher

161 temperature<sup>20</sup> and lower humidity<sup>21</sup> but these important potential effect modifiers have not  
162 been investigated.

163 Given the above gaps in knowledge, we investigated the associations between grass pollen  
164 exposure on the day of clinical testing and the preceding three days and lung function and  
165 airway inflammation in a general population cohort at overall at potentially greater risk for  
166 asthma and allergies. We examined for susceptible groups and periods by exploring effect  
167 modifications by current asthma, pollen sensitization, current hay fever, age and other  
168 environmental factors.

169

170

171

172

## 173 **Methods**

### 174 Study design and population

175 The study sample consisted of participants of the Melbourne Atopy Cohort Study (MACS)<sup>22</sup>.  
176 The original cohort consisted of 620 'high risk' babies born between 24 March 1990 and 1  
177 November 1994 in Melbourne, Australia. Each had at least one first-degree relative with a  
178 history of allergic diseases i.e. eczema, asthma, hay fever or severe food allergy. The  
179 recruitment process and study methodology has been described in detail elsewhere<sup>22, 23</sup>.

180 The present work is based on cross-sectional data from the 18-year follow-up visit. During  
181 this follow-up, parents and siblings were also invited to participate. However, only the  
182 participants who attended the laboratory between September 2009 and December 2011 were  
183 included in the analysis as daily outdoor grass pollen counts were only available during this  
184 period. These participants underwent a single episode of clinical testing for lung function  
185 (spirometry), airway inflammation (exhaled NO and exhaled breath condensate), and skin-  
186 prick tests (SPTs) during the study period. These one-time visits were random in terms of  
187 when the individuals attended the clinic relative to the phase of the pollen season.

188 Environmental pollen data were obtained for the day of lab testing (lag 0) and the three days  
189 prior to attendance (designated lag day 1, lag day 2 and lag day 3).

190 Ethics approval was granted by The University of Melbourne and Royal Children's Hospital  
191 Human Research Ethics Committees. Participants also provided written informed consent.

#### 192 Environmental data collection

193 Daily ambient outdoor grass pollen and fungi levels were collected between September 2009  
194 and December 2011 for the Melbourne Air Pollen Children and Adolescent Health  
195 (MAPCAH) study<sup>24</sup>. These aeroallergens were collected using a volumetric spore trap  
196 (Burkard, UK) located on the rooftop of the Earth Sciences building at The University of  
197 Melbourne. For this study, we analyzed grass pollen, *Cladosporium*, *Leptosphaeria*,  
198 *Alternaria* and smuts as they were found to be most abundant in Melbourne<sup>25, 26</sup>. The  
199 collection method is detailed in the Supplementary Material.

200 Air pollutants including the daily maximum 4-hour average ozone (O<sub>3</sub>), the daily maximum  
201 1-hour average nitrogen oxide (NO<sub>2</sub>) and the daily 24-hour average concentrations of  
202 particulate matter up to 2.5µm in diameter (PM<sub>2.5</sub>) and up to 10µm in diameter (PM<sub>10</sub>) during  
203 the study period were obtained from the Victorian Environmental Protection Authority. The  
204 daily maximum temperatures, rainfall and relative humidity data were obtained from The  
205 Bureau of Meteorology.

#### 206 Clinical data collection

207 Lung function was measured following ATS/ERS guidelines<sup>27</sup> using the EasyOne Spirometer  
208 (ndd Medical Technologies Inc., Andover, MA). Associations with the following pre-  
209 bronchodilator (pre-BD) spirometry parameters were assessed: forced expiratory volume in  
210 the first second (FEV<sub>1</sub>), forced vital capacity (FVC), mid-forced expiratory flow (FEF<sub>25-75%</sub>)  
211 and FEV<sub>1</sub>/FVC ratio. Bronchodilator response (BDR) after 400µg inhaled salbutamol was  
212 measured as a continuous and dichotomous variable. "Absolute BDR" was defined as the  
213 difference between post- and pre-bronchodilator FEV<sub>1</sub> values, while a "positive BDR"  
214 response was defined as having ≥ 12% change and ≥ 200ml of absolute change in FEV<sub>1</sub>  
215 from pre-bronchodilator to post-bronchodilator measurements<sup>28, 29</sup>. Pollen sensitization to  
216 mixed grass and/or ryegrass (Bayer, Spokane, WA, USA) allergen was determined by a skin-  
217 prick test (SPT) on the forearm. The measurement details are provided in the Supplementary  
218 Material.

219 Fractional exhaled nitric oxide (FeNO) was measured following ATS/ERS guidelines<sup>30</sup> using  
220 the HypAir™ FeNO machine (Medisoft, Sorinnes, Belgium). Exhaled breath condensate  
221 (EBC) was collected only in probands (but not relatives) and assayed according to the

222 ATS/ERS guidelines<sup>31, 32</sup> for acidity (pH) and nitrogen oxides (NO<sub>x</sub>) using a glass-  
223 condensing chamber in wet ice. The methodology of EBC collection is detailed elsewhere<sup>33</sup>.

#### 224 Clinical definitions

225 Participants were considered to be pollen-sensitized if they had a wheal diameter of at least  
226 3mm greater than saline for any of the grasses tested (i.e. mixed grass and/or ryegrass).  
227 Current asthma and hay fever status were identified using a valid ISAAC questionnaire<sup>34</sup>.  
228 The participants were considered to have current asthma if they had a history of doctor-  
229 diagnosed asthma and one or more of the following in the 12 months prior to the follow-up:  
230 (a) wheeze; (b) at least one episode of asthma; (c) use of asthma medication. This definition  
231 of current asthma is in line with previous publications from this cohort and did not include  
232 BDR, since this might not be a reliable marker for asthma<sup>35</sup>. Current hay fever was defined as  
233 having at least one episode of hay fever and/or the use of medication for the symptoms in the  
234 12 months prior to the follow-up. Age was stratified into two groups: (a) children and  
235 adolescents (< 18 years old) and (b) adults (≥ 18 years old).

#### 236 Statistical analysis

237 The associations between grass pollen exposure on the same day of clinical testing (lag 0)  
238 and lagged grass pollen exposure (lag day 1 up to lag day 3 and cumulative exposure 3 days  
239 prior) and lung function outcomes (i.e. FEV<sub>1</sub>, FEF<sub>25-75%</sub>, FEV<sub>1</sub>/FVC ratio, absolute BDR);  
240 and airway inflammation outcomes (i.e. FeNO and EBC pH) were assessed using generalized  
241 linear models. Positive BDR was assessed using generalized estimating equations. EBC NO<sub>x</sub>  
242 was found to be strongly right skewed, so it was stratified into none (below level of  
243 detection), low (≥ 2.5 to 20 μmol) or high (> 20 μmol)<sup>36</sup> and analyzed using ordinal logistic  
244 regression (OLR). All models were accounted for familial clustering. In addition, we  
245 conducted a sensitivity analysis investigating grass pollen exposure at cumulative lag 7  
246 (average exposure over the prior 7 days).

247 The multivariable models were adjusted for age, sex, height, maximum temperature, relative  
248 humidity, ozone and season of testing as *a priori* confounders<sup>37-41</sup>. Models assessing EBC pH  
249 and EBC NO<sub>x</sub> were also adjusted for storage time to account for possible sample  
250 degradation. Potential effect modifications by current asthma (irrespective of hay fever and  
251 pollen sensitization status), current hay fever (irrespective of asthma and pollen sensitization  
252 status), pollen sensitization (irrespective of hay fever and asthma status), age, air pollutants,  
253 ambient fungal levels, temperature, humidity and rainfall were explored separately using

254 Wald's test. We also conducted an exploratory analyses of the relevant associations by the  
255 combined phenotypes of current asthma and hay fever (current asthma only, hay fever only or  
256 both), pollen sensitization and current hay fever (pollen sensitization only, hay fever only or  
257 both), and pollen sensitization and current asthma (pollen sensitization only, current asthma  
258 only or both), separately using Wald's test .

259 The coefficients and odds ratios (OR) were reported as the effect size per increase in 29 grass  
260 pollen grains/m<sup>3</sup>. Significant associations were defined as  $p \leq 0.05$  and  $p\text{-int} \leq 0.1$ . Strata-  
261 specific associations were only reported if the  $p$ -value for interaction term was  $\leq 0.1$ . All  
262 statistical analyses were performed using Stata IC 14.2 (StataCorp, College Station, Texas).

263

264

265

266

267

## 268 **Results**

### 269 Sample characteristics

270 This study included altogether 936 participants, consisting of 264 probands, plus some  
271 parents (n=398) and siblings (n=274). The median (range) age of the study participants was  
272 21 years (6-71). Of the total sample, 45% were males, 30% reported they had current asthma,  
273 13% reported they had current hay fever, and 47% were sensitized to at least one grass  
274 pollen. These results are summarized in Table 1. Of those with current asthma, 64.8% were  
275 sensitized to at least one grass pollen extract and 19.6% reported current hay fever. The  
276 proportion of people with current asthma who attended the clinic each month did not vary  
277 (Table S1).

### 278 Grass pollen distribution

279 The 2009 to 2011 study period captured three grass pollen seasons, from September to  
280 February, peaking in October to January<sup>42</sup>. There were no distinct variations in the grass  
281 pollen seasons between these years. Although a minor thunderstorm asthma event was  
282 reported on the 25th of November 2010<sup>4</sup>, none of the participants attended the clinic on this  
283 day and up to 12 days later. Grass pollen levels were negatively correlated with relative

284 humidity and nitrogen dioxide levels; and positively correlated with fungal levels  
285 (*Cladosporium*, *Leptosphaeria*, *Alternaria*, Smuts), maximum temperature, PM<sub>10</sub> and ozone.  
286 The summary statistics and the Spearman's correlations between grass pollen and other  
287 environmental factors are presented in Table S2.

#### 288 Adjusted analyses

289 The unadjusted (Table S3) and adjusted results (Table 2) were consistent. For adjusted lung  
290 function analyses, there were significant inverse associations between grass pollen exposure  
291 at lag 2 and lag 3 and FEF<sub>25-75%</sub> and FEV<sub>1</sub>/FVC ratio. For adjusted airway inflammation  
292 analyses (Table 2), there were significant positive associations between grass pollen exposure  
293 at lag 1 and lag 2 and FeNO. The associations at cumulative lag 7 were not significant (data  
294 not shown).

#### 295 Effect modification

296 Some of the observed associations were modified by current asthma and age (Table 3).  
297 Exposure to higher levels of grass pollen at lag 0, lag 1, lag 2 and cumulative lag 3 were  
298 associated with an increase in absolute BDR and FeNO in subjects with current asthma  
299 (irrespective of hay fever status) but not those without current asthma (p-values for  
300 interaction range from 0.004 to 0.06). Furthermore, exposure to higher levels of grass pollen  
301 was associated with a significantly lower FEF<sub>25-75%</sub> at lag 2, lag 3 and cumulative lag 3; and  
302 FEV<sub>1</sub>/FVC ratio at all time points in adults aged 18 and above, but not in children or  
303 adolescents (p-values for interaction range from 0.009 to 0.1).

304 The interactions with current hay fever were not significant but the observed associations in  
305 those with current hay fever were stronger compared to those without current hay fever  
306 (Table S4). Similar can be observed for pollen sensitization (Table S4). While there were  
307 some associations observed between grass pollen and lung function in the presence of little to  
308 no fungi, the number of associations seen were consistent with chance given the number of  
309 interactions tested in this analysis (Table S5). The interactions between grass pollen and air  
310 pollutants (PM<sub>2.5</sub>, PM<sub>10</sub>, ozone, nitrogen dioxide), temperature, rainfall, and humidity were  
311 not significant at all (p-values for interaction  $\geq 0.2$ , data not shown).

312 Interactions between grass pollen and the combined variable of current asthma and hay fever  
313 were significant for FEF<sub>25-75%</sub>, FeNO and absolute BDR (range: 0.086 to 0.099, see Table  
314 S6). In those with hay fever alone (Table 4), increased grass pollen exposure was associated  
315 with decreased FEF<sub>25-75%</sub> at all lag periods except lag 0, increased FeNO at all lag periods

316 except lag 2 and increased absolute BDR at lag 2. In those with hay fever and asthma (Table  
317 4), increased grass pollen exposure was associated with increased absolute BDR at all lag  
318 periods. In those with asthma alone (Table 4), increased grass pollen exposure was associated  
319 with increased absolute BDR at lag 1, lag 2 and cumulative lag 3.

320 Interactions between grass pollen and the combined variable of pollen sensitization and  
321 current hay fever were significant for FEV<sub>1</sub>, FVC and FeNO (range: 0.01 to 0.09, see Table  
322 S7). The associations were stronger in pollen-sensitized individuals with hay fever compared  
323 to pollen-sensitized individuals without hay fever. In pollen-sensitized individuals with hay  
324 fever, increased grass pollen exposure was associated with decreased FEV<sub>1</sub> at lag 3 and  
325 higher FeNO at lag 0 and lag 1 (Table 5). The direction of association for FVC was  
326 consistent, such that pollen-sensitized individuals with hay fever were observed to have lower  
327 FVC at all lags associated with increased grass pollen exposure (Table S8).

328 Interactions between grass pollen and the combined variable of pollen sensitization and  
329 current asthma were significant for FeNO (range: 0.0001 to 0.003, see Table S9). Pollen-  
330 sensitized individuals with asthma were observed to have higher FeNO at all lags associated  
331 with increased grass pollen exposure (Table S10).

## 332 Discussion

333 This analysis has shown that short-term exposure (up to 3 days) to grass pollen was  
334 associated with reduced lung function and increased airway inflammation, particularly for  
335 FEF<sub>25-75%</sub>, FEV<sub>1</sub>/FVC ratio and FeNO, but with different lag patterns. These associations  
336 were stronger in specific groups of individuals – those with current asthma or hay fever, grass  
337 pollen-sensitized individuals and adults. All associations were lagged, which is consistent  
338 with grass pollen exposure inducing a late inflammatory response<sup>43</sup>. An inflammatory  
339 cascade resulting in the recruitment of immune cells such as eosinophils is suggested to peak  
340 approximately 24-72 hours after grass pollen exposure<sup>44</sup> which could be the reason why we  
341 did not observe significant associations at cumulative lag 7 days. An alternative explanation  
342 for the lack of association found at cumulative lag 7 days is the nature of grass pollen seasons  
343 in Melbourne. It consists of short periods of 2-3 days with hot/sunny weather with continental  
344 anticyclones, interspersed with westerly cold/wet low-pressure fronts. This intermittent  
345 weather coupled with the different variations of grass pollen levels makes it an ideal place for  
346 such a natural experiment investigating short-term effects of up to 3 days. Interestingly, we  
347 observed a more delayed effect on lung function compared to airway inflammation. Lung  
348 function was observed to be lower 2-3 days after grass pollen exposure, but airway  
349 inflammation was observed to increase 1-2 days after grass pollen exposure. These findings  
350 suggest a temporal association between lung function deficits and airway inflammation.

351 Our observation that grass pollen exposure was associated with FEF<sub>25-75%</sub> but not  
352 significantly with FEV<sub>1</sub> may indicate that the greatest impact is on medium-sized to small  
353 airways. This is consistent with observations by Marseglia et al.<sup>45</sup> in a population with  
354 allergic rhinitis. Reduction in FVC in pollen-sensitized individuals with hay fever is  
355 consistent with narrowing of the smallest airways affecting air-trapping and so raising  
356 Residual Volume (RV) and impinging on FVC. Unfortunately, we did not have RV data to  
357 confirm this. Smaller airways have in aggregate far greater cross-sectional surface areas and  
358 airway volumes compared to large airways, but they only contribute to 10% of airway  
359 resistance<sup>46</sup>. FEV<sub>1</sub> is less sensitive to smaller airway changes<sup>46</sup> so this could be a reason why  
360 past studies<sup>47-49</sup> found no association with lung function when they measured FEV<sub>1</sub> alone.  
361 While there have been contradictory views on FEF<sub>25-75%</sub> as a reliable marker of small airway  
362 obstruction and variability in this measure can occur as a result of its close relationship to  
363 FEV<sub>1</sub> and FVC<sup>50</sup>, FEF<sub>25-75%</sub> has been suggested to be a marker of a more distal airway  
364 narrowing<sup>51</sup>. We did find small reductions in FEV<sub>1</sub>/FVC ratio which were statistically

365 significant but might not be clinically meaningful. In addition, we also observed a significant  
366 increase in FeNO, which would be consistent with eosinophilic airway inflammation<sup>52, 53</sup>.

367 Moreover, we observed that the associations between grass pollen exposure, FeNO and  
368 absolute BDR were stronger in people with current asthma compared to those without. This is  
369 consistent with asthmatics being more susceptible to reversible airflow limitation during peak  
370 grass pollen seasons<sup>12</sup>. In asthma, the lung elasticity is further lost, and exposure to grass  
371 pollen would lead to increased inflammation. Additionally, we observed a significant  
372 association with FEF<sub>25-75%</sub>, FeNO and absolute BDR among those with hay fever alone, but  
373 only with absolute BDR among those with both asthma and hay fever or asthma alone. This  
374 was not what we expected because one would speculate that people with current asthma  
375 would have more pulmonary function changes to allergen exposure. Nevertheless, the  
376 strongest associations in absolute BDR were observed among those with both asthma and hay  
377 fever, followed by those with asthma alone. For those with hay fever alone, the changes in  
378 absolute BDR were smaller but significant at lag 2, indicating a degree of sub-clinical asthma  
379 making them potentially quite vulnerable to a large surge of atmospheric pollen such as in a  
380 thunderstorm.

381 Indeed, grass pollen-sensitized individuals irrespective of current hay fever and asthma status  
382 were observed to have increased airway inflammation with increased grass pollen exposure,  
383 but pollen-sensitized individuals with either current hay fever or asthma were shown to have  
384 greater associations in terms of both lung function and airway inflammation. There was also a  
385 modifying effect of age with adults having stronger associations with increased grass pollen  
386 exposure, which is consistent with adults being more affected by thunderstorm asthma<sup>4, 54</sup>.

387 Although the response rate for mothers was slightly higher compared to fathers, this matched  
388 the distribution of males to females in the study (42% and 58% respectively) and so is  
389 unlikely to have biased the findings. Furthermore, reporting bias was unlikely because we  
390 used objective lung function measures to assess the outcomes.

#### 391 Strengths and limitations

392 To the best of our knowledge, this is the first large, community-based study to investigate the  
393 association between grass pollen levels and objective measures of lung function and airway  
394 inflammation. Furthermore, clinical and exposure measurements were carried out using gold  
395 standard techniques and validated questionnaires to limit measurement bias. The wide range

396 of data collected from this study also allowed investigation into effect modifiers, which had  
397 not been previously documented.

398 We did not find evidence to support the hypothesis of interactions between exposures to  
399 fungi and pollen. Nevertheless, this should be examined in future studies, especially given the  
400 isolated findings that we observed in relation to few specific fungi taxa. There was also no  
401 evidence of interaction by air pollutants and meteorological variables. This could be due to  
402 limited power, lack of individual level data (e.g. proximity of residence to major roads) and  
403 the relatively low pollution levels in this study setting. Thunderstorm asthma events occur  
404 during extreme weather conditions such as an increase in humidity, a sudden drop in  
405 temperature and presence of thunderstorm outflows, which result in the increase of pollen  
406 allergen load in the air. Unfortunately, we did not have patient data to assess the effects of  
407 these extreme weather conditions.

408 Several other study limitations also need to be considered. There was potential for spurious  
409 findings due to multiple hypothesis testing. Grass pollen counts were measured at a single  
410 site and this might not have accurately reflected exposure at an individual level. Nonetheless,  
411 measurements were made within approximately 50km of the central business district, where  
412 all participants resided. Data such as the relative distance between residence and collection  
413 site, time spent outdoors and indoor pollen measurements were also not collected, so these  
414 would have limited exposure assessment. However, these misclassifications were likely to be  
415 non-differential, would bias towards the null and so unlikely to explain the positive results.  
416 Furthermore, the cross-sectional approach of this analysis did not consider daily changes in  
417 allergen exposure on lung function and airway inflammation as the participants were assessed  
418 only once during the study period. This would have provided stronger evidence for the  
419 associations found in this study. Moreover, we had limited data to explore grass pollen at the  
420 species level and sensitization to specific grass pollen taxa. This would have provided us with  
421 a better understanding on the relationship between grass pollen, lung function and airway  
422 inflammation. Lastly, the MACS cohort consisted of individuals with a family history of  
423 allergic diseases. While this may limit generalizability, approximately 70% of the Australian  
424 population has a family history of allergies<sup>55</sup>.

425 In conclusion, the findings from our study highlight the relationship between short-term grass  
426 pollen exposure (within 3 days) and subsequent lung function and airway inflammation  
427 changes. Our data suggest that adults and individuals with current asthma, current hay fever

428 and grass pollen-sensitization are especially vulnerable. These findings are fundamental to  
429 inform individual and public health preventive strategies for asthma exacerbation and  
430 hospital admissions due to grass pollen exposure, such as a preventive asthma therapy or  
431 allergen immunotherapy. In addition, the use of modern technologies such as early warning  
432 systems on smartphones and social media sites for impending high pollen days, as well as  
433 broader information technology systems (ITS)-based public education to the adverse effects  
434 of grass pollen exposure could increasingly play an important preventive role.

Author Manuscript

436 **References**

- 437 1. Network GA. The Global Asthma Report 2018. Auckland, New Zealand. 2016.
- 438 2. Ordoñez GA, Phelan PD, Olinsky A, Robertson CF. Preventable factors in hospital  
439 admissions for asthma. *Archives of Disease in Childhood*. 1998;78(2):143.
- 440 3. Erbas B, Chang JH, Dharmage S, Ong EK, Hyndman R, Newbiggin E, et al. Do levels  
441 of airborne grass pollen influence asthma hospital admissions? *Clinical & Experimental*  
442 *Allergy*. 2007;37(11):1641-7.
- 443 4. Thien F, Beggs PJ, Csutoros D, Darvall J, Hew M, Davies JM, et al. The Melbourne  
444 epidemic thunderstorm asthma event 2016: an investigation of environmental triggers, effect  
445 on health services, and patient risk factors. *The Lancet Planetary Health*. 2018;2(6):e255-e63.
- 446 5. Idrose NS, Dharmage SC, Lowe AJ, Lambert KA, Lodge CJ, Abramson MJ, et al. A  
447 systematic review of the role of grass pollen and fungi in thunderstorm asthma.  
448 *Environmental Research*. 2020;181:108911.
- 449 6. Mohapatra SS, Lockey RF, Shirley S. Immunobiology of grass pollen allergens.  
450 *Current Allergy and Asthma Reports*. 2005;5(5):381.
- 451 7. Suphioglu C, Blaher B, Rolland JM, McCluskey J, Schäppi G, Kenrick J, et al.  
452 Molecular basis of IgE-recognition of Lol p 5, a major allergen of rye-grass pollen.  
453 *Molecular Immunology*. 1998;35(5):293-305.
- 454 8. Suphioglu C, Singh MB, Taylor P, Bellomo R, Holmes P, Puy R, et al. Mechanism of  
455 grass-pollen-induced asthma. *Lancet (London, England)*. 1992;339(8793):569-72.
- 456 9. Schmidt CW. Pollen Overload: Seasonal Allergies in a Changing Climate.  
457 *Environmental health perspectives*. 2016;124(4):A70-5.
- 458 10. Silver JD, Spriggs K, Haberle S, Katelaris CH, Newbiggin EJ, Lampugnani ER.  
459 Crowd-sourced allergic rhinitis symptom data: The influence of environmental and  
460 demographic factors. *Science of The Total Environment*. 2020;705:135147.
- 461 11. Di Gioacchino M, Cavallucci E, Di Stefano F, Paolini F, Ramondo S, Di Sciascio  
462 MB, et al. Effect of natural allergen exposure on non-specific bronchial reactivity in  
463 asthmatic farmers. *The Science of the Total Environment*. 2001;270(1-3):43-8.
- 464 12. Djukanović R, Feather I, Gratziau C, Walls A, Peroni D, Bradding P, et al. Effect of  
465 natural allergen exposure during the grass pollen season on airways inflammatory cells and  
466 asthma symptoms. *Thorax*. 1996;51(6):575-81.

- 467 13. Epton MJ, Martin IR, Graham P, Healy PE, Smith H, Balasubramaniam R, et al.  
468 Climate and aeroallergen levels in asthma: a 12 month prospective study. *Thorax*.  
469 1997;52(6):528-34.
- 470 14. Steerenberg PA, Bischoff EW, de Klerk A, Verlaan AP, Jongbloets LM, van Loveren  
471 H, et al. Acute effect of air pollution on respiratory complaints, exhaled NO and biomarkers  
472 in nasal lavages of allergic children during the pollen season. *International Archives of*  
473 *Allergy and Immunology*. 2003;131(2):127-37.
- 474 15. Gruzieva O, Pershagen G, Wickman M, Melen E, Hallberg J, Bellander T, et al.  
475 Exposure to grass pollen--but not birch pollen--affects lung function in Swedish children.  
476 *Allergy*. 2015;70(9):1181-3.
- 477 16. Packe GE, Ayres JG. Asthma outbreak during a thunderstorm. *Lancet (London,*  
478 *England)*. 1985;2(8448):199-204.
- 479 17. Venables KM, Allitt U, Collier CG, Emberlin J, Greig JB, Hardaker PJ, et al.  
480 Thunderstorm-related asthma--the epidemic of 24/25 June 1994. *Clinical and experimental*  
481 *allergy : Journal of the British Society for Allergy and Clinical Immunology*. 1997;27(7):725-  
482 36.
- 483 18. Knox RB. Grass pollen, thunderstorms and asthma. *Clinical & Experimental Allergy*.  
484 1993;23(5):354-9.
- 485 19. Amato G, Liccardi G, Amato M, Cazzola M. Outdoor air pollution, climatic changes  
486 and allergic bronchial asthma. *European Respiratory Journal*. 2002;20(3):763.
- 487 20. Krämer U, Weidinger S, Darsow U, Möhrenschrager M, Ring J, Behrendt H.  
488 Seasonality in Symptom Severity Influenced by Temperature or Grass Pollen: Results of a  
489 Panel Study in Children with Eczema. *Journal of Investigative Dermatology*.  
490 2005;124(3):514-23.
- 491 21. Bartková-Ščevková J. The influence of temperature, relative humidity and rainfall on  
492 the occurrence of pollen allergens (*Betula*, *Poaceae*, *Ambrosia artemisiifolia*) in the  
493 atmosphere of Bratislava (Slovakia). *International Journal of Biometeorology*. 2003;48(1):1-  
494 5.
- 495 22. Lowe AJ, Lodge CJ, Allen KJ, Abramson MJ, Matheson MC, Thomas PS, et al.  
496 Cohort Profile: Melbourne Atopy Cohort study (MACS). *International Journal of*  
497 *Epidemiology*. 2017;46(1):25-6.
- 498 23. Lowe AJ, Hosking CS, Bennett CM, Allen KJ, Axelrad C, Carlin JB, et al. Effect of a  
499 partially hydrolyzed whey infant formula at weaning on risk of allergic disease in high-risk

- 500 children: a randomized controlled trial. *The Journal of Allergy and Clinical immunology*.  
501 2011;128(2):360-5.e4.
- 502 24. Erbas B, Dharmage S, Sullivan M, Akram M, Newbiggin E, Taylor P, et al. A case-  
503 crossover design to examine the role of aeroallergens and respiratory viruses on childhood  
504 asthma exacerbations requiring hospitalization : The MAPCAH study. *Journal of Biometrics  
505 and Biostatistics*. 2012.
- 506 25. Tham R, Vicendese D, Dharmage SC, Hyndman RJ, Newbiggin E, Lewis E, et al.  
507 Associations between outdoor fungal spores and childhood and adolescent asthma  
508 hospitalizations. *The Journal of Allergy and Clinical immunology*. 2017;139(4):1140-7.e4.
- 509 26. Erbas B, Akram M, Dharmage SC, Tham R, Dennekamp M, Newbiggin E, et al. The  
510 role of seasonal grass pollen on childhood asthma emergency department presentations.  
511 *Clinical and Experimental Allergy : Journal of the British Society for Allergy and Clinical  
512 Immunology*. 2012;42(5):799-805.
- 513 27. Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, et al.  
514 Standardisation of spirometry. *European Respiratory Journal*. 2005;26(2):319.
- 515 28. Sim YS, Lee J-H, Lee W-Y, Suh DI, Oh Y-M, Yoon J-S, et al. Spirometry and  
516 Bronchodilator Test. *Tuberc Respir Dis (Seoul)*. 2017;80(2):105-12.
- 517 29. Tan WC, Bourbeau J, Hernandez P, Chapman KR, Cowie R, FitzGerald JM, et al.  
518 Bronchodilator Responsiveness and Reported Respiratory Symptoms in an Adult Population.  
519 *PLOS ONE*. 2013;8(3):e58932.
- 520 30. ATS/ERS recommendations for standardized procedures for the online and offline  
521 measurement of exhaled lower respiratory nitric oxide and nasal nitric oxide, 2005. *American  
522 Journal of Respiratory and Critical Care Medicine*. 2005;171(8):912-30.
- 523 31. Silkoff PE, Erzurum SC, Lundberg JO, George SC, Marczin N, Hunt JF, et al. ATS  
524 workshop proceedings: exhaled nitric oxide and nitric oxide oxidative metabolism in exhaled  
525 breath condensate. *Proceedings of the American Thoracic Society*. 2006;3(2):131-45.
- 526 32. Horvath I, Hunt J, Barnes PJ, Alving K, Antczak A, Baraldi E, et al. Exhaled breath  
527 condensate: methodological recommendations and unresolved questions. *The European  
528 Respiratory Journal*. 2005;26(3):523-48.
- 529 33. Aldakheel FM, Bourke JE, Thomas PS, Matheson MC, Abramson MJ, Hamilton GS,  
530 et al. NO<sub>x</sub> in exhaled breath condensate is related to allergic sensitization in young and  
531 middle-aged adults. *Clinical & Experimental Allergy*. 2019;49(2):171-9.

- 532 34. Asher MI, Keil U, Anderson HR, Beasley R, Crane J, Martinez F, et al. International  
533 Study of Asthma and Allergies in Childhood (ISAAC): rationale and methods. *The European*  
534 *Respiratory Journal*. 1995;8(3):483-91.
- 535 35. Fingleton J, Weatherall M, Beasley R. Bronchodilator responsiveness: interpret with  
536 caution. *Thorax*. 2012;67(8):667.
- 537 36. Tham R, Erbas B, Dharmage SC, Tang MLK, Aldakheel F, Lodge CJ, et al. Outdoor  
538 fungal spores and acute respiratory effects in vulnerable individuals. *Environmental*  
539 *Research*. 2019;178:108675.
- 540 37. Todea A, Suatean I, Coman A, Rosca L. The Effect of Climate Change and Air  
541 Pollution on Allergenic Potential of Pollens. *Notulae Botanicae Horti Agrobotanici Cluj-*  
542 *Napoca*. 2013;41:646-50.
- 543 38. Khatri SB, Holguin FC, Ryan PB, Mannino D, Erzurum SC, Teague WG. Association  
544 of ambient ozone exposure with airway inflammation and allergy in adults with asthma. *J*  
545 *Asthma*. 2009;46(8):777-85.
- 546 39. Frank U, Ernst D. Effects of NO<sub>2</sub> and Ozone on Pollen Allergenicity. *Front Plant Sci*.  
547 2016;7:91-.
- 548 40. Li S, Baker PJ, Jalaludin BB, Marks GB, Denison LS, Williams GM. Ambient  
549 temperature and lung function in children with asthma in Australia. *European Respiratory*  
550 *Journal*. 2014;43(4):1059.
- 551 41. Noh SR, Paek D. Weather and Airway Inflammation. *Epidemiology*. 2011;22(1):S18.
- 552 42. Erbas B, Dharmage SC, Tang ML, Akram M, Allen KJ, Vicendese D, et al. Do  
553 human rhinovirus infections and food allergy modify grass pollen-induced asthma hospital  
554 admissions in children? *The Journal of Allergy and Clinical Immunology*. 2015;136(4):1118-  
555 20.e2.
- 556 43. Roberts G, Hurley C, Bush A, Lack G. Longitudinal study of grass pollen exposure,  
557 symptoms, and exhaled nitric oxide in childhood seasonal allergic asthma. *Thorax*.  
558 2004;59(9):752-6.
- 559 44. Boldogh I, Bacsı A, Choudhury BK, Dharajiya N, Alam R, Hazra TK, et al. ROS  
560 generated by pollen NADPH oxidase provide a signal that augments antigen-induced allergic  
561 airway inflammation. *The Journal of Clinical Investigation*. 2005;115(8):2169-79.
- 562 45. Marseglia GL, Cirillo I, Vizzaccaro A, Klersy C, Tosca MA, La Rosa M, et al. Role  
563 of forced expiratory flow at 25-75% as an early marker of small airways impairment in  
564 subjects with allergic rhinitis. *Allergy and Asthma Proceedings*. 2007;28(1):74-8.

- 565 46. Thien F. Measuring and imaging small airways dysfunction in asthma. *Asia Pac*  
566 *Allergy*. 2013;3(4):224-30.
- 567 47. Bake B, Viklund E, Olin AC. Effects of pollen season on central and peripheral nitric  
568 oxide production in subjects with pollen asthma. *Respiratory Medicine*. 2014;108(9):1277-  
569 83.
- 570 48. Larsson P, Larstad M, Bake B, Hammar O, Bredberg A, Almstrand AC, et al. Exhaled  
571 particles as markers of small airway inflammation in subjects with asthma. *Clinical*  
572 *Physiology and Functional Imaging*. 2017;37(5):489-97.
- 573 49. Panzner P, Malkusova I, Vachova M, Liska M, Brodska P, Ruzickova O, et al.  
574 Bronchial inflammation in seasonal allergic rhinitis with or without asthma in relation to  
575 natural exposure to pollen allergens. *Allergologia et immunopathologia*. 2015;43(1):3-9.
- 576 50. Barreiro TJ, Perillo I. An approach to interpreting spirometry. *Am Fam Physician*.  
577 2004;69(5):1107-14.
- 578 51. Riley CM, Wenzel SE, Castro M, Erzurum SC, Chung KF, Fitzpatrick AM, et al.  
579 Clinical Implications of Having Reduced Mid Forced Expiratory Flow Rates (FEF25-75),  
580 Independently of FEV1, in Adult Patients with Asthma. *PLoS One*. 2015;10(12):e0145476.
- 581 52. George L, Brightling CE. Eosinophilic airway inflammation: role in asthma and  
582 chronic obstructive pulmonary disease. *Ther Adv Chronic Dis*. 2016;7(1):34-51.
- 583 53. Berry MA, Shaw DE, Green RH, Brightling CE, Wardlaw AJ, Pavord ID. The use of  
584 exhaled nitric oxide concentration to identify eosinophilic airway inflammation: an  
585 observational study in adults with asthma. *Clinical and Experimental allergy : Journal of the*  
586 *British Society for Allergy and Clinical Immunology*. 2005;35(9):1175-9.
- 587 54. Silver JD, Sutherland MF, Johnston FH, Lampugnani ER, McCarthy MA, Jacobs SJ,  
588 et al. Seasonal asthma in Melbourne, Australia, and some observations on the occurrence of  
589 thunderstorm asthma and its predictability. *PLOS ONE*. 2018;13(4):e0194929.
- 590 55. Koplin JJ, Allen KJ, Gurrin LC, Peters RL, Lowe AJ, Tang MLK, et al. The impact of  
591 family history of allergy on risk of food allergy: a population-based study of infants. *Int J*  
592 *Environ Res Public Health*. 2013;10(11):5364-77.

Table 1. Characteristics of MACS participants during the study period.

Variables		All	Probands	Mothers	Fathers	Siblings
N (%)		936	264 (28%)	238 (25%)	160 (17%)	274 (29%)
Median age (range) (years)		21 (65)	18 (5)	49 (20)	52 (30)	16 (35)
Mean height $\pm$ SD (cm)		169.9 $\pm$ 10.2	172.6 $\pm$ 1.5	164.2 $\pm$ 6.0	177.7 $\pm$ 6.4	167.8 $\pm$ 11.8
		<b>n (% of all)</b>	<b>n (% of subgroup)</b>	<b>n (% of subgroup)</b>	<b>n (% of subgroup)</b>	<b>n (% of subgroup)</b>
Gender	Male	424 (45%)	127 (48%)	-	160 (100%)	137 (50%)
	Female	512 (55%)	137 (52%)	238 (100%)	-	137 (50%)
Mixed grass sensitisation	Tested	n=895	n=253	n=228	n=154	n=258
	Yes	391 (44%)	105 (42%)	103 (45%)	60 (39%)	122 (47%)
Ryegrass sensitisation	Tested	n=896	n=253	n=227	n=154	n=260
	Yes	410 (46%)	109 (43%)	110 (48%)	66 (43%)	124 (48%)
Pollen sensitisation (Any†)	Tested	n=890	n=252	n=224	n=154	n=258
	Yes	422 (47%)	112 (44%)	111 (50%)	68 (44%)	130 (50%)
Current asthma	Total	n=811	n=229	n=207	n=146	n=229
	Yes	247 (30%)	66 (29%)	74 (36%)	28 (19%)	79 (34%)

Current hay fever	Total	n=815	232	205	141	237
	Yes	108 (13%)	23 (9.9%)	38 (19%)	14 (9.9%)	33 (14%)
Spirometry test		873 (93%)	262 (99%)	218 (92%)	146 (91%)	244 (89%)
FeNO test		797 (85%)	222 (84%)	205 (86%)	139 (87%)	231 (84%)
EBC test		231 (24.6%)	231 (88%)	-	-	-

†Sensitisation to mixed grass and/or ryegrass

Table 2. **Adjusted analysis** of grass pollen and various outcomes on same day, lag 1, lag 2, lag 3 and cumulative 3-day lag.

Outcomes	Grass pollen									
	Lag 0		Lag 1		Lag 2		Lag 3		Lag 0-3	
	Coef. (95% CI)	p-value	Coef. (95% CI)	p-value	Coef. (95% CI)	p-value	Coef. (95% CI)	p-value	Coef. (95% CI)	p-value
<b>FEV<sub>1</sub> (ml)</b>	1.45 (-70, 74)	0.97	-16.0 (-70, 38)	0.56	-29.0 (-81, 21)	0.25	-36.0 (-97, 26)	0.26	-7.3 (-23, 8.7)	0.37
<b>FVC (ml)</b>	0.87 (-65, 67)	0.98	-2.0 (-55, 51)	0.94	6.1 (-48, 60)	0.82	-2.9 (-71, 65)	0.93	0.29 (-16, 17)	0.96
<b>FEF<sub>25-75%</sub> (ml/second)</b>	-10.1 (-45, 25)	0.58	-66.4 (-171, 38)	0.21	-119 (-226, -11)*	0.03*	-122.1 (-225, -20)*	0.02*	-26 (-56, 3.8)	0.086
<b>FEV<sub>1</sub>/FVC ratio (%)</b>	-0.3 (-2.0, 1.0)	0.45	-0.6 (-1.0, 0.3)	0.21	-1.0 (-3.0, -0.03)*	0.04*	-2.0 (-3.0, -0.3)*	0.02*	-0.2 (-0.4, 0.06)	0.15
<b>Absolute BDR (ml)</b>	-1.2 (-19, 17)	0.90	9.0 (-11, 29)	0.37	16 (-4.6, 36)	0.13	13.1 (-7.3, 34)	0.21	2.7 (-2.9, 8.4)	0.34
<b>FeNO (ppb)</b>	-2.9 (-2.0, 8.1)	0.25	4.35 (-0.1, 8.7)*	0.05*	4.35 (-0.001, 8.7)*	0.05*	1.74 (-2.0, 5.5)	0.39	1.0 (-0.18, 2.2)	0.096
<b>EBC (pH)</b>	0.09 (-0.06, 0.3)	0.17	0.12 (-0.01, 0.3)	0.07	0.15 (0.06, 0.3)	0.04	0.15 (-0.01, 0.3)	0.07	0.03 (-0.0003, 0.07)	0.05
	<b>OR (95% CI)</b>	<b>p-value</b>	<b>OR (95% CI)</b>	<b>p-value</b>	<b>OR (95% CI)</b>	<b>p-value</b>	<b>OR (95% CI)</b>	<b>p-value</b>	<b>OR (95% CI)</b>	<b>p-value</b>
<b>EBC NO<sub>x</sub> (</b>	0.75 (0.31, 1.0)	0.31	0.75 (0.41, 1.3)	0.45	0.89 (0.56, 1.8)	0.72	1.0 (0.56, 1.0)	0.92	1.0 (0.7, 1.0)	0.65

<b><math>\mu\text{mol}</math></b>										
<b>Positive BDR</b>	0.75 (0.56, 1.0)	0.50	1.0 (0.75, 1.0)	0.998	1.0 (0.75, 1.78)	0.27	1.0 (0.75, 1.78)	0.72	1.0 (0.94, 1.0)	0.56

N.B. FEV<sub>1</sub>, FEF<sub>25-75%</sub>, absolute BDR and FeNO associations were adjusted for age, sex, height, relative humidity, temperature, ozone and season of testing. FEV<sub>1</sub>/FVC ratio associations were adjusted for relative humidity, temperature, ozone and season of testing. EBC associations were adjusted for storage time in addition to the other confounders. All results are interpreted as per increase in 29 grass pollen grains/m<sup>3</sup>. \*p ≤ 0.05.

Author Manuscript

Table 3. **Significant associations** between grass pollen and different outcomes stratified by current asthma and age.

Grass pollen	Outcomes					
	Absolute BDR (ml)			FeNO (ppb)		
	No current asthma	Current asthma		No current asthma	Current asthma	
	n=517	n=235		n=517	n=235	
Coef. (95% CI)	Coef. (95% CI)	p-int	Coef. (95% CI)	Coef. (95% CI)	p-int	
Lag 0	-16 (-37, 4.6)	50 (6.4, 93)	0.004*	-2.9 (-5.8, 2.9)	8.7 (-0.9, 17.4)	0.04*
Lag 1	-15 (-40, 11)	52 (10, 95)	0.004*	-1.2 (-4.4, 2.0)	8.7 (-0.6, 18.0)	0.048*
Lag 2	-5.5 (-33, 22)	46 (6.1, 86)	0.03*	-1.74 (-4.9, 1.4)	8.4 (-0.3, 17.1)	0.03*
Lag 3	-2.6 (-30, 25)	35 (-11, 81)	0.17*	-1.7 (-4.9, 1.7)	2.6 (-5.5, 10.7)	0.31
Lag 0-3	-2.6 (-9.3, 4.1)	13 (1.3, 24)	0.016*	-0.4 (-1.4, 0.5)	2.0 (-0.4, 4.6)	0.06*
	FEF <sub>25-75%</sub> (ml/second)			FEV <sub>1</sub> /FVC ratio (%)		
	< 18 years old	≥ 18 years old		< 18 years old	≥ 18 years old	
	n=246	n=627		n=246	n=627	
	Coef. (95% CI)	Coef. (95% CI)	p-int	Coef. (95% CI)	Coef. (95% CI)	p-int
Lag 0	-30 (-115, 176)	-83 (-202, 36)	0.13	0.6 (-0.3, 1.5)	-0.9 (-1.5, -0.0003)	0.009*
Lag 1	43 (-107, 192)	-104 (-221, 13)	0.07*	0.4 (-0.6, 1.5)	-0.9 (-1.7, -0.1)	0.02*
Lag 2	-7.9 (-181, 165)	-142 (-255, -29)	0.10*	0.09 (-1.2, 1.5)	-1.2 (-1.7, -0.3)	0.04*
Lag 3	-22 (-211, 167)	-168 (-280, -57)	0.10*	-0.1 (-1.5, 1.2)	-1.2 (-2.0, -0.3)	0.08*
Lag 0-3	3.2 (-45, 51)	-36 (-67, -4.9)	0.09*	0.06 (-0.3, 0.3)	-0.3 (-0.6, -0.09)	0.02*

N.B. FEF<sub>25-75%</sub>, FeNO and absolute BDR associations were adjusted for age, sex, height, relative humidity, temperature, ozone and season of testing. FEV<sub>1</sub>/FVC ratio associations were adjusted for relative humidity, temperature, ozone and season of testing. All results are interpreted as per increase in 29 grass pollen grains/m<sup>3</sup>. p-int= p-value for interaction term; \*p-int ≤ 0.1.

Author Manuscript

Table 4. **Significant associations** between grass pollen and lung function outcomes; and FeNO on same day, lag 1, lag 2, lag 3 and cumulative 3-day lag in those with current hay fever alone, current hay fever and current asthma, and current asthma alone.

Outcomes	Grass pollen									
	Lag 0		Lag 1		Lag 2		Lag 3		Lag 0-3	
	Coef. (95% CI)	p-value	Coef. (95% CI)	p-value	Coef. (95% CI)	p-value	Coef. (95% CI)	p-value	Coef. (95% CI)	p-value
In those with current hay fever and no asthma (N=44 for spirometry; N=48 for FeNO test)										
<b>FEF<sub>25-75%</sub></b> <b>(ml/second)</b>	-306 (-736, 124)	0.16	-451 (-845, -56)*	0.03*	-508 (-856, -160)*	0.004*	-525 (-907, -142)*	0.007*	-120 (-220, -19)*	0.02*
<b>Absolute BDR (ml)</b>	20 (-40, 80)	0.52	39 (-21, 99)	0.21	63 (-0.003, 125)*	0.05*	63 (-1.5, 128)	0.06	12.5 (-3.8, 29)	0.13
<b>FeNO (ppb)</b>	12.5 (1.5, 23)*	0.025*	16 (4.6, 27)*	0.006*	11.3 (-0.7, 23)	0.066	12.2 (1.0, 23)*	0.03*	3.5 (0.6, 6.1)*	0.02*
In those with current hay fever and current asthma (N=43 for spirometry)										
<b>Absolute BDR (ml)</b>	98 (44, 153)*	<0.001*	95 (44, 146)*	<0.001*	92 (39, 146)*	<0.001*	94 (39, 148)*	0.001*	25 (11, 38)*	<0.001*
In those with current asthma and no current hay fever (N=166 for spirometry)										

<b>Absolute BDR (ml)</b>	56 (-6.4, 119)	0.08	69 (7.8, 131)*	0.03*	63 (7.5, 119)*	0.03*	43.5 (-8.7, 96)	0.10	17.1 (2.6, 32)*	0.02*
--------------------------	----------------	------	----------------	-------	----------------	-------	-----------------	------	-----------------	-------

N.B. FEF<sub>25-75%</sub> absolute BDR and FeNO associations were adjusted for age, sex, height, relative humidity, temperature, ozone and season of testing. All results are interpreted as per increase in 29 grass pollen grains/m<sup>3</sup>. \*p ≤ 0.05.

Author Manuscript

Table 5. **Adjusted analysis** of grass pollen and FEV<sub>1</sub>; and FeNO stratified by grass pollen sensitisation, with and without current hay fever on same day, lag 1, lag 2, lag 3 and cumulative 3-day lag.

Grass pollen	FEV <sub>1</sub> (ml)				FeNO (ppb)			
	Sensitized to $\geq 1$ grass pollen extract without current hay fever		Sensitized to $\geq 1$ grass pollen extract with current hay fever		Sensitized to $\geq 1$ grass pollen extract without current hay fever		Sensitized to $\geq 1$ grass pollen extract with current hay fever	
	n=373		n=426		n=385		n=396	
	Coef. (95% CI)	p-value	Coef. (95% CI)	p-value	Coef. (95% CI)	p-value	Coef. (95% CI)	p-value
Lag 0	34 (-53, 122)	0.44	43 (-120, 34)	0.28	1.3 (-3.5, 6.1)	0.60	5.5 (0.09, 11)*	0.047*
Lag 1	53 (-39, 146)	0.26	-37 (-105, 31)	0.28	3.2 (-1.8, 7.8)	0.22	5.2 (-0.09, 11)*	0.05*
Lag 2	57 (-44, 159)	0.27	-67 (-136, 2.3)	0.06	2.9 (-1.9, 7.8)	0.23	5.2 (-0.2, 11)	0.06
Lag 3	30 (-91, 149)	0.63	-78 (-157, -2.5)*	0.04*	2.0 (-1.97, 6.4)	0.31	2.3 (-2.8, 7.8)	0.37
Lag 0-3	13 (-15, 42)	0.35	-16 (-35, 3.8)	0.10	0.6 (-0.6, 2.0)	0.29	1.2 (-0.1, 2.6)	0.07

N.B. FEV<sub>1</sub> and FeNO associations were adjusted for age, sex, height, relative humidity, temperature, ozone and season of testing. All results are interpreted as per increase in 29 grass pollen grains/m<sup>3</sup>. \*p $\leq$ 0.05.