

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Jaegle, I;Adachi, I;Aihara, H;Al Said, S;Asner, DM;Aushev, T;Ayad, R;Bakich, AM;Bansal, V;Barrett, M;Bhuyan, B;Bozek, A;Bračko, M;Browder, TE;Červenkov, D;Chang, MC;Cheon, BG;Chilikin, K;Cho, K;Chobanova, V;Choi, SK;Choi, Y;Cinabro, D;Dalseno, J;Doležal, Z;Drásal, Z;Drutskoy, A;Dutta, D;Eidelman, S;Epifanov, D;Farhat, H;Fast, JE;Ferber, T;Frost, O;Gaur, V;Gabyshev, N;Ganguly, S;Garmash, A;Getzkow, D;Gillard, R;Goh, YM;Golob, B;Grzymkowska, O;Hayasaka, K;Hayashii, H;He, XH;Hedges, M;Hou, WS;Iijima, T;Inami, K;Ishikawa, A;Iwasaki, Y;Julius, T;Kang, KH;Kato, E;Kawasaki, T;Kim, DY;Kim, JB;Kim, JH;Kim, SH;Kinoshita, K;Ko, BR;Kodyš, P;Korpar, S;Križan, P;Krokovny, P;Kuzmin, A;Kwon, YJ;Lange, JS;Lee, IS;Lewis, P;Li Gioi, L;Libby, J;Liventsev, D;Matvienko, D;Miyata, H;Mizuk, R;Mohanty, GB;Moll, A;Mussa, R;Nakano, E;Nakao, M;Nisar, NK;Nishida, S;Ogawa, S;Pakhlov, P;Pakhlova, G;Park, H;Pedlar, TK;Pesántez, L;Petrič, M;Piilonen, LE;Ritter, M;Rostomyan, A;Sakai, Y;Sandilya, S;Santelj, L;Sanuki, T;Sato, Y;Savinov, V

Title:

Search for the dark photon and the dark higgs boson at belle

Date:

2015-05-27

Citation:

Jaegle, I., Adachi, I., Aihara, H., Al Said, S., Asner, D. M., Aushev, T., Ayad, R., Bakich, A. M., Bansal, V., Barrett, M., Bhuyan, B., Bozek, A., Bračko, M., Browder, T. E., Červenkov, D., Chang, M. C., Cheon, B. G., Chilikin, K., Cho, K., ... Savinov, V. (2015). Search for the dark photon and the dark higgs boson at belle. *Physical Review Letters*, 114 (21), <https://doi.org/10.1103/PhysRevLett.114.211801>.

Persistent Link:

<https://hdl.handle.net/11343/116024>

Search for the dark photon and the dark Higgs boson at Belle

I. Jaegle,¹³ I. Adachi,^{14, 10} H. Aihara,⁵⁵ S. Al Said,^{50, 25} D. M. Asner,⁴⁴ T. Aushev,^{35, 22} R. Ayad,⁵⁰ A. M. Bakich,⁴⁹ V. Bansal,⁴⁴ M. Barrett,¹³ B. Bhuyan,¹⁶ A. Bozek,⁴¹ M. Bračko,^{31, 23} T. E. Browder,¹³ D. Červenkov,⁴ M.-C. Chang,⁷ B. G. Cheon,¹² K. Chilikin,²² K. Cho,²⁶ V. Chobanova,³² S.-K. Choi,¹¹ Y. Choi,⁴⁸ D. Cinabro,⁵⁹ J. Dalseno,^{32, 52} Z. Doležal,⁴ Z. Drásal,⁴ A. Drutskey,^{22, 34} D. Dutta,¹⁶ S. Eidelman,³ D. Epifanov,⁵⁵ H. Farhat,⁵⁹ J. E. Fast,⁴⁴ T. Ferber,⁶ O. Frost,⁶ V. Gaur,⁵¹ N. Gabyshev,³ S. Ganguly,⁵⁹ A. Garmash,³ D. Getzkow,⁸ R. Gillard,⁵⁹ Y. M. Goh,¹² B. Golob,^{30, 23} O. Grzymkowska,⁴¹ K. Hayasaka,³⁷ H. Hayashii,³⁸ X. H. He,⁴⁵ M. Hedges,¹³ W.-S. Hou,⁴⁰ T. Iijima,^{37, 36} K. Inami,³⁶ A. Ishikawa,⁵⁴ Y. Iwasaki,¹⁴ T. Julius,³³ K. H. Kang,²⁸ E. Kato,⁵⁴ T. Kawasaki,⁴² D. Y. Kim,⁴⁷ J. B. Kim,²⁷ J. H. Kim,²⁶ S. H. Kim,¹² K. Kinoshita,⁵ B. R. Ko,²⁷ P. Kodyš,⁴ S. Korpar,^{31, 23} P. Krizán,^{30, 23} P. Krokovny,³ A. Kuzmin,³ Y.-J. Kwon,⁶¹ J. S. Lange,⁸ I. S. Lee,¹² P. Lewis,¹³ L. Li Gioi,³² J. Libby,¹⁷ D. Liventsev,¹⁴ D. Matvienko,³ H. Miyata,⁴² R. Mizuk,^{22, 34} G. B. Mohanty,⁵¹ A. Moll,^{32, 52} R. Mussa,²¹ E. Nakano,⁴³ M. Nakao,^{14, 10} N. K. Nisar,⁵¹ S. Nishida,^{14, 10} S. Ogawa,⁵³ P. Pakhlov,^{22, 34} G. Pakhlova,²² H. Park,²⁸ T. K. Pedlar,⁶² L. Pesántez,² M. Petrič,²³ L. E. Piilonen,⁵⁸ M. Ritter,³² A. Rostomyan,⁶ Y. Sakai,^{14, 10} S. Sandilya,⁵¹ L. Santelj,¹⁴ T. Sanuki,⁵⁴ Y. Sato,³⁶ V. Savinov,⁴⁶ O. Schneider,²⁹ G. Schnell,^{1, 15} C. Schwanda,¹⁹ D. Semmler,⁸ K. Senyo,⁶⁰ O. Seon,³⁶ I. Seong,¹³ M. E. Sevir,³³ V. Shebalin,³ T.-A. Shibata,⁵⁶ J.-G. Shiu,⁴⁰ B. Shwartz,³ F. Simon,^{32, 52} R. Sinha,²⁰ Y.-S. Sohn,⁶¹ M. Starič,²³ M. Sumihama,⁹ K. Sumisawa,^{14, 10} U. Tamponi,^{21, 57} G. Tatishvili,⁴⁴ Y. Teramoto,⁴³ F. Thorne,¹⁹ M. Uchida,⁵⁶ S. Uehara,^{14, 10} Y. Unno,¹² S. Uno,^{14, 10} S. E. Vahsen,¹³ C. Van Hulse,¹ P. Vanhoefer,³² G. Varner,¹³ A. Vinokurova,³ M. N. Wagner,⁸ C. H. Wang,³⁹ M.-Z. Wang,⁴⁰ P. Wang,¹⁸ X. L. Wang,⁵⁸ M. Watanabe,⁴² Y. Watanabe,²⁴ K. M. Williams,⁵⁸ E. Won,²⁷ J. Yamaoka,⁴⁴ S. Yashchenko,⁶ Y. Yook,⁶¹ Y. Yusa,⁴² V. Zhilich,³ V. Zhulanov,³ and A. Zupanc²³

(The Belle Collaboration)

¹University of the Basque Country UPV/EHU, 48080 Bilbao

²University of Bonn, 53115 Bonn

³Budker Institute of Nuclear Physics SB RAS and Novosibirsk State University, Novosibirsk 630090

⁴Faculty of Mathematics and Physics, Charles University, 121 16 Prague

⁵University of Cincinnati, Cincinnati, Ohio 45221

⁶Deutsches Elektronen-Synchrotron, 22607 Hamburg

⁷Department of Physics, Fu Jen Catholic University, Taipei 24205

⁸Justus-Liebig-Universität Gießen, 35392 Gießen

⁹Gifu University, Gifu 501-1193

¹⁰The Graduate University for Advanced Studies, Hayama 240-0193

¹¹Gyeongsang National University, Chinju 660-701

¹²Hanyang University, Seoul 133-791

¹³University of Hawaii, Honolulu, Hawaii 96822

¹⁴High Energy Accelerator Research Organization (KEK), Tsukuba 305-0801

¹⁵IKERBASQUE, Basque Foundation for Science, 48011 Bilbao

¹⁶Indian Institute of Technology Guwahati, Assam 781039

¹⁷Indian Institute of Technology Madras, Chennai 600036

¹⁸Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049

¹⁹Institute of High Energy Physics, Vienna 1050

²⁰Institute of Mathematical Sciences, Chennai 600113

²¹INFN - Sezione di Torino, 10125 Torino

²²Institute for Theoretical and Experimental Physics, Moscow 117218

²³J. Stefan Institute, 1000 Ljubljana

²⁴Kanagawa University, Yokohama 221-8686

²⁵Department of Physics, Faculty of Science, King Abdulaziz University, Jeddah 21589

²⁶Korea Institute of Science and Technology Information, Daejeon 305-806

²⁷Korea University, Seoul 136-713

²⁸Kyungpook National University, Daegu 702-701

²⁹École Polytechnique Fédérale de Lausanne (EPFL), Lausanne 1015

³⁰Faculty of Mathematics and Physics, University of Ljubljana, 1000 Ljubljana

³¹University of Maribor, 2000 Maribor

³²Max-Planck-Institut für Physik, 80805 München

³³School of Physics, University of Melbourne, Victoria 3010

³⁴Moscow Physical Engineering Institute, Moscow 115409

³⁵Moscow Institute of Physics and Technology, Moscow Region 141700

³⁶Graduate School of Science, Nagoya University, Nagoya 464-8602

- ³⁷Kobayashi-Maskawa Institute, Nagoya University, Nagoya 464-8602
³⁸Nara Women's University, Nara 630-8506
³⁹National United University, Miao Li 36003
⁴⁰Department of Physics, National Taiwan University, Taipei 10617
⁴¹H. Niewodniczanski Institute of Nuclear Physics, Krakow 31-342
⁴²Niigata University, Niigata 950-2181
⁴³Osaka City University, Osaka 558-8585
⁴⁴Pacific Northwest National Laboratory, Richland, Washington 99352
⁴⁵Peking University, Beijing 100871
⁴⁶University of Pittsburgh, Pittsburgh, Pennsylvania 15260
⁴⁷Soongsil University, Seoul 156-743
⁴⁸Sungkyunkwan University, Suwon 440-746
⁴⁹School of Physics, University of Sydney, NSW 2006
⁵⁰Department of Physics, Faculty of Science, University of Tabuk, Tabuk 71451
⁵¹Tata Institute of Fundamental Research, Mumbai 400005
⁵²Excellence Cluster Universe, Technische Universität München, 85748 Garching
⁵³Toho University, Funabashi 274-8510
⁵⁴Tohoku University, Sendai 980-8578
⁵⁵Department of Physics, University of Tokyo, Tokyo 113-0033
⁵⁶Tokyo Institute of Technology, Tokyo 152-8550
⁵⁷University of Torino, 10124 Torino
⁵⁸CNP, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061
⁵⁹Wayne State University, Detroit, Michigan 48202
⁶⁰Yamagata University, Yamagata 990-8560
⁶¹Yonsei University, Seoul 120-749
⁶²Luther College, Decorah, Iowa 52101
(Dated: March 31, 2015 / version 39)

The dark photon, A' , and the dark Higgs boson, h' , are hypothetical constituents featured in a number of recently proposed Dark Sector Models. Assuming prompt decays of both dark particles, we search for their production in the so-called Higgs-strahlung channel, $e^+e^- \rightarrow A'h'$, with $h' \rightarrow A'A'$. We investigate ten exclusive final-states with $A' \rightarrow e^+e^-$, $\mu^+\mu^-$, or $\pi^+\pi^-$, in the mass ranges $0.1 \text{ GeV}/c^2 < m_{A'} < 3.5 \text{ GeV}/c^2$ and $0.2 \text{ GeV}/c^2 < m_{h'} < 10.5 \text{ GeV}/c^2$. We also investigate three inclusive final-states, $2(e^+e^-)X$, $2(\mu^+\mu^-)X$, and $(e^+e^-)(\mu^+\mu^-)X$, where X denotes a dark photon candidate detected via missing mass, in the mass ranges $1.1 \text{ GeV}/c^2 < m_{A'} < 3.5 \text{ GeV}/c^2$ and $2.2 \text{ GeV}/c^2 < m_{h'} < 10.5 \text{ GeV}/c^2$. Using the entire 977 fb^{-1} data set collected by Belle, we observe no significant signal. We obtain individual and combined 90% credibility level upper limits on the branching fraction times the Born cross section, $\mathcal{B} \times \sigma_{\text{Born}}$, on the Born cross section, σ_{Born} , and on the dark photon coupling to the dark Higgs boson times the kinetic mixing between the Standard Model photon and the dark photon, $\alpha_D \times \epsilon^2$. These limits improve upon and cover wider mass ranges than previous experiments. The limits from the final-states $3(\pi^+\pi^-)$ and $2(e^+e^-)X$ are the first placed by any experiment. For α_D equal to $1/137$, $m_{h'} < 8 \text{ GeV}/c^2$, and $m_{A'} < 1 \text{ GeV}/c^2$, we exclude values of the mixing parameter, ϵ , above $\sim 8 \times 10^{-4}$.

PACS numbers: 12.60.-i, 14.80.Ec, 14.60.-z, 14.40.Aq

Recent results from dedicated dark-matter searches [1–3], muon-spin precession measurements [4], and space-based particle observatories [5–7] may be interpreted as deviations from the Standard Model (SM) of particle physics. Attempts at devising unified explanations have led to Dark Sector Models (DSM) that introduce a new hidden or dark U(1) interaction that imbues dark matter with a novel charge [8–28]. A possible mediator of this new Abelian force is the dark photon, which has an expected mass of the order of $\text{MeV}/c^2 - \text{GeV}/c^2$ and has a very small kinetic mixing with the Standard Model photon, ϵ , of the order of $10^{-5} - 10^{-2}$ [13]. The dark U(1) symmetry group could be spontaneously broken, by a Higgs mechanism, adding a dark Higgs boson h' (or several of these) to such models [24].

Due to the small coupling to SM particles and the low expected mass of the dark photon, the ideal tools to discover the dark photon and the dark Higgs boson are low energy and high-luminosity experiments such as Belle at KEKB, Belle II at SuperKEKB [24], BaBar at PEP-II [29, 30], and dedicated fixed target and beam dump experiments, several of which are planned or under construction [31–37]. This article focuses on the Higgs-strahlung channel, $e^+e^- \rightarrow A'h'$. Generally, the dark photon A' can decay into lepton pairs, hadrons, or invisible particles while the dark Higgs boson h' can decay into either $A'A'^{(*)}$, leptons pairs, or hadrons, where $A'^{(*)}$ is a virtual dark photon. The decay modes of the A' and h' depend on their masses and decay lengths [24, 38]. There are three main cases: (a) $m_{h'} < m_{A'}$: h' is

long-lived and decays to lepton pairs or hadrons, (b) $m_{A'} < m_{h'} < 2m_{A'}$: $h' \rightarrow A'A'^*$, where A'^* decays into leptons, and (c) $m_{h'} > 2m_{A'}$: $h' \rightarrow A'A'$. This article is concerned with case (c); in particular, we investigate ten exclusive final-states of type $3(l^+l^-)$, $2(l^+l^-)(\pi^+\pi^-)$, $2(\pi^+\pi^-)(l^+l^-)$, and $3(\pi^+\pi^-)$, where l^+l^- is an electron or muon pair but not a tau pair, and three inclusive final-states of type $2(l^+l^-)X$, where X is a dark photon candidate detected via missing mass.

The Higgs-strahlung channel involves the effective coupling of the dark photon to SM particles, α' , induced via kinematic mixing with the SM photon, and the coupling of the dark-photon to the dark Higgs boson, α_D . KLOE and BaBar have reported searches for the dark photon and the dark Higgs boson [29, 39]: KLOE focused on $m_{h'} < m_{A'}$ and BaBar on $m_{h'} > 2m_{A'}$ (assuming prompt decays of the A' and h'), but no signal was found in either case. BaBar set limits on the product $\alpha_D \times \epsilon^2$ (where $\epsilon^2 = \alpha'/\alpha_{\text{em}}$ and α_{em} is the SM electromagnetic coupling constant) for dark photon and dark Higgs boson mass ranges of $0.25 - 3.0 \text{ GeV}/c^2$ and $0.8 - 10.0 \text{ GeV}/c^2$, respectively. Beam dump experiments [40–47] have placed 90% confidence level upper limits on ϵ for the processes $e^-p \rightarrow A'X'$ and $pp \rightarrow A'X'$ (where X' is not identified) of $\epsilon < 10^{-4}$ for a dark photon mass range of $1 - 300 \text{ MeV}/c^2$. Recently, BaBar [30] set an upper limit of $\epsilon < 3 \times 10^{-3}$ for a dark photon mass range of $0.3 - 10 \text{ GeV}/c^2$ for the radiative decay process $e^+e^- \rightarrow \gamma A'$. The advantage of the Higgs-strahlung channel compared to the radiative decay is that the Quantum Electrodynamics (QED) background is expected to be much smaller. If, in addition, the coupling between the dark photon and the dark Higgs boson is of order unity, then the Higgs-strahlung channel is the most sensitive probe for the dark photon.

Here, we report individual upper limits on the branching fraction times the Born cross section, $\mathcal{B} \times \sigma_{\text{Born}}$, for the thirteen aforementioned Higgs-strahlung final states as well as combined upper limits on σ_{Born} and on the product $\alpha_D \times \epsilon^2$ for these final-states, in the mass ranges $0.1 \text{ GeV}/c^2 < m_{A'} < 3.5 \text{ GeV}/c^2$ and $0.2 \text{ GeV}/c^2 < m_{h'} < 10.5 \text{ GeV}/c^2$, assuming prompt decays of the dark particles. We use data collected with the Belle detector [48] at the KEKB e^+e^- collider [49], amounting to 977 fb^{-1} at center-of-mass energies corresponding to the $\Upsilon(1S)$ to $\Upsilon(5S)$ resonances and in the nearby continuum.

We optimize the selection criteria and determine the $e^+e^- \rightarrow A'h'$ signal detection efficiency using a Monte Carlo (MC) simulation where the interaction kinematics and detector response are simulated with the packages MadGraph [50] and GEANT3 [51], respectively. There is no suitable background simulation available, so background samples are taken from data sidebands.

We choose loose particle identification criteria to enhance the detection efficiency of final states with leptons. To ensure that only prompt decays are selected, *i.e.*, that




FIG. 1: Example $m_{A'_{\text{cand}}}^1 - m_{A'_{\text{cand}}}^3$ distribution for the $A'h' \rightarrow A'A'A' \rightarrow 6\pi$ channel, for $m_{A'_{\text{cand}}}^1 = 2.0 \pm 0.1 \text{ GeV}/c^2$, where $m_{A'_{\text{cand}}}^1$ and $m_{A'_{\text{cand}}}^3$ are the dark photon candidates with the highest and lowest mass, respectively. The “same-sign” distributions (blue), where at least one A' candidate is reconstructed from $\pi^+\pi^+$ or $\pi^-\pi^-$, are normalized to the “opposite-sign” $3(\pi^+\pi^-)$ distributions (red) in the sidebands, and are used to predict the background in the signal region.

the decay of each A' candidate occurs near the e^+e^- interaction point (IP), we require that the vertex fit of all tracks detected in the event be consistent with an origin at the IP, and that each track have impact parameters $|dz| < 1.5 \text{ cm}$ and $dr < 0.2 \text{ cm}$, where dz is measured along the positron beam (collinear with the z axis) and dr is measured in the transverse r - ϕ plane. We also require that the second-order Fox-Wolfman moment [52] satisfy $R_2 < 0.9$, and that the electron helicity angle, α_e , in the A' rest frame satisfy $\cos(\alpha_e) < 0.9$, as in Ref. [29].

For exclusive channels, we select final states with exactly three pairs of oppositely charged particles. For inclusive channels, we select final states of the type $2(l^+l^-)X$, where X is constrained by the missing mass of the event and contains zero, one, or two reconstructed tracks that are not identified as leptons or pions. We require that both $m_{l^+l^-}$ and m_X be greater than $1.1 \text{ GeV}/c^2$. Above this mass, the branching fraction of A' to hadronic final-states other than charged pion pairs is dominant [24]. We refer to events selected according to these criteria as “opposite-sign” to distinguish them from the “same-sign” events used for background estimation.

For exclusive final states, we select candidate events with final-state masses between 98% and 105% of the initial-state mass. For inclusive channels, where this condition cannot be applied, we perform a missing-mass analysis: X is treated as an unobserved particle whose missing four-momentum is given by

$$P_X = P_{e^+e^-} - P_{A'_{\text{cand}} \rightarrow l^+l^-}^1 - P_{A'_{\text{cand}} \rightarrow l^+l^-}^2, \quad (1)$$

where $P_{e^+e^-}$ and $P_{A'_{\text{cand}} \rightarrow l^+l^-}^{1,2}$ are the four-momenta of




FIG. 2: (a) : Signal candidates observed versus dark photon candidate mass, $m_{A'_{cand}}$, and dark Higgs boson candidate mass, $m_{A'_{cand} A'_{cand}}$, for the 13 final states. There are three entries per event. (b) and (c): Projection of signal candidates onto $m_{A'_{cand} A'_{cand}}$ and $m_{A'_{cand}}$ (red points) with the predicted background (blue squares) from the scaled same-sign distributions for comparison. The dark photon candidate mass distribution has been scaled by $1/3$. (d): Normalized residuals between the signal candidate distribution and predicted background, versus dark photon candidate mass (red points) and dark Higgs boson candidate mass (black squares). The same-sign error bars contain statistical and systematic errors. For empty bins, the systematic error is one event.

the initial-state and the two fully reconstructed dark photon candidates, respectively. The mass m_X of the missing four-momentum P_X is then compared to the reconstructed masses of dark photon candidates 1 and 2 using:

$$\Delta m = m_X - (m_{A'_{cand} \rightarrow l^+ l^-}^1 + m_{A'_{cand} \rightarrow l^+ l^-}^2)/2. \quad (2)$$

We select inclusive final-states by requiring

$$\Delta m_{\min} < \Delta m < \Delta m_{\max}, \quad (3)$$

where the optimized limits $\Delta m_{\min}$ and $\Delta m_{\max}$ each depend on the measured mean mass of dark photon candidates 1 and 2 and on the particular final state.

For exclusive (inclusive) final-states, we then require the invariant masses of dark photon candidates, $m_{A'_{cand}}$, to be consistent with three (two) distinct $A' \rightarrow l^+ l^-$ or $\pi^+ \pi^-$ decays. Signal candidates with three (two) consistent dark photon masses are kept by requiring

$$m_{A'_{cand}}^{\min} < m_{A'_{cand}} < m_{A'_{cand}}^{\max}, \quad (4)$$

where the optimized limits $m_{A'_{cand}}^{\min}$ and $m_{A'_{cand}}^{\max}$ each depend on the measured mean mass of the three (two) fully

reconstructed dark photon candidates and on the simulated width of the invariant mass distribution of the dark photon for that mass.

For each event, if there is more than one signal candidate that fulfills the selection criteria for a given final state, we select the candidate with the smallest Δm . For exclusive channels, we use: $\Delta m = \sum_1^3 \Delta m_i^2$ with

$$\Delta m_i = m_{A'_{cand}}^i - (m_{A'_{cand}}^1 + m_{A'_{cand}}^2 + m_{A'_{cand}}^3)/3. \quad (5)$$

If an event satisfies the selection criteria for multiple final states, we allocate the event to a single final state to ensure that the datasets for each final state are statistically independent. This is accomplished by selecting the lowest numbered final-state category from the following list: (1) exclusive with 6 leptons, (2) exclusive with four leptons, (3) exclusive with two leptons, (4) exclusive with six pions, and (5) inclusive final-states. For the signal MC simulation, the fraction of events with multiple signal candidates ranges from 7% to 15% in the channels where we need to apply this ordering. For data, the fraction is below 0.5% in all final states.

We optimize the event selection, including particle identification, the final-state mass requirements, and the parameters $\Delta m_{\min}$, $\Delta m_{\max}$, $m_{A'_{cand}}^{\min}$ and $m_{A'_{cand}}^{\max}$ using the signal MC simulation only. Events reconstructed as described above are used for signal. Background distributions are derived from the same event sample, by using events where at least one dark photon candidate is reconstructed from two tracks with charges of the same sign, enforcing all selection criteria except charge conservation. We refer to these as “same-sign” events. We verify that the background estimation is consistent with data as shown in Fig. 1. We generate MC with specific dark photon and dark Higgs boson masses and interpolate between samples where necessary. The detection efficiencies are 20% and 30%, on average, for the $3(e^+e^-)$ and $3(\mu^+\mu^-)$ final-states, respectively.

For setting limits, we also estimate the background using “same-sign” events, but in this case they are from experimental data. We sort the dark photon candidates by mass in descending order, $m_{A'_{cand}}^1 > m_{A'_{cand}}^2 > m_{A'_{cand}}^3$, and calculate the mass difference $m_{A'_{cand}}^1 - m_{A'_{cand}}^3$. We divide the data into different bins of $m_{A'_{cand}}^1$, with each bin analyzed separately. We divide the $m_{A'_{cand}}^1 - m_{A'_{cand}}^3$ distribution into two regions: signal and sideband. The signal region size is determined by equation (4). The sideband region starts at 1.5 times and ends at 5.0 times the signal-region upper limit. Figure 1 shows the distribution of the mass difference $m_{A'_{cand}}^1 - m_{A'_{cand}}^3$ for the bin $m_{A'_{cand}}^1 = 2.0 \pm 0.1$ GeV/c² for the six-pion final-state. We assume that, in the absence of signal, the same-sign and the opposite-sign distributions have the same shape (but different yields) in both the signal region and the sideband. Therefore, for each $m_{A'_{cand}}^1$ bin, the same-sign distribu-




FIG. 3: Left: 90% CL upper limit on the product $\mathcal{B} \times \sigma_{\text{Born}}$ for each of the 13 final states considered versus dark photon mass for different hypotheses for the dark Higgs boson mass. Black, red, green, blue and yellow curves correspond to $m_{h'} = 1, 3, 5, 7$, and 9 GeV/c², respectively, for exclusive channels and $m_{h'} = 3, 4, 5, 7$, and 9 GeV/c², respectively, for inclusive channels. Right: 90% CL upper limit on the cross section of $e^+e^- \rightarrow A'h'$, $h' \rightarrow A'A'$ versus dark photon and dark Higgs boson mass.

TABLE I: Number of events observed after all selection criteria are applied.

Final-state	Events	Final-state	Events
$3(e^-e^+)$	1	$2(\mu^+\mu^-)(e^+e^-)$	1
$3(\mu^+\mu^-)$	2	$2(\mu^+\mu^-)(\pi^+\pi^-)$	1
$3(\pi^+\pi^-)$	147	$2(\pi^+\pi^-)(e^+e^-)$	5
$2(e^+e^-)(\mu^+\mu^-)$	7	$2(\pi^+\pi^-)(\mu^+\mu^-)$	6
$2(e^+e^-)(\pi^+\pi^-)$	2	$(e^+e^-)(\mu^+\mu^-)(\pi^+\pi^-)$	7
$2(e^+e^-)X$	572	$(e^+e^-)(\mu^+\mu^-)X$	30
$2(\mu^+\mu^-)X$	20		

tion (blue squares) is scaled so that the number of events in the sideband agree with the number of opposite-sign events (red points) in the sideband. The expected background in the signal region is then the scaled number of events of the same-sign distribution in that region. This procedure is illustrated by Fig. 1. The opposite-sign and scaled same-sign distributions are consistent in the signal region and the sideband. In the presence of signal, we would expect an excess of opposite-sign events over the predicted background in the signal region, as can be seen for the signal MC distribution. Figure 2 summarizes the background estimation. Figure 2 (a) shows the distribution of events measured as a function of the dark photon candidate mass, $m_{A'_{\text{cand}}}$, and the dark Higgs boson candidate mass, $m_{A'_{\text{cand}}A'_{\text{cand}}}$. Table I shows the number of events observed after all selection criteria are applied.

Figures 2 (b) and (c) show the projections on the mass axis of the dark Higgs boson and dark photon, respec-

tively. The number of events observed in the signal region, N_{obs} , and the number of predicted background events, N_{bkg} , are in good agreement. Their differences are quantified by the normalized residuals, shown in Fig. 2 (d) and defined as $(N_{\text{obs}} - N_{\text{bkg}})/\sqrt{\sigma_{\text{obs}}^2 + \sigma_{\text{bkg}}^2}$, where σ_{obs} and σ_{bkg} are the standard deviations of the distributions. In all cases, the number of events observed is consistent with the background estimate. For exclusive final states, the background is mostly due to processes with ρ and ω resonance particles, such as SM 2γ processes. The discontinuity at 1.1 GeV/c² in Fig. 2 (c) is an artifact of the selection criteria.

The upper limits on $\mathcal{B} \times \sigma_{\text{Born}}$ and σ_{Born} are calculated for ranges of $m_{A'}$ and $m_{h'}$, based on the signal MC mass resolution, with a Bayesian inference method with the use of Markov Chain Monte Carlo [53]. The number of observed events can be expressed as:

$$N_{\text{obs}} = \sigma_{\text{Born}} \cdot (1 + \delta) \cdot |1 - \Pi|^2 \cdot \mathcal{L} \cdot \mathcal{B} \cdot \varepsilon + N_{\text{bkg}}, \quad (6)$$

where $1 + \delta$ is an initial-state radiative correction factor, $|1 - \Pi|^2$ is the vacuum polarization factor, $\mathcal{L}$ is the luminosity, ε is the detection efficiency, and N_{bkg} is the number of predicted background events. We calculate, for the exclusive (inclusive) channels, $1 + \delta$ using the formulae in Ref. [54] and assuming the theoretical cross section is proportional to $1/s$ [24], where s is the square of the initial-state mass, and also assuming a cut-off value corresponding to 98% (a value between 20% and 90%) of the initial-state mass. $1 + \delta$ varies from 0.804 (0.93) to 0.807 (1.17) depending on s and for the inclusive chan-




FIG. 4: 90% CL upper limit on the product $\alpha_D \times \epsilon^2$ versus dark photon mass (top row) and dark Higgs boson mass (bottom row) for Belle (solid red curve) and BaBar [29] (dashed black curve). BaBar limits should be divided by $(1 + \delta)$ before being compared with Belle limits. The blue dotted curve, which coincides more or less with the solid red curve, shows the expected Belle limit.

nels also the effective cut-off value. We use $1 + \delta = 0.8055$ (1.0) and include the variation as a systematic error in the upper limit calculation. The value of $|1 - \Pi(s)|^2$ is taken from Ref. [55, 56] and varies between 0.9248 and 1.072 depending on s . For $\mathcal{B} \times \sigma_{\text{Born}}$ and σ_{Born} , logarithmic priors are used, and for $1 + \delta$, $|1 - \Pi|^2$, $\mathcal{L}$, $\mathcal{B}$, ϵ , and N_{bkg} Gaussian priors are used to take into account the systematic uncertainty. In Fig. 3, the left panel shows the 90% credibility level (CL)¹ upper limits on $\mathcal{B} \times \sigma_{\text{Born}}$ versus the dark photon mass, for different hypotheses of the dark Higgs boson mass, for each of the 13 final states considered, while the right panel shows the combined upper limit on σ_{Born} for $e^+e^- \rightarrow Ah'$ versus the dark photon and dark Higgs boson mass. For the combined limit, compared to BaBar, we use two extra channels, $3(\pi^+\pi^-)$ and $2(e^+e^-)X$, which contribute 91% of our background. The branching fractions were taken from Ref. [24].

The combined limit can also be expressed as a limit on the product $\alpha_D \times \epsilon^2$ by using the equations described in Ref. [24]. Figure 4 shows the 90% CL upper limits on $\alpha_D \times \epsilon^2$ for Belle, expected and measured, and for BaBar, for five different mass hypotheses for the dark Higgs boson (top row) and dark photon (bottom row) masses. Note that the BaBar limits were based on the visible cross section, rather than the Born cross section. For the expected limit, we assume: $N_{\text{obs}} = N_{\text{bkg}}$.

The inclusion of the $3(\pi^+\pi^-)$ final state dramatically improves the limit around the ρ and ω resonances. The dominant sources of systematic uncertainties are: the integrated luminosity (1%), branching fractions (4%), track identification (6%), particle identification efficiency (5%), detection efficiency (15%), background estimation (10%) and initial-state radiation (15%). All systematic uncertainties added in quadrature amount to 25%.

In summary, we search for the dark photon and the dark Higgs boson in the mass ranges $0.1 - 3.5 \text{ GeV}/c^2$ and $0.2 - 10.5 \text{ GeV}/c^2$, respectively. No significant signal is observed. We obtain individual and combined 90% CL upper limits on the product of branching fraction times the Born cross section, $\mathcal{B} \times \sigma_{\text{Born}}$, on the Born cross section, σ_{Born} , and on the product of the dark photon coupling to the dark Higgs boson and the kinetic mixing between the Standard Model photon and the dark photon, $\alpha_D \times \epsilon^2$. These limits improve upon and cover wider mass ranges than previous experiments and the limits in the final-states $3(\pi^+\pi^-)$ and $2(e^+e^-)X$, where X is a dark photon candidate detected via missing mass, are the first limits placed by any experiment. For α_D equal to $1/137$, $m_{h'} < 8 \text{ GeV}/c^2$, and $m_{A'} < 1 \text{ GeV}/c^2$, we exclude values of the mixing parameter, ϵ , above $\sim 8 \times 10^{-4}$. In the mass ranges, and for modes, where previous measurements from BaBar exist, the limits reported here are almost a factor of two smaller. Since the backgrounds are very low to non-existent, the improvement scales nearly linearly with the integrated luminosity. This bodes well for future searches with Belle II.

We thank the KEKB group for excellent operation of the accelerator; the KEK cryogenics group for effi-

¹ In common High Energy Physics usage, this credibility level has been reported as “confidence level”, which is a frequentist-statistics term.

cient solenoid operations; and the KEK computer group, the NII, and PNNL/EMSL for valuable computing and SINET4 network support. We acknowledge support from MEXT, JSPS and Nagoya's TLPSC (Japan); ARC (Australia); FWF (Austria); NSFC (China); MSMT (Czechia); CZF, DFG, and VS (Germany); DST (India); INFN (Italy); MOE, MSIP, NRF, GSDC of KISTI, and BK21Plus (Korea); MNiSW and NCN (Poland); MES and RFAAE (Russia); ARRS (Slovenia); IKERBASQUE and UPV/EHU (Spain); SNSF (Switzerland); NSC and MOE (Taiwan); and DOE and NSF (USA). We would also like to thank Bertrand Echenard for discussing the BaBar analysis and Rouven Essig for providing the models for MadGraph.

-
- [1] R. Bernabei *et al.* [DAMA Collaboration], *Eur. Phys. J. C* **56**, 333 (2008).
 - [2] Z. Ahmed *et al.* [CDMS-II Collaboration], arXiv:0912.3592 (2010).
 - [3] C.E. Aalseth *et al.* [CoGeNT Collaboration], *Phys. Rev. Lett.* **106**, 131301 (2011).
 - [4] G.W. Bennett *et al.* [Muon g-2 Collaboration], *Phys. Rev. D* **73**, 072003 (2006).
 - [5] O. Adriani *et al.* [PAMELA Collaboration], *Nature* **458**, 607-609 (2009).
 - [6] M. Ackermann *et al.* [Fermi LAT. Collaboration], *Phys. Rev. D* **82**, 092004 (2010).
 - [7] M. Aguilar *et al.* [AMS Collaboration], *Phys. Rev. Lett.* **110**, 141102 (2013).
 - [8] P. Fayet, *Phys. Lett. B* **95**, 285 (1980).
 - [9] P. Fayet, *Nucl. Phys. B* **187**, 184 (1981).
 - [10] P. Fayet and J. Kaplan, *Phys. Lett. B* **269**, 213 (1991).
 - [11] C. Boehm and P. Fayet, *Nucl. Phys. B* **683**, 219 (2004).
 - [12] P. Fayet, *Phys. Rev. D* **70**, 023514 (2004).
 - [13] N. Arkani-Hamed, D.P. Finkbeiner, T.R. Slatyer and N. Weiner, *Phys. Rev. D* **79**, 015014 (2009).
 - [14] M. Pospelov, A. Ritz and M. Voloshin, *Phys. Lett. B* **662**, 53 61b (2008).
 - [15] E.J. Chun and J.C. Park, *JCAP* **0902**, 026 (2009).
 - [16] C. Cheung, J.T. Ruderman, L.T. Wang, and I. Yavin, *Phys. Rev. D* **80**, 035008 (2009), arXiv:0902.3246.
 - [17] A. Katz and R. Sundrum, *JHEP* **0906**, 003 (2009).
 - [18] D. Morrissey, D. Poland and K. Zurek, *JHEP* **0907**, 050 (2009).
 - [19] M. Goodsell, J. Jaeckel, J. Redondo, and A. Ringwald, *JHEP* **0911**, 027 (2009).
 - [20] M. Baumgart, C. Cheung, L.-T. Wang, J. Ruderman, I. Yavin, *JHEP* **0904**, 014 (2009).
 - [21] Y. Nomura and J. Thaler, *Phys. Rev. D* **79**, 075008 (2009).
 - [22] D. Alves, S. Behbabani, P. Schuster, and J. Wacker, [hep-ph] SU-ITP-09/13 (2009), arXiv:0903.3945.
 - [23] J. Jaeckel, A. Ringwald, *Ann. Rev. Nucl. Part. Sci.* **60**, 405 (2010).
 - [24] B. Batell, M. Pospelov, and A. Ritz, *Phys. Rev. D* **79**, 115008 (2009).
 - [25] M. Reece and L. T. Wang, *JHEP* **0907**, 051 (2009).
 - [26] R. Essig, P. Schuster, and N. Toro, *Phys. Rev. D* **80**, 015003 (2009).
 - [27] F. Bossi, arXiv:0904.3815 (2009).
 - [28] P.-f. Yin, J. Liu, and S.-h. Zhu, *Phys. Lett. B* **679**, 362-368 (2009).
 - [29] J. P. Lees *et al.* [BaBar Collaboration], *Phys. Rev. Lett.* **108**, 211801 (2012).
 - [30] J.P. Lees *et al.* [BaBar Collaboration], arXiv:1406.2980 (2014).
 - [31] S. Abrahamyan *et al.* [APEX Collaboration], *Phys. Rev. Lett.* **107**, 191804 (2011).
 - [32] J.L. Hewett *et al.* [HPS Collaboration], ANL-HEP-TR-12-25, SLAC-R-991, arXiv:1205.2671 (2012).
 - [33] J. Balewski *et al.* [DarkLight Collaboration], (2013), arXiv:1307.4432.
 - [34] M. Battaglieri *et al.* [BDX Collaboration], arXiv:1406.3028 (2014).
 - [35] H. Merkel *et al.* [A1 Collaboration], *Phys. Rev. Lett.* **106**, 251802 (2011).
 - [36] H. Merkel *et al.* [A1 Collaboration], *Phys. Rev. Lett.* **112**, 221802 (2014).
 - [37] T. Beranek, H. Merkel and M. Vanderhaeghen, *Phys. Rev. D* **88**, 015032 (2013).
 - [38] R. Essig, R. Harnik, J. Kaplan, N. Toro, *Phys. Rev. D* **82**, 113008 (2010).
 - [39] S. Giovannella "KLOE Collaboration", *J. Phys. Conf. Ser.* **335**, 012067 (2011).
 - [40] J. Blümlein *et al.*, *Z. Phys. C* **51**, 341 (1991).
 - [41] J. Blümlein *et al.*, *Int. J. Mod. Phys. A* **7**, 3835 (1992).
 - [42] J. Blümlein and J. Brunner, arXiv:1104.2747 (2011).
 - [43] J. Blümlein and J. Brunner, *Phys. Lett. B* **731** (2014) 320, arXiv:1311.3870.
 - [44] L. Barabash *et al.*, *Phys. Lett. B* **295**, 154 (1992).
 - [45] J. D. Bjorken *et al.* [E137 Collaboration], *Phys. Rev. D* **38**, 3375 (1988) .
 - [46] E. M. Riordan *et al.* [E141 Collaboration], *Phys. Rev. Lett.* **59**, 755 (1987) .
 - [47] A. Bross *et al.* [E774 Collaboration] *Phys. Rev. Lett.* **67**, 2942 (1991).
 - [48] A. Abashian *et al.* [Belle Collaboration], *Nucl. Instrum. Methods Phys. Res. Sect. A* **479**, 117 (2002); also see detector section in J.Brodzicka *et al.*, *Prog. Theor. Exp. Phys.* **2012**, 04D001 (2012).
 - [49] S. Kurokawa and E. Kikutani, *Nucl. Instrum. Methods Phys. Res. Sect. A* **499**, 1 (2003), and other papers included in this Volume; T. Abe *et al.*, *Prog. Theor. Exp. Phys.* **2013**, 03A001 (2013) and following articles up to 03A011.
 - [50] J. Alwall *et al.*, *J. HEP* **07**, 079 (2014).
 - [51] R. Brun *et al.*, GEANT, Cern/DD/ee/84-1, (1986).
 - [52] G.C. Fox and S. Wolfram, *Phys. Rev. Lett.* **41**, 1581 (1978).
 - [53] A. Caldwell, D. Kollar, K. Kröninger, BAT -The Bayesian Analysis Toolkit Computer Physics Communications, **180**, 2197-2209 (2009).
 - [54] E.A. Kuraev and V.S. Fadin, *Sov. J. Nucl. Phys.* **41**, 466 (1985) [*Yad. Fiz.* **41**, 733 (1985)].
 - [55] S. Actis *et al.*, *Eur. Phys. J. C* **66**, 585 (2010).
 - [56] F. Ignatov, <http://cmd.inp.nsk.su/~ignatov/vpl/>