



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Jiang, F;Liu, GS;Dusting, GJ;Chan, EC

Title:

NADPH oxidase-dependent redox signaling in TGF- β -mediated fibrotic responses

Date:

2014-01-01

Citation:

Jiang, F., Liu, G. S., Dusting, G. J. & Chan, E. C. (2014). NADPH oxidase-dependent redox signaling in TGF- β -mediated fibrotic responses. *Redox Biology*, 2 (1), pp.267-272. <https://doi.org/10.1016/j.redox.2014.01.012>.

Persistent Link:

<https://hdl.handle.net/11343/268843>

License:

[CC BY-NC-ND](#)



ELSEVIER

Contents lists available at ScienceDirect

Redox Biology

journal homepage: www.elsevier.com/locate/redox

Graphical Review

NADPH oxidase-dependent redox signaling in TGF- β -mediated fibrotic responses [☆]Fan Jiang ^{a,*}, Guei-Sheung Liu ^{b,c}, Gregory J. Dusting ^{b,c}, Elsa C. Chan ^{b,**}^a Key Laboratory of Cardiovascular Remodeling and Function Research, Qilu Hospital, Shandong University, Jinan, Shandong 250012, China^b Centre for Eye Research Australia, University of Melbourne, VIC 3002, Australia^c Department of Ophthalmology, University of Melbourne, VIC 3002, Australia

ARTICLE INFO

Article history:

Received 13 January 2014

Accepted 14 January 2014

Available online 20 January 2014

Keywords:

NADPH oxidase

Nox4

Transforming growth factor- β

Fibrosis

Redox signaling

ABSTRACT

Uncontrolled fibrosis in organs like heart, kidney, liver and lung is detrimental and may lead to end-stage organ failure. Currently there is no effective treatment for fibrotic disorders. Transforming growth factor (TGF)- β has a fundamental role in orchestrating the process of fibrogenesis; however, interventions directly targeting TGF- β would have undesired systemic side effects due to the multiple physiological functions of TGF- β . Further characterization of the downstream signaling pathway(s) involved in TGF- β -mediated fibrosis may lead to discovery of novel treatment strategies for fibrotic disorders. Accumulating evidence suggests that Nox4 NADPH oxidase may be an important downstream effector in mediating TGF- β -induced fibrosis, while NADPH oxidase-dependent redox signaling may in turn regulate TGF- β /Smad signaling in a feed-forward manner. It is proposed that pharmacological inhibition of the Nox4 function may represent a novel approach in treatment of fibrotic disorders.

© 2014 The Authors. Published by Elsevier B.V. All rights reserved.

Contents

TGF- β and fibrotic disorders	267
Involvement of NADPH oxidase in TGF- β -mediated profibrotic effects	268
Redox modulation of TGF- β signaling	269
Mechanisms of TGF- β regulation of Nox4 expression	271
Acknowledgements	271
References	271

TGF- β and fibrotic disorders

Fibrosis is caused by uncontrolled wound healing responses during tissue repair in pathological conditions such as myocardial infarction, idiopathic pulmonary fibrosis, hepatitis, and chronic kidney disease [1]. Fibrotic scars are characterized by accumulation

of contractile matrix proteins and tend to compromise normal tissue functions. One of the major profibrotic cytokines orchestrating fibrogenesis is transforming growth factor (TGF)- β (see Fig. 1). More than two decades ago it was shown that an exogenous application of TGF- β to incisional wounds accelerated production of collagen and improved the wound healing efficiency [2]. TGF- β is a multifunctional protein, and one of its major biological effects is to promote the recruitment of fibroblasts to wound site and synthesis of structural matrix proteins [3]; both processes are crucial to fibrogenesis (Fig. 1). TGF- β is produced by infiltrating inflammatory cells (for example macrophages), parenchymal cells, and platelets during tissue repair [4,5]. The profibrotic activity of TGF- β can be further regulated by other endogenous factors such as angiotensin II [6].

A major source of fibroblasts identified in scarred tissues is proliferation and migration of resident fibroblasts to the injured

[☆]This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-No Derivative Works License, which permits non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

* Corresponding author. Tel.: +86 531 8216 9257; fax: +86 531 8616 9356.

** Corresponding author. Centre for Eye Research Australia (CERA), 1/32, Gisborne Street, East Melbourne, VIC 3002, Australia. Tel.: +61 3 9929 8360; fax: +61 3 9416 0926.

E-mail addresses: fjiang@sdu.edu.cn (F. Jiang), elsa.chan@unimelb.edu.au (E.C. Chan).




Fig. 1. How TGF- β stimulates fibrotic responses during tissue repair. TGF- β is released by platelets, parenchymal cells and macrophages or its activity stimulated by endogenous factors such as angiotensin II. Under the influence of TGF- β , proliferation and migration of resident fibroblasts and the process of epithelial to mesenchymal transition (EMT) occur, while fibroblasts differentiate into myofibroblasts in the wound site. Differentiated myofibroblasts have a higher efficiency in synthesizing extracellular matrix proteins. Accumulation of extracellular matrix is enhanced by the inhibitory and stimulatory effects of TGF- β on expression of matrix metalloproteinases and their inhibitors respectively.

site, partially stimulated by TGF- β [7,8], while other sources may include epithelial to mesenchymal transition (EMT) and cells derived from the bone marrow [9]. Iwano et al. used bone marrow chimeras and transgenic reporter mice to trace origins of fibroblasts in fibrotic kidneys; two populations of fibroblasts have been identified, with the majority being derived from local EMT and a minor proportion being from the bone marrow [9]. TGF- β mediates EMT in a variety of organs including lung, liver, kidney, heart and eye [10]. Endothelial to mesenchymal transition mediated by TGF- β has also been observed and may play a role in cardiac fibrosis [11]. Hence, TGF- β is a crucial mediator for the generation and mobilization of fibroblasts required for fibrogenesis. Fibroblasts identified in scarred tissues are responsible for the production of matrix proteins [12]. TGF- β promotes accumulation of these proteins by inducing a phenotypic change of fibroblast to a highly efficient matrix-producing type termed myofibroblast [3]. To prevent degradation of the newly synthesized matrix proteins, TGF- β also inhibits expression of matrix catabolizing enzymes such as matrix metalloproteinases, and induces expression of matrix metalloproteinase inhibitors such as the tissue inhibitor of metalloproteinase [4].

Interventions directly targeting TGF- β would have undesired systemic side effects due to the multiple physiological functions of TGF- β [13,14]. Therefore, further characterization of the downstream signaling pathway(s) involved in TGF- β -induced fibrotic process may provide useful information in the discovery of novel treatment strategies for various fibrotic disorders.

Involvement of NADPH oxidase in TGF- β -mediated profibrotic effects

Accumulative evidence highlights the involvement of NADPH oxidase-dependent redox signaling in the profibrotic responses mediated by TGF- β [15]. Five isoforms of the Nox catalytic subunit, namely Nox1, Nox2, Nox3, Nox4 and Nox5, have been identified [16]. The prototypical NADPH oxidase is composed of the membrane-bound Nox and p22phox, as well as the cytosolic subunits p40phox, p47phox, p67phox and a small GTPase Rac (Fig. 2 insert) [16]. Whereas all of the Nox isoforms except for Nox3 have been shown to be expressed in fibroblasts from different organs (including blood vessels, heart, lung and kidney), current evidence suggests that the Nox4 type NADPH oxidase may have a pivotal role in mediating TGF- β -induced profibrotic responses [15,17–22].

TGF- β increased Nox4 gene expression without effects on Nox1, Nox2 or Nox5 in human cardiac fibroblasts [18]. Treatment with siRNA against Nox4 suppressed expression of TGF- β target genes including fibronectin, collagen I, α -smooth muscle actin (α -SMA, a marker of myofibroblast), and connective tissue growth factor, indicating that Nox4 was involved in TGF- β -induced differentiation of cardiac fibroblasts to myofibroblasts [18]. In renal tubular epithelial cells, Rhyu et al. showed that inhibition of NADPH oxidase activity with the non-selective Nox inhibitor DPI suppressed EMT and matrix protein production [23,24]. In a recent study, we showed that in mouse cardiac fibroblasts, TGF- β -stimulated collagen synthesis and myofibroblast differentiation




Fig. 2. Involvement of Nox4 type NADPH oxidase in TGF- β -induced profibrotic responses. Nox4-derived ROS facilitate TGF- β -mediated fibrosis by inducing differentiation of fibroblasts into myofibroblasts and synthesis of extracellular matrix proteins, while inhibiting their degradation. Nox4-derived ROS also promote apoptosis of epithelial cells. It remains to be confirmed whether Nox4 signaling is also involved in TGF- β -mediated epithelial to mesenchymal transition (EMT). It is proposed that selective inhibitors of Nox4 such as GKT-136901 and compound 88 may represent a novel strategy to combat fibrotic disorders. The inset illustrates the structure of the prototype Nox2.

were abrogated by EUK-134 (a scavenger of both superoxide and hydrogen peroxide) or a dominant negative form of Nox4 [17]. Such modulating effects of Nox4 in TGF- β -mediated collagen expression have also been observed in fetal lung mesenchymal cells [19], pulmonary fibroblasts [25], kidney fibroblasts [20] and liver stellate cells [26]; all of these cells are engaged in the development of fibrosis in the affected organs. Moreover, there is evidence showing that Nox4 mRNA and protein are increased in pulmonary fibroblasts isolated from patients with idiopathic pulmonary fibrosis, and the Nox4 expression level correlates with expressions of procollagen I and α -SMA [25].

Amara et al. showed that targeted gene silencing of Nox4 with tracheal administration of siRNA suppressed bleomycin-induced pulmonary fibrosis in mice [25]. Similarly, Carnesecchi et al. reported that bleomycin-induced pulmonary fibrosis was prevented in Nox4 knockout mice, and this protective effect of Nox4 deficiency was attributable to a decrease in epithelial cell apoptosis induced by TGF- β [27]. Moreover, we used a mouse model of subcutaneous sponge implantation and demonstrated that local inhibition of Nox4 expression with RNA interference suppressed the stimulatory effects of TGF- β on collagen accumulation in vivo [17]. These in vitro and in vivo findings support the notion that Nox4 has a pivotal role in mediating the profibrotic actions of TGF- β (Fig. 2).

Currently, a challenge is that we lack isoform-specific NADPH oxidase inhibitors. GKT-136901 is an orally active selective inhibitor of Nox1 and Nox4 [28], which has been shown to be able to suppress liver fibrosis induced by bile duct ligation [29]. Jarman et al. recently showed that oral treatment with a putative Nox4 inhibitor known as compound 88 attenuated bleomycin-induced lung fibrosis in rats and TGF- β -induced induction of procollagen and α -SMA expression in human pulmonary fibroblasts [30]. These studies suggest that pharmacological inhibition of the Nox4 function may represent a novel strategy in treatment of fibrotic disorders.

Redox modulation of TGF- β signaling

In addition to a potential role of NADPH oxidase as downstream mediators in TGF- β -induced profibrotic effects, there is also evidence suggesting that redox pathways may regulate TGF- β /Smad signaling in a feed-forward manner. For example, using cell-free assays people have shown that irradiation or ion-catalyzed reactive oxygen species (ROS) formation elicits conversion of the latent form of TGF- β to its active form, which is one of the most important mechanisms that regulate activation of the TGF- β signaling [31]. ROS may also regulate TGF- β signaling indirectly via certain oxidative intermediate products. In cultured macrophages, treatment with 4-hydroxy nonenal (4-HNE), a major reactive aldehyde formed during lipid peroxidation, upregulated TGF- β expression and release [32], although it was not clear whether the same effect was also present in fibroblasts. Moreover, ROS may have modulating effects on TGF- β -induced Smad2/3 activation. In cardiac fibroblasts, it was observed that TGF- β -induced Smad2/3 phosphorylation was significantly suppressed by various antioxidant agents and by Nox4 gene silencing [18]. In a model of bleomycin-induced pulmonary fibrosis, the level of Smad2 phosphorylation was significantly decreased in Nox4 knockout mice [27]. In a separate study in kidney proximal tubular epithelial cells, it was reported that antioxidant treatment effectively inhibited TGF- β -induced phosphorylation of Smad2, although exogenous H_2O_2 failed to modulate Smad2 phosphorylation [24]. Nox4 modulation of Smad2/3 phosphorylation was also observed in cultured pulmonary fibroblasts [25]. These findings together indicate that NADPH oxidase and ROS molecules may have critical roles in facilitating TGF- β -induced Smad2/3 activation.

The mechanism of the enhancing effect of ROS on TGF- β -induced Smad phosphorylation is currently unclear. However, several possibilities that may explain this phenomenon are proposed. Activation of TGF- β signaling requires phosphorylation of the type I receptor (ALK5) on serine and threonine residues in the





Fig. 3. Potential redox-dependent mechanisms that may be involved in modulating TGF- β -induced intracellular signaling. ROS may facilitate TGF- β signaling by (1) promoting conversion of latent TGF- β to its active form; (2) enhancing phosphorylation of the TGF- β receptor I (ALK5); (3) enhancing phosphorylation of Smad2/3; (4) enhancing JNK and p38 activation, leading to increased transcriptional activities of Smad proteins. PP1, protein phosphatase 1.



B

mouse	---CACAACAACAGGCTCGCGC	AGACAAAGGGGCTGGCGCAG	-----	-221
rat	----CGGC---CACAACAACAGGCTCGCGC	AGACAAAGGGGCTGGCGCGG	-----	-221
human	ACAGCAGCAGCCACAACAACAGGCTCGCCCC	TAGACAAAGGGGCCGGCGCGGCGGAGCAG		-298
	* * *	*****	*****	*

Fig. 4. Potential mechanisms of TGF- β -induced upregulation of Nox4 gene expression. (A) TGF- β may stimulate Nox4 expression via the classical Smad2/3 pathway and activation of the PI3K pathway. (B) A putative Smad binding motif (boxed) is present in the Nox4 gene promoter region, which is conserved in human, mouse and rat.

GS domain upon ligand binding, while phosphorylated ALK5 in turn induces phosphorylation of Smad2/3 in their C-terminal SXS motif and thereby initiates downstream signaling events [33]. ALK5 phosphorylation is negatively regulated (dephosphorylated) by protein phosphatase 1 (PP1) and PP2A [33]. There is evidence showing that PP1 and PP2A are redox-sensitive, with ROS molecules such as H₂O₂ being able to inactivate the activity of these enzymes [34], leading to enhanced ALK5 activation and Smad phosphorylation. Moreover, people have demonstrated that the phosphatase PTEN may induce a decrease in Smad2/3 phosphorylation, although this effect of PTEN appears to require another phosphatase PPM1A [33]. It is well documented that PTEN is sensitive to ROS-induced oxidation and inactivation [35]. In this scenario, ROS may facilitate Smad phosphorylation by inhibiting PTEN/PPM1A-mediated dephosphorylation. Nevertheless, the importance of these potential mechanisms in modulating TGF- β signaling in vivo is currently not understood, which needs to be further corroborated by direct experimental evidence.

In addition to regulating Smad phosphorylation, ROS may also modulate TGF- β signaling via non-canonical, Smad-independent mechanisms. In addition to Smad-transduced signals, TGF- β may activate other signaling pathways including the mitogen-activated protein kinase (MAPK) members c-Jun N-terminal kinase (JNK) and p38 [36]. Of note, it is clear that both JNK and p38 are redox sensitive, which can be activated by ROS in the cytoplasm [35]. Interestingly, experimental evidence has indicated that JNK and p38 may in turn enhance the transcriptional activities of Smad proteins by direct phosphorylation of Smad3 or indirectly by promoting Smad3 association with the transcriptional co-activator p300 [36]. Hence, ROS may facilitate TGF- β -induced signaling by enhancing JNK and p38 activation. These redox mechanisms possibly involved in modulating TGF- β functions are summarized in Fig. 3.

Mechanisms of TGF- β regulation of Nox4 expression

TGF- β upregulates Nox4 expression in a variety of cells [17,18,20,37–40]. Two major mechanisms potentially involved in TGF- β -induced Nox4 expression are the classical Smad2/3 pathway and the phosphoinositide 3-kinase (PI3K) pathway (Fig. 4). We and others have demonstrated that TGF- β -induced Nox4 upregulation and ROS generation can be inhibited by inhibitors of ALK5 and Smad3, or dormant negative mutants of Smad2/3 [17,37,40,41], suggesting that Nox4 expression occurs downstream of Smad2/3 activation. Similarly, overexpression of the inhibitory Smad protein Smad7 reversed the stimulating effects of TGF- β on Nox4 expression [41]. In addition to the Smad pathway, there is evidence suggesting that activation of PI3K may also be involved in TGF- β -induced regulation of Nox4 expression. For example, using a pharmacological inhibitor of PI3K, Michaeloudes et al. showed that in airway smooth muscle cells, PI3K inhibition abrogated Nox4 expression in response to TGF- β treatment [41]. Moreover, Sturrock et al. also demonstrated that TGF- β -stimulated Nox4 expression in airway smooth muscle cells was sensitive to PI3K inhibition [42]. However, the role of PI3K in TGF- β -induced modulation of Nox4 expression in fibroblasts is currently unclear, which needs to be further clarified.

Acknowledgements

This work was supported by the National 973 Basic Research Program (2010CB732605), Natural Science Foundation of China (81271269), Ophthalmic Research Institute of Australia (ORIA), and the Early Career Researcher grant (from University of

Melbourne). G.J.D receives a Principal Research Fellowship from NHMRC. G.S.L. and E.C.C. are supported by The Ansell Ophthalmology Foundation. The Centre for Eye Research Australia receives Operational Infrastructure Support from the Victorian Government.

References

- [1] A.K. Ghosh, S.E. Quaggin, D.E. Vaughan, Molecular basis of organ fibrosis: potential therapeutic approaches, *Exp. Biol. Med.* (Maywood) 238 (2013) 461–481.
- [2] T.A. Mustoe, G.F. Pierce, A. Thomason, P. Gramates, M.B. Sporn, et al., Accelerated healing of incisional wounds in rats induced by transforming growth factor-beta, *Science* 237 (1987) 1333–1336.
- [3] A. Leask, D.J. Abraham, TGF-beta signaling and the fibrotic response, *FASEB J.* 18 (2004) 816–827.
- [4] M.H. Branton, J.B. Kopp, TGF-beta and fibrosis, *Microb. Infect.* 1 (1999) 1349–1365.
- [5] A. Meyer, W. Wang, J. Qu, L. Croft, J.L. Degen, et al., Platelet TGF-beta1 contributions to plasma TGF-beta1, cardiac fibrosis, and systolic dysfunction in a mouse model of pressure overload, *Blood*. 119 (2012) 1064–1074.
- [6] B.D. Uhal, J.K. Kim, X. Li, M. Molina-Molina, Angiotensin-TGF-beta 1 crosstalk in human idiopathic pulmonary fibrosis: autocrine mechanisms in myofibroblasts and macrophages, *Curr. Pharm. Des.* 13 (2007) 1247–1256.
- [7] S.H. Phan, Biology of fibroblasts and myofibroblasts, *Proc. Am. Thorac. Soc.* 5 (2008) 334–337.
- [8] R. Kalluri, E.G. Neilson, Epithelial-mesenchymal transition and its implications for fibrosis, *J. Clin. Invest.* 112 (2003) 1776–1784.
- [9] M. Iwano, D. Plieth, T.M. Danoff, C. Xue, H. Okada, et al., Evidence that fibroblasts derive from epithelium during tissue fibrosis, *J. Clin. Invest.* 110 (2002) 341–350.
- [10] J.C. Kovacic, N. Mercader, M. Torres, M. Boehm, V. Fuster, Epithelial-to-mesenchymal and endothelial-to-mesenchymal transition: from cardiovascular development to disease, *Circulation* 125 (2012) 1795–1808.
- [11] E.M. Zeisberg, O. Tarnavski, M. Zeisberg, A.L. Dorfman, J.R. McMullen, et al., Endothelial-to-mesenchymal transition contributes to cardiac fibrosis, *Nat. Med.* 13 (2007) 952–961.
- [12] T.A. Wynn, Common and unique mechanisms regulate fibrosis in various fibroproliferative diseases, *J. Clin. Invest.* 117 (2007) 524–529.
- [13] E. Pardali, M.J. Goumans, P. ten Dijke, Signaling by members of the TGF-beta family in vascular morphogenesis and disease, *Trends Cell Biol.* 20 (2010) 556–567.
- [14] M. Tian, J.R. Neil, W.P. Schiemann, Transforming growth factor-beta and the hallmarks of cancer, *Cell Signalling* 23 (2011) 951–962.
- [15] R. Samarakoon, J.M. Overstreet, P.J. Higgins, TGF-beta signaling in tissue fibrosis: redox controls, target genes and therapeutic opportunities, *Cell Signalling* 25 (2013) 264–268.
- [16] E.C. Chan, F. Jiang, H.M. Peshavariya, G.J. Dusting, Regulation of cell proliferation by NADPH oxidase-mediated signaling: potential roles in tissue repair, regenerative medicine and tissue engineering, *Pharmacol. Ther.* 122 (2009) 97–108.
- [17] E.C. Chan, H.M. Peshavariya, G.S. Liu, F. Jiang, S.Y. Lim, et al., Nox4 modulates collagen production stimulated by transforming growth factor beta1 in vivo and in vitro, *Biochem. Biophys. Res. Commun.* 430 (2013) 918–925.
- [18] I. Cucoranu, R. Clempus, A. Dikalova, P.J. Phelan, S. Ariyan, et al., NAD(P)H oxidase 4 mediates transforming growth factor-beta1-induced differentiation of cardiac fibroblasts into myofibroblasts, *Circ. Res.* 97 (2005) 900–907.
- [19] L. Hecker, R. Vittal, T. Jones, R. Jagirdar, T.R. Luckhardt, et al., NADPH oxidase-4 mediates myofibroblast activation and fibrogenic responses to lung injury, *Nat. Med.* 15 (2009) 1077–1081.
- [20] C.D. Bondi, N. Manickam, D.Y. Lee, K. Block, Y. Gorin, et al., NAD(P)H oxidase mediates TGF-beta1-induced activation of kidney myofibroblasts, *J. Am. Soc. Nephrol.* 21 (2010) 93–102.
- [21] N. Sampson, P. Berger, C. Zenzmaier, Therapeutic targeting of redox signaling in myofibroblast differentiation and age-related fibrotic disease, *Oxid. Med. Cell. Longev.* 2012 (2012) 458276.
- [22] J.L. Barnes, Y. Gorin, Myofibroblast differentiation during fibrosis: role of NAD(P)H oxidases, *Kidney Int.* 79 (2011) 944–956.
- [23] D.Y. Rhyu, J. Park, B.R. Sharma, H. Ha, Role of reactive oxygen species in transforming growth factor-beta1-induced extracellular matrix accumulation in renal tubular epithelial cells, *Transplant. Proc.* 44 (2012) 625–628.
- [24] D.Y. Rhyu, Y. Yang, H. Ha, G.T. Lee, J.S. Song, et al., Role of reactive oxygen species in TGF-beta1-induced mitogen-activated protein kinase activation and epithelial-mesenchymal transition in renal tubular epithelial cells, *J. Am. Soc. Nephrol.* 16 (2005) 667–675.
- [25] N. Amara, D. Goven, F. Prost, R. Muloway, B. Crestani, et al., NOX4/NADPH oxidase expression is increased in pulmonary fibroblasts from patients with idiopathic pulmonary fibrosis and mediates TGFbeta1-induced fibroblast differentiation into myofibroblasts, *Thorax* 65 (2010) 733–738.
- [26] P. Sancho, J. Mainez, E. Crosas-Molist, C. Roncero, C.M. Fernandez-Rodriguez, et al., NADPH oxidase NOX4 mediates stellate cell activation and hepatocyte cell death during liver fibrosis development, *PLoS One* 7 (2012) e45285.
- [27] S. Carnesecci, C. Deffert, Y. Donati, O. Basset, B. Hinz, et al., A key role for NOX4 in epithelial cell death during development of lung fibrosis, *Antioxid Redox Signalling* 15 (2011) 607–619.

- [28] B. Laleu, F. Gaggini, M. Orchard, L. Fioraso-Cartier, L. Cagnon, et al., First in class, potent, and orally bioavailable NADPH oxidase isoform 4 (Nox4) inhibitors for the treatment of idiopathic pulmonary fibrosis, *J. Med. Chem.* 53 (2010) 7715–7730.
- [29] J.X. Jiang, X. Chen, N. Serizawa, C. Szyndralewicz, P. Page, et al., Liver fibrosis and hepatocyte apoptosis are attenuated by GKT137831, a novel NOX4/NOX1 inhibitor in vivo, *Free Radic. Biol. Med.* 53 (2012) 289–296.
- [30] E.R. Jarman, V.S. Khambata, C. Cope, P. Jones, J. Roger, et al., An inhibitor of NADPH oxidase-4 attenuates established pulmonary fibrosis in a rodent disease model, *Am. J. Respir. Cell Mol. Biol.* (2013).
- [31] M.H. Barcellos-Hoff, T.A. Dix, Redox-mediated activation of latent transforming growth factor-beta 1, *Mol. Endocrinol.* 10 (1996) 1077–1083.
- [32] G. Leonarduzzi, A. Scavazza, F. Biasi, E. Chiarotto, S. Camandola, et al., The lipid peroxidation end product 4-hydroxy-2,3-nonenal up-regulates transforming growth factor beta1 expression in the macrophage lineage: a link between oxidative injury and fibrosclerosis, *FASEB J.* 11 (1997) 851–857.
- [33] K.H. Wrighton, X. Lin, X.H. Feng, Phospho-control of TGF-beta superfamily signaling, *Cell. Res.* 19 (2009) 8–20.
- [34] H.S. Kim, M.C. Song, I.H. Kwak, T.J. Park, I.K. Lim, Constitutive induction of p-Erk1/2 accompanied by reduced activities of protein phosphatases 1 and 2A and MKP3 due to reactive oxygen species during cellular senescence, *J. Biol. Chem.* 278 (2003) 37497–37510.
- [35] F. Jiang, Y. Zhang, G.J. Dusting, NADPH oxidase-mediated redox signaling: roles in cellular stress response, stress tolerance, and tissue repair, *Pharmacol. Rev.* 63 (2011) 218–242.
- [36] A. Moustakas, C.H. Heldin, Non-Smad TGF-beta signals, *J. Cell Sci.* 118 (2005) 3573–3584.
- [37] A. Sturrock, B. Cahill, K. Norman, T.P. Huecksteadt, K. Hill, et al., Transforming growth factor-beta1 induces Nox4 NAD(P)H oxidase and reactive oxygen species-dependent proliferation in human pulmonary artery smooth muscle cells, *Am. J. Physiol. Lung Cell. Mol. Physiol.* 290 (2006) L661–L673.
- [38] I. Carmona-Cuenca, C. Roncero, P. Sancho, L. Caja, N. Fausto, et al., Upregulation of the NADPH oxidase NOX4 by TGF-beta in hepatocytes is required for its proapoptotic activity, *J. Hepatol.* 49 (2008) 965–976.
- [39] A. Martin-Garrido, D.I. Brown, A.N. Lyle, A. Dikalova, B. Seidel-Rogol, et al., NADPH oxidase 4 mediates TGF-beta-induced smooth muscle alpha-actin via p38MAPK and serum response factor, *Free Radic. Biol. Med.* 50 (2011) 354–362.
- [40] H.E. Boudreau, B.W. Casterline, B. Rada, A. Korzeniowska, T.L. Leto, Nox4 involvement in TGF-beta and SMAD3-driven induction of the epithelial-to-mesenchymal transition and migration of breast epithelial cells, *Free Radic. Biol. Med.* 53 (2012) 1489–1499.
- [41] C. Michaeloudes, M.B. Sukkar, N.M. Khorasani, P.K. Bhavsar, K.F. Chung, TGF-beta regulates Nox4, MnSOD and catalase expression, and IL-6 release in airway smooth muscle cells, *Am. J. Physiol. Lung Cell. Mol. Physiol.* 300 (2011) L295–L304.
- [42] A. Sturrock, T.P. Huecksteadt, K. Norman, K. Sanders, T.M. Murphy, et al., Nox4 mediates TGF-beta1-induced retinoblastoma protein phosphorylation, proliferation, and hypertrophy in human airway smooth muscle cells, *Am. J. Physiol. Lung Cell. Mol. Physiol.* 292 (2007) L1543–L1555.