



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Li, C;Hu, HW;Chen, QL;Chen, D;He, JZ

Title:

Growth of comammox Nitrospira is inhibited by nitrification inhibitors in agricultural soils

Date:

2020-02-01

Citation:

Li, C., Hu, H. W., Chen, Q. L., Chen, D. & He, J. Z. (2020). Growth of comammox Nitrospira is inhibited by nitrification inhibitors in agricultural soils. *Journal of Soils and Sediments*, 20 (2), pp.621-628. <https://doi.org/10.1007/s11368-019-02442-z>.

Persistent Link:

<https://hdl.handle.net/11343/282514>

[Click here to view linked References](#)

SOILS, SEC 1 • SOIL ORGANIC MATTER DYNAMICS AND NUTRIENT CYCLING •  
RESEARCH ARTICLE

**Growth of comammox *Nitrospira* is inhibited by nitrification inhibitors in agricultural  
soils**

**Chaoyu Li<sup>1</sup> • Hang-Wei Hu<sup>1</sup> • Qing-Lin Chen<sup>1</sup> • Deli Chen<sup>1</sup> • Ji-Zheng He<sup>1</sup>**

Received: 24 June 2019 / Accepted: 24 August 2019

© Springer-Verlag GmbH Germany, part of Springer Nature 2019

---

Responsible editor: Huaiying Yao

---

<sup>1</sup>School of Agriculture and Food, Faculty of Veterinary and Agricultural Sciences, The  
University of Melbourne, Parkville, Victoria 3010, Australia

✉ Ji-Zheng He

jizheng.he@unimelb.edu.au

## Abstract

**Purpose** The discovery of comammox *Nitrospira* being capable of complete oxidizing ammonia to nitrate radically challenged the conventional concept of two-step nitrification. However, the response of comammox *Nitrospira* to nitrification inhibitors (NIs) and their role in soil nitrification remain largely unknown, which has hindered our ability to predict the efficiency of NIs in agroecosystems.

**Materials and methods** We evaluated the effect of four NIs, 2-chloro-6-(trichloromethyl)pyridine (nitrapyrin), 3,4-dimethylpyrazole phosphate (DMPP), allylthiourea (ATU) and dicyandiamide (DCD) on the growth of comammox *Nitrospira*, ammonia-oxidizing archaea (AOA) and ammonia-oxidizing bacteria (AOB) in two pasture and arable soils.

**Results and discussion** The amendment of nitrogen fertiliser significantly increased soil nitrate concentrations over time, indicating a sustaining nitrification activity in both soils. The addition of all the four NIs effectively reduced the production of nitrate in both soils, but to varying degrees during incubation. The abundances of comammox *Nitrospira* clade A were significantly increased by addition of nitrogen fertilisers and significantly impeded by the four NIs in the pasture soil, but their abundances were only remarkably hindered by nitrapyrin in the arable soil. All the four NIs obviously inhibited the AOB abundances in both soils. Except for DMPP, the other three NIs effectively suppressed the AOA abundances in both soils.

**Conclusions** We provided new evidence that growth of comammox *Nitrospira* clade A can be stimulated by nitrogen fertilisers and inhibited by various nitrification inhibitors, suggesting their potential role in nitrification of agricultural soils.

**Keywords** Agricultural soil • Comammox *Nitrospira* • Nitrification inhibitor • Nitrogen use efficiency

## 1 Introduction

1  
2 The intensive use of nitrogen (N) fertilizers in agro-ecosystems is driven by a growing  
3 demand of food for a global human population of more than 11 billion by 2100 (Gerland et al.  
4  
5 2014; Gruber and Galloway 2008). However, nitrogen use efficiency (NUE) is rarely  
6  
7 exceeding 50%, as majority of N fertilizers is lost to the environment through nitrate leaching,  
8  
9 nitrous oxide (N<sub>2</sub>O) emission, and ammonia volatilization (Wang et al. 2018a; Zhang et al.  
10  
11 2019). Application of nitrification inhibitors (NIs) together with N-based fertilizers is a  
12  
13 practical tool to reduce N losses and to promote plant NUE in order to increase crop yields in  
14  
15 agricultural systems ( Dinnes et al. 2002; Subbarao et al. 2006; Di and Cameron 2012; Shi et  
16  
17 al. 2016b). Dicyandiamide (DCD), 3,4-dimethylpyrazole phosphate (DMPP) and 2-chloro-6-  
18  
19 (trichloromethyl) pyridine (nitrapyrin, NP) are commercial NIs widely applied in agricultural  
20  
21 soils (Shi et al. 2016a; Subbarao et al. 2006) and allylthiourea (ATU) is frequently used in  
22  
23 laboratory experiments to differentiate the relative contribution of ammonia oxidizers to  
24  
25 nitrification (Lehtovirta-Morley et al. 2013; Shen et al. 2013; Wang and Gu 2014). The  
26  
27 underlying mechanisms for NIs are generally attributed to their deactivation of ammonia  
28  
29 monooxygenase (AMO), which catalyses the ammonia oxidation process, conversion from  
30  
31 ammonia to nitrite, mediated by ammonia-oxidizing archaea (AOA) and ammonia oxidizing  
32  
33 bacteria (AOB) (Subbarao et al. 2009; Shen et al. 2013). However, the outcomes of these NIs  
34  
35 are reported to be variable across soils (Shi et al. 2016a, b), primarily owing to their largely  
36  
37 unknown effects on nitrogen-cycling microorganisms (Di et al. 2009; Shi et al. 2017).  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 Nitrification was traditionally considered as a two-step process including ammonia  
49  
50 oxidation catalysed by AOA and AOB, and nitrite oxidation catalysed by nitrite-oxidizing  
51  
52 bacteria (NOB). This long-held perspective was radically challenged by the recent discovery  
53  
54 of complete ammonia oxidizers, namely comammox organisms within the genus *Nitrospira*,  
55  
56 which are capable of oxidizing ammonia to nitrate in a single organism (Daims et al. 2015; van  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 Kessel et al. 2015). The assembled complete genomes of comammox *Nitrospira* harbour the  
2 full set of genes encoding AMO and hydroxylamine dehydrogenase for ammonia oxidation,  
3  
4 and genes encoding nitrite oxidoreductase for nitrite oxidation (Daims et al. 2015; van Kessel  
5  
6 et al. 2015). Since the discovery of comammox *Nitrospira* as a third group of ammonia  
7  
8 oxidizers, numerous studies have been carried out to investigate their environmental prevalence  
9  
10 in various environmental settings, including agricultural soils, forest soils, paddy soils,  
11  
12 freshwater sediments, and wastewater treatment plants (Bartelme et al. 2017; Hu and He 2017;  
13  
14 Pjevac et al. 2017; Fowler et al. 2018; Wang et al. 2018b; Yu et al. 2018; Beach and Noguera  
15  
16 2019). It was found that comammox *Nitrospira* are widely prevalent in terrestrial ecosystems  
17  
18 and can be more abundant than canonical ammonia oxidizers in some soils (Hu and He 2017;  
19  
20 Pjevac et al. 2017), but their role in soil nitrification and responses to NIs remains unknown.  
21  
22  
23  
24  
25

26  
27 Given the fact that genomes of comammox *Nitrospira* contain the *amoA* gene encoding  
28  
29 the enzyme AMO, which is the target of NIs, we hypothesized that growth of comammox  
30  
31 organisms can be inhibited by the currently available NIs, but to varying degrees. The study of  
32  
33 van Kessel et al. (2015) reported that the addition of ATU significantly inhibited ammonia-  
34  
35 dependent carbon fixation of comammox *Nitrospira* in an incubation experiment. As far as we  
36  
37 know, no studies have compared the responses of comammox *Nitrospira*, AOA and AOB to  
38  
39 NIs in agroecosystems. The objective of this study was therefore to assess the impacts of four  
40  
41 NIs on nitrification and the abundance of comammox *Nitrospira*, AOA and AOB in two  
42  
43 agricultural soils during a 28-day microcosm incubation. The results of this study can improve  
44  
45 our understanding of the possible mechanisms governing the efficacy of NIs on nitrification by  
46  
47 including comammox *Nitrospira* as a third group of nitrifiers.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 2 Materials and methods

### 2.1 Site description and soil sampling

Soil samples were collected during May 2018 (autumn in Australia) from two long-term research farms: a dairy farm site (pasture soil) at Dookie (36°33'S, 145°69'E) and a vegetable farm site (arable soil) at Clyde (38°13'S, 145°33'E), Victoria, Australia. The mean annual rainfall at Dookie is 554 mm, and the mean annual maximum and minimum temperatures are 21 °C and 9 °C, respectively. The Dookie site has been used as a dairy farm for more than 30 years, including around 43 hectares of irrigated pastures with a feeding capacity for 180 cows. The pastures at the Dookie site are a mixture of perennial ryegrass (*Lolium perenne*) and white clover (*Trifolium repens*). The mean annual rainfall at Clyde is 816 mm and the mean annual maximum and minimum temperatures are 20 °C and 10 °C, respectively. Approximately 5 kg soils were collected from the upper 10 cm of soils from each site, placed into plastic buckets, and shipped on ice to the laboratory. The soil samples were sieved through a 2 mm sieve to remove roots and stones, mixed thoroughly and stored at 4 °C prior to construction of soil microcosms.

### 2.2 Soil physicochemical analysis

Soil physicochemical properties were analysed through the Melbourne Trace Analysis for Chemical, Earth and Environmental Sciences platform. Soil moisture content was measured by oven drying  $10 \pm 0.05$  g of soil at 105 °C for 24 h. Soil pH and electrical conductivity were estimated with a soil to water ratio of 1: 5 using an Orion Star A211 pH meter (Thermo Scientific Inc., Melbourne, Australia). Total C and N were determined via the Dumas combustion method on an isotope ratio mass spectrometer (Sercon Hydra, Crewe, UK). Soil  $\text{NH}_4^+$ -N and  $\text{NO}_3^-$ -N were extracted by 1 M potassium chloride (KCl) solution with a soil to solution ratio of 1:5, filtered through a Whatman Grade 42 filter paper (Sigma-Aldrich Co., St.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Louis, MO, USA), and analysed with a segmented flow analyzer (SAN++; Skalar, Breda, Holland). Available phosphorus (P) were measured via the Colwell P test on a segmented flow analyzer (SAN++; Skalar, Breda, Holland). Available sulphur (S) and potassium (K) were extracted with a soil to Mehlich 3 ratio of 1:10 and estimated by Optima 8300 ICP-OES spectrometer (PerkinElmer Inc., Waltham, WA, USA). More detailed information about the soil properties is shown in Table 1.

### 2.3 Soil microcosm incubation

The soil incubation was performed in 250-ml plastic vials containing 20 g soil samples (oven dry weight equivalent). Six treatments were established with three replicates: (i) control; (ii)  $(\text{NH}_4)_2\text{SO}_4$  (100 mg  $\text{NH}_4^+\text{-N kg}^{-1}$  soil); (iii)  $(\text{NH}_4)_2\text{SO}_4$  plus nitrapyrin (50  $\mu\text{mol kg}^{-1}$  soil); (iv)  $(\text{NH}_4)_2\text{SO}_4$  plus DMPP (5  $\mu\text{mol kg}^{-1}$  soil); (v)  $(\text{NH}_4)_2\text{SO}_4$  plus ATU (1000  $\mu\text{mol kg}^{-1}$  soil); and (vi)  $(\text{NH}_4)_2\text{SO}_4$  plus DCD (10  $\mu\text{mol kg}^{-1}$  soil). The selected NIs, including DCD, DMPP, nitrapyrin (NP) and ATU, have been widely used in agricultural soils or laboratory experiments to inhibit the conversion of ammonium to nitrate, and the concentrations of NIs used were based on the findings from previous studies (Lehtovirta-Morley et al. 2013; Shen et al. 2013; Wang and Gu 2014; Shi et al. 2016a). All microcosms were incubated in a temperature-controlled incubator at 25 °C in the dark for 28 days. Soil moisture content was maintained at 55% water filled pore space (WFPS) throughout the incubation by weekly adding water to the incubation vials based on the weight changes. Aerobic conditions were maintained twice a week by opening the vials and replenishing. Destructive samples were collected for molecular and mineral N analysis on days 0, 3, 7, 14, 21 and 28.

## 2.4 DNA extraction and quantitative PCR assay

1  
2 Nucleic acids were extracted from 0.25 g of soil using MoBio PowerSoil DNA isolation  
3 kits (MoBio Laboratories, Carlsbad, CA, USA) according to the manufacturer's instructions,  
4 with slight modifications as previously described (Hu et al. 2015). The concentrations of  
5 extracted DNA were determined using the NanoDrop ND2000c spectrophotometer (NanoDrop  
6 Technologies).

7  
8  
9  
10  
11  
12  
13 All quantitative PCR (qPCR) assays were performed on a Bio-Rad CFX384 optical  
14 real-time PCR detection system (Bio-Rad, Laboratories Inc., Hercules, CA, USA). The primer  
15 sets *comaA*-244F/*comaA*-659R, *comaB*-244F/*comaB*-659R (Pjevac et al. 2017),  
16  
17 *CrenamoA23f*/*CrenamoA616r* (Tourna et al. 2008) and *amoA1F*/*amoA2R* (Rotthauwe et al.  
18 1997) were used to qualify the *amoA* gene abundances of comammox clade A, clade B, AOA  
19 and AOB, respectively (Table S1, Electronic Supplementary Material - ESM). Standard curves  
20 were generated using 10-fold serial dilutions of plasmids containing correct inserts of the target  
21 genes. Melting curve analysis was performed at the end of each qPCR run to check the  
22 specificity of amplification products, before identification by agarose gel electrophoresis.  
23  
24 Based on the agarose gel results, the qPCR products of comammox clade A yielded single and  
25 bright bands with the expected 415 bp target size in both soils. The resultant PCR products  
26 were cloned into the pGEM<sup>®</sup>-T Easy vector and transformed into *Escherichia coli* JM109  
27 competent cells based on the manufacturer's instructions. Positive clones were sent to  
28 Macrogen Sequencing Department, South Korea to analyse the sequences containing the  
29 correct insert. The obtained sequences were checked for chimeras and aligned with reference  
30 sequences using MUSCLE in MEGA 7.0 (Kumar et al. 2016). The results showed that all  
31 obtained sequences (based on 54 positive clones) have high similarity (> 95%) with reference  
32 sequences of comammox *Nitrospira*. The sequence for comammox *Nitrospira* clade A has been  
33 deposited into GenBank with the accession number MK581036. However, we did not obtain  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 any specific amplification for comammox *Nitrospira* clade B in either pasture or arable soil,  
2 and therefore comammox *Nitrospira* clade B were excluded from the downstream analysis.  
3  
4 The amplification efficiencies of all *amoA* gene were 85%-99% and the  $r^2$  values were  $> 0.99$ .  
5  
6  
7  
8  
9

## 10 **2.5 Statistical analysis**

11  
12 Two-way analysis of variance (ANOVA) based on the Duncan test was conducted to  
13 analyse the differences in ammonium and nitrate concentrations, the abundance of comammox  
14 *Nitrospira* clade A, AOA and AOB across different treatments and sampling times with SPSS  
15 Statistics 25 (IBM, USA). Differences at a P value of  $< 0.05$  were statistically significant.  
16  
17  
18  
19  
20  
21  
22  
23

## 24 **3 Results**

### 25 **3.1 Changes in soil mineral N in microcosm incubation**

26  
27 The  $\text{NH}_4^+$ -N concentrations remained largely unchanged during incubation in the  
28 control treatments of both soils (Fig. 1A and 1B). The  $\text{NH}_4^+$ -N concentrations in the  $(\text{NH}_4)_2\text{SO}_4$   
29 treatment rapidly reduced from  $55.43 (\pm 4.71)$  to  $0.65 (\pm 0.10)$   $\text{mg kg}^{-1}$  and from  $78.83 (\pm 2.68)$   
30 to  $0.88 (\pm 0.04)$   $\text{mg kg}^{-1}$  soil in the control treatment from day 0 to day 3 in the pasture and  
31 arable soils, respectively (Fig. 1A and 1B). The addition of nitrapyrin and DMPP significantly  
32 ( $P < 0.001$ ) increased the  $\text{NH}_4^+$ -N concentrations in the pasture soil and slowed down the  
33 reduction of the  $\text{NH}_4^+$ -N concentrations in the arable soil. No significant differences ( $P > 0.05$ )  
34 in the temporal changes of the  $\text{NH}_4^+$ -N concentrations in the  $(\text{NH}_4)_2\text{SO}_4$ ,  $(\text{NH}_4)_2\text{SO}_4 + \text{ATU}$   
35 and  $(\text{NH}_4)_2\text{SO}_4 + \text{DCD}$  was observed (Table 2).  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

51 In the control microcosms, the  $\text{NO}_3^-$ -N concentrations continuously increased over time,  
52 indicating a sustaining nitrification activity in both soils (Fig. 1C and 1D). In the  $(\text{NH}_4)_2\text{SO}_4$   
53 treatment, the  $\text{NO}_3^-$ -N concentrations sharply increased and had significantly ( $P < 0.001$ )  
54 higher net nitrification rates of  $\sim 9.36$  and  $\sim 4.64$   $\text{mg NO}_3^-$ -N  $\text{kg}^{-1}$  soil per day than the control  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 treatments of ~5.86 and ~ 1.44 mg NO<sub>3</sub><sup>-</sup>-N kg<sup>-1</sup> soil per day in the pasture and arable soils,  
2 respectively. The addition of all the four NIs effectively reduced the production of NO<sub>3</sub><sup>-</sup>-N in  
3 both soils, but to varying degrees during incubation. The addition of nitrapyrin showed the  
4 most significant effect on the inhibition of nitrification in both soils. Net nitrification rates in  
5 the other three treatments with NIs also had significantly ( $P < 0.001$ ) lower values than that in  
6 the (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> treatment (Table 2).  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16

### 17 **3.2 Changes of the comammox *Nitrospira amoA* gene abundance**

18  
19 In the control treatments, the comammox *Nitrospira* clade A *amoA* gene abundance  
20 ranged between 4.32 to 9.26 × 10<sup>7</sup> copies g<sup>-1</sup> soil in the pasture soil (Fig. 2A), and 1.77 to 3.28  
21 × 10<sup>7</sup> copies g<sup>-1</sup> soil in the arable soil (Fig. 2B). Addition of (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> significantly ( $P < 0.05$ )  
22 increased the comammox *Nitrospira* clade A *amoA* gene abundance from 6.08 × 10<sup>7</sup> at day 7  
23 to 5.50 × 10<sup>7</sup> at day 28 in the pasture soil (Fig. 2A), and from 2.37 × 10<sup>7</sup> at day 7 to 3.06 × 10<sup>7</sup>  
24 at day 28 in the arable soil (Fig. 2B), compared to the control treatment (Table 2).  
25  
26  
27  
28  
29  
30  
31  
32  
33

34 The comammox *Nitrospira* clade A abundance showed different responses to the four  
35 NIs in both soils. Compared with the (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> treatment, the comammox *Nitrospira* clade A  
36 abundance in the (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> + nitrapyrin treatment of the pasture and arable soils significantly  
37 ( $P < 0.05$ ) declined from 5.59 to 3.91 × 10<sup>7</sup> copies g<sup>-1</sup> soil and from 1.56 to 1.71 × 10<sup>7</sup> copies  
38 g<sup>-1</sup> soil, respectively (Fig. 2A and 2B). The other three NIs showed different effects on the  
39 comammox *Nitrospira* clade A in the pasture and arable soils. In the pasture soil, comammox  
40 *Nitrospira* clade A abundance was significantly ( $P < 0.05$ ) reduced in the presence of ATU and  
41 DCD, and significantly ( $P < 0.05$ ) decreased from day 14 to day 28 after the addition of DMPP  
42 (Fig. 2A). In the arable soil, no obvious inhibitory effect of DCD on the growth of comammox  
43 *Nitrospira* clade A was observed, and comammox *Nitrospira* clade A abundance even tended  
44 to increase in the DMPP and ATU treatments (Fig. 2B).  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

### 3.3 Changes of the AOA and AOB *amoA* gene abundance

The AOA *amoA* gene abundance was significantly increased ( $P < 0.05$ ) after adding  $(\text{NH}_4)_2\text{SO}_4$  in both soils (Fig. 3A and 3B). Compared with the  $(\text{NH}_4)_2\text{SO}_4$  treatment, only DMPP had no significant ( $P > 0.05$ ) inhibitory effect on the AOA abundance in both soils (Table 2). The addition of nitrapyrin significantly ( $P < 0.05$ ) impeded the growth of AOA from day 14 to day 28 in the pasture soil (Fig. 3A) and the AOA abundance was significantly ( $P < 0.05$ ) lower in the  $(\text{NH}_4)_2\text{SO}_4$  + nitrapyrin treatment than in the  $(\text{NH}_4)_2\text{SO}_4$  treatment during the first three weeks of incubation in the arable soil (Fig. 3B). The other two NIs (ATU and DCD) significantly ( $P < 0.05$ ) reduced the AOA abundance in both soils throughout the incubation (Table 2).

The  $(\text{NH}_4)_2\text{SO}_4$  application in both soils significantly ( $P < 0.05$ ) increased the AOB *amoA* gene abundances, compared with the control treatments, in both soils during the 28-day incubation. All the four NIs significantly ( $P < 0.05$ ) inhibited the growth of AOB in both soils but to varying degrees during the incubation (Fig. 3C and 3D, Table 2).

## 4 Discussion

This study, for the first time, demonstrated that comammox *Nitrospira* clade A can be effectively impeded by diverse NIs (nitrapyrin, DMPP, ATU and DCD) in agricultural soils. Previous studies focused on the responses of AOA and AOB to various NIs without taking comammox *Nitrospira* into consideration, because comammox *Nitrospira* were recently discovered in 2015 (Daims et al. 2015; van Kessel et al. 2015). A large body of studies suggested that most NIs are non-selective and can suppress either AOA or AOB *amoA* gene abundances depending on which nitrifying group is more functionally dominant during the nitrification process (Jia and Conrad 2009; Offre et al. 2009; Zhang et al. 2012). For example, AOB were effectively inhibited by acetylene in agricultural soils functionally dominated by

1 AOB (Jia and Conrad 2009), while DCD significantly inhibited the growth and activity of AOA  
2 in acidic soils thought to be dominated by AOA (Zhang et al. 2012). Although the responses  
3 of comammox *Nitrospira* clade A to some NIs (DMPP, ATU and DCD) in the two tested soils  
4 were not consistent due to different soil properties, all the NIs were effective in impeding the  
5 growth of comammox *Nitrospira* clade A, accompanied by significant reduction in net  
6 nitrification rates in the treatments with NIs, compared to the (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> treatment. The results  
7 indicated that comammox *Nitrospira* might be involved in the active nitrification of  
8 agricultural soils with high N input, and are sensitive to the currently available NIs, albeit the  
9 inhibitory effects on comammox *Nitrospira* are soil dependent. However, the main soil factors  
10 affecting the responses of comammox *Nitrospira* to NIs are required to be further identified by  
11 incubating multiple types of agricultural soils.

12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
Previous studies revealed that soil ammonia availability is an important factor affecting  
the abundances and activities of AOA and AOB, and further influencing their responses to NIs  
(He et al. 2012; Prosser and Nicol 2012; Hu et al. 2014). Based on the ionization equilibrium  
(NH<sub>4</sub><sup>+</sup> ↔ NH<sub>3</sub> + H<sup>+</sup>; pK<sub>a</sub> = 9.25 at 25°C) (He et al. 2012; Hu et al. 2014), we found that  
comammox *Nitrospira* clade A might be involved in nitrification in the soils with theoretical  
ammonia concentrations ranging from 159 nM to 893 nM in the pasture soil and from 408 nM  
to 781 nM in the arable soil. To date, comammox *N. inopinata*, isolated from aquatic systems,  
was the only kinetically characterized comammox species and its apparent half-saturation  
constant of K<sub>m</sub>(NH<sub>3</sub>) is around 63 nM (Kits et al. 2017), which is remarkably lower than the  
theoretical soil ammonia concentrations in this study. This finding suggests that, contrasting to  
the previous perception that comammox *Nitrospira* live an oligotrophic lifestyle (Daims et al.  
2015; van Kessel et al. 2015), comammox *Nitrospira* in the soil environment might be more  
favoured under nitrogen-rich conditions, which was also supported by recent findings that the  
abundance of comammox *Nitrospira* increased in response to added N sources in acidic forest

1 soils (Shi et al. 2018). Thus, further research regarding the  $K_m$  (NH<sub>3</sub>) of comammox *Nitrospira*  
2 is required to obtain a better understanding of the ecological niches of this new group of  
3 ammonia oxidizers in terrestrial ecosystems.  
4  
5

6  
7 Apart from nitrapyrin, the other three NIs did not inhibit the abundance of comammox  
8 *Nitrospira* clade A in the arable soil. The concentrations of some NIs which can inhibit AOA  
9 or AOB in most environments probably have little influence on comammox *Nitrospira* clade  
10 A, and the inhibitory effects are dependent on soil properties. Acid soil may be a favourable  
11 habitat for the growth of comammox *Nitrospira* clade A, which influences their responses to  
12 diverse NIs. The pH value rapidly decreased from 6.90 to 6.01 and even to 5.49 after the  
13 addition of (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> fertilizer in the pasture soil, whereas the pH values remained neutral  
14 levels between 7.13 and 6.94 in the arable soil (Fig. S1, Electronic Supplementary Material -  
15 ESM). Previous studies considered that soil pH is a main factor driving the niche differentiation  
16 between AOA and AOB (Hu et al. 2014; Li et al. 2018) and the range of soil pH directly affects  
17 the response of canonical ammonia oxidisers to inhibitors (Shi et al. 2016a, b). Therefore,  
18 comammox *Nitrospira* may be not very sensitive to some inhibitors in neutral arable soils.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

36 The NIs ATU and DCD significantly inhibited AOA and AOB abundance but cannot  
37 inhibit comammox *Nitrospira* clade A in the arable soil. The different responses between  
38 comammox *Nitrospira* and canonical nitrifiers to NIs could be ascribed to the unique genomic  
39 structures and metabolic versatility of comammox *Nitrospira*. For example, besides the inputs  
40 of ammonium from soils, comammox *Nitrospira* may use an alternative substrate to catalyse  
41 ammonia oxidation. Comammox *Nitrospira* harbour the genes encoding enzymes to generate  
42 energy from cyanate and urea degradation, indicative of their genetic potential to utilize  
43 cyanate and/or urea as the substrates for nitrification (Hu and He 2017; Koch et al. 2019).  
44 Comammox *Nitrospira* harbour enzymes that regulate the pathways involved in degradation  
45 of various carbon compounds suggestive of their potential to grow mixotrophically (Palomo et  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 al. 2018). Therefore, we assume that comammox *Nitrospira* may also use substrates other than  
2 ammonia to support the process of nitrification, which could explain why they are not sensitive  
3  
4 to NIs in the arable soil as the currently NIs are mainly targeting the AMO of ammonia  
5 oxidizers. Apart from soil pH, ammonia availability and genomic characteristics, other factors,  
6  
7 such as soil clay which can adsorb a certain amount of NI (Roco and Blu 2006), may also affect  
8  
9 the NI efficacy to hamper the abundance of comammox *Nitrospira*.  
10  
11  
12  
13  
14  
15  
16

## 17 **5 Conclusions**

18  
19 Our studies provided new evidence that growth of comammox *Nitrospira* clade A can be  
20 stimulated by nitrogen fertilisers and inhibited by various NIs, suggesting their potential role  
21 in nitrification of agricultural soils. Notably, the inhibitory efficacy of different NIs on  
22 comammox *Nitrospira* is soil dependent, which may be attributed to the different soil  
23 properties that can influence the absorption and degradation of NIs. These findings are essential  
24 to a better mechanistic understanding of the variable efficiency of NIs in different soils, and  
25 have implications for refining agricultural management practices to promote N use efficiency  
26 and reduce N losses in agriculture.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 **Acknowledgements** This work was supported by Australian Research Council (DP160101028,  
42 LP160101134). The authors would like to thank Ms Zhenzhen Yan and Dr. Bing Han for their  
43 assistance with soil sampling and Mrs Qing Xie for her help during the incubation trial.  
44  
45  
46  
47  
48  
49

## 50 **References**

51  
52  
53 Bartelme RP, McLellan SL, Newton RJ (2017) Freshwater recirculating aquaculture system  
54 operations drive biofilter bacterial community shifts around a stable nitrifying  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 consortium of ammonia-oxidizing archaea and comammox Nitrospira. Front Microbiol  
2 8:101  
3

4 Beach NK, Noguera DR (2019) Design and assessment of species-level qPCR primers  
5 targeting comammox. Front Microbiol 10:35  
6

7  
8  
9 Daims H, Lebedeva EV, Pjevac P, Han P, Herbold C, Albertsen M, Jehmlich N, Palatinszky  
10 M, Vierheilig J, Bulaev A (2015) Complete nitrification by Nitrospira bacteria. Nature  
11  
12 528:504-509  
13  
14

15  
16  
17 Di H, Cameron K (2012) How does the application of different nitrification inhibitors affect  
18 nitrous oxide emissions and nitrate leaching from cow urine in grazed pastures? Soil  
19 use. Manage 28:54-61  
20  
21

22  
23  
24 Di H, Cameron K, Shen JP, Winefield C, O'Callaghan M, Bowatte S, He J (2009) Nitrification  
25 driven by bacteria and not archaea in nitrogen-rich grassland soils. Nat Geosci 2:621-  
26  
27 624  
28  
29

30  
31 Dinnes DL, Karlen DL, Jaynes DB, Kaspar TC, Hatfield JL, Colvin TS, Cambardella CA (2002)  
32 Nitrogen management strategies to reduce nitrate leaching in tile-drained Midwestern  
33 soils. Agron J 94:153-171  
34  
35

36  
37  
38 Fowler SJ, Palomo A, Dechesne A, Mines PD, Smets BF (2018) Comammox Nitrospira are  
39 abundant ammonia oxidizers in diverse groundwater-fed rapid sand filter communities.  
40  
41 Environ Microbiol 20:1002-1015  
42  
43  
44

45  
46  
47 Gerland P, Raftery AE, Ševčíková H, Li N, Gu D, Spoorenberg T, Alkema L, Fostick BK,  
48 Chunn J, Lalic N (2014) World population stabilization unlikely this century. Science  
49  
50 346:234-237  
51  
52

53  
54  
55 Gruber N, Galloway JN (2008) An Earth-system perspective of the global nitrogen cycle.  
56  
57 Nature 451:293-296  
58  
59  
60  
61  
62  
63  
64  
65

- 1 He JZ, Hu HW, Zhang LM (2012) Current insights into the autotrophic thaumarchaeal  
2 ammonia oxidation in acidic soils. *Soil Biol Biochem* 55:146-154  
3  
4 Hu HW, He JZ (2017) Comammox—a newly discovered nitrification process in the terrestrial  
5 nitrogen cycle. *J Soils Sediments* 17:2709-2717  
6  
7 Hu HW, Macdonald CA, Trivedi P, Holmes B, Bodrossy L, He JZ, Singh BK (2015) Water  
8 addition regulates the metabolic activity of ammonia oxidizers responding to  
9 environmental perturbation in dry subhumid ecosystems. *Environ Microbiol* 17:444-461  
10  
11 Hu HW, Xu ZH, He JZ (2014) Ammonia-oxidizing archaea play a predominant role in acidic  
12 soil nitrification. *Adv Agron* 125:261-302  
13  
14 Jia Z, Conrad R (2009) Bacteria rather than Archaea dominate microbial ammonia oxidation  
15 in an agricultural soil. *Environ Microbiol* 11:1658-1671  
16  
17 Kits KD, Sedlacek CJ, Lebedeva EV et al (2017) Kinetic analysis of a complete nitrifier reveals  
18 an oligotrophic lifestyle. *Nature* 54:269-272  
19  
20 Koch H, van Kessel MA, Lüscher S (2019) Complete nitrification: insights into the  
21 ecophysiology of comammox *Nitrospira*. *Appl Microbiol Biotechnol* 103:177-189  
22  
23 Kumar S, Stecher G, Tamura K (2016) MEGA7: molecular evolutionary genetics analysis  
24 version 7.0 for bigger datasets. *Mol Biol Evol* 33:1870-1874  
25  
26 Lehtovirta-Morley LE, Verhamme DT, Nicol GW, Prosser JI (2013) Effect of nitrification  
27 inhibitors on the growth and activity of *Nitrosotalea devanattera* in culture and soil.  
28 *Soil Biol Biochem* 62:129-133  
29  
30 Li Y, Chapman SJ, Nicol GW, Yao H (2018) Nitrification and nitrifiers in acidic soils. *Soil*  
31 *Biol Biochem* 116:290-301  
32  
33 Offre P, Prosser JI, Nicol GW (2009) Growth of ammonia-oxidizing archaea in soil  
34 microcosms is inhibited by acetylene. *FEMS Microbiol Ecol* 70:99-108  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Palomo A, Pedersen AG, Fowler SJ, Dechesne A, Sicheritz-Ponten T, Smets BF (2018) Comparative genomics sheds light on niche differentiation and the evolutionary history of comammox *Nitrospira*. ISME J 12:1779-1793
- Pjevac P, Schauburger C, Poghosyan L, Herbold CW, van Kessel MA, Daebeler A, Steinberger M, Jetten MS, Lüscher S, Wagner M (2017) AmoA-targeted polymerase chain reaction primers for the specific detection and quantification of comammox *Nitrospira* in the environment. Front Microbiol 8: 1508
- Prosser JI, Nicol GW (2012) Archaeal and bacterial ammonia-oxidisers in soil: the quest for niche specialisation and differentiation. Trends Microbiol 20:523-531
- Roco MM, Blu RO (2006) Evaluation of the nitrification inhibitor 3, 4-dimethylpyrazole phosphate in two Chilean soils. J Plant Nutr 29:521-534
- Shen T, Stieglmeier M, Dai J, Urich T, Schleper C (2013) Responses of the terrestrial ammonia-oxidizing archaeon *Ca. Nitrososphaera viennensis* and the ammonia-oxidizing bacterium *Nitrospira multiformis* to nitrification inhibitors. FEMS Microbiol Lett 344:121-129
- Shi X, Hu HW, He JZ, Chen D, Suter HC (2016a) Effects of 3, 4-dimethylpyrazole phosphate (DMPP) on nitrification and the abundance and community composition of soil ammonia oxidizers in three land uses. Biol Fertil Soils 52:927-939
- Shi X, Hu HW, Müller C, He JZ, Chen D, Suter HC (2016b) Effects of the nitrification inhibitor 3, 4-dimethylpyrazole phosphate on nitrification and nitrifiers in two contrasting agricultural soils. Appl Environ Microbiol 82:5236-5248
- Shi X, Hu HW, Wang J, He JZ, Zheng C, Wan X, Huang Z (2018) Niche separation of comammox *Nitrospira* and canonical ammonia oxidizers in an acidic subtropical forest soil under long-term nitrogen deposition. Soil Biol Biochem 126:114-122

- 1 Shi X, Hu HW, Zhu-Barker X, Hayden H, Wang J, Suter H, Chen D, He JZ (2017) Nitrifier-  
2  
3 induced denitrification is an important source of soil nitrous oxide and can be inhibited  
4  
5  
6 by a nitrification inhibitor 3, 4 - dimethylpyrazole phosphate. *Environ Microbiol*  
7  
8  
9 19:4851-4865  
10
- 11 Subbarao G, Ito O, Sahrawat K, Berry W, Nakahara K, Ishikawa T, Watanabe T, Suenaga K,  
12  
13 Rondon M, Rao IM (2006) Scope and strategies for regulation of nitrification in  
14  
15 agricultural systems—challenges and opportunities. *Crit Rev Plant Sci* 25:303-335  
16  
17
- 18 Subbarao G, Nakahara K, Hurtado MdP, Ono H, Moreta D, Salcedo AF, Yoshihashi A,  
19  
20  
21 Ishikawa T, Ishitani M, Ohnishi-Kameyama M (2009) Evidence for biological  
22  
23 nitrification inhibition in *Brachiaria* pastures. *Proc Natl Acad Sci* 106:17302-17307  
24  
25
- 26 van Kessel MA, Speth DR, Albertsen M, Nielsen PH, den Camp HJO, Kartal B, Jetten MS,  
27  
28  
29 Lucker S (2015) Complete nitrification by a single microorganism. *Nature* 528:555-  
30  
31 559  
32  
33
- 34 Wang YY, Cheng YH, Chen KE, Tsay YF (2018a) Nitrate transport, signaling, and use  
35  
36 efficiency. *Annu Rev Plant Biol* 69:85-122  
37  
38
- 39 Wang YF, Gu JD (2014) Effects of allylthiourea, salinity, and pH on ammonia/ammonium-  
40  
41 oxidizing prokaryotes in mangrove sediment incubated in laboratory microcosms. *Appl*  
42  
43 *Microbiol Biotechnol* 98:3257-3274  
44  
45
- 46 Wang M, Huang G, Zhao Z, Dang C, Liu W, Zheng M (2018b) Newly designed primer pair  
47  
48 revealed dominant and diverse comammox amoA gene in full-scale wastewater  
49  
50 treatment plants. *Bioresour Technol* 270:580-587  
51  
52
- 53 Zhang LM, Hu HW, Shen JP, He JZ (2012) Ammonia-oxidizing archaea have more important  
54  
55 role than ammonia-oxidizing bacteria in ammonia oxidation of strongly acidic soils.  
56  
57 *The ISME J* 6:1032-1045  
58  
59  
60  
61  
62  
63  
64  
65

Zhang S, Zheng Q, Noll L, Hu Y, Wanek W (2019) Environmental effects on soil microbial  
nitrogen use efficiency are controlled by allocation of organic nitrogen to microbial  
growth and regulate soil gross N mineralization. *Soil Biol Biochem* 135:304-315

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Figure legends

**Fig. 1** Changes in the ammonium and nitrate concentrations during the microcosm incubation of the pasture soil (A and C) and the arable soil (B and D). Error bars indicate standard errors of triplicate samples

**Fig. 2** Changes in the comammox *Nitrospira* clade A *amoA* gene abundance during the microcosm incubation of the pasture soil (A) and arable soil (B). Error bars indicate standard errors of triplicate samples

**Fig. 3** Changes in the AOA and AOB *amoA* gene abundance during the microcosm incubation of the pasture soil (A and C) and arable soil (B and D). Error bars indicate standard errors of triplicate samples

**Fig. 1**






Fig. 2



**Fig. 3**



**Table 1** Basic characteristics of the two soils in this study

<b>Property</b>	<b>Pasture soil (Dookie)</b>	<b>Arable soil (Clyde)</b>
Water content (%)	29.4	10.2
Soil pH (H <sub>2</sub> O)	6.90	7.13
Total C (g kg <sup>-1</sup> )	35	25
Total N (g kg <sup>-1</sup> )	3.8	2.9
NH <sub>4</sub> <sup>+</sup> -N (mg kg <sup>-1</sup> )	0.82	0.17
NO <sub>3</sub> <sup>-</sup> -N (mg kg <sup>-1</sup> )	38.60	21.83
Sulphur (mg kg <sup>-1</sup> )	22	45
Phosphorus (mg kg <sup>-1</sup> )	196	675
Potassium (mg kg <sup>-1</sup> )	348	129
Texture	Silty loam	Loamy sand

**Table 2** Two-way ANOVA analysis of mineral N concentrations and gene abundances

Factor	Source of variation	NH <sub>4</sub> <sup>+</sup> -N conc.		NO <sub>3</sub> <sup>-</sup> -N conc.		Coma clade A <i>amoA</i> copies		AOA <i>amoA</i> copies		AOB <i>amoA</i> copies	
		PS	AS	PS	AS	PS	AS	PS	AS	PS	AS
		Control	T	127.94**	35.77**	1161.83**	4685.68**	5.27*	3.21	15.38*	68.87**
VS (NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	D	119.27**	34.18**	687.27**	615.95**	11.14**	3.46*	3.78*	39.81**	12.65**	0.33
	T×D	124.23**	34.21**	49.54**	195.06**	1.76	2.43	1.63	7.62**	3.06*	4.51*
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	T	1591.35**	293.01**	640.42**	1092.99**	5.93*	17.91**	3.21	0.60	44.47**	23.55**
VS	D	60.40**	96.16**	451.04**	837.85**	7.54**	3.04*	0.47	5.42*	5.84*	0.49
NP	T×D	95.98**	16.51**	49.87**	69.77**	2.05	1.39	1.25	1.92	9.23**	3.20*
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	T	79.66**	30.73**	283.15**	175.72**	3.99	6.29*	0.08	0.16	12.96*	11.47*
VS	D	60.42**	92.23**	620.07**	611.22**	2.53	1.51	4.45*	33.22**	2.82*	1.39
DMPP	T×D	0.66	7.86**	18.65**	19.74**	2.15	0.55	0.90	5.52*	3.66*	2.21
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	T	0.00	1.22	87.37**	166.74**	6.22*	10.78*	6.79*	26.30**	18.35**	2.01
VS	D	346.02**	104.99**	556.97**	1043.04**	14.10**	2.16	3.70*	12.68**	16.39**	1.25
ATU	T×D	0.10	1.86	7.57**	19.89**	3.77*	1.18	2.04	0.92	3.41*	2.30
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	T	0.27	0.14	104.36**	107.04**	15.02*	2.51	12.66*	5.47*	35.66**	0.76
VS	D	168.52**	113.18**	611.73**	391.51**	10.49**	1.96	2.32	6.81**	23.17**	2.29
DCD	T×D	0.63	0.25	12.54**	17.10**	1.85	2.01	0.93	0.91	6.17*	1.34

The data indicates F value. Significance levels (P): \* $P < 0.05$ ; \*\* $P < 0.001$ . T, treatment; D, day; PS, pasture soil; AS, vegetable soil

## ELECTRONIC SUPPLEMENTARY MATERIAL

SOILS, SEC 1 • SOIL ORGANIC MATTER DYNAMICS AND NUTRIENT CYCLING •  
RESEARCH ARTICLE

**Growth of comammox *Nitrospira* is inhibited by nitrification inhibitors in agricultural  
soils**

**Chaoyu Li<sup>1</sup> • Hang-Wei Hu<sup>1</sup> • Qing-Lin Chen<sup>1</sup> • Deli Chen<sup>1</sup> • Ji-Zheng He<sup>1</sup>**

Received: 24 June 2019 / Accepted: 24 August 2019

© Springer-Verlag GmbH Germany, part of Springer Nature 2019

---

Responsible editor: Huaiying Yao

---


<sup>1</sup>School of Agriculture and Food, Faculty of Veterinary and Agricultural Sciences, The  
University of Melbourne, Parkville, Victoria 3010, Australia

✉ Ji-Zheng He

jizheng.he@unimelb.edu.au

**Table S1** Polymerase chain reaction (PCR) primer sets used in this study

Target gene	Primer name	Primer sequence (5'-3')	Amplification conditions	Length (bp)	References
Comammox <i>Nitrospira</i> clade A <i>amoA</i>	comaA-244f_a	TACAACTGGGTGAACTA	95°C for 10 min, 40 cycles of 94°C for 30 s, 54°C for 45 s and 72°C for 1 min	415	Pjevac et al. (2017)
	comaA-244f_b	TATAACTGGGTGAACTA			
	comaA-244f_c	TACAATTGGGTGAACTA			
	comaA-244f_d	TACAACTGGGTCAACTA			
	comaA-244f_e	TACAACTGGGTCAATTA			
	comaA-244f_f	TATAACTGGGTCAATTA			
	comaA-659r_a	AGATCATGGTGCTATG			
	comaA-659r_b	AAATCATGGTGCTATG			
	comaA-659r_c	AGATCATGGTGCTGTG			
	comaA-659r_d	AAATCATGGTGCTGTG			
	comaA-659r_e	AGATCATCGTGCTGTG			
comaA-659r_f	AAATCATCGTGCTGTG				
Comammox <i>Nitrospira</i> clade B <i>amoA</i>	comaB-244f_a	TAYTTCTGGACGTTCTA	95°C for 10 min, 40 cycles of 94°C for 30 s, 54°C for 45 s and 72°C for 1 min	415	Pjevac et al. (2017)
	comaB-244f_b	TAYTTCTGGACATTCTA			
	comaB-244f_c	TACTTCTGGACTTTCTA			
	comaB-244f_d	TAYTTCTGGACGTTTTA			
	comaB-244f_e	TAYTTCTGGACATTTTA			
	comaB-244f_f	TACTTCTGGACCTTCTA			
	comaB-659r_a	ARATCCAGACGGTGTG			
	comaB-659r_b	ARATCCAAACGGTGTG			
	comaB-659r_c	ARATCCAGACAGTGTG			
	comaB-659r_d	ARATCCAAACAGTGTG			
	comaB-659r_e	AGATCCAGACTGTGTG			
comaB-659r_f	AGATCCAAACAGTGTG				
AOA <i>amoA</i>	CrenamoA23f CrenamoA616r	ATGGTCTGGCTWAGACG GCCATCCATCTGTATGTCCA	95°C for 10 min, 35 cycles of 94°C for 30 s, 55°C for 30 s and 72°C for 1 min	629	Tourna et al. (2008)
AOB <i>amoA</i>	amoA-1F amoA-2R	GGGGTTTCTACTGGTGGT CCCCTCKGSAAAGCCTTCTTC	95°C for 10 min, 35 cycles of 94°C for 20 s, 56°C for 30 s and 72°C for 1 min	491	Rotthauwe et al. (1997)



**Fig. S1** Changes in soil pH during incubation trial of the pasture soil (A) and the arable soil (B). Error bars indicate standard errors of triplicate sample