



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Chen, ZY;Sukee, T;Koehler, AV;Webster, BL;Gasser, RB;Ponder, WF;Young, ND

Title:

Mitogenome and nuclear rRNA gene cluster of *Austropeplea subaquatilis* (Tate, 1880) from South Australia, with molecular and morphological comparison of *A. cf. brazieri* (Smith, 1882) from Victoria (Gastropoda, Hygrophila, Lymnaeidae)

Date:

2025-01-01

Citation:

Chen, Z. Y., Sukee, T., Koehler, A. V., Webster, B. L., Gasser, R. B., Ponder, W. F. & Young, N. D. (2025). Mitogenome and nuclear rRNA gene cluster of *Austropeplea subaquatilis* (Tate, 1880) from South Australia, with molecular and morphological comparison of *A. cf. brazieri* (Smith, 1882) from Victoria (Gastropoda, Hygrophila, Lymnaeidae). *Zookeys*, 1255 (1255), pp.41-62. <https://doi.org/10.3897/zookeys.1255.164109>.

Persistent Link:

<https://hdl.handle.net/11343/367694>

License:

CC BY

# Mitogenome and nuclear rRNA gene cluster of *Austropeplea subaquatilis* (Tate, 1880) from South Australia, with molecular and morphological comparison of *A. cf. brazieri* (Smith, 1882) from Victoria (Gastropoda, Hygrophila, Lymnaeidae)

Zhe-Yu Chen<sup>1</sup>, Tanapan Sukee<sup>1</sup>, Anson V. Koehler<sup>1</sup>, Bonnie L. Webster<sup>2</sup>, Robin B. Gasser<sup>1</sup>, Winston F. Ponder<sup>3</sup>, Neil D. Young<sup>1</sup>

<sup>1</sup> Department of Veterinary Biosciences, Melbourne Veterinary School, Faculty of Science, The University of Melbourne, Parkville, VIC 3010, Australia

<sup>2</sup> Department of Life Science, Natural History Museum, London, SW7 5BD, UK

<sup>3</sup> Australian Museum Research Institute, Australian Museum, Sydney, NSW 2010, Australia

Corresponding authors: Zhe-Yu Chen ([chenzheyu1998@163.com](mailto:chenzheyu1998@163.com)); Neil D. Young ([nyoung@unimelb.edu.au](mailto:nyoung@unimelb.edu.au))



Academic editor: Thierry Backeljau

Received: 4 July 2025

Accepted: 12 September 2025

Published: 9 October 2025

ZooBank: <https://zoobank.org/602AF124-3022-4B8C-BE69-C9F9006DE8CF>

**Citation:** Chen Z-Y, Sukee T, Koehler AV, Webster BL, Gasser RB, Ponder WF, Young ND (2025) Mitogenome and nuclear rRNA gene cluster of *Austropeplea subaquatilis* (Tate, 1880) from South Australia, with molecular and morphological comparison of *A. cf. brazieri* (Smith, 1882) from Victoria (Gastropoda, Hygrophila, Lymnaeidae). ZooKeys 1255: 41–62. <https://doi.org/10.3897/zookeys.1255.164109>

**Copyright:** © Zhe-Yu Chen et al. This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

## Abstract

Species of *Austropeplea* are lymnaeid snails endemic to Australia and New Zealand, and most are intermediate hosts of parasitic trematodes. Their taxonomy has long been uncertain due to the high phenotypic plasticity of most species. In this study, we used Oxford Nanopore sequencing technology to characterise the mitogenome and nuclear ribosomal RNA (rRNA) gene cluster of *Austropeplea subaquatilis* from South Australia to support comparative taxonomic investigations of this species. Then, *A. subaquatilis* was compared with *A. cf. brazieri* at both morphological and molecular levels. Morphologically, *A. subaquatilis* and *A. cf. brazieri* can be distinguished by shell morphometric indices, mantle edge morphology, pigmentation, and reproductive and neural anatomy. The two taxa differed by 1.9% in both the mitogenome and nuclear rRNA gene cluster. Sequence divergence was pronounced in the internal transcribed spacer (ITS) regions of the latter gene cluster, with nucleotide differences of 13.8% in ITS1 and 8.2% in ITS2. Phylogenetic analyses of sequence data for the mitochondrial 16S gene and ITS2 placed the two taxa in distinct groups. Taken together, the integrative evidence presented herein supported species-level divergence between *A. subaquatilis* and *A. cf. brazieri*.

**Key words:** Australia, freshwater snails, integrative taxonomy, mitochondrial genome, molecular phylogeny, nuclear rRNA gene cluster

## Introduction

Lymnaeidae Rafinesque, 1815 (commonly known as pond snails) are a globally distributed group of freshwater hygrophilid snails, which have attracted widespread attention as intermediate hosts of various species of parasitic trematodes (e.g., Vinarski et al. 2019, 2023; Vázquez et al. 2023). The Australian native lymnaeid genus *Austropeplea* Cotton, 1942, is one of the vectors of *Fasciola hepatica* (liver fluke) disease in Australia and New Zealand (Boray and McMichael 1961; Boray 1964a, 1969, 1978; Ponder and Waterhouse 1997).

Boray and McMichael (1961) synonymised all 23 previously named *Austropeplea* species to just one species, *Austropeplea tomentosa* (Pfeiffer, 1885), a name based on New Zealand specimens, although they recognised two distinct phenotypes A and B. The type A snails were darker with a smaller mantle border, and had a more robust and opaque shell with a distinct spire, while type B snails were lighter with a much larger mantle border, and with a fragile and transparent shell with a short spire (Boray and McMichael 1961; Boray 1964b). Despite the existence of two phenotypes, Boray and McMichael (1961) argued that they were all one species, based on assumed environment-mediated differences in snail phenotype and they cited evidence of successful hybridisation of the two phenotypes under laboratory conditions (Boray 1964b). In contrast, Puslednik et al. (2009), using morphological and molecular evidence, indicated that Australian *Austropeplea* represented a distinct lineage from the New Zealand *A. tomentosa*, and could be divided into different operational taxonomic units (OTUs). Based on these OTUs, Ponder et al. (2024) divided the Australian *Austropeplea* into four species in two subgenera, *Austropeplea* (*Austropeplea*) *brazieri* (Smith, 1882) which is widely distributed in eastern Australia, *Austropeplea* (*Austropeplea*) *subaquatilis* (Tate, 1880) in South Australia, and *Austropeplea* (*Austropeplea*) *huonensis* (Tenison Woods, 1876) and *Austropeplea* (*Kutikina*) *hispida* (Ponder & Waterhouse, 1997) in Tasmania. Nevertheless, due to limited defined morphological characters and the low phylogenetic resolution of current molecular markers, comprehensive species delimitation remains lacking for the nominal species other than *A. (K.) hispida*. As a result, the current classification of species (or OTUs) within the Australian members of typical *Austropeplea* largely relies on geographic origin rather than diagnostic features of the specimens themselves. Therefore, detailed morphological and molecular characterisation is needed to improve the taxonomic resolution within this genus.

The aim of this study was to characterise *A. subaquatilis* from South Australia morphologically and molecularly, and then compare this species with *A. cf. brazieri* from Victoria, Australia to establish their taxonomic and phylogenetic relationship.

## Materials and methods

### Sample collection and preservation

*Austropeplea subaquatilis* were collected from “Drain M” near Princes Highway in Thornlea, South Australia, Australia (latitude -37.36891397, longitude 140.2052126). *Austropeplea cf. brazieri* were previously collected from a roadside irrigation channel in Werribee South, Victoria, Australia (latitude -37.944706, longitude 144.698857) (see Sukee et al. 2024).

Both species of *Austropeplea* were then cultured in aquaria within a designated laboratory at The University of Melbourne, Victoria, Australia. Snails from different sources were maintained in strict isolation in separate tanks containing clean artificial pond water with aeration. Water was changed regularly and the snails were fed a commercial fish diet. A small section (~5 mm) of the foot muscle was excised from adult specimens and preserved in RNAlater at 4 °C for 24 h, -20 °C for one month and then stored at -80 °C until further processing. The remainder of each adult specimen was then placed in 70% ethanol for subsequent dissection and collection of morphological features.

## Gross morphology of mature snails

Animal external features in the living state were observed in the field and in individuals maintained in culture. Ten and 13 individuals of *A. subaquatilis* and *A. cf. brazieri* were dissected for internal morphological observation respectively, all individuals had attained a body size sufficient for oviposition. Terminology for the characters followed Ponder and Waterhouse (1997) and Vinarski and Pointier (2023). Images of living animals were captured using a Canon® 5D Mark IV camera with a Canon® EF 100 mm f/2.8L Macro IS USM lens. Preserved specimens were gently cleaned with soft brushes to remove the coagulated mucus and dissected under an Olympus SZ30 stereomicroscope. Detailed images of anatomical features were taken with the same Canon® 5D Mark IV camera with a Laowa® 25 mm f/2.8 2.5-5X Ultra Macro lens. The final high depth-of-field images were produced by a WeMacro® Rail System and stacked from 20–30 single photos using Zerene Stacker® 1.04. Anatomical illustrations were prepared using a Wild M5 stereomicroscope attached with a drawing tube. All images were modified and assembled using Adobe Photoshop 2023.

## Scanning electron microscopy

The radular sac was removed from three adult snails of each species and the soft tissue removed using a 10% potassium hydroxide (KOH) solution. After complete dissolution of soft tissues, each radula was washed extensively in MilliQ water. While still soft, each radula ribbon was transferred onto a round coverslip (Ø 13 mm), air-dried and fixed in place. The coverslip was then mounted onto an aluminium stub using conductive carbon tape (ProSciTech Pty Ltd). A 4-µm thick gold coating was deposited on the radula surface using SafeMatic® CCU-010 coater. Radulae were then scanned using a Hitachi® SU7000 scanning electron microscope under a low vacuum mode (5 kV). Images generated from the middle detector were used in this study.

## DNA isolation

Foot tissue of a specimen of *A. subaquatilis* (voucher number: AB291) was removed from the RNAlater and washed extensively in nuclease-free water (Qiagen). DNA was then isolated from each section of tissue using the E.Z.N.A. Mollusc DNA Kit according to manufacturer's instructions (Omega Bio-tek Inc.). The quantity of DNA isolated from each tissue was determined using the Qubit 1X dsDNA HS Assay and a Qubit 2.0 Fluorometer 2 (Invitrogen, ThermoFisher).

## Target-enriched amplification of mitochondrial DNA

Amplification of the *A. subaquatilis* mitochondrial DNA was performed using the REPLI-g mitochondrial DNA kit (Qiagen) following the manufacturer's protocol using a custom primer mix that was designed to match conserved regions of the lymnaeid 12S and 16S mitochondrial ribosomal subunits and the *cox1* gene. Primers were 11–14 nt in length and incorporated phosphorothioate links between the last three bases at the 3' end of the primers (Table 1). A total of 8 primers were combined into a 100 µM stock. Template DNA sample was diluted with water (supplied

**Table 1.** Modified primers used to amplify lymnaeid mitochondrial DNA using the REPLI-g mitochondrial DNA amplification kit (Qiagen). Asterisks represent the incorporated phosphorothioate links.

| Primer name   | Primer sequence   | Targeted region | Direction |
|---------------|-------------------|-----------------|-----------|
| RepGS_Aust16F | TACCTGTTTATC*A*A  | 16S             | Forward   |
| RepGS_16SBRis | AACTCAGATCAT*G*T  | 16S             | Reverse   |
| RepGS_12sF    | CAACGGCAATAT*A*T  | 12S             | Forward   |
| RepGS_12sR    | CTAGGATTAGAT*A*C  | 12S             | Reverse   |
| RepGT_JB3     | ATCCT GAGGTTT*A*T | cox1            | Forward   |
| RepGT_JB4.5   | ACATAATGAAAA*T*G  | cox1            | Reverse   |
| RepGT_16sF    | CCTTTTGCATCA*T*G  | 16S             | Forward   |
| RepGT_16sR    | CGGTCTTAACTC*A*A  | 16S             | Reverse   |

from the kit) to 150 ng in a total volume of 20 µL. A fresh amplification mix containing the custom primers was prepared as per manufacturer's instructions (Qiagen). In brief, 29 µL of DNA template and primer mix was denatured at 75 °C for 5 min then cooled to room temperature. Next, 1 µL of REPLI-g Midi Polymerase was added, and the sample was incubated at 33 °C for 8 h. Finally, the polymerase was deactivated by raising the temperature to 65 °C for 3 min. Amplified DNA was quantified as described above and stored at 4 °C until further processing.

### Library preparation and long-read sequencing

For *A. subaquatilis*, a barcode was assigned to amplified DNA template using the RAPID 24 (SQK-RBK114) barcoding library kit (Oxford Nanopore Technologies) and loaded onto a R10.4.1 flow cell and sequenced for 4 h on the PromethION 2 Solo (Oxford Nanopore Technologies) sequencing platform. Post-sequencing base-calling of POD5 data was performed using the program Dorado v. 0.7.2 (Oxford Nanopore Technologies) in super-accurate mode and reads were stored in FASTQ format. Nanopore long read sequence data for *A. cf. brazieri* was available from a previous study (Sukee et al. 2024).

### Clustering analyses, mitochondrial genome assembly, and gene annotations

Long reads with homology to reference mitochondrial genomes or ribosomal RNA subunit of freshwater molluscs were identified using pblat v. 2.5.1 (Wang and Kong 2019) and extracted from the raw long read data file using seqtk v. 1.4-r122 (<https://github.com/lh3/seqtk/>). A consensus mitochondrial sequence or nuclear rRNA gene cluster sequence was assembled using canu v. 2.3 (<https://github.com/marbl/canu>) with minimum read length set to 1,000 bp and genome size set to 20,000 bp (mitochondrial genome) or 8,000 bp (nuclear rRNA gene cluster). The *A. subaquatilis* mitochondrial genome was primarily annotated using Geneious Prime v. 2024.0.7 and MITOS v. 2.0.2 (Bernt et al. 2013), then manually curated following the suggested roles for molluscs (Fourdrilis et al. 2018; Ghiselli et al. 2021). The published *A. cf. brazieri* mitochondrial genome (GenBank accession number [PP100270](#)) were also further curated according to the same criteria. All sequence annotations and GC content plots were visualised using the Proksee online server (Grant et al. 2023).

## Nucleotide diversity comparison

Nucleotide pairwise distances (p-distance) of complete mitochondrial genome, the nuclear rRNA gene cluster and each gene were calculated in Geneious Prime v. 2024.0.7 after alignment with Clustal Omega v. 1.2.3. Sliding window analyses of nucleotide diversity (300-bp windows with 10-bp steps for mitochondrial genome and 50-bp windows with 10-bp steps for nuclear rRNA gene cluster) were performed on the aligned mitogenomes and nuclear rRNA gene cluster of *A. cf. brazieri* and *A. subaquatilis* using the PopGenome package (Pfeifer et al. 2014) in R. For each comparison, nucleotide diversity values were plotted using the R package ggplot2 (Wickham 2016). The GC/AT content and skew values were calculated using a custom Python v. 3.9.13 script.

## Phylogenetic analyses

To compare *A. cf. brazieri* and *A. subaquatilis* samples with available molecular data for *Austropeplea* spp. (see Puslednik et al. 2009), phylogenies were reconstructed using only the mitochondrial 16S and nuclear ITS2 regions (Suppl. material 1: table S1). *Orientogalba viridis* was used as outgroup (Liu et al. 2012; Suwancharoen et al. 2023). The 16S and ITS2 sequences were aligned separately using MUSCLE v. 3.7 in AliView v. 1.28 and concatenated using catfasta2phyml (<https://github.com/nylander/catfasta2phyml>). Numbers of variable and parsimony informative sites of the nucleotide were calculated using MEGA v. 11.0.13. Partitioned Maximum Likelihood (ML) analyses with the majority-rule consensus tree were performed in IQ-TREE v. 1.6.12 (Minh et al. 2013) using Ultrafast fast bootstrap approach (Minh et al. 2013) with 10,000 reiterations. Bayesian inference (BI) was conducted in MrBayes v. 3.2.6 (Ronquist et al. 2012), with four independent runs, each of which was performed for 1,000,000 generations and sampled every 1000 generations with the first 25% samples discarded as burn-in. Convergence of the Markov chain Monte Carlo simulations was assessed to ensure that the average standard deviation of split frequencies was < 0.01 and the potential scale reduction factors (PSRFs) were ~1. Additionally, Tracer v. 1.7.2 (Rambaut et al. 2018) was used to verify that all effective sample size (ESS) values exceeded 200. The most appropriate model of sequence evolution (ML: K2P+G4 for ITS2 and K3Pu+F+G4 for 16S; BI: K2P+G4 for ITS2 and HKY+F+G4 for 16S) was selected using ModelFinder (Kalyaanamoorthy et al. 2017).

## Results

### Mitochondrial genome composition and comparison

The mitochondrial genomes (mitogenome) of *A. subaquatilis* (GenBank accession number [PV749633](#)) and *A. cf. brazieri* (GenBank accession number [PP100270](#)) are 13,768 and 13,757 bp, respectively (Fig. 1A, B). Both genomes contain 37 genes: 13 protein-coding genes (PCGs), 2 ribosomal RNA genes (rRNAs), and 22 transfer RNA genes (tRNAs) (Fig. 1A, B, Table 2). Eight (*nad4L*, *cytb*, *cox1*, *cox2*, *atp8*, *atp6*, *nad3*, *nad2*) of the PCGs terminate with a truncated stop codon, which is assumed to be completed as




**Figure 1.** Reference mitochondrial genome and nuclear rRNA gene cluster of *subaquatilis* (A, C) and *Austropeplea cf. brazieri* (B, D). The direction of gene transcription is shown with an arrow. In each panel GC content is displayed via the observed skew patterns.

TAA by the addition of 3' A residues to the mRNA during transcription. The arrangement of genes within both mitogenomes are identical with only minor differences in inferred lengths of non-coding tRNA (tRNA-D, F, W, C, G, H, Q, L2, M, T and K) and 12S genes (Table 2). The mitogenome nucleotide composition of *A. subaquatilis* is 32.8% A, 40.7% T, 12.4% C and 14.1% G, resulting in an AT content of 73.4%, which is nearly same as the 73.3% observed in *A. cf. brazieri* (with 32.8% A, 40.5% T, 12.5% C and 14.2% G). The AT/GC skew values for *A. subaquatilis* and *A. cf. brazieri* are -0.1072/0.0624 and -0.1051/0.0612, respectively. The overall plots of GC content in *A. subaquatilis* closely resembles those observed in *A. cf. brazieri*, with no substantial differences in content or distribution patterns across the mitogenomes (Fig. 1A, B). Both species exhibit higher GC content in the *cox3* and *cox1* regions and reduced GC content in *nad2*, *nad6*, and 16S rRNA. The overall mitochondrial genome p-distance between *A. subaquatilis* and *A. cf. brazieri* is 1.9%. The p-distance of each of the PCGs range from 3.1% (*nad5*) to 0.6% (16S) (Fig. 2A, Table 2).




**Figure 2.** Sliding window analysis of the pairwise differences in the nucleotide identity of *Austropeplea subaquatilis* and *Austropeplea cf. brazieri* mitochondrial genomes (A) and nuclear rRNA gene cluster (B). Gene boundaries are indicated by vertical dotted lines. The horizontal dotted line indicates the average nucleotide diversity between the two sequences.

### Nuclear rRNA gene cluster composition and comparison

The completed nuclear rRNA gene cluster of *A. subaquatilis* (GenBank accession no. [PV593739](#)) and *A. cf. brazieri* (GenBank accession no. [PV593740](#)) span 6,712 bp and 6,747 bp respectively (Fig. 1C, D). The two ribosomal DNA sequences exhibit similar overall structures, comprising the 18S rRNA, 5.8S rRNA, 28S rRNA genes and two internal transcribed spacer regions (ITS1 and ITS2) (Fig. 1C, D, Table 2). Both sequences share identical lengths for the 5.8S rRNA (158 bp). The 18S rRNA and 28S rRNA regions measure 1,865 bp and 3,815 bp in *A. subaquatilis*, and 1,866 bp and 3,832 bp in *A. cf. brazieri*, respectively. Notable differences are observed in the ITS regions: *A. subaquatilis* contained a 509 bp ITS1 and a 368 bp ITS2, whereas *A. cf. brazieri* possesses a 504 bp ITS1 and a 388 bp ITS2 (Table 2). The nuclear rRNA gene cluster of *A. subaquatilis* and *A. cf. brazieri* show similar nucleotide compositions: 22.8% A, 21.8% T, 25.7% C, and 29.7% G in *A. subaquatilis*; 22.5% A, 22.0% T, 25.7% C, and 29.8% G in *A. cf. brazieri*. Both sequences have consistent GC content (55.4% and 55.5%, respectively) and similar GC skew values, while *A. subaquatilis* shows slightly higher AT skew than *A. cf. brazieri* (AT/GC skew: 0.0227/0.0715 and 0.0123/0.0737). The GC content plots of the nuclear rRNA gene cluster in *Austropeplea subaquatilis* and *A. cf. brazieri* are similar (Fig. 1C, D). In both species, the GC content plots exhibit several local maxima in GC proportion within the central region of the 28S rRNA gene, while pronounced decreases in GC proportion were observed at the boundaries of the internal transcribed spacers (ITS1 and ITS2), particularly at their junctions with the adjacent 5.8S and 28S rRNA genes. The overall nuclear rRNA gene cluster p-distance between *A. subaquatilis* and *A. cf. brazieri* was 1.9%. While the rRNA genes are largely identical (99%–100%), the ITS regions exhibit substantial sequence divergence, with the p-distance of ITS1 and ITS2 regions being 13.8% and 8.2%, respectively (Fig. 2B, Table 2).

**Table 2.** Location, lengths, and directions of annotated genes within the mitochondrial and nuclear rRNA gene clusters of *Austropeplea subaquatilis*, with comparisons to *Austropeplea cf. brazieri* (values after slash). Nucleotide pairwise identity of each gene between the two species is shown.

| Gene designations         | Location start | Location end | Length (bp) | Direction | Pairwise Identity | p-distance |
|---------------------------|----------------|--------------|-------------|-----------|-------------------|------------|
| Mitochondrial genes       |                |              |             |           |                   |            |
| 16S                       | 1/1            | 986/986      | 986/986     | forward   | 99.4%             | 0.6%       |
| tRNA-L1(tag)              | 988/988        | 1051/1051    | 64/64       | forward   | N/A               | N/A        |
| tRNA-P(tgg)               | 1047/1047      | 1106/1106    | 60/60       | forward   | N/A               | N/A        |
| tRNA-A(tgc)               | 1107/1107      | 1170/1170    | 64/64       | forward   | N/A               | N/A        |
| nad6                      | 1171/1171      | 1629/1629    | 459/459     | forward   | 97.6%             | 2.4%       |
| nad5                      | 1631/1631      | 3277/3277    | 1647/1647   | forward   | 96.9%             | 3.1%       |
| nad1                      | 3279/3279      | 4154/4154    | 876/876     | forward   | 98.3%             | 1.7%       |
| nad4L                     | 4155/4155      | 4452/4452    | 298/298     | forward   | 98.3%             | 1.7%       |
| cytb                      | 4453/4453      | 5533/5533    | 1081/1081   | forward   | 98.3%             | 1.7%       |
| tRNA-D(gtc)               | 5536/5536      | 5588/5587    | 53/52       | forward   | N/A               | N/A        |
| tRNA-F(gaa)               | 5589/5588      | 5651/5650    | 63/63       | forward   | N/A               | N/A        |
| cox2                      | 5652/5651      | 6294/6293    | 643/643     | forward   | 98.3%             | 1.7%       |
| tRNA-Y(gta)               | 6297/6296      | 6346/6345    | 50/50       | forward   | N/A               | N/A        |
| tRNA-W(tca)               | 6347/6346      | 6405/6405    | 59/60       | forward   | N/A               | N/A        |
| tRNA-C(gca)               | 6410/6410      | 6468/6468    | 59/59       | forward   | N/A               | N/A        |
| tRNA-G(tcc)               | 6471/6470      | 6524/6522    | 54/53       | forward   | N/A               | N/A        |
| tRNA-H(gtg)               | 6527/6525      | 6583/6582    | 57/58       | forward   | N/A               | N/A        |
| tRNA-Q(ttg)               | 6592/6591      | 6650/6649    | 59/59       | reverse   | N/A               | N/A        |
| tRNA-L2(taa)              | 6651/6650      | 6703/6701    | 53/52       | reverse   | N/A               | N/A        |
| atp8                      | 6705/6703      | 6855/6853    | 151/151     | reverse   | 98.2%             | 1.8%       |
| tRNA-N(gtt)               | 6857/6854      | 6920/6917    | 64/64       | reverse   | N/A               | N/A        |
| atp6                      | 6921/6918      | 7560/7557    | 640/640     | reverse   | 98.6%             | 1.4%       |
| tRNA-R(tcg)               | 7561/7558      | 7623/7620    | 63/63       | reverse   | N/A               | N/A        |
| tRNA-E(gaa)               | 7624/7621      | 7675/7672    | 52/52       | reverse   | N/A               | N/A        |
| 12S                       | 7676/7673      | 8393/8388    | 718/716     | reverse   | 98.2%             | 1.8%       |
| tRNA-M(cat)               | 8394/8389      | 8466/8457    | 73/69       | reverse   | N/A               | N/A        |
| nad3                      | 8467/8458      | 8806/8797    | 340/340     | reverse   | 99.1%             | 0.9%       |
| tRNA-S2(tga)              | 8817/8808      | 8871/8862    | 55/55       | reverse   | N/A               | N/A        |
| tRNA-S1(gct)              | 8872/8863      | 8926/8917    | 55/55       | reverse   | N/A               | N/A        |
| nad4                      | 8927/8918      | 10252/10243  | 1326/1326   | forward   | 98%               | 2%         |
| tRNA-T(tgt)               | 10253/10244    | 10319/10311  | 67/68       | reverse   | N/A               | N/A        |
| cox3                      | 10321/10313    | 11100/11092  | 780/780     | reverse   | 98.8%             | 1.2%       |
| tRNA-I(gat)               | 11141/11133    | 11205/11197  | 65/65       | forward   | N/A               | N/A        |
| nad2                      | 11206/11198    | 12109/12101  | 904/904     | forward   | 97.8%             | 2.2%       |
| tRNA-K(ttt)               | 12110/12102    | 12191/12180  | 82/79       | forward   | N/A               | N/A        |
| cox1                      | 12203/12192    | 13694/13683  | 1492/1492   | forward   | 97.7%             | 2.3%       |
| tRNA-V(tac)               | 13695/13684    | 13755/13744  | 61/61       | forward   | N/A               | N/A        |
| Nuclear rRNA gene cluster |                |              |             |           |                   |            |
| 18S                       | 1/1            | 1865/1865    | 1865/1865   | forward   | 100%              | 0%         |
| ITS1                      | 1866/1866      | 2374/2369    | 509/504     | forward   | 86.2%             | 13.8%      |
| 5.8S                      | 2375/2370      | 2532/2527    | 158/158     | forward   | 100%              | 0%         |
| ITS2                      | 2533/2528      | 2900/2915    | 368/388     | forward   | 91.8%             | 8.2%       |
| 28S                       | 2901/2916      | 6731/6747    | 3831/3832   | forward   | 99.3%             | 0.7%       |

## Phylogenetic analysis

Phylogenetic trees were constructed from a dataset consisting of 29 16S + ITS2 sequences of *Austropeplea*, 27 of them were obtained from previous study (Puslednik et al. 2009), two were generated for this study and one available out-group taxon (Liu et al. 2012; Suwancharoen et al. 2023). The aligned lengths of 16S and ITS2 genes were 433 and 452 nucleotides, respectively. Within these sequences, 79 and 87 sites were variable, while 69 and 46 sites were parsimony informative. The Bayesian-inferred (BI) and maximum likelihood (ML) phylogenetic trees of *Austropeplea* were not fully resolved. Nevertheless, both Bayesian posterior probabilities (BPP) and ML bootstrap (BS) values supported separation between Australian and New Zealand species. *Austropeplea hispida* was recovered as sister to the remaining Australian species, which together formed a trichotomy in the inferred topology (Fig. 3). The sequences generated here for *Austropeplea subaquatilis* and *A. cf. brazieri* group with the southern Australian and eastern Australian samples from previous study, respectively.

## Taxonomic account

### *Austropeplea subaquatilis* (Tate, 1880)

Figs 4A, C, E, 5A, C, 6A, C, 7A

**Material examined.** Specimens from “Drain M” near Princes Highway in Thornlea, South Australia, Australia, and their artificially bred offspring.

**Description. Shell** (Fig. 4A) medium in size (up to 12.5 mm in height), ovate, with low, narrow conical spire and strongly inflated last whorl. Shell wall thin, fragile in some specimens. Whorls (4.0–4.5 in number) rounded, slightly convex, separated by a shallow, slightly oblique to nearly straight suture. Last whorl comprises ~0.9 of shell height. Shell surface smooth, somewhat shiny, light brown to nearly colourless, covered by collabral growth lines. Aperture pyriform, with evenly rounded basal and palatal margins, posterior corner forming angle with last whorl. Peristome sharp, not expanded but columellar lip reflexed and attached to back of last whorl. Parietal callus thin but distinct, extending to last whorl far beyond inner lip. Columellar fold weakly developed. Umbilicus covered by inner lip, closed or very narrow (slot-like).

**Head-foot** (Fig. 4C, E) typical of family. Foot broad, reaching 1.5–2× shell height when fully extended, light grey with sparse white freckles (observed when living). When stimulated, considerable quantities of mucus produced and covers entire body. Tentacles shield-shaped (Fig. 4E), twice as long as wide (observed when living). Mantle light grey with large black blotches on pallial roof. Mantle collar (Fig. 4E) reflexed and attached to shell, extended as thin flap on both sides to enclose shell fully or largely in mature individuals. Closed edge of mantle collar situated along midline of animal, forming marginal fold near shell apex. Mantle covering visceral coil with band-like black pigment in mature individuals (Fig. 4C); disconnected from pigmentation on pallial roof, and readily lost in preserved specimens.

**Central nervous system** typical of family (Fig. 5A). Cerebral ganglia with regular borders, pale yellow (fresh). Commissural lobule distinct, white, approximately equal to cerebral ganglia in size.




**Figure 3.** Bayesian phylogenetic tree of *Austropeplea* based on combined 16S and ITS2 sequences. Red tip labels represent the sequences generated in this study. Coloured shading and the corresponding inset map summarise the geographic distributions of the principal lineages. Maximum likelihood (ML) bootstrap (BS) and Bayesian posterior probability (BPP) support values for shown for each node. Scale bar represents substitutions per site.

**Pulmonary roof** (pallial complex) (Fig. 2C) with heart and kidney in their typical positions for family. Kidney spindle-shaped, thin-walled, with transversely pleated lining of sinuate tube visible through transparent wall, proximal part opposite anterior pericardium. Ureter short, urinary opening not observed.

**Prostate** pear-shaped, with single internal fold. Sperm duct long and thick, equal to or slightly longer than oothecal gland in length. Praeputium (Fig. 6A) light greyish-white, cylindrical, tapers towards proximal end, distal part folded near male genital opening. Bulbous termination of praeputium distinct in lighter colour to white, equal to or wider than narrowest part of praeputium. Penis sheath narrow, shorter than praeputium, proximal part slightly inflated. ICA 1.34 to 2.02. Spermatheca (Fig. 6C) spherical, duct short, not exceeding length of spermatheca, width approx. 0.1 of its length.

**Radula** of the haplolateral multidentate type (Fig. 7A). Radular formula 28-C-28 to 32-C-32. Teeth in same row bend upward to margin. Central tooth small, bicuspid, asymmetrical, right cusp significantly larger than left. Lateral teeth pairs 1–8~11 tricuspid, middle cusp largest, left cusp larger than right, rarely with denticle situated on the right basal side; pairs 9~12–28~32 with four or five cusps.

**Remarks.** The name *A. subaquatilis* (Tate, 1880: 103, pl. 4, Fig. 5A–C) is here tentatively considered to be the species name for South Australian *Austropeplea* populations following Ponder et al. (2024). This species was described based on type material collected from the River Torrens in Ade-




**Figure 4.** Shells (A, B), mantle pigmentations (C, D) and living individuals (E, F) of *Austropeplea* spp. A, C, E. *Austropeplea subaquatilis*; B, D, F. *Austropeplea* cf. *brazieri*. G. Schematic representation of mantle extension ranges (green areas), *A. subaquatilis* shown above, *A. cf. brazieri* below. The red arrows (in C) indicate the mantle pigmentation on the visceral coil. Scale bar: 5 mm (except G not to scale).

laide and are not aware of it having been found from type locality in recent years (Z.-Y. Chen, unpublished data). A currently recognised synonym, *A. aruntalis* Cotton & Godfrey, 1938 (replacement name for *Limnaea papyracea* Tate, 1880: 103, pl. 4, Fig. 6A, B), may represent a valid species name for the South Australian Limestone Coast population, from which the specimens in this study were collected, if future studies reveal consistent differences between populations of the two forms.

**Distribution.** South-eastern South Australia and western Victoria (Ponder et al. 2024).




**Figure 5.** Dorsal side of central nervous system (A, B) and pallial complex (C, D) of *Austropeplea* spp. A, C. *Austropeplea subaquatilis*; B, D. *Austropeplea* cf. *brazieri*. Abbreviations: an – anus, au – auricle, cc – cerebral commissure, cg – cerebral ganglia, cl – commissural lobule, k – kidney, pn – cut wall of pneumostome, pv – pulmonary vein, re – rectum, ve – ventricle. The red-dotted line shows the additional lobe from the cerebral ganglion with a clear boundary. Scale bars: 1 mm.

### ***Austropeplea* cf. *brazieri* (E. A. Smith, 1882)**

Figs 4B, D, F, 5B, D, 6B, D, 7B

**Material examined.** Artificially bred specimens in the lab, which were originally from Werribee South, Victoria, Australia.

**Description. Shell** (Fig. 4B) medium in size ( $\leq 14.7$  mm in height in lab condition), high-conical, with relatively narrow and high spire and moderately inflated body whorl. Shell thin but somewhat solid, in some specimens almost translucent. Whorls (4–4.5 in number) rounded, convex, slowly increasing, separated by deep and oblique suture. Shell surface smooth, light brown, covered by collabral




**Figure 6.** Male (A, B) and female (C, D) copulatory apparatus of *Austropeplea* spp. A, C. *Austropeplea subaquatilis*; B, D. *Austropeplea* cf. *brazieri*. Abbreviations: bt – bulbous termination of praeputium, prep – praeputium, prer – praeputium retractor muscles, ps – penis sheath, sp – spermatheca, srd – spermatheca duct, vag – vagina, vd – vas deferens. Scale bars: 1 mm.

growth lines. Aperture pyriform, with evenly rounded basal and palatal margins. Peristome sharp, not expanded but columellar lip slightly reflexed. Parietal callus thin but distinct, extending a little over the parietal wall. Columellar fold weakly developed. Umbilicus covered by inner lip, closed or very narrow (slot-like).

**Head-foot** (Fig. 4D, F) typical of family. Foot broad, fully extended approximately equal to shell height, light grey with dense white freckles (observed when living). Tentacle elongated triangular, length equal to width (observed when living). Mantle light grey with large black blotches on pallial roof. Mantle collar slightly reflexed and attached to aperture margin (Fig. 4F). Mantle covering visceral coil unpigmented (Fig. 4D).

**Central nervous system** (Fig. 5B) typical of family. Cerebral ganglia with irregular-borders, pale yellow in fresh material. Commissural lobule distinct, white, notably smaller in size than cerebral ganglia.

**Pulmonary roof** (Fig. 5D) with heart and kidney in typical positions for family. Kidney spindle-shaped, thin-walled, with transversely pleated lining of sinuate tube visible through translucent wall, proximal part opposite to anterior pericardium. Ureter short, urinary opening not observed.

**Prostate** fusiform, with single internal fold. Sperm duct short, almost invisible in natural position. Praeputium (Fig. 6B) light greyish white, cylindrical, tapers




**Figure 7.** Radulae of *Austropeplea* spp. A. *Austropeplea subaquatilis*; B. Normal form of *Austropeplea* cf. *brazieri*; C. Variant form of *Austropeplea* cf. *brazieri*. Scale bars: 100 µm.

towards proximal end, distal part near opening folded. Bulbous termination of praeputium distinct in lighter colour to white, narrowest across praeputium. Penis sheath narrow, shorter than praeputium, proximal part slightly inflated. Index of copulatory apparatus (ICA) 1.12 to 2.21. Spermatheca (Fig. 6D) ellipsoid. Spermatheca duct shorter than length of spermatheca, width approx. quarter of length, proximal external side somewhat adherent to the vaginal duct.

**Radula** of the haplolateral multidentate type (Fig. 7B). Radular formula 28-C-28 to 38-C-38. Teeth in same row approximately aligned horizontally. Central tooth small, bicuspid, asymmetrical, right cusp significantly larger than left. Lateral teeth pairs 1–11~13 tricuspid, middle cusp largest, left cusp larger than right, rarely with denticle situated on the right basal side; pairs 12~14–28~38 with four or five cusps. A variant individual observed among examined specimens (Fig. 7C), with lateral teeth pairs 1–13 tricuspid, middle cusp largest, left cusp larger than right, rarely left cusp absent; pairs 14–17 tricuspid, left cusp largest, right two cusps gradually reduced; pairs 18–28 with 3–6 cusps.

**Remarks.** We use the species-group name *A. cf. brazieri* for our Victorian specimens because both molecular and morphological data indicate clear differences from typical *Austropeplea brazieri* (E. A. Smith, 1882: 274, pl. 5, fig. 15, from Glebe Point, Sydney, New South Wales). The currently recognised distribution of *A. brazieri* is broad (Ponder et al. 2024), and our analysis and that done previously by

Puslednik et al. (2009) have shown that it does not form a monophyletic group. To reflect this taxonomic uncertainty, we follow our earlier usage (Sukee et al. 2024) in applying the provisional name *A. cf. brazieri* to the Victorian population.

There are three additional names available for this taxon, all from New South Wales (*Glacilimnaea gelida* Iredale, 1943: 214, Blue Lake, Mt Kosciusko, NSW; *Simlimnea morbida* Iredale, 1944: 119, figs 5-4, Walcha, NSW; and *Simlimnea aegrifer* Iredale, 1944: 119, fig. 5-5, Bombala, NSW), but based on the molecular data of species from representative location of above (Fig. 3), none are applicable to the Victorian taxon.

**Distribution.** Victoria, Australia.

## Morphological remarks

Due to its short apex and very large aperture, *A. subaquatilis* can be distinguished from *A. cf. brazieri* by shell morphometrics such as the Index Spire height/Shell height and Index Aperture height/Shell height (Table 3). A distinguishing feature that allows mature individuals of *A. subaquatilis* to be readily separated from those of *A. cf. brazieri* is the markedly expanded and extended mantle edge in *A. subaquatilis*. Additionally, the shell of *A. subaquatilis* is fragile and relatively small compared to the broad foot of the animal. Under conditions of high population density and limited food availability in aquaria, *A. subaquatilis* often entered a state of developmental arrest (diapause), producing dwarf forms lacking mantle extension, yet remaining reproductively functional. When these dwarf individuals were transferred to low-density environments with adequate food supply, they resumed growth and eventually developed an expanded mantle that enclosed the shell, similar to wild-type adults. This phenomenon has not been observed in *A. cf. brazieri* under same laboratory conditions.

**Table 3.** Morphometric characteristics of studied specimens of *Austropeplea subaquatilis* and *Austropeplea cf. brazieri*. ICA: Index of the copulatory apparatus, ratio of praeputium length and penis sheath length. Parenthetical values represent the mean of the group, \* indicates  $p < 0.001$ .

| Characters and Indexes      | Species                         |                                 | p |
|-----------------------------|---------------------------------|---------------------------------|---|
|                             | <i>A. subaquatilis</i> (n = 10) | <i>A. cf. brazieri</i> (n = 13) |   |
| Number of whorls            | 3.75–4.5                        | 4–4.25                          |   |
| Shell height (SH), mm       | 9.02 – (10.37) – 12.5           | 9.71 – (11.30) – 14             |   |
| Shell width (SW), mm        | 6.04 – (6.87) – 8.41            | 5.92 – (7.14) – 8.98            |   |
| Spire height (SpH), mm      | 1.45 – (2.07) – 2.63            | 2.86 – (3.79) – 4.66            | * |
| Body whorl height (BWH), mm | 8.58 – (9.54) – 11.45           | 8.63 – (9.95) – 12.19           |   |
| Aperture height (AH), mm    | 7.27 – (8.20) – 9.75            | 6.36 – (7.46) – 9.5             |   |
| Aperture width (AW), mm     | 4.29 – (5.05) – 6.48            | 3.84 – (4.72) – 6.12            |   |
| Index SW/SH                 | 0.63 – (0.66) – 0.70            | 0.55 – (0.63) – 0.69            |   |
| Index SpH/SH                | 0.16 – (0.20) – 0.21            | 0.28 – (0.34) – 0.41            | * |
| Index BWH/SH                | 0.87 – (0.92) – 0.95            | 0.86 – (0.88) – 0.92            |   |
| Index AH/SH                 | 0.75– (0.79) – 0.85             | 0.59 – (0.66) – 0.71            | * |
| Index AW/AH                 | 0.44 – (0.49) – 0.52            | 0.34– (0.42) – 0.45             | * |
| ICA                         | 1.34 – (1.69) – 2.02            | 1.12 – (1.51) – 2.21            |   |

The mantle pigment on the visceral coil of mature individuals of *A. subaquatilis* does not occur in *A. cf. brazieri*. The internal pigmentation of *A. subaquatilis* is generally lighter than that of *A. cf. brazieri*. The difference in the relative size of the bulbous termination of the praeputium is a consistent distinction in the male reproductive system of the two species, as are the shape of the spermatheca and the width of the spermathecal duct in the female system. Apart from the larger commissural lobule in *A. cf. brazieri*, mature *A. subaquatilis* formed a distinctly demarcated and ellipsoid lobe from each cerebral ganglion situated opposite the commissural lobule (namely adjacent to the buccal mass).

## Discussion

This study used an integrative approach, combining morphological data, mitogenome and nuclear rRNA gene cluster comparisons and phylogenetic analyses to assess the relationship between *A. subaquatilis* and *A. cf. brazieri*. Based on morphological features, these two snail taxa could be readily distinguished. The complete mitochondrial and nuclear rRNA gene clusters sequenced and compared herein are some of the first such data sets for the family Lymnaeidae and are among the very few available for the order Hygrophila (McQuirk et al. 2025). Analysis of mitochondrial and nuclear markers showed no significant difference in gene order or structure, although sufficient phylogenetically informative sites were reported to suggest that their divergence corresponded to species-level differentiation, and thus supported our morphological findings.

The patterns of nucleotide divergence in mitochondrial genes and the nuclear rRNA clusters of *A. subaquatilis* and *A. cf. brazieri* are distinct. Most mitochondrial genes exhibit low pairwise nucleotide divergence between the two species. For instance, the nucleotide p-distance for the *cox1* gene is 2.3%, a value that was at the threshold between intra- and interspecific variation in other lymnaeid taxonomic studies (Bolotov et al. 2014; Vinarski et al. 2016, 2022; Aksenova et al. 2017, 2024; Lounnas et al. 2018; Ferreira et al. 2021; Falniowski et al. 2023), and is comparable to the minimal genetic divergence (p-distance = 2.4%) observed between *Ladislavella tumrokensis* and *L. elodes*, and between *Galba cubensis* and *G. neotropica* (see Vinarski et al. 2016; Ferreira et al. 2021). Moreover, the mitochondrial 16S rRNA gene shows the lowest p-distance (0.6%) among all mitochondrial genes between the two species, further supporting previous findings that mitochondrial 16S rRNA alone lacked sufficient phylogenetic resolution to delimit Australian *Austropeplea* species (Puslednik et al. 2009; Sukee et al. 2024). In contrast, although the sequences of the three nuclear rRNA genes (18S, 5.8S, 28S) of the two species are nearly identical, the observed differences in ITS regions (p-distance = 13.8% for ITS1; p-distance = 8.2% for ITS2) fall within the range of interspecific divergence commonly used in current lymnaeid taxonomic practices (see Vinarski et al. 2016; Ferreira et al. 2021; Aksenova et al. 2024). The higher divergence in ITS regions compared with mitochondrial genes may suggest the presence of introgression and incomplete lineage sorting within Australian *Austropeplea* (see Davis and Nixon 1992; Doyle 1992; Harrison and Larson 2014). Laboratory experiments showed that mating between species of *Austropeplea* was possible (Boray 1964b), although the fertility of hybrid offspring had not been demonstrated.

Despite the limitations of using 16S gene and ITS2 markers alone for this group, we proceeded to use only these regions to make a phylogenetic compari-

son with data available for *Austropeplea*. The topology of the 16S + ITS2 phylogenetic trees in this study and that of Puslednik et al. (2009) were largely consistent.

The newly sequenced *A. subaquatilis* and *A. cf. brazieri* were placed in two geographically structured lineages within *Austropeplea* (southern versus eastern Australia), consistent with regional structuring rather than constituting definitive evidence of species level divergence. Nevertheless, the limitations of the current markers were evident, as the main Australian *Austropeplea* lineage and several within group relationships showed polytomies, indicating unresolved relationships. These unresolved nodes may have reflected limited phylogenetic signal, rapid radiation, or incomplete lineage sorting, and thus warrant further investigation using additional genetic markers. Future studies employing complete mitochondrial genome sequences and nuclear rRNA gene clusters are warranted to address the unresolved relationships within the genus.

The morphological features of *A. subaquatilis* and *A. cf. brazieri* generally corresponded to the type B and type A morphs, respectively, as described by Boray and McMichael (1961). Morphologically, the differences between the two species were primarily focused on the shell, mantle extension, nervous system, and the reproductive system. The differences in the reproductive system between *A. subaquatilis* and *A. cf. brazieri* were distinct. The morphological difference in the bulbous termination of the praeputium may suggested underlying internal structural differences, which warrant further investigation in future studies. The nervous systems of *A. subaquatilis* and *A. cf. brazieri* differed, particularly in the size of the commissural lobule and the presence of additional lobes arising from the cerebral ganglia in *A. subaquatilis*. These differences were not reported in previous studies. The lack of comparative neuroanatomical studies on this group currently limited our ability to assess whether these features were structurally significant or how they related to other members of the Lymnaeidae. Further research is needed to clarify their nature and taxonomic relevance. The two unique features mentioned by Puslednik et al. (2009) for the South Australian samples (*A. subaquatilis* in this study), namely the longer cephalic tentacles (twice as long as wide) and the prostate (tube) being much longer than the female reproductive system, were also observed in this study. However, the recognition of these traits is largely depended on how much contraction had occurred due to preservation, as these features were less pronounced in contracted material compared to the other distinguishing features noted above.

The external features of the head-foot and mantle of *A. subaquatilis* included a broad foot with the shell enveloped by the mantle. In its natural habitat, *A. subaquatilis* was observed living in extremely dense water milfoil (*Myriophyllum* spp.) environments, which presumably necessitated moving between tightly packed branches. The reduction of shell-related hindrance may have enhanced the animal's ability to survive in such environments. The abnormally high mucus production in *A. subaquatilis* when stimulated may have served as an alternative defence mechanism against predators, perhaps compensating for the fragile shell. This trait was commonly observed in land slugs and semi-slugs (Luchtel and Deyrup-Olsen 2001). The mantle extension may also have functioned to enhance cutaneous respiration, thereby reducing the frequency of surfacing for air or adapting to hypoxic conditions caused by intensified respiration of aquatic plants at night (Russell-Hunter 1978). The degree of mantle extension has not been well characterised across species in this family, with only the Europe-

an *Myxas glutinosa* (O. F. Müller, 1774) explicitly documented as having a fully shell-enveloping mantle, similar to what was observed in *A. subaquatilis* (Hubendick 1951; Stadnichenko 2004). However, based on the phylogenetic relationships of *Austropeplea* from Tasmania and South Australia reconstructed in this study and previous research (Puslednik et al. 2009), it appeared that the development of mantle extensions in *Austropeplea* was not an isolated occurrence. Boray and McMichael (1961) noted that mantle extension in their type B morphs tended to diminish or disappear over successive laboratory generations. This observation closely resembled the dwarf forms we observed. However, these dwarf individuals regained mantle extension once placed in favourable environmental conditions, in contrast to *A. cf. brazieri*, in which this trait appeared to be permanently absent. Further studies are required to clarify the mechanism of mantle extension and to determine its biological or ecological relevance.

Taken together, the integrative evidence presented herein supported species-level divergence between *A. subaquatilis* and *A. cf. brazieri*. While mitochondrial divergence alone approached the threshold of interspecific separation, the pronounced differences in ITS regions, along with the morphological traits, provided a coherent framework for species delimitation. Nevertheless, the phylogeny of this group remains poorly resolved. Continued efforts involving additional phylogenetically informative genetic datasets and comparative anatomical studies will be essential for resolving outstanding taxonomic uncertainties and understanding the evolutionary dynamics of this group.

## Acknowledgements

We are grateful to Ms Christine Andersen, Prof. Ian Beveridge, and Mr Kasem Driver for their invaluable support during the fieldwork.

## Additional information

### Conflict of interest

The authors have declared that no competing interests exist.

### Ethical statement

No ethical statement was reported.

### Use of AI

No use of AI was reported.

### Funding

This study was supported by the Australian Research Council Discovery Project no. DP230100270.

### Author contributions

All authors have contributed equally.

### Author ORCIDs

Zhe-Yu Chen  <https://orcid.org/0000-0002-4150-8906>

Tanapan Sukee  <https://orcid.org/0000-0003-3181-5045>

Anson V. Koehler  <https://orcid.org/0000-0001-8330-6416>  
Bonnie L. Webster  <https://orcid.org/0000-0003-0930-9314>  
Robin B. Gasser  <https://orcid.org/0000-0002-4423-1690>  
Winston F. Ponder  <https://orcid.org/0000-0002-8600-3952>  
Neil D. Young  <https://orcid.org/0000-0001-8756-229X>

## Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

## References

- Aksenova O, Vinarski M, Bolotov I, Kondakov A, Beshpalaya Y, Tomilova A, Paltser I, Gofarov M (2017) Two *Radix* spp. (Gastropoda: Lymnaeidae) endemic to thermal springs around Lake Baikal represent ecotypes of the widespread *Radix auricularia*. *Journal of Zoological Systematics and Evolutionary Research* 55(4): 298–309. <https://doi.org/10.1111/jzs.12174>
- Aksenova OV, Vinarski MV, Itagaki T, Ohari Y, Oshida T, Kim SK, Lee JH, Kondakov AV, Khrebtova IS, Soboleva AA, Travina OV, Sokolova SE, Palatov DM, Beshpalaya YV, Vikhrev IV, Gofarov MY, Bolotov IN (2024) Taxonomy and trans-Beringian biogeography of the pond snails (Gastropoda: Lymnaeidae) of East Asia: an integrative view. *Zoological Journal of the Linnean Society* 201(4): zlae083. <https://doi.org/10.1093/zoolinnean/zlae083>
- Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsche G, Pütz J, Middendorff M, Stadler PF (2013) MITOS: Improved *de novo* metazoan mitochondrial genome annotation. *Molecular Phylogenetics and Evolution* 69(2): 313–319. <https://doi.org/10.1016/j.ympev.2012.08.023>
- Bolotov I, Beshpalaya Y, Aksenova O, Aksenov A, Bolotov N, Gofarov M, Kondakov A, Paltser I, Vikhrev I (2014) A taxonomic revision of two local endemic *Radix* spp. (Gastropoda: Lymnaeidae) from Khodutka geothermal area, Kamchatka, Russian Far East. *Zootaxa* 3869(5): 585–593. <https://doi.org/10.11646/zootaxa.3869.5.9>
- Boray JC (1964a) Studies on the ecology of *Lymnaea tomentosa*, the intermediate host of *Fasciola hepatica*. 1. History, geographical distribution, and environment. *Australian Journal of Zoology* 12(2): 217–230. <https://doi.org/10.1071/ZO9640217>
- Boray JC (1964b) Studies on the ecology of *Lymnaea tomentosa*, the intermediate host of *Fasciola hepatica*. 2. The sexual behaviour of *Lymnaea tomentosa*. *Australian Journal of Zoology* 12(2): 231–238. <https://doi.org/10.1071/ZO9640231>
- Boray JC (1969) Experimental Fascioliasis in Australia. In: Dawes B (Ed.) *Advances in Parasitology*. Academic Press, 95–210. [https://doi.org/10.1016/S0065-308X\(08\)60435-2](https://doi.org/10.1016/S0065-308X(08)60435-2)
- Boray JC (1978) The potential impact of exotic *Lymnaea* spp. on fascioliasis in Australasia. *Veterinary Parasitology* 4(2): 127–141. [https://doi.org/10.1016/0304-4017\(78\)90004-3](https://doi.org/10.1016/0304-4017(78)90004-3)
- Boray JC, McMichael DF (1961) The identity of the Australian lymnaeid snail host of *Fasciola hepatica* L. and its response to environment. *Marine and Freshwater Research* 12(2): 150–163. <https://doi.org/10.1071/MF9610150>
- Cotton BC (1942) Some Australian freshwater Gastropoda. *Transactions of the Royal Society of South Australia* 66: 75–82.
- Cotton BC, Godfrey FK (1938) A systematic list of the Gastropoda, the marine freshwater and land univalve Mollusca of South and Central Australia. *Malacological Society of South Australia Publication No. 1*, 44 pp.

- Davis JI, Nixon KC (1992) Populations, genetic variation, and the delimitation of phylogenetic species. *Systematic Biology* 41(4): 421–435. <https://doi.org/10.1093/sysbio/41.4.421>
- Doyle JJ (1992) Gene trees and species trees: Molecular systematics as one-character taxonomy. *Systematic Botany* 17(1): 144–163. <https://doi.org/10.2307/2419070>
- Falniowski A, Jaszczyńska A, Hofman S (2023) *Radix rufescens* (J. E. Gray, 1822) (Gastropoda: Lymnaeidae), a new species for Oman and Arabian Peninsula. *Acta Zoologica Academiae Scientiarum Hungaricae* 69(3): 303–312. <https://doi.org/10.17109/AZH.69.3.303.2023>
- Ferreira APPN, Costa ALO, Becattini RM, Ferreira MAND, da Paixão HPR, Coscarelli D, Vidigal THDA, Lima W dos S, Pereira CA de J (2021) Integrative taxonomy: Combining molecular and morphological characteristics to identify *Lymnaea (Galba) cubensis*, intermediate host of *Fasciola hepatica*. *Revista Brasileira de Parasitologia Veterinária = Brazilian Journal of Veterinary Parasitology : Órgão Oficial do Colégio Brasileiro de Parasitologia Veterinária* 30(2): e026320. <https://doi.org/10.1590/s1984-29612021052>
- Fourdrilis S, de Frias Martins AM, Backeljau T (2018) Relation between mitochondrial DNA hyperdiversity, mutation rate and mitochondrial genome evolution in *Melanhaphe neritoides* (Gastropoda: Littorinidae) and other Caenogastropoda. *Scientific Reports* 8(1): 17964. <https://doi.org/10.1038/s41598-018-36428-7>
- Ghiselli F, Gomes-dos-Santos A, Adema CM, Lopes-Lima M, Sharbrough J, Boore JL (2021) Molluscan mitochondrial genomes break the rules. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 376(1825): 20200159. <https://doi.org/10.1098/rstb.2020.0159>
- Grant JR, Enns E, Marinier E, Mandal A, Herman EK, Chen C, Graham M, Van Domselaar G, Stothard P (2023) Proksee: In-depth characterization and visualization of bacterial genomes. *Nucleic Acids Research* 51(W1): W484–W492. <https://doi.org/10.1093/nar/gkad326>
- Harrison RG, Larson EL (2014) Hybridization, introgression, and the nature of species boundaries. *The Journal of Heredity* 105: 795–809. <https://doi.org/10.1093/jhered/esu033>
- Hubendick B (1951) Recent Lymnaeidae: Their variation, morphology, taxonomy, nomenclature and distribution. *Kungliga Svenska Vetenskapsakademiens Handlingar. Fjärde Serien* 3: 1–223.
- Iredale T (1943) A basic list of the freshwater Mollusca of Australia. *Australian Zoologist* 10(2): 188–230.
- Iredale T (1944) Guide to the freshwater shells of New South Wales. Pt II. *Australian Naturalist* 11: 113–127.
- Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermin LS (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods* 14(6): 587–589. <https://doi.org/10.1038/nmeth.4285>
- Liu G-H, Wang S-Y, Huang W-Y, Zhao G-H, Wei S-J, Song H-Q, Xu M-J, Lin R-Q, Zhou D-H, Zhu X-Q (2012) The complete mitochondrial genome of *Galba pervia* (Gastropoda: Mollusca), an intermediate host snail of *Fasciola* spp. *PLoS ONE* 7(7): e42172. <https://doi.org/10.1371/journal.pone.0042172>
- Lounnas M, Correa AC, Alda P, David P, Dubois M-P, Calvopiña M, Caron Y, Celi-Erazo M, Dung BT, Jarne P, Loker ES, Noya O, Rodríguez-Hidalgo R, Toty C, Uribe N, Pointier J-P, Hurtrez-Boussès S (2018) Population structure and genetic diversity in the invasive freshwater snail *Galba schirazensis* (Lymnaeidae). *Canadian Journal of Zoology* 96(5): 425–435. <https://doi.org/10.1139/cjz-2016-0319>
- Luchtel DL, Deyrup-Olsen I (2001) Body wall: form and function. In: Barker GM (Ed.) *The biology of terrestrial molluscs*. CABI Publishing, Oxon, 147–178. <https://doi.org/10.1079/9780851993188.0147>

- McQuirk KA, DeCore JM, Castillo MG, Adema CM (2025) Rewilding shows differential fitness of sympatric *Physella acuta* (Draparnaud, 1805) snail lineages. *Aquatic Ecology* 59(2): 435–454. <https://doi.org/10.1007/s10452-025-10172-3>
- Minh BQ, Nguyen MAT, von Haeseler A (2013) Ultrafast approximation for phylogenetic bootstrap. *Molecular Biology and Evolution* 30(5): 1188–1195. <https://doi.org/10.1093/molbev/mst024>
- Müller OF (1774) Vermium terrestrium et fluviatilium, seu animalium infusorium, Helminthicorum, et testaceorum, non marinorum, succincta historia. vol 2. Havniae et Lipsiae, apud Heineck et Faber, ex officina Mölleriana. I-XXXVI, 1–214 pp. [10 unnumbered pages] <https://doi.org/10.5962/bhl.title.46299>
- Pfeifer B, Wittelsbürger U, Ramos-Onsins SE, Lercher MJ (2014) PopGenome: An efficient Swiss Army knife for population genomic analyses in R. *Molecular Biology and Evolution* 31(7): 1929–1936. <https://doi.org/10.1093/molbev/msu136>
- Pfeiffer L (1855) Descriptions of fifty-seven new species of Helicea, from Mr Cunningham's collection. *Proceedings of the Zoological Society of London* 1854: 286–298. <https://doi.org/10.1111/j.1469-7998.1854.tb07277.x>
- Ponder WF, Waterhouse JH (1997) A new genus and species of Lymnaeidae from the lower Franklin River, south western Tasmania, Australia. *The Journal of Molluscan Studies* 63(3): 441–468. <https://doi.org/10.1093/mollus/63.3.441>
- Ponder WF, Hallan A, Shea ME, Clark SA, Richards K, Klunzinger MW, Kessner V (2024) Australian Freshwater Molluscs. Revision 2A. [https://keys.lucidcentral.org/keys/v3/freshwater\\_molluscs/](https://keys.lucidcentral.org/keys/v3/freshwater_molluscs/)
- Puslednik L, Ponder WF, Dowton M, Davis AR (2009) Examining the phylogeny of the Australasian Lymnaeidae (Heterobranchia: Pulmonata: Gastropoda) using mitochondrial, nuclear and morphological markers. *Molecular Phylogenetics and Evolution* 52(3): 643–659. <https://doi.org/10.1016/j.ympev.2009.03.033>
- Rafinesque CS (1815) Analyse de la nature ou Tableau de l'univers et des corps organisés. Published by the author, Palermo, 224 pp. <https://doi.org/10.5962/bhl.title.106607>
- Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA (2018) Posterior Summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* 67(5): 901–904. <https://doi.org/10.1093/sysbio/syy032>
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Russell-Hunter WD (1978) Ecology of freshwater pulmonates. In: Fretter V, Peake J (Eds) *Pulmonates*, vol 2A. Academic Press, London, 335–384.
- Smith EA (1882) On the freshwater shells of Australia. *Journal of the Linnean Society of London, Zoology* 16(92): 255–316. <https://doi.org/10.1111/j.1096-3642.1882.tb02283.x>
- Stadnichenko AP (2004) Pond snails and limpet snails (Lymnaeidae and Acroloxidae) of Ukraine. Tsentr uchebnoi literatury, Kiev, 327 pp. [in Russian]
- Sukee T, Koehler AV, Webster BL, Gauci CG, Fogarty CE, Ponder WF, Gasser RB, Young ND (2024) Mitochondrial genome of the fluke pond snail, *Austropeplea* cf. *brazieri* (Gastropoda: Lymnaeidae). *Parasites & Vectors* 17(1): 283. <https://doi.org/10.1186/s13071-024-06358-7>
- Suwancharoen C, Phuangsri C, Siriwechviriyi P, Bunsong T, Japa O (2023) Diversity of trematode cercariae among naturally infected lymnaeid snails from Phayao, Thailand. *Parasitology Research* 122(11): 2691–2708. <https://doi.org/10.1007/s00436-023-07971-8>

- Tate R (1880) Descriptions of some new species of south Australian Pulmonifera. Transactions of the Royal Society of South Australia 3: 102–104.
- Tenison Woods JE (1876) On the freshwater shells of Tasmania. Papers and Proceedings of the Royal Society of Tasmania 1875: 66–82. <https://doi.org/10.5962/bhl.part.9781>
- Vázquez AA, Alba A, Alda P, Vittecoq M, Chapuis E, Faugère D, Pointier J-P, Hurtrez-Boussès S (2023) Lymnaeid snails and the transmission of Fasciolosis: Understanding the differential risks from local to global scale. In: Vinarski MV, Vázquez AA (Eds) The Lymnaeidae: A Handbook on Their Natural History and Parasitological Significance. Springer International Publishing, Cham, 359–394. [https://doi.org/10.1007/978-3-031-30292-3\\_13](https://doi.org/10.1007/978-3-031-30292-3_13)
- Vinarski MV, Pointier J-P (2023) Conchological and anatomical identification of the lymnaeid snails. In: Vinarski MV, Vázquez AA (Eds) The Lymnaeidae: A Handbook on Their Natural History and Parasitological Significance. Springer International Publishing, Cham, 103–120. [https://doi.org/10.1007/978-3-031-30292-3\\_4](https://doi.org/10.1007/978-3-031-30292-3_4)
- Vinarski M, Aksenova O, Bespalaya Y, Bolotov I, Gofarov M, Kondakov A (2016) *Ladislavella tumrokensis*: The first molecular evidence of a Nearctic clade of lymnaeid snails inhabiting Eurasia. Systematics and Biodiversity 14(3): 276–287. <https://doi.org/10.1080/14772000.2016.1140244>
- Vinarski MV, Clewing C, Albrecht C (2019) Lymnaeidae Rafinesque, 1815. In: Lydeard C, Cummings S (Eds) Freshwater Mollusks of the World. Johns Hopkins University Press, Baltimore, 158–162.
- Vinarski MV, Von Oheimb PV, Aksenova OV, Gofarov MY, Kondakov AV, Nekhaev IO, Bolotov IN (2022) Trapped on the roof of the world: taxonomic diversity and evolutionary patterns of Tibetan Plateau endemic freshwater snails (Gastropoda: Lymnaeidae: Tibetoradix). Integrative Zoology 17(5): 825–848. <https://doi.org/10.1111/1749-4877.12600>
- Vinarski MV, Aksenova OV, Bolotov IN, Vázquez AA, Alda P, Pointier J-P, Hurtrez-Boussès S (2023) Biogeography of the living Lymnaeidae. In: Vinarski MV, Vázquez AA (Eds) The Lymnaeidae: A Handbook on Their Natural History and Parasitological Significance. Springer International Publishing, Cham, 183–206. [https://doi.org/10.1007/978-3-031-30292-3\\_7](https://doi.org/10.1007/978-3-031-30292-3_7)
- Wang M, Kong L (2019) pblat: A multithread blat algorithm speeding up aligning sequences to genomes. BMC Bioinformatics 20(1): 28. <https://doi.org/10.1186/s12859-019-2597-8>
- Wickham H (2016) ggplot2. Elegant Graphics for Data Analysis. Second Edition. Springer International Publishing, Cham, 260 pp. <https://doi.org/10.1007/978-3-319-24277-4>

## Supplementary material 1

### Location, code, and NCBI GenBank accession numbers of the lymnaeid snail nucleotide sequences used in this study

Authors: Zhe-Yu Chen, Tanapan Sukee, Anson V. Koehler, Bonnie L. Webster, Robin B. Gasser, Winston F. Ponder, Neil D. Young

Data type: pdf

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1255.164109.suppl1>