

Literature Review of Benign Müllerian Papilloma contrasted with Vaginal
Rhabdomyosarcoma

Dr. Sarah K. McQuillan, MD, FRCSC
Obstetrics & Gynecology
University of Melbourne
The Royal Children's Hospital Melbourne

Dr. Sonia R. Grover
MBBS, MD, FRANZCOG
Department of Paediatric Gynaecology
University of Melbourne
The Royal Children's Hospital Melbourne

Dr. Jan Pyman
MBBS
Department of Anatomical Pathology
University of Melbourne
The Royal Women's Hospital

Dr. Yasmin L. Jayasinghe
MBBS, FRANZCOG, PhD
Department of Obstetrics & Gynaecology Royal Women's Hospital
University of Melbourne
The Royal Children's Hospital Melbourne

Corresponding Author:

Dr. Sarah K. McQuillan

Sarah.mcquillan@medportal.ca

The Royal Children's Hospital

50 Flemington Rd., Parkville, Vic, Australia, 3052

(03) 9345 5522 ext: 55310

Fax number: (03) 9345 5585

No financial support to disclose.

Abstract:

Study Objectives: Benign Müllerian papillomas of the genital tract are rare hence can be mistaken for vaginal rhabdomyosarcoma on initial clinical review. This review of the literature will consolidate the previous cases of Müllerian papilloma reported and look for clues to differentiate the 2 entities.

Design: Case report and literature review

Setting: Pediatric Adolescent Gynecology Clinic in Tertiary Center

Data Source: A search of English language publications from January 2014 until 1951 (the first case report) was performed using the search words “Müllerian papilloma” and “pre-pubertal bleeding”. References from previous published reports were also obtained for completeness.

Main Outcome: Literature review of Benign Müllerian papilloma

Results: Since 1951, fifty-six cases of Müllerian papilloma were reported, including 4 cases at our own institution. Co-morbid conditions were found in 31.5% of cases (with 3 cases associated with mesenchymal tumors). The average length of time from onset of symptoms (primarily vaginal bleeding) to diagnosis was 6.7 months (range 1 day to 3 years) with only one case diagnosed incidentally. Median age of presentation was 5 years (range 1day-52 years). Most cases were localized and resected with ease. Histology reveals complex papillary lesions without cytological atypia.

Conclusion: Benign Müllerian papilloma is distinguished from the more significant diagnosis of vaginal rhabdomyosarcoma by initial length of vaginal bleeding at presentation, lack of vaginal wall extension, ease of resection and histopathology. This is compared with vaginal rhabdomyosarcoma, commonly exhibit localized or distant spread at presentation.

Keywords: Benign Müllerian papilloma, rhabdomyosarcoma, prepubescent

1 INTRODUCTION:

2 Pre-pubertal vaginal bleeding is most frequently caused by local non-hormonal causes
3 such as trauma, a foreign body or severe vulvovaginitis (1). Rarely, causes such as
4 rhabdomyosarcoma of the vagina can present as vaginal bleeding. Rhabdomyosarcoma is
5 the most common soft tissue sarcoma in children accounting for 4-6% of all malignancies
6 in this age group (0-19). The most common primary sites for rhabdomyosarcoma are the
7 head, the genitourinary tract, and the extremities(2). Younger age at diagnosis (1-9) and
8 non-bladder/non-prostate genitourinary tract subtype have more favorable prognoses (3).
9 The mean age of diagnosis of vaginal rhabdomyosarcoma is 2 years with almost all
10 occurring in children under the age of 5 (4) compared with cervical rhabdomyosarcoma
11 which is more commonly diagnosed in adolescences (5). Benign Müllerian papillomas
12 (BMP) are rare polypoid masses, which also present with pre-pubertal bleeding, and on
13 first inspection at vaginoscopy, may have an appearance very similar to either vaginal or
14 cervical rhabdomyosarcoma. Therefore, this review aims to characterize any
15 distinguishing features that may aid in the differential diagnosis and thus guide
16 appropriate intraoperative management prior to final histopathology assessment. We will
17 present 4 case reports of BMP along with a review of fifty-two reports in the literature,
18 which may shed further light on clinical presentation and pathogenesis of these rare
19 tumors.

20

21 **Case 1:** A 5-year-old girl presented with a 2-year history of intermittent painless bright
22 red vaginal bleeding of varying amounts. She had attended previous health providers
23 who had initially diagnosed vulvovaginitis, however with ongoing symptoms she was

24 referred to a Pediatric Gynecology Centre. At the initial consultation, a full history and
25 physical examination using the frog leg position were performed. There was no personal
26 history of trauma, irritation, bleeding diathesis, and no family history of gynecologic
27 cancer. Examination was unremarkable for irritation, or vulvar dermatoses such as lichen
28 sclerosis, and there was no obvious vaginal bleeding. Breast and pubic hair were Tanner
29 stage 1. A bleeding profile was undertaken in view of the unusual duration of bleeding
30 and the activated partial thromboplastin time (aPTT) was slightly increased however a
31 bleeding disorder was excluded. Hormonal testing revealed a normal pre-pubertal
32 profile. Ultrasound showed no significant abnormalities with normal urinary bladder,
33 prepubescent ovaries and uterus. Vaginoscopy revealed a 3cm polypoid friable but solid
34 mass arising from the ectocervix with no obvious extension into the anterior or posterior
35 vaginal walls (Figure 1). A biopsy was performed and 2 weeks later a debulking
36 procedure removed the entire mass to the base of the lesion using forceps and electro-
37 cautery. The pathology described BMP as a complex papillary lesion covered by
38 cuboidal or low columnar surface epithelium without cytological atypia supported by
39 mature fibrous cores.

40

41 **Case 2:** A 7-year-old girl presented with a history of intermittent painless vaginal
42 bleeding. She had a past history of osteosarcoma of the jaw 3 years earlier, treated with
43 surgery, chemotherapy and radiation and considered cured. Physical examination
44 confirmed Tanner Stage 1 breast/pubertal hair with no evidence of vulvovaginitis.
45 Vaginoscopy revealed a small polypoid mass with no extension into the surrounding
46 tissue, originating from the region of the cervical os. This was removed entirely at initial

47 excision/biopsy surgery. At her most recent follow up at age 31, she has had no
48 recurrence.

49

50 **Case 3:** A 9-year-old girl presented with a three-week history of bright red painless
51 vaginal bleeding on a background history of constipation. Her physical examination was
52 unremarkable. Because of the multiple recurrences of bleeding despite treatment for
53 constipation she underwent a vaginoscopy, which showed a frond-like polypoid mass
54 originating from the cervix with the final pathology confirming BMP.

55

56 **Case 4:** A 10-year-old girl with a history of insulin dependent diabetes presented with a
57 one-year history of intermittent painless vaginal bleeding. She was pre-pubertal on
58 examination and investigation. A vaginoscopy showed a very small polypoid lesion
59 originating from the cervix. An excisional biopsy to remove the lesion was undertaken
60 using hysteroscopic forceps.

61

62 In summary, all patients presented with intermittent painless bright red prepubescent
63 vaginal bleeding with Tanner Stage 1 breast/pubertal hair and no evidence of vulvovaginitis
64 with polypoid masses on vaginoscopy. Vaginoscopy was used to visualize the source of
65 bleeding and the polypoid mass was subsequently removed ensuring the entire lesion was
66 removed from the base of the stalk. The diagnosis of BMP was confirmed in all cases by
67 2 independent pathologists both specializing in gynecology pathology (CWC and JP).
68 Cytogenetics of the lesion confirmed no clonal abnormalities in case 1 (Figure 3). All 4
69 patients have been followed up postoperatively with no more episodes of vaginal

70 bleeding. Case 1, 3 and 4 have had 1-2 years of follow-up. Case 2 has had yearly
71 Papanicolaou testing in adulthood with no visible evidence of recurrence, and has had
72 one hysteroscopy for abnormal uterine bleeding at age 26 (with benign proliferative
73 endometrium on pathology).

74

75 METHODS:

76 A MEDLINE search was conducted of all articles published using the search terms
77 “Müllerian papilloma” and “prepubertal vaginal bleeding” until January 2014. Standard
78 reference tracing was also performed for completeness. Nineteen articles fitting the
79 search criteria were initially obtained using MEDLINE and a further seven were found
80 using reference tracing. All articles were read and non-English articles were excluded.
81 The term “Müllerian papilloma” was first employed in 1981(6) and prior to that the
82 terminology was found to be inconsistent. All the articles included were case reports,
83 mostly of 1 or 2 cases with 1 case series including both pre-pubertal and adult cases of
84 BMP that included 14 cases (7).

85

86 RESULTS:

87 Benign Müllerian papilloma (BMP) is a rare tumor of the female genital tract first
88 reported in 1951(8). Since that first case, there have been 55 further reports from 23
89 different articles (including our case series of 4, summarized in this review) in the
90 literature involving the cervix, vagina or both. These are summarized in Table 1(6, 9-29).
91 The majority of cases (32/56, 57.1%) occurred in pre-pubertal girls with a median age of
92 presentation being 5 years (range 1 day – 52 years). Similar to our cases, most subjects

93 presented with a history of painless intermittent bright red vaginal bleeding except for
94 one report where the BMP was found incidentally(21). In the literature, vaginal bleeding
95 was often first managed as vaginitis with review of vulvar hygiene and then only with
96 subsequent diagnostic vaginoscopy if there was no resolution. On vaginoscopy, the
97 presentation of BMP can parallel more sinister diagnoses such as rhabdomyosarcoma,
98 endodermal sinus tumor, clear cell adenocarcinoma and papillary carcinomas because of
99 its polypoid appearance. Diagnosis relies on histopathology. Definitive management
100 requires excision of the tumor. The biopsy procedure and the excision may be done as a
101 two-step procedure, which is the standard of care unless the excision is unavoidable with
102 a biopsy (as the treatment for a rhabdomyosarcoma may require a conservative wide local
103 excision with the possible addition of either one or both of chemotherapy and
104 radiotherapy (30, 31)). Frequently by the time a rhabdomyosarcoma presents with vaginal
105 bleeding, there is already distant metastasis (in 28.4%) or direct/regional infiltration to
106 adjacent organs (in 16.2%) (32).

107

108 Currently, 5 cases of recurrent BMP have been reported in the English literature (7, 9, 12,
109 16, 21, 27) with 1 of these 5 patients experiencing multiple recurrences and eventually
110 being diagnosed with a malignant transformation (9, 12). That case involved a woman
111 with an intellectual disability and severe cerebral palsy who, at the age of 42 years,
112 underwent partial excision of the mass, with pathology showing borderline changes in the
113 BMP of the vagina with a normal cervix. Three years later she was diagnosed with clear
114 cell carcinoma of the upper one third of the anterior vagina, which was managed
115 conservatively (without surgery) due to the patient's other medical conditions. The

116 associated use of diethylstilbestrol (DES) in this patient in utero could not be determined
117 and would be useful to know due to its established association with vaginal adenosis and
118 vaginal clear cell carcinoma (33). Three of the 5 patients with recurrences documented in
119 the literature, had recurrences on more than 1 occasion (7, 9, 16, 21, 27). Given the rarity
120 and limited number of recurrences and only 1 documented case of both borderline and
121 subsequently malignant transformation, radical surgery on initial presentation in
122 childhood should be avoided.

123 DISCUSSION:

124 Given the concern regarding the alternative diagnosis of rhabdomyosarcoma in the
125 context of vaginal bleeding and a polypoid lesion on vaginoscopy, the main
126 distinguishing features of BMP may include, a relatively long duration of prepubertal
127 bleeding prior to presentation, lack of systemic or abdominal physical examination
128 findings, frond-like structure on vaginoscopy originating from either the vaginal wall or
129 cervix with lack of extension into the vaginal walls (rather than the grape-like vesicular
130 appearance of vaginal or cervical rhabdomyosarcoma), ease of resection and
131 characteristic histopathology showing complex arborizing papillae with fibrovascular
132 cores, and lined by bland cuboidal or low columnar epithelium with no cytological atypia
133 or mitotic activity. Our literature review demonstrates that the mean time to diagnosis for
134 BMP from initial onset of symptoms was 6.7 months (range 1 day to 36 months). In
135 comparison, genital rhabdomyosarcoma has a more fulminant course, with 16.2%
136 presenting with local/regional metastasis at the time of clinical presentation with
137 mucosanguinous discharge (32, 34-36) (Table 2).

138

139 It is important to note that the majority of cases of BMP reported in the literature
140 particularly in the young, have been associated with another medical presentation ranging
141 from physiological conditions or benign complaints (chronic abdominal pain, pregnancy)
142 to more concerning pathology (history of osteosarcoma). Therefore, in some of these
143 patients, BMP may have been discovered incidentally during management of other
144 conditions, or may have been diagnosed earlier due to co-existing medical care.

145 Furthermore, only cases that cause vaginal bleeding will be investigated by
146 vaginoscopy/hysteroscopy, thus, it is entirely possible that asymptomatic cases are not
147 detected, or may even spontaneously resolve, as suggested by Norris *et al.* in 1966(7).

148

149 BMP are thought to develop from maternal hormones that induce intrauterine
150 development of the tumors according to Norris *et al.*(7). However the presence of
151 mesenchymal tumors/conditions [osteosarcoma (current report), Proteus syndrome(22),
152 Wilms tumor (15)] reported in our review is interesting and may shed light on a separate
153 pathophysiology. Vaginal bleeding in a prepubertal girl with a history of mesenchymal
154 tumor may raise further suspicion regarding the presence of a BMP. We await further
155 reports, which will enable exploration of such a potential association.

156

157 Less than 10% (5/56) of reported cases of BMP presenting with abnormal uterine
158 bleeding eventually have recurred, with only 1 borderline lesion and subsequent
159 malignant transformation occurring in the same patient (0.9%), Given the paucity of
160 cases reported in the literature, and the low possibility of recurrence the question of long
161 term follow up remains a challenge, particularly in the pediatric population. When these

162 young women reach the age for cervical screening according to respective country
163 guidelines, it may be useful to include the clinical history of previous BMP diagnosis
164 both to assist the pathologist in review of the slides or to flag the potential for
165 cytopathologist review rather than automated review or HPV DNA testing alone (37).
166 This has been recommended for cases associated with recurrences (21). In 1 case report
167 published in 2007, BMP was confirmed on routine cytological smear as well as on the
168 Thin Prep™ PreservCy™ slide (cytyc Corp., Boxborough, Massachusetts, MA)(13). The
169 one previous report of transformation to clear cell carcinoma raises the question of
170 whether human papillomavirus (HPV), may be a potential etiological agent. Clear cell
171 carcinoma of the cervix has been associated with HPV positivity (38), and has also been
172 reported in very young patients who have never been sexually active (39). However in a
173 report by Hollowell *et al.*, BMP did not demonstrate positivity for either low or high risk
174 Human Papilloma Virus (HPV) genotypes (13). HPV testing of the biopsy specimen is
175 not recommended in the pediatric population, as it does not change management in that
176 age group.

177

178 In conclusion, BMP is a very rare benign tumor of the vagina and cervix, which may
179 present with vaginal bleeding and initially raise concern regarding rhabdomyosarcoma.
180 BMP may be recurrent with only one case of malignant transformation in an older
181 woman reported, therefore invasive surveillance in children is not recommended.
182 Surveillance during routine cytological screening in older asymptomatic women is
183 recommended. BMP in children has been reported in association with mesenchymal
184 tumors, which may shed new light on pathophysiology of these tumors.

Acknowledgement: Dr. D. MacGregor for microscopic photography, and Dr. CW Chow for discussing the pathology results

REFERENCES

1. Sanfilippo JS, Wakim NG. Bleeding and vulvovaginitis in the pediatric age group. *Clinical obstetrics and gynecology*. 1987;30(3):653-61. Epub 1987/09/01.
2. Crist W, Gehan EA, Ragab AH, Dickman PS, Donaldson SS, Fryer C, et al. The Third Intergroup Rhabdomyosarcoma Study. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 1995;13(3):610-30. Epub 1995/03/01.
3. Sultan I, Qaddoumi I, Yaser S, Rodriguez-Galindo C, Ferrari A. Comparing adult and pediatric rhabdomyosarcoma in the surveillance, epidemiology and end results program, 1973 to 2005: an analysis of 2,600 patients. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2009;27(20):3391-7. Epub 2009/04/29.
4. Kumar RJ, Ellenson L, Ronnett BM. *Blaustein's Pathology of the Female Genital Tract 6th Edition*. Springer. Pg.291-293.
5. Zeisler H, Mayerhofer K, Joura EA, Bancher-Todesca D, Kainz C, Breitenecker G, et al. Embryonal rhabdomyosarcoma of the uterine cervix: case report and review of the literature. *Gynecologic oncology*. 1998;69(1):78-83. Epub 1998/05/08.
6. Ulbright TM, Alexander RW, Kraus FT. Intramural papilloma of the vagina: evidence of Mullerian histogenesis. *Cancer*. 1981;48(10):2260-6. Epub 1981/11/15.
7. Norris HJ, Taylor HB. Polyps of the vagina. A benign lesion resembling sarcoma botryoides. *Cancer*. 1966;19(2):227-32. Epub 1966/02/01.
8. James T. A benign polypoid tumour of the cervix uteri in a girl three years old. *J Obstet Gynaecol Br Emp*. 1951;58(5):762-3. Epub 1951/10/01.
9. Abu J, Nunns D, Ireland D, Brown L. Malignant progression through borderline changes in recurrent Mullerian papilloma of the vagina. *Histopathology*. 2003;42(5):510-1. Epub 2003/04/26.
10. Arbo E, dos Reis R, Uchoa D, Acetta SG, Rivoire WA, Capp E. Vaginal Mullerian papilloma in a 2-year-old child. *Gynecologic and obstetric investigation*. 2004;58(1):55-6. Epub 2004/05/19.
11. Cohen M, Pedemonte L, Drut R. Pigmented mullerian papilloma of the vagina. *Histopathology*. 2001;39(5):541-3. Epub 2001/12/12.
12. Dobbs SP, Shaw PA, Brown LJ, Ireland D. Borderline malignant change in recurrent mullerian papilloma of the vagina. *Journal of clinical pathology*. 1998;51(11):875-7. Epub 1999/04/08.
13. Hollowell ML, Goulart RA, Gang DL, Otis CN, Prior J, Sachs BF, et al. Cytologic features of mullerian papilloma of the cervix: mimic of malignancy. *Diagnostic cytopathology*. 2007;35(9):607-11. Epub 2007/08/19.
14. Kumar A, Kumar A. Benign mullerian papilloma of the cervix. *Journal of minimally invasive gynecology*. 2012;19(5):541-2. Epub 2012/09/01.
15. Lane BR, Ross JH, Hart WR, Kay R. Mullerian papilloma of the cervix in a child with multiple renal cysts. *Urology*. 2005;65(2):388. Epub 2005/02/15.
16. Luttgies JE, Lubke M. Recurrent benign Mullerian papilloma of the vagina. Immunohistological findings and histogenesis. *Archives of gynecology and obstetrics*. 1994;255(3):157-60. Epub 1994/01/01.

17. McCluggage WG, Nirmala V, Radhakumari K. Intramural mullerian papilloma of the vagina. *International journal of gynecological pathology : official journal of the International Society of Gynecological Pathologists*. 1999;18(1):94-5. Epub 1999/01/19.
18. McCluggage WG, O'Rourke D, McElhenney C, Crooks M. Mullerian papilloma-like proliferation arising in cystic pelvic endosalpingiosis. *Human pathology*. 2002;33(9):944-6. Epub 2002/10/16.
19. Mierau GW, Lovell MA, Wyatt-Ashmead J, Goin L. Benign mullerian papilloma of childhood. *Ultrastructural pathology*. 2005;29(3-4):209-16. Epub 2005/07/23.
20. Reck-Burneo CA, Villanueva J, Velcek FT. Vaginal mullerian papilloma: an unusual cause of vaginal bleeding in a toddler. *Journal of pediatric and adolescent gynecology*. 2009;22(5):e124-6. Epub 2009/07/21.
21. Smith YR, Quint EH, Hinton EL. Recurrent benign mullerian papilloma of the cervix. *Journal of pediatric and adolescent gynecology*. 1998;11(1):29-31. Epub 1998/04/04.
22. Smrkolj S, Sorc L, Sinkovec J, Rakar S. Mullerian papilloma in a patient with Proteus syndrome: case report and review of the literature. *European journal of gynaecological oncology*. 2012;33(4):428-32. Epub 2012/10/25.
23. Tumini S, Carinci S, Anzellotti MT, Chiesa PL, Rossi C, Stuppia L, et al. Genital sanguineous discharge in prepuberty: a case of mullerian papilloma of vagina in a nine-year-old girl. *Journal of pediatric endocrinology & metabolism : JPEM*. 2010;23(8):831-2. Epub 2010/11/16.
24. Steelman J, Redondo MJ, Lovell MA, Koyle M, Kappy M. Pathology teach and tell: vaginal mesonephric papilloma. *Pediatr Pathol Mol Med*. 2001;20(4):341-4. Epub 2001/08/07.
25. Schmedding A, Zense M, Fuchs J, Gluer S. Benign papilloma of the cervix in childhood: immunohistochemical findings and review of the literature. *Eur J Pediatr*. 1997;156(4):320-2. Epub 1997/04/01.
26. Andrews CF, Jourdain L, Damjanov I. Benign cervical mesonephric papilloma of childhood. Report of a case studied by light and electron microscopy. *Diagn Gynecol Obstet*. 1981;3(1):39-43. Epub 1981/01/01.
27. Janovski NA, Kasdon EJ. Benign Mesonephric Papillary and Polypoid Tumors of the Cervix in Childhood. *J Pediatr*. 1963;63:211-6. Epub 1963/08/01.
28. Selzer I, Nelson HM. Benign papilloma (polypoid tumor) of the cervix uteri in children. Report of 2 cases. *Am J Obstet Gynecol*. 1962;84:165-9. Epub 1962/07/15.
29. Novak E, Woodruff JD, Novak ER. Probable mesonephric origin of certain female genital tumors. *Am J Obstet Gynecol*. 1954;68(5):1222-42. Epub 1954/11/01.
30. Ghaemmaghami F, Karimi Zarchi M, Ghasemi M. Lower genital tract rhabdomyosarcoma: case series and literature review. *Archives of gynecology and obstetrics*. 2008;278(1):65-9. Epub 2008/04/23.
31. Arndt CA, Donaldson SS, Anderson JR, Andrassy RJ, Laurie F, Link MP, et al. What constitutes optimal therapy for patients with rhabdomyosarcoma of the female genital tract? *Cancer*. 2001;91(12):2454-68. Epub 2001/06/20.
32. Kirsch CH, Goodman M, Esiashvili N. Outcome of Female Pediatric Patients Diagnosed With Genital Tract Rhabdomyosarcoma Based on Analysis of Cases

Literature review of Benign Müllerian papilloma

Registered in SEER Database Between 1973 and 2006. *Am J Clin Oncol*. 2012. Epub 2012/11/01.

33. Herbst AL, Ulfelder H, Poskanzer DC. Adenocarcinoma of the vagina. Association of maternal stilbestrol therapy with tumor appearance in young women. *The New England journal of medicine*. 1971;284(15):878-81. Epub 1971/04/15.
34. Manor E, Bodner L, Kachko P, Kapelushnik J. Trisomy 8 as a sole aberration in embryonal rhabdomyosarcoma (sarcoma botryoides) of the vagina. *Cancer Genet Cytogenet*. 2009;195(2):172-4. Epub 2009/12/08.
35. Andrassy RJ, Hays DM, Raney RB, Wiener ES, Lawrence W, Lobe TE, et al. Conservative surgical management of vaginal and vulvar pediatric rhabdomyosarcoma: a report from the Intergroup Rhabdomyosarcoma Study III. *J Pediatr Surg*. 1995;30(7):1034-6; discussion 6-7. Epub 1995/07/01.
36. Khan S, McDowell H, Upadhyaya M, Fryer A. Vaginal rhabdomyosarcoma in a patient with Noonan syndrome. *J Med Genet*. 1995;32(9):743-5. Epub 1995/09/01.
37. Money DM, Roy M. Canadian Consensus Guidelines on Human Papillomavirus. *J Obstet Gynecol Canada* 2007; 29(8): supplement 3.
38. Guo YF, Liu AJ, Wang XL, Wu XZ, Song L, Liu HT. [Human papillomavirus detection in clear cell carcinoma of the cervix]. *Zhonghua Shi Yan He Lin Chuang Bing Du Xue Za Zhi*. 2009;23(2):82-4. Epub 2010/01/29.
39. Fernandez-Pineda I, Spunt SL, Parida L, Krasin MJ, Davidoff AM, Rao BN. Vaginal tumors in childhood: the experience of St. Jude Children's Research Hospital. *J Pediatr Surg*. 2011;46(11):2071-5. Epub 2011/11/15.

Figure Legend:

Figure 1A: Vaginoscopy photograph with 10x magnification of a polypoid cluster of friable tissue protruding from the cervical os with normal prepubescent vaginal hypoestrogenic mucosa and no extension to the anterior or posterior aspects of the vaginal walls.

Figure 1B Demonstration of the vascularity of the pedicle within the mass.

Figure 2: Complex papillary lesion covered by cuboidal or low columnar surface epithelium without cytological atypia supported by mature fibrous cores with a variably dense mixed inflammatory infiltrate including focally prominent mast cells.

Haematoxyllin &Eosin staining. The original magnification of the image was 1.6x (picture courtesy of DM).

Table 1: Review of Published English Language Müllerian Papilloma Literature (1951-2013)

N/A – Not Available

Table 2: Comparison of Müllerian Papilloma and Rhabdomyosarcoma. Genital Rhabdomyosarcoma presents on average at an earlier age, can present with metastatic disease and is more likely to be located on the anterior wall of the vagina compared with MP (23).

Literature review of Benign Müllerian papilloma

Table 1: Review of Published English Language Müllerian Papilloma Literature

Article	Age	Length of vaginal bleeding (months)	Co-morbidities	Location	Follow up (months)	Recurrence
McQuillan (Unpublished)	5 years	24	Prolonged Activated Partial Thromboplastin Time (APTT)	Cervix	16	
McQuillan (Unpublished)	7years	<3	Osteosarcoma	Cervix	300	
McQuillan (Unpublished)	9 years	1	Constipation	Cervix	12	
McQuillan (Unpublished)	10 years	12	Insulin Dependent Diabetes	Cervix	12	
Smrkolj 2012(22)	19years	NA	Proteus Syndrome	Cervix	60	
Kumar 2012 (14)	4 years	NA	None	Cervix	0	
Tumini 2010 (23)	9 years	NA	NA	Vagina	NA	
Reck-Burneo 2009 (20)	2 years	4	None	Anterior Vaginal Wall	36	
Hollowell 2007 (13)	15 months	3	None	Cervix	6	

Mierau 2005 (19) (includes report from Steelman 2001 (24))	4 years 9 days	36 9 days	None None	Posterior vaginal wall	4 48	
Lane 2005 (15)	18 months	9	Multiple renal cysts & Wilms Tumour)	Cervix	8	
Arbo 2004 (10)	2 years	6	None	NA	NA	
Abu 2003(9) (includes Dobbs 1998 (12))	4 years	NA	Cerebral Palsy	Vagina	46 years	10 recurrences including borderline malignant changes in 1998 and clear cell malignant changes in 2002
Cohen 2001 (11)	13 years	NA	Post tylenol use	Vagina	NA	
McCluggage 1999 (17)	24 years	NA	Pregnant	Vagina	NA	
Smith 1998 (21)	4 years	None	Chronic abdominal pain	Cervix	35	2 recurrences at 12 & 24 months
Schmedding 1997 (25)	2 years	1.5	NA	Cervix	6	
Luttges 1994 (16)	5years	NA	Whooping cough	Vagina		recurrence at 24 months
Ulbright 1981 (6)	5 years	NA	NA	Posterior vaginal wall	1	

Andrews 1981 (26)	NA	NA	NA	Cervix	NA	
Norris & Taylor (7) 1966	1 day 1 day 36, 30, 20, 52, 20, 51, 29, 41, 18, 17, 40, 31	1 day 1 day Median 2 to 3 months (longer if pregnant)	None 8 pregnant	Vagina Vagina Vagina	96 9 Patient 12 (age 31): 100	Patient 12: 2 recurrences at 4 & 6 months
Janovski & Kasdon 1963 (27)	5 years	12	NA	Cervix (then vagina)	20	recurrence at 4 months
Selzer & Nelson 1962 (28)	3 years 3 years	NA	NA	Cervix Cervix	8.5 2	
Novak 1954 (29)	14 months	<3	NA	Vagina	7	
James 1951 (8)	3 years	<3	NA	Cervix	120	

NA - not available

Table 2: Comparison of Müllerian Papilloma and Rhabdomyosarcoma.

	Müllerian Papilloma	Rhabdomyosarcoma
Median Age	5 years, (range 1 day to age 52)	3.7 years (range 0.16 to age 15) (31) but on average 2 years (2) for vaginal and slightly older for cervical rhabdomyosarcoma in the second decade of life (5).
Presentation	Vaginal bleeding (average duration 6.71 months, range 1 day to 3 years)	Vaginal bleeding (61%) with 44.6% having local, regional or distant spread (mainly lung and bone) and a further 28.4% unstaged (27)
Past Medical Conditions	19 with Associated Co-morbid condition (including 8 pregnancy and 3 mesenchymal tumors)	One case associated with trisomy 8 (28); other genetic conditions increase the risk of childhood rhabdomyosarcoma (30)
Location	Vagina (but both anterior and posterior walls) or less	Anterior Wall of the Vagina

	commonly cervix	
Physical examination	Tanner stage 1	Tanner stage 1
Vaginoscopy	Well demarcated friable polypoid cluster with no local infiltration	Gray or tan edematous friable polypoid cluster with possible local infiltration in the vagina or protruding from the cervix.
Resection	Easy with minimal blood loss	Risk of hemorrhage and worse prognosis with incomplete resection (29)
Microscopic Findings	Papillae covered by bland epithelium without cytological atypia or mitotic activity. No cambium layer. Fibrovascular stromal cores.	Densely cellular cambium layer underlying intact epithelium and the presence of associated spindled stromal cells showing rhabdomyoblastic differentiation.