

Article Type: Research letter

Article begins on page three of this document.

<b>Title</b>	Increased incidence of community-associated <i>Staphylococcus aureus</i> bloodstream infections in Victoria and Western Australia, 2011–2016
--------------	--

**Authors:**

	Title	First name	Mid inits	Last name	Postnom (eg, PhD)	Position1	Address1	Position2	Address2	Tel	Email
1	Dr.	Nabeel		Imam	MBBS, MPH	Biostatistician/Epidemiologist	1			93429333	nabeel.imam@mh.org.au
2	Ms.	Simone		Tempone	Masters of Infectious Diseases	Program Officer	2				Simone.Tempone@health.wa.gov.au
3	Dr.	Paul	K	Armstrong	MBBS, FRACP, MAppEpid	Director	2			893884801	Paul.Armstrong@health.wa.gov.au
4	Ms.	Rebecca		McCann	BSc (Nursing)	Manager	2			0439920819	rebecca.mccann@health.wa.gov.au
5	Ms.	Sandra		Johnson	Bachelor of Science, Masters of Public Health	Epidemiologist	1				Sandra.Johnson@mh.org.au
6	Assoc. Prof.	Leon	J	Worth	MBBS, FRACP, PhD, Grad Dip Epi	Infectious diseases physician	1				leon.worth@mh.org.au
7	Prof.	Michael	J	Richards	MB, BS, FRACP, MD	Infectious Diseases Physician	3			0393427481	Michael.Richards@mh.org.au

Number of corresponding author:	1
Number of alternative corresponding author:	

**Addresses:**

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/mja2.12057](https://doi.org/10.1002/mja2.12057)

This article is protected by copyright. All rights reserved

	Institution	Postal address	City	State	Post Code	Nation
1	Victorian Healthcare Associated Infection Surveillance System (VICNISS) Coordinating Centre, The Peter Doherty Institute for Infection and Immunity	792 Elizabeth Street	Melbourne	VIC	3000	
2	Western Australia Department of Health	Perth	Perth	WA	6000	
3	Royal Melbourne Hospital	C/- Post Office	Melbourne	VIC	3050	
4						
5						
6						
7						
8						
9						
10						

Postal address of first corresponding author (if different from the institutional address given above)	
--	--

Primary Keywords [Office use only]	Infectious diseases; Statistics, epidemiology and research design
Secondary keywords [Office use only]	Staphylococcus; Sepsis; Community acquired infections; Surveillance
Notes:	

**Article details** (press ctrl – 9 to enter details):

Article type	Research letter
Blurb	Characterising the isolates responsible for infection would help identify virulence factors and the relatedness of isolates
Pullquote	

**Office use**

<i>Ms. Number</i>	mja18.00731. R2
<i>Medical editor</i>	Robyn Godding
<i>Medical editor email</i>	rgodding@mja.com.au
<i>Structural editor</i>	Paul Foley
<i>Structural editor email</i>	pfoley@mja.com.au
<i>Section/Category</i>	Research letter
<i>Strapheading</i>	Research letter
<i>Substrap</i>	

**Elsevier – file data:**

Filename for copyediting	ima_mja18.00731_ms.docx
Accompanying graphics	ima_mja18.00731_gr.eps ima_mja18.00731_gr.jpg
Stock images	None
Appendices	None

**Office use – history:**

Event	Date
Original submission received	05/07/2018

Event	Date
Accept	13/09/2018

Proof sent to author	
Proof returned by author	
Published (date format xx/xx/xx)	18/02/19
Issue	3
Vol	210
DOI	10.5694/mja18.00731
Journal	The Medical Journal of Australia
Original article DOI (for response)	

Author Manuscript

# Increased incidence of community-associated *Staphylococcus aureus* bloodstream infections in Victoria and Western Australia, 2011–2016

Standardised national surveillance of health care-associated *Staphylococcus aureus* bloodstream infections (HA-SABs)<sup>1</sup> has found that rates are declining in Australia.<sup>2</sup> The incidence of community-associated SABs (CA-SABs), however, has not been investigated. These infections frequently have complicated courses (eg, metastatic sites of infection)<sup>3</sup> and high mortality (about 20%).<sup>4</sup>

During calendar years 2016 and 2017, the Victorian Healthcare Associated Infection Surveillance System (VICNISS) and Healthcare Infection Surveillance Western Australia (HISWA) received reports from affiliated hospitals about increases in the frequency of CA-SABs. We sought to formally review trends in their incidence and to determine whether populations at particular risk could be identified. We analysed quarterly data for the period 1 January 2011 – 31 December 2016 from 93 Victorian public hospitals (VICNISS) and 58 Western Australian public hospitals (HISWA). As a quality assurance evaluation of surveillance data, no patient-identifying or hospital-level data were captured; formal ethics review was therefore not required.

A total of 10 320 SAB events were reported (7262 in Victoria, 3058 in WA); 6800 infections (65.9%) were community-associated, corresponding to an aggregate crude rate of 13.3 CA-SABs per 100 000 person-years (Victoria, 12.9; WA, 14.3 CA-SABs per 100 000 person-years). The incidence in each state increased significantly during the study period: 8% (95% confidence interval [CI], 6–10%) per year in Victoria and 6% (95% confidence interval [CI], 4–9%) per year in WA (Box 1).

Most CA-SABs (5698, 83.8%) were caused by methicillin-susceptible *S. aureus* (MSSA). There was a small but statistically significant annual increase in the proportion of MSSA CA-SABs in Victoria (4% per year; 95% CI, 3–8%), but not in WA; that is, the incidence of CA-SABs caused by MSSA rose more rapidly in Victoria than the overall incidence of CA-SABs.

The incidence of CA-SABs was higher among older patients and in men, and was particularly high for men over 60: the standardised incidence in this group was 50.9 cases

(95% CI, 48.5–53.4) in Victoria and 56.7 cases (95% CI, 52.6–61.0) per 100 000 person-years in WA, twice the incidence among women of the same age (Victoria, 24.7 [95% CI, 23.2–26.4]; WA, 24.9 [95% CI, 22.4–27.7] cases per 100 000 person-years).

Our study is the first to evaluate the epidemiology of CA-SAB in more than one Australian state. Other investigators have similarly reported a greater disease burden for older people and men.<sup>4</sup> It is notable that residents of aged care facilities have been identified as being at particular risk;<sup>5</sup> infections in this group would be classified as CA-SABs by our analysis. One limitation of our study was that only infections in patients presenting to public hospitals are captured by surveillance systems, perhaps missing some community-onset infections, but the vast majority of patients with SABs would require admission to hospital. Some infected persons would have been managed entirely in private health care, meaning that we will have underestimated the incidence of CA-SAB.

The increases in incidence we describe may reflect the emergence of virulent *S. aureus* strains in the community or changes in host risk factors. Characterising the isolates responsible for infection would assist in identifying virulence factors and the relatedness of isolates. Further evaluation of infection risks in people over 60 years of age is also needed for developing targeted prevention strategies.

**Acknowledgements:** We acknowledge the infection control consultants in Victorian and Western Australian hospitals who collected the data we analysed. VICNISS is fully funded by the Victorian Department of Health and Human Services; HISWA is a program in the Public Health Division of the Department of Health Western Australia.

**Competing interests:** No relevant disclosures.

Received 5 July 2018, accepted 13 September 2018.

## Author details

Nabeel Imam<sup>1</sup>

Simone Tempone<sup>2</sup>

Paul K Armstrong<sup>2</sup>

Rebecca McCann<sup>2</sup>

Sandra Johnson<sup>1</sup>

Leon J Worth<sup>1</sup>

Michael J Richards<sup>3</sup>

1 Victorian Healthcare Associated Infection Surveillance System (VICNISS) Coordinating Centre, The Peter Doherty Institute for Infection and Immunity, Melbourne, VIC.

2 Western Australia Department of Health, Perth, WA.

3 Royal Melbourne Hospital, Melbourne, VIC.

**nabeel.imam@mh.org.au**

doi: 10.5694/mja18.00731

## References

- 1 Australian Commission on Safety and Quality in Health Care. Implementation guide for surveillance of *Staphylococcus aureus* bacteraemia. Sydney: ACSQHC, 2013. [https://www.safetyandquality.gov.au/wp-content/uploads/2012/02/SAQ019\\_Implementation\\_guide\\_SAB\\_v10.pdf](https://www.safetyandquality.gov.au/wp-content/uploads/2012/02/SAQ019_Implementation_guide_SAB_v10.pdf) (viewed June 2018).
- 2 Worth LJ, Spelman T, Bull AL, Richards MJ. *Staphylococcus aureus* bloodstream infection in Australian hospitals:

findings from a Victorian surveillance system. *Med J Aust* 2014; 200: 282-284. <https://www.mja.com.au/journal/2014/200/5/staphylococcus-aureus-bloodstream-infection-australian-hospitals-findings>

- 3 Tong SYC, Davis JS, Eichenberger E, et al. *Staphylococcus aureus* infections: epidemiology, pathophysiology, clinical manifestations, and management. *Clin Microbiol Rev* 2015; 28: 603-661.
- 4 Turnidge JD, Kotsanas D, Munckhof W, et al. *Staphylococcus aureus* bacteraemia: a major cause of mortality in Australia and New Zealand. *Med J Aust* 2009; 191: 368-373. <https://www.mja.com.au/journal/2009/191/7/staphylococcus-aureus-bacteraemia-major-cause-mortality-australia-and-new>
- 5 Agostino JW, Ferguson JK, Eastwood K, Kirk MD. The increasing importance of community-acquired methicillin-resistant *Staphylococcus aureus* infections. *Med J Aust* 2017; 207: 388-393. <https://www.mja.com.au/journal/2017/207/9/increasing-importance-community-acquired-methicillin-resistant-staphylococcus>

**Box. Quarterly incidence of community-associated *Staphylococcus aureus* bloodstream infections (CA-SABs) in Victoria and Western Australia, 2011–2016, with trend lines and 95% confidence envelopes**

