

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Hayhurst, H;de Coo, IFM;Piekutowska-Abramczuk, D;Alston, CL;Sharma, S;Thompson, K;Rius, R;He, L;Hopton, S;Ploski, R;Ciara, E;Lake, NJ;Compton, AG;Delatycki, MB;Verrips, A;Bonnen, PE;Jones, SA;Morris, AA;Shakespeare, D;Christodoulou, J;Wesol-Kucharska, D;Rokicki, D;Smeets, HJM;Pronicka, E;Thorburn, DR;Gorman, GS;McFarland, R;Taylor, RW;Ng, YS

Title:

Leigh syndrome caused by mutations in *MTFMT* is associated with a better prognosis (vol 6, pg 515, 2019)

Date:

2019-04

Citation:

Hayhurst, H., de Coo, I. F. M., Piekutowska-Abramczuk, D., Alston, C. L., Sharma, S., Thompson, K., Rius, R., He, L., Hopton, S., Ploski, R., Ciara, E., Lake, N. J., Compton, A. G., Delatycki, M. B., Verrips, A., Bonnen, P. E., Jones, S. A., Morris, A. A., Shakespeare, D., ... Ng, Y. S. (2019). Leigh syndrome caused by mutations in *MTFMT* is associated with a better prognosis (vol 6, pg 515, 2019). ANNALS OF CLINICAL AND TRANSLATIONAL NEUROLOGY, 6 (4), pp.821-821. <https://doi.org/10.1002/acn3.780>.

Persistent Link:

<https://hdl.handle.net/11343/271489>

License:

CC BY-NC-ND

ERRATUM

Leigh syndrome caused by mutations in *MTFMT* is associated with a better prognosis

In Hayhurst et al. (2019),¹ the authors' affiliation were incorrectly published on page 515.

The authors' affiliation should read as:

Hannah Hayhurst¹, Irenaeus F. M. de Co^{2,11}, Dorota Piekutowska-Abramczuk³, Charlotte L. Alston¹, Sunil Sharma¹, Kyle Thompson¹, Rocio Rius^{4,5}, Langping He¹, Sila Hopton¹, Rafal Ploski³, Elzbieta Ciara³, Nicole J. Lake^{4,5}, Alison G. Compton^{4,5}, Martin B. Delatycki^{4,5}, Aad Verrips⁶, Penelope E. Bonnen⁷, Simon A. Jones⁸, Andrew A. Morris⁸, David Shakespeare⁹, John Christodoulou^{4,5}, Dorota Wesol-Kucharska¹⁰, Dariusz Rokicki¹⁰, Hubert J. M. Smeets¹¹, Ewa Pronicka^{3,10}, David R. Thorburn^{4,5}, Gráinne S. Gorman¹, Robert McFarland¹, Robert W. Taylor¹, Yi Shiao Ng¹

¹Wellcome Centre for Mitochondrial Research, Institute of Neuroscience, Newcastle University, Newcastle upon Tyne, United Kingdom

²Department of Neurology, Erasmus Medical Centre, Rotterdam, Netherlands

³Department of Medical Genetics, Warsaw Medical University, 02-106 Warsaw, Poland

⁴Victorian Clinical Genetics Service and Murdoch Children's Research Institute, Parkville, Victoria 3052, Australia

⁵Department of Paediatrics, University of Melbourne, Melbourne, Victoria 3052, Australia

⁶Department of Neurology, Canisius Wilhelmina Hospital, Nijmegen, The Netherlands

⁷Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, Texas, USA

⁸Manchester Centre for Genomic Medicine, Manchester University NHS Foundation Trust, Manchester Academic Health Sciences Centre, Manchester, UK

⁹Neuro-rehabilitation unit, Royal Preston Hospital, Preston, United Kingdom

¹⁰Department of Paediatrics, Nutrition and Metabolic Diseases, The Children's Memorial Health Institute, 04-730 Warsaw, Poland

¹¹Department of Clinical Genomics, Research Schools GROW and MHeNS, Maastricht University, Maastricht, The Netherlands

We apologize for this error.

Reference

1. Hayhurst H, de Co IF, Piekutowska-Abramczuk D, et al. Leigh syndrome caused by mutations in *MTFMT* is associated with a better prognosis. *Ann Clin Transl Neurol* 2019;6:515–524.